

NCERT EXERCISES

- 3.1 Arrange the following metals in the order in which they displace each other from the solution of their salts.
Al, Cu, Fe, Mg and Zn.
- 3.2 Given the standard electrode potentials,
 $K^+/K = -2.93V$, $Ag^+/Ag = 0.80V$,
 $Hg^{2+}/Hg = 0.79V$
 $Mg^{2+}/Mg = -2.37 V$, $Cr^{3+}/Cr = - 0.74V$
 Arrange these metals in their increasing order of reducing power.
- 3.3 Depict the galvanic cell in which the reaction $Zn(s)+2Ag^+(aq) \rightarrow Zn^{2+}(aq)+2Ag(s)$ takes place. Further show:
 (i) Which of the electrode is negatively charged?
 (ii) The carriers of the current in the cell.
 (iii) Individual reaction at each electrode.
- 3.4 Calculate the standard cell potentials of galvanic cell in which the following reactions take place:
 (i) $2Cr(s) + 3Cd^{2+}(aq) \rightarrow 2Cr^{3+}(aq) + 3Cd$
 (ii) $Fe^{2+}(aq) + Ag^+(aq) \rightarrow Fe^{3+}(aq) + Ag(s)$
 Calculate the $\Delta_r G^\ominus$ and equilibrium constant of the reactions.
- 3.5 Write the Nernst equation and emf of the following cells at 298 K:
 (i) $Mg(s) | Mg^{2+}(0.001M) || Cu^{2+}(0.0001 M) | Cu(s)$
 (ii) $Fe(s) | Fe^{2+}(0.001M) || H^+(1M) | H_2(g)(1bar) | Pt(s)$
 (iii) $Sn(s) | Sn^{2+}(0.050 M) || H^+(0.020 M) | H_2(g) (1 bar) | Pt(s)$
 (iv) $Pt(s) | Br_2(l) | Br^-(0.010 M) || H^+(0.030 M) | H_2(g) (1 bar) | Pt(s)$.
- 3.6 In the button cells widely used in watches and other devices the following reaction takes place:
 $Zn(s) + Ag_2O(s) + H_2O(l) \rightarrow Zn^{2+}(aq) + 2Ag(s) + 2OH^-(aq)$
 Determine $\Delta_r G^\ominus$ and E^\ominus for the reaction.
- 3.7 Define conductivity and molar conductivity for the solution of an electrolyte. Discuss their variation with concentration.
- 3.8 The conductivity of 0.20 M solution of KCl at 298 K is $0.0248 S cm^{-1}$. Calculate its molar conductivity.
- 3.9 The resistance of a conductivity cell containing 0.001M KCl solution at 298 K is 1500Ω . What is the cell constant if conductivity of 0.001M KCl solution at 298 K is $0.146 \times 10^{-3} S cm^{-1}$.
- 3.10 The conductivity of sodium chloride at 298 K has been determined at different concentrations and the results are given below:

Concentration/M	0.001	0.010	0.020	0.050	0.100
$10^2 \times \kappa/S m^{-1}$	1.237	11.85	23.15	55.53	106.74

 Calculate Λ_m for all concentrations and draw a plot between Λ_m and $c^{1/2}$. Find the value of Λ_m^0 .
- 3.11 Conductivity of 0.00241 M acetic acid is $7.896 \times 10^{-5} S cm^{-1}$. Calculate its molar conductivity and if Λ_m^0 for acetic acid is $390.5 S cm^2 mol^{-1}$, what is its dissociation constant?
- 3.12 How much charge is required for the following reductions:
 (i) 1 mol of Al^{3+} to Al.
 (ii) 1 mol of Cu^{2+} to Cu.
 (iii) 1 mol of MnO_4^- to Mn^{2+} .
- 3.13 How much electricity in terms of Faraday is required to produce
 (i) 20.0 g of Ca from molten $CaCl_2$.
 (ii) 40.0 g of Al from molten Al_2O_3 .

- 3.14 How much electricity is required in coulomb for the oxidation of
- * (i) 1 mol of H_2O to O_2 .
 - * (ii) 1 mol of FeO to Fe_2O_3 .
- 3.15 A solution of $\text{Ni}(\text{NO}_3)_2$ is electrolysed between platinum electrodes using a current of 5 amperes for 20 minutes. What mass of Ni is deposited at the cathode?
- 3.16 Three electrolytic cells A,B,C containing solutions of ZnSO_4 , AgNO_3 and CuSO_4 , respectively are connected in series. A steady current of 1.5 amperes was passed through them until 1.45 g of silver deposited at the cathode of cell B. How long did the current flow? What mass of copper and zinc were deposited?
- 3.17 Using the standard electrode potentials given in Table 3.1, predict if the reaction between the following is feasible:
- (i) $\text{Fe}^{3+}(\text{aq})$ and $\text{I}^-(\text{aq})$
 - (ii) $\text{Ag}^+(\text{aq})$ and $\text{Cu}(\text{s})$
 - (iii) $\text{Fe}^{3+}(\text{aq})$ and $\text{Br}^-(\text{aq})$
 - (iv) $\text{Ag}(\text{s})$ and $\text{Fe}^{3+}(\text{aq})$
 - (v) $\text{Br}_2(\text{aq})$ and $\text{Fe}^{2+}(\text{aq})$.
- 3.18 Predict the products of electrolysis in each of the following:
- * (i) An aqueous solution of AgNO_3 with silver electrodes.
 - * (ii) An aqueous solution of AgNO_3 with platinum electrodes.
 - * (iii) A dilute solution of H_2SO_4 with platinum electrodes.
 - * (iv) An aqueous solution of CuCl_2 with platinum electrodes.