


# GPLUS EDUCATION

Date :  
Time :  
Marks :


CHEMISTRY

## COORDINATION COMPOUNDS

### Single Correct Answer Type

- The IUPAC name of  $\text{Na}_3[\text{Co}(\text{ONO})_6]$  is:  
a) Sodium cobaltinitrite  
b) Sodium hexanitritocobaltate(III)  
c) Sodium hexanitrocobalt(III)  
d) Sodium hexanitritocobaltate(II)
- $\text{CuSO}_4$  decolourises on addition of KCN, the product is:  
a)  $\text{Cu}(\text{CN})_4^{2-}$                       b)  $[\text{Cu}(\text{CN})_4]^{3-}$                       c)  $\text{Cu}(\text{CN})_2$                       d)  $\text{CuCN}$
- Exchange of coordination group by a water molecule in complex molecule results in:  
a) Ionization isomerism  
b) Ligand isomerism  
c) Hydration isomerism  
d) Geometrical isomerism
- The type of isomerism found in urea molecule is  
a) Chain  
b) Position  
c) Tautomerism  
d) None of these
- The IUPAC name of the compound  is  
a) Butane-2-aldehyde                      b) 2-methyl butanal                      c) 2-ethyl propanal                      d) None of the above
- Anisol is a product obtained from phenol by the reaction known as:  
a) Coupling                      b) Etherification                      c) Oxidation                      d) Esterification
- Which of the following is diamagnetic in nature?  
a)  $[\text{Fe}(\text{CN})_6]^{3-}$                       b)  $[\text{NiCl}_4]^{2-}$                       c)  $[\text{Ni}(\text{CO})_4]$                       d)  $[\text{MnCl}_4]^{2-}$
- Which is the strongest field ligand?  
a)  $\text{CN}^-$                       b)  $\text{NO}_2^-$                       c)  $\text{NH}_3$                       d) en
- Nitrobenzene on reduction with Zn and aq.  $\text{NH}_4\text{Cl}$  gives:  
a) Aniline  
b) Nitrosobenzene  
c) *N*-phenyl hydroxylamine  
d) Hydrazobenzene
- The IUPAC name of  $[\text{Co}(\text{NH}_3)_5\text{ONO}]^{2+}$  ion is  
a) Pentaammine nitrito cobalt (IV) ion                      b) Pentaammine nitro cobalt (III) ion  
c) Pentaammine nitrito cobalt (III) ion                      d) Pentaammine nitro cobalt (IV) ion
- The compound which does not show paramagnetism is  
a)  $\text{NO}_2$                       b)  $\text{NO}$                       c)  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$                       d)  $[\text{Cu}(\text{NH}_3)_4\text{Cl}_2]$
- Which of the following is expected to undergo nitration more easily and readily to furnish the corresponding nitro derivatives employing the usual nitrating mixture?  
a)  $\text{C}_6\text{H}_6$                       b)  $\text{C}_6\text{H}_5\text{NO}_2$                       c)  $\text{C}_6\text{H}_5\text{CH}_3$                       d)  $\text{C}_6\text{H}_5 \cdot \text{CCl}_3$
- The number of unpaired electrons calculated in  $[\text{Co}(\text{NH}_3)_6]^{3+}$  and  $[\text{Co}(\text{F}_6)]^{3-}$  are  
a) 4 and 4                      b) 0 and 2                      c) 2 and 4                      d) 0 and 4

14. The IUPAC name of



- a) 4-hydroxy amino benzene carboxylic acid      b) 4-(N-hydroxy) imino benzene carboxylic acid  
c) 4-hydroxy imino cyclohexanoic acid      d) 4-(N-hydroxy) imino cyclohexane-1  
-carboxylic acid

15. The IUPAC name of the coordination compound  $K_2[Zn(OH)_4]$  is

- a) Potassium tetrahydroxozine (II)      b) Dipotassium tetrahydroxo(II)  
c) Potassium tetrahydroxozincate (II)      d) Potassium tetrahydroxozincate (III)

16. Arrange in order of decreasing trend towards  $S_E$  reactions,  
Chlorobenzene, Benzene, Anilium chloride, Toluene:

- I.      (II)      (III)      (IV)  
a)  $II > I > III > IV$       b)  $III > I > II > IV$       c)  $IV > II < I > III$       d)  $I > II > III > IV$

17. Toluene is nitrated and the resulting product is reduced with tin and hydrochloric acid. The product so obtained is diazotised and then heated with cuprous bromide. The reaction mixture so formed contains:

- a) Mixture of *o*- and *m*-bromotoluenes  
b) Mixture of *o*- and *p*-bromotoluenes  
c) Mixture of *o*- and *p*-dibromobenzenes  
d) Mixture of *o*- and *p*-bromoanilines

18. A positive carbylamine test is given by:

- a) *N,N*-dimethylaniline  
b) 2,4-dimethylaniline  
c) *N*-methyl-*o*-methylaniline  
d) *p*-methyl benzylamine

19.  $CN^-$  is strong field ligand. This is due to the fact that

- a) It carries negative charge  
b) It is a pseudohalide  
c) It can accept electrons from metal species  
d) It forms high spin complexes with metal species.

20. Which of the following is not true for ligand metal complex?

- a) Highly charged ligand forms strong bond  
b) Greater the ionization potential of central metal, the stronger is the bond  
c) Larger the permanent dipole moment of ligand, the more stable is the bond  
d) Larger the ligand, the more stable is the metal-ligand bond

21. The nitration of nitrobenzene with fuming  $HNO_3$  will give:

- a) TNB      b) 1,3-dinitrobenzene      c) Picric acid      d) 1,4-dinitrobenzene

22. A ligand can also be regarded as

- a) Lewis acid      b) Bronsted base      c) Lewis base      d) Bronsted acid


23. The correct statement with respect to the complexes  $Ni(CO)_4$  and  $[Ni(CN)_4]^{2-}$  is

- a) Nickel is in the same oxidation state in both  
b) Both have tetrahedral geometry  
c) Both have square planar geometry  
d) Have tetrahedral and square planar geometry respectively

24. Which one of the following has lowest value of paramagnetic behaviour?





- a)  $[Cr(CN)_6]^{3-}$       b)  $[Mn(CN)_6]^{3-}$       c)  $[Fe(CN)_6]^{3-}$       d)  $[Co(CN)_6]^{3-}$

25. In the reaction;



the products are:








26. An octahedral complex is formed when central metal atom undergoes hybridization amongst the....orbitals.

a)  $sp^3$

b)  $dsp^2$

c)  $sp^3d$

d)  $sp^3d^2$



28. Biological oxidation of  $\text{C}_6\text{H}_6$  taking place in body of dog, gives:

a) Benzoic acid

b) Toluic acid

c) Maleic acid

d) Muconic acid

29. Ammonia forms the complex ion  $[\text{Cu}(\text{NH}_3)_4]^{2+}$  with copper ions in the alkaline solutions but not in acidic solutions. What is the reason for it?

a) In acidic solutions hydration protects copper ions

b) In acidic solutions protons coordinate with ammonia molecules forming  $\text{NH}_4^+$  ions and  $\text{NH}_3$  molecules are not available

c) In alkaline solutions insoluble  $\text{Cu}(\text{OH})_2$  is precipitated which is soluble in excess of any alkali

d) Copper hydroxide is an amphoteric substance

30. Which of the following has the highest molar conductivity in solution?

a)  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$

b)  $[\text{Pt}(\text{NH}_3)_5\text{Cl}]\text{Cl}_3$

c)  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}_2$

d)  $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]\text{Cl}$

31. Which of the following is not *meta* directing group?

a)  $-\text{SO}_3\text{H}$

b)  $-\text{NO}_2$

c)  $-\text{CN}$

d)  $-\text{NH}_2$

32. Which of the following is an organometallic compound?

a) Lithium methoxide

b) Lithium acetate

c) Lithium dimethylamine

d) Methyl lithium

33. Which among the following is very strong *o*-, *p*-directing group?





a)  $-\text{Cl}$

b)  $-\text{OR}$




c)  $-\text{NH}_2$

d)  $-\text{NHR}$


34. The type of hybridisation in tetrahedral complexes of metal atom is

- a)  $dsp^2$                       b)  $d^2sp$                       c)  $sp^3$                       d)  $sp^2$
35. Chlorobenzene on heating with NaOH at 300°C under pressure gives:  
a) Phenol                      b) Benzaldehyde                      c) Chlorophenol                      d) None of these
36. The coordination number of Fe in  $[\text{Fe}(\text{CN})_6]^{4-}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$  and  $[\text{FeCl}_4]^-$  are respectively.  
a) 2, 3, 3                      b) 6, 6, 4                      c) 6, 3, 3                      d) 6, 4, 6
37. Consider the following statements  
I. Chain and position isomerism are not possible together between two isomers  
II. Tautomerism is a chemical phenomenon which is catalysed by acid as well as base  
III. Tautomers are always metamers  
IV. Tautomers are always functional isomers  
Select the correct answer by using the codes given below  
a) Only III is correct                      b) III and IV are correct  
c) I, II and III are correct                      d) I, II and IV are correct
38. What is the EAN of nickel in  $[\text{Ni}(\text{CN})_4]^{2-}$ ?  
a) 32                      b) 35                      c) 34                      d) 36
39. Which of the following alcohols is dehydrated most readily with conc.  $\text{H}_2\text{SO}_4$ ?  
a)  $p\text{-O}_2\text{NC}_6\text{H}_4\text{CH}(\text{OH})\text{CH}_3$   
b)  $p\text{-ClC}_6\text{H}_4\text{CH}(\text{OH})\text{CH}_3$   
c)  $p\text{-CH}_3\text{OC}_6\text{H}_4\text{CH}(\text{OH})\text{CH}_3$   
d)  $\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{CH}_3$
40. The compound having tetrahedral geometry is  
a)  $[\text{Ni}(\text{CN})_4]^{2-}$                       b)  $[\text{Pd}(\text{CN})_4]^{2-}$                       c)  $[\text{PdCl}_4]^{2-}$                       d)  $[\text{NiCl}_4]^{2-}$
41. Identify 'Z' in the change;  
$$\text{C}_6\text{H}_5\text{NH}_2 \xrightarrow[280\text{ K}]{\text{NaNO}_2/\text{HCl}} \text{X} \xrightarrow{\text{CuBr/HBr}} \text{Z:}$$
- a) 
- b) 
- c) 
- d) 
42. Which of the following is most acidic?  
a)  $p$ -cresol                      b)  $p$ -chlorophenol                      c)  $p$ -nitrophenol                      d)  $p$ -aminophenol
43. Benzoylacetone beryllium exhibit isomerism of the type  
a) Structural                      b) Geometrical                      c) Optical                      d) Conformational
44. Which one of the following has a square planar geometry?  
(At. No. Fe=26, Co=27, Ni=28, Pt=78)  
a)  $[\text{CoCl}_4]^{2-}$                       b)  $[\text{FeCl}_4]^{2-}$                       c)  $[\text{NiCl}_4]^{2-}$                       d)  $[\text{PtCl}_4]^{2-}$
45. The number of ions formed on dissolving one molecule of  $\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$  in water is:  
a) 4                      b) 5                      c) 3                      d) 6
46. A solution of potassium ferrocyanide would contains-ions  
a) 2                      b) 3                      c) 4                      d) 5
47. Which of the following is not considered as an organometallic compound?  
a) Grignard reagent                      b)  $\text{cis-platin}$                       c) Zeise's salt                      d) Ferrocene
48. When phenol is reacted with  $\text{CHCl}_3$  and NaOH followed by acidification, salicylaldehyde is obtained. Which of the following species are involved in the above mentioned reaction as intermediates?

d) Both (a) and (b)



49. Number of geometrical isomers for the molecule



are

a) 2

b) 3

c) 6

d) 5

50. Which statement about coordination number of a cation is true?

a) Most metal ions exhibit only a single characteristic coordination number

b) The coordination number is equal to the number of ligands bonded to the metal atom

c) The coordination number is determined solely by the tendency to surround the metal atom with the same number of electrons as one of the rare gases

d) For most cations, the coordination number depends on the size, structure and charge of the ligands

51. Among the following, the strongest base is:

a)  $C_6H_5NH_2$

b)  $p\text{-NO}_2\text{—C}_6\text{H}_4\text{NH}_2$

c)  $m\text{-NO}_2\text{—C}_6\text{H}_4\text{NH}_2$

d)  $C_6H_5CH_2NH_2$

52. General formula for arenes is:

a)  $C_nH_{2n+6}$

b)  $C_nH_{2n+6y}$

c)  $C_nH_{2n}$

d)  $C_nH_{2n-6y}$

53. Which of the following doesn't have a metal-carbon bond?

a)  $Al(OC_2H_5)_3$

b)  $C_2H_5MgBr$

c)  $K[Pt(C_2H_4)Cl_3]$

d)  $Ni(CO)_4$

54. How many isomers are possible in  $[Co(en)_2Cl_2]^+$ ?

a) 2

b) 4

c) 6

d) 1

55. How many carbon atoms in the molecule  $HOOC - (CHOH)_2 - COOH$  are asymmetric?

a) 1

b) 2

c) 3

d) None of these

56. In benzene, there is a delocalisation of  $\pi$ -electrons. Hence, each  $\pi$ -electron is attached by....carbon nuclei.

a) 2

b) 3

c) 6

d) 4

57. Which can be used to distinguish  $C_6H_5NH_2$  and  $C_6H_5CH_2NH_2$ ?

a) Diazotisation followed with coupling with phenol

b) Carbylamine reaction

c) Reimer-Tiemann reaction

d) None of the above

58. When  $RCOCl$  and  $AlCl_3$  are used in Friedel-Craft's reaction, the electrophile is:

a)  $Cl^+$

b)  $RCOCl$

c)  $R^+CO$

d)  $R^+$

59. Thiophene is separated from benzene by:

a) Chlorination of thiophene

b) Sulphonation of thiophene

c) Nitration of thiophene

d) Oxidation of thiophene

60. A complex compound of  $CO^{3+}$  with molecular formula  $CoCl_x \cdot yNH_3$  gives a total of 3 ions when dissolved in water. How many  $Cl^-$  ions satisfy both primary and secondary valencies in this complex?

a) 3

b) 1

c) 4

d) Zero

61. The correct IUPAC name of alcohol  $[(CH_3)_2CH]_3COH$  is

a) Tri isopropyl carbinol

b) 2, 4-dimethyl-3-isopropyl pentan-3-ol

c) 2,4-dimethyl-3-(1-methyl) ethyl pentan-3-ol

d) None of the above

62. Colour of transition metal complexes can be explained by:





a) Completely filled  $d$ -orbitals

b) Vacant  $d$ -orbitals

c)  $d - d$  transition

d) None of the above

63. Which is most reactive towards electrophilic reagents:



64. Pick a poor electrolytic conductor complex in solution:

a)  $K_2[PtCl_6]$

b)  $[Co(NH_3)_3](NO_2)_3$

c)  $K_4[Fe(CN)_6]$

d)  $[Co(NH_3)_4]SO_4$

65. Benzene reacts with sulphuric acid only when the acid is:


a) Dilute and cold

b) Dilute and hot

c) Hot and concentrated

d) Mixed with  $HNO_3$

66. In the following reaction the catalyst used is:



a)  $Cr_2O_3$

b)  $Al_2O_3$

c) Zn dust

d)  $Cr_2O_3$  and  $Al_2O_3$

67. The alkane which has only primary hydrogen atom is


a) Pentane

b) isopentane

c) neopentane

d) 2, 2-dimethyl butane

68. The correct IUPAC name of the complex;



a) Dichlorodimethylglyoximate cobalt(II)




b) Bis(dimethylglyoxime) dichloro cobalt(II)

c) Dimethylglyoxime cobalt(II) chloride

d) Dichlorodimethylglyoxime-N,N-cobalt(II)

69. Which of the following nitroalkane will not show tautomerism?

a)  $CH_3CH_2CH_2CH_2NO_2$



70. Which is low spin complex?

a)  $[Fe(CN)_6]^{3-}$

b)  $[Co(NO_2)_6]^{3-}$

c)  $[Mn(CN)_6]^{3-}$

d) All of these

71. The probable formula for Prussian blue is:



a)  $Fe_3[Fe(CN)_6]_2$


b)  $Fe_2[Fe(CN)_6]_3$


c)  $Fe_4[Fe(CN)_6]_3$

d)  $Fe_3[Fe(CN)_6]_4$


72. Which represents Reimer-Tiemann reaction?













73. The complex ion which has no 'd'-electron in the central metal atom is :  
 a)  $[\text{MnO}_4]^-$                       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                       c)  $[\text{Fe}(\text{CN})_6]^{3-}$                       d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
74. The shape of cobalt hexaammine cation, which has its central cobalt atom surrounded by six ammonia molecules is:  
 a) Tetrahedral                      b) Octahedral                      c) Square planar                      d) Trigonal
75. Which ligand is capable of forming low spin as well as high spin complexes?  
 a) CO                      b)  $\text{NO}_2^-$                       c)  $\text{CN}^-$                       d)  $\text{NH}_3$
76.   
 The IUPAC name of \_\_\_\_\_ is  
 a) 7-ethyl-2, 4, 5, 6-tetramethyl-deca-1, 9-diene                      b) 7-ethyl-2, 4, 5, 6-tetramethyl-deca-1, 8-diene  
 c) 4-ethyl-4, 5, 6, 7-tetramethyl-deca-1, 9-diene                      d) 7-(1-propenyl)-2, 3, 4, 5-tetramethyl-non-1-ene
77. IUPAC name of  $[\text{Pt}(\text{NH}_3)_3\text{Br}(\text{NO}_2)\text{Cl}]\text{Cl}$  is  
 a) Triamminechlorobromonitro platinum (IV) chloride  
 b) Triamminebromonitrochloro platinum (IV) chloride  
 c) Triamminebromochloronitro platinum (IV) chloride  
 d) Triamminenitrochlorobromo platinum (IV) chloride
78. An aromatic ether is not cleaved by HI even at 525 K. The compound is:

- a)  $\text{C}_6\text{H}_5\text{OCH}_3$                       b)  $\text{C}_6\text{H}_5\text{OC}_6\text{H}_5$                       c)  $\text{C}_6\text{H}_5\text{OC}_3\text{H}_7$











79. Phenol does not react with:  
 a)  $\text{Na}_2\text{CO}_3$                       b) NaOH                      c)  $\text{NaHCO}_3$                       d) KOH
80.  $[\text{EDTA}]^{4-}$  is a  
 a) Monodentate ligand                      b) Bidentate ligand  
 c) Quadridentate ligand                      d) Hexadentate ligand
81.  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Br}_2$  and  $[\text{Pt}(\text{NH}_3)_4\text{Br}_2]\text{Cl}_2$  are related to  
 a) Optical isomer                      b) Linkage isomers                      c) Coordinate isomers                      d) Ionization isomers
82. Ferrocene is an example of  
 a) Sand-wiched complex  
 b) Pi-bonded complex  
 c) A complex in which all the five carbon atoms of cyclopentadiene anion are bonded to the metal  
 d) All of the above
83. Which compound is zero valent metal complex?  
 a)  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$                       b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$                       c)  $[\text{Ni}(\text{CO})_4]$                       d)  $\text{K}_3[\text{Fe}(\text{CN})_6]$
84. Which of the following compounds is 2, 2, 3-trimethyl hexane?  
 a)  $(\text{CH}_3)_3\text{CCH}(\text{CH}_3)\text{CH}_2\text{CH}_3$                       b)  $(\text{CH}_3)_3\text{CCH}_2(\text{CH}_3)_2$   
 c)  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$                       d)  $(\text{CH}_3)_3\text{CCH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$
85. The formula of phenoxy benzene is:  
 a)  $\text{C}_6\text{H}_5\text{C}_6\text{H}_5$                       b)  $\text{C}_6\text{H}_5-\text{O}-\text{C}_6\text{H}_5$                       c)  $\text{C}_6\text{H}_5-\text{O}-\text{C}_6\text{H}_6$                       d) None of these

86. Ziegler-Natta catalyst is an organometallic compound containing  
 a) Iron                                      b) Titanium                                      c) Rhodium                                      d) Zirconium
87. Ziegler-Natta catalyst is  
 a)  $(\text{Ph}_3\text{P})_3\text{RhCl}$                                       b)  $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$                                       c)  $[\text{Al}_2(\text{C}_2\text{H}_6)_6 + \text{TiCl}_4]$                                       d)  $[\text{Fe}(\text{C}_2\text{H}_5)_2]$
88. The tendency to show complex formation is maximum in ....elements.  
 a) s-block                                      b) p-block                                      c) d-block                                      d) f-block
89. EDTA has coordination number  
 a) 3                                      b) 4                                      c) 5                                      d) 6
90. Which of the following is used in Friedel-Craft's acylation reaction?  
 a)                                       b)  $\text{CH}_3\text{CH}_2\text{Cl}$                                       c)  $\text{CH}_3\text{COOCH}_3$                                       d)  $\text{CH}_3\text{Cl}$
91. The correct IUPAC name of  $\text{Mn}_3(\text{CO})_{12}$  is  
 a) Dodacacarbonyl maganate (0)                                      b) Dodacacarbonyl maganate (II)  
 c) Didacacarbonyl trimaganese (0)                                      d) Manganic dodecacarbonyl (0)
92. The  $\pi$  -bonded organometallic compound which has ethene as one of its component is  
 a) Zeise's salt                                      b) Ferrocene                                      c) Dibenzene chromium                                      d) Tetraethyl tin
93. IUPAC name of the compound  
 is  
 a) Ethyl-2-methyl-2-(m-nitro) phenyl propanoate                                      b) Ethyl-2-methyl-2-(o-nitro) phenyl propanoate  
 c) Ethyl-2-methyl-2-(3-nitro) phenyl propanoate                                      d) Ethyl-2-methyl-2-(3-nitro) phenyl propanoic acid
94. What is the product obtained in the following reaction:  
  
 a)                                       b)   
 c)                                       d) 
95.  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is called:  
 a) Hexaammine cobalt (III) chloride  
 b) Amino cobalt chloride (III)  
 c) Cobalt chloride hexaammine  
 d) Hexaammine tricobalt chloride
96. The complexes  $[\text{PtCl}_2(\text{NH}_3)_4]\text{Br}_2$  and  $[\text{PtBr}_2(\text{NH}_3)_4]\text{Cl}_2$  are example for isomerism  
 a) Geometrical                                      b) Optical                                      c) Ionization                                      d) Linkage
97. Geometrical shapes of the complexes formed by the reaction of  $\text{Ni}^{2+}$  with  $\text{Cl}^-$ ,  $\text{CN}^-$  and  $\text{H}_2\text{O}$ , respectively, are  
 a) Octahedral, tetrahedral and square planar                                      b) Tetrahedral, square planar and octahedral  
 c) Square planar, tetrahedral and octahedral                                      d) Octahedral, square planar and octahedral
98. Identify the correct order of reactivity in electrophilic substitution reactions of the following compounds:  
  
 a)  $1 > 2 > 3 > 4$                                       b)  $4 > 3 > 2 > 1$                                       c)  $2 > 1 > 3 > 4$                                       d)  $2 > 3 > 1 > 4$

99. The centric formula for benzene was proposed by:  
 a) Dewar  
 b) Armstrong and Baeyer  
 c) Ladenberg  
 d) Kekule
100. Which is the correct statement?  
 a) Benzyl alcohol is more acidic than phenol  
 b) Ethanol is a powerful oxidizing agent  
 c) Phenol is more acidic than propanol  
 d) Ethane has high boiling point than ethanol
101. Phenol on sulphonation gives:  
 a) *o*-phenol sulphonic acid  
 b) *p*-phenol sulphonic acid  
 c) *m*-phenol sulphonic acid  
 d) Mixture of *o*-and *p*-phenol sulphonic acids
102. Which of the following organometallic compound is  $\sigma$  and  $\pi$  bonded?  
 a)  $\text{Fe}(\text{CH}_3)_3$                       b)  $[\text{Co}(\text{CO})_5\text{NH}_3]^{2+}$                       c)  $[\text{Fe}(\eta^5 - \text{C}_5\text{H}_5)_2]$                       d)  $\text{K}[\text{PtCl}_3(\eta^2 - \text{C}_2\text{H}_4)]$
103. The number of double bonds in BHC (gammexane) is:  
 a) 1                                      b) 2                                      c) 3                                      d) Zero
104. Given the molecular formula of the hexa coordinated complexes (A)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  (B)  $\text{CoCl}_3 \cdot 5\text{NH}_3$  (C)  $\text{CoCl}_3 \cdot 4\text{NH}_3$ . If the number of coordinated  $\text{NH}_3$  molecules in A, B and C respectively are 6, 5 and 4, primary valency in (A), (B) and (C) are  
 a) 0, 1, 2                              b) 3, 2, 1                              c) 6, 5, 4                              d) 3, 3, 3
105. Type of isomerism shown by  $[\text{Cr}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$  is  
 a) Optical                              b) Ionisation                              c) Geometrical                              d) Linkage
106.  $[\text{Sc}(\text{H}_2\text{O})_6]^{3+}$  ion is  
 a) Colourless and diamagnetic                              b) Coloured and octahedral  
 c) Colourless and paramagnetic                              d) Coloured and paramagnetic
107. Which one of the following octahedral complexes will not show geometrical isomerism? (A and B are monodentate ligands)  
 a)  $[\text{MA}_4\text{B}_2]$                               b)  $[\text{MA}_5\text{B}]$                               c)  $[\text{MA}_2\text{B}_4]$                               d)  $[\text{MA}_3\text{B}_3]$
108. The IUPAC name of the following compound is  

$$\begin{array}{c} \text{O}=\text{C}-\text{CH}-\text{CH}_2 \\ | \quad | \quad | \\ \text{OH} \quad \text{NH}_2 \quad \text{OH} \end{array}$$
  
 a) 3-amino-2-hydroxy propanoic acid                              b) 2-aminopropan-3-ol-1-oic acid  
 c) 2-amion-3-hydroxy propanoic acid                              d) Aminohydroxy propanoic acid
109. Which of the following complex ion is not expected to absorb visible light?  
 a)  $[\text{Ni}(\text{CN})_4]^{2-}$   
 b)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$   
 c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$   
 d)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$
110. The correct sequence of activating power of a group in benzene is:  
 a)  $-\text{NH}_2 > -\text{NHCOCH}_3 > -\text{CH}_3$   
 b)  $-\text{NH}_2 < -\text{NHCOCH}_3 < -\text{CH}_3$   
 c)  $-\text{NH}_2 > -\text{NHCOCH}_3 < -\text{CH}_3$   
 d)  $-\text{NH}_2 < -\text{NHCOCH}_3 > -\text{CH}_3$
111. The pair of compounds having metals in their highest oxidation state is  
 a)  $\text{MnO}_2, \text{FeCl}_3$                               b)  $[\text{MnO}_4]^- , \text{CrO}_2\text{Cl}_2$   
 c)  $[\text{Fe}(\text{CN})_6]^{3-} , [\text{Co}(\text{CN})_3]$                               d)  $[\text{NiCl}_4]^{2-} , [\text{CoCl}_4]^-$
112. Total number of geometrical isomers for the complex  $[\text{RhCl}(\text{CO})(\text{PPh}_3)(\text{NH}_2)]$  is



- a) 1                                      b) 2                                      c) 3                                      d) 4
113. The reaction of chloroform with alc. KOH and *p*-toluidine forms:
- a) 
- b) 
- c) 
- d) 
114. Which order is correct in spectrochemical series of ligands?
- a)  $\text{Cl}^- < \text{F}^- < [\text{C}_2\text{O}_4]^{2-} < \text{NO}_2^- < \text{CN}^-$
- b)  $\text{CN}^- < [\text{C}_2\text{O}_4]^{2-} < \text{Cl}^- > \text{NO}_2^- < \text{F}^-$
- c)  $[\text{C}_2\text{O}_4]^{2-} < \text{F}^- < \text{Cl}^- > \text{NO}_2^- < \text{CN}^-$
- d)  $\text{F}^- < \text{Cl}^- < \text{NO}_2^- < \text{CN}^- < [\text{C}_2\text{O}_4]^{2-}$
115. The IUPAC name of compound  $\text{K}_3[\text{Fe}(\text{CN})_5\text{NO}]$  is
- a) Pentacyano nitrosyl potassium ferrate(II)                                      b) Potassium cyano pentanitrosyl ferrate(II)
- c) Potassium pentacyanonitrosyl ferrate (III)                                      d) Potassium pentacyanonitrosyl ferrate (II)
116. The colour of  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$  is due to:
- a) Transfer of an electron from one Ti to another
- b) Presence of water molecule
- c) Excitation of electrons from *d* – *d*
- d) Intramolecular vibration
117. The oxidation number of Fe in  $\text{K}_4[\text{Fe}(\text{CN})_6]$  is
- a) +3                                      b) +4                                      c) +2                                      d) –2
118. Correct structures of [E][S]-5-bromo-2,7-dimethyl, non-4-ene is
- a) 
- b) 
- c) 
- d) 
119. Name the metal *M* which is extracted on the basis of following reactions,
- $4\text{M} + 8\text{CN}^- + 2\text{H}_2\text{O} + \text{O}_2 \rightarrow 4[\text{M}(\text{CN})_2]^- + 4\text{OH}^-$
- $2[\text{M}(\text{CN})_2]^- + \text{Zn} \rightarrow [\text{Zn}(\text{CN})_4]^{2-} + 2\text{M}$ :
- a) Nickel                                      b) Silver                                      c) Copper                                      d) Mercury
120. EAN of Cr in  $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$  is:
- a) 32                                      b) 33                                      c) 34                                      d) 35
121. The complex  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$  furnishes:
- a) 5 ions                                      b) 6 ions                                      c) 4 ions                                      d) 2 ions
122. Ammoniacal solution of  $\text{Ni}(\text{CN})_2$  reacts with  $\text{C}_6\text{H}_6$  to produce a light violet coloured crystalline compound of the formula:
- a)  $\text{Ni}(\text{CN})_2 \cdot \text{C}_6\text{H}_5$                                       b)  $\text{C}_6\text{H}_5\text{CH}_3$                                       c)  $\text{Ni}(\text{CN})_2\text{C}_6\text{H}_6$                                       d)  $\text{Ni}(\text{CN})_2\text{NH}_3 \cdot \text{C}_6\text{H}_6$
123. Ammonia forms the complex ion  $[\text{Cu}(\text{NH}_3)_4]^{2+}$  with copper ions in alkaline solution but not in acidic solution. What is the reason for it?
- a) In acidic solutions, hydration protects copper ions

- b) In alkaline solution, insoluble  $\text{Cu}(\text{OH})_2$  is precipitated which in excess of any alkali  
 c) Copper hydroxide is an amphoteric substance  
 d) In acidic solutions, protons coordinate with ammonia molecules forming  $\text{NH}_4^+$  ions and  $\text{NH}_3$  molecules are not available

124. Which of the following shows geometrical isomerism?

- a) 1, 2-dichloroethane  
 b) 1, 2-dimethylcyclopropane  
 c)  $\text{CH}_3\text{CH} \begin{array}{c} \text{CO—NH} \\ \text{NH—CO} \end{array} \text{CHCH}_3$   
 d) All of the above


125. The shape of the complex  $[\text{Ag}(\text{NH}_3)_2]^+$  is:

- a) Octahedral  
 b) Square planar  
 c) Tetrahedral  
 d) Linear

126. The  $\pi$ -bonded organometallic compound which has ethane as one of its component is

- a) Dibenzene chromium  
 b) Zeise salt  
 c) Ferrocene  
 d) Tetraethyl tin

127. The major product of the following reaction is:




- a)
- b)
- c)
- d)

128. Which is true in the case of  $\text{Ni}(\text{CO})_4$  complex?


- a) Hybridization of Ni is  $sp^3$   
 b) Tetrahedral shape of the molecule  
 c) Diamagnetic  
 d) All are correct


129. The reaction,  $C_6H_5N_2Cl \xrightarrow{Cu_2Cl_2/HCl} C_6H_5Cl + N_2$  is called:  
 a) Etard's reaction      b) Sandmeyer's reaction      c) Wurtz-Fittig reaction      d) Perkin's reaction
130. Which of the following does not show optical isomerism?  
 a)  $[Co(en)_3]^{3+}$       b)  $[Co(en)_2Cl_2]^+$       c)  $[Co(NH_3)_3Cl_3]^0$       d)  $[Co(en)Cl_2(NH_3)_2]^+$



Having the IUPAC name as





- a) 1, 2-dimethyl cyclobutane      b) 2, 3-dimethyl cyclobutene  
 c) 2, 3-dimethyl butane      d) 1, 2-dimethyl cyclobut-1-ene
132. Which of the following ions is produced when we prepare nitrating mixture by mixing together concentrated  $HNO_3$  and concentrated  $H_2SO_4$ ?  
 a)  $NO_2^-$       b)  $NO_2^+$       c)  $NO_3^-$       d)  $SO_3^+H$
133. The correct IUPAC name of




- a) 1-bromo-2-chloro-6-fluoro-4-iodobenzene      b) 1-bromo-6-chloro-2-fluoro-4-iodobenzene  
 c) 2-bromo-1-chloro-3-fluoro-5-iodobenzene      d) 2-bromo-3-chloro-1-fluoro-5-iodobenzene
134.  $[Co(NH_3)_4(NO_2)_2]Cl$  exhibits:  
 a) Ionization isomerism, geometrical isomerism and optical isomerism  
 b) Linkage isomerism, geometrical isomerism and optical isomerism  
 c) Linkage isomerism, ionization isomerism and optical isomerism  
 d) Linkage isomerism, ionization isomerism and geometrical isomerism
135. Which of the following complexes are not correctly matched with hybridisation of their central metal ion?  
 1.  $[Ni(CO)_4]$        $sp^3$   
 2.  $[Ni(CO)_4]^{2-}$        $sp^3$   
 3.  $[CoF_6]^{3-}$        $d^2sp^3$   
 4.  $[Fe(CN)_6]^{3-}$        $sp^3d^2$
- Select the correct answer using the codes given below  
 a) 1 and 2      b) 1 and 3      c) 2 and 4      d) 2, 3 and 4
136. Which of the following is an explosive?  
 a)  $PCl_5$       b)  $HNO_3$       c)  $C_6H_5OH$       d) 2,4,6-trinitrophenol
137. The coordination number of Cr in  $[Cr(NH_3)_3(H_2O)_3]Cl_3$  is:  
 a) 3      b) 4      c) 6      d) 2
138. The major product obtained when 3-phenyl-1, 2-propane-diol is heated with  $H_2SO_4$  is:  
 a)  $C_6H_5-CH_2-CO-CH_3$   
 b)  $C_6H_5-CH_2-CH_2-CHO$   
 c)  $C_6H_5-CH_2-CH=CH_2$   
 d) 

139. Rate of substitution in phenol is:  
 a) Slower than as in benzene  
 b) Faster than as in benzene  
 c) Equal to that as in benzene  
 d) None of the above

140. Magnetic moment of  $[Ag(CN)_2]^-$  is zero. How many unpaired electrons are there?

- a) Zero                                      b) 4                                      c) 3                                      d) 1
141. Chlorophyll is a coordination compound having central atom of:  
a) Ca                                      b) Mg                                      c) Na                                      d) K
142. Which of the following statements is incorrect?  
a) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied only the secondary valency of ferric ion.  
b) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferric ion.  
c) In  $K_4[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferrous ion.  
d) In  $[Cu(NH_3)_4]SO_4$ , the ligand has satisfied only the secondary valency of copper.
143. Maximum number of open chain isomers that an alkene can have with the molecular formula  $C_4H_8$  is  
a) 5                                      b) 4                                      c) 3                                      d) 2
144. Which one is the wrong statement?  
a) Open chain compounds are called aliphatic  
b) Unsaturated compounds contain multiple bonds in them  
c) Saturated hydrocarbons are called alkene  
d) Aromatic compounds possess a characteristic aroma
145. According to postulates of Werner's theory for coordination compounds, which of the following is true?  
a) Primary valencies are ionizable                                      b) Secondary valencies are ionizable  
c) Only primary valencies are non-ionizable                                      d) Primary and secondary valencies are non-ionizable
146. Atomic numbers of Cr and Fe are respectively 24 and 26. Which of the following is paramagnetic with the spin of the electron?  
a)  $[Cr(CO)_6]$                                       b)  $[Fe(CO)_5]$                                       c)  $[Fe(CN)_6]^{4-}$                                       d)  $[Cr(NH_3)_6]^{3+}$
147. Which of the following structures correspond to the product expected, when excess of  $C_6H_6$  reacts with  $CH_2Cl_2$  in presence of anhy.  $AlCl_3$  ?
- a)       b)       c)       d) 
148. Which of the following will give a pair of enantiomorphs?  
a)  $[Co(en)_2Cl_2]Cl$                                       b)  $[Cr(NH_3)_6][Co(CN)_6]$   
c)  $[Pt(NH_3)_4][PtCl_6]$                                       d)  $[Co(NH_3)_4Cl_2]NO_2$
149. The crystal field splitting energy for octahedral ( $\Delta_0$ ) and tetrahedral ( $\Delta_t$ ) complexes is related to  
a)  $\Delta_t = \frac{4}{9} \Delta_0$                                       b)  $\Delta_t = \frac{1}{2} \Delta_0$                                       c)  $\Delta_0 = 2\Delta_t$                                       d)  $\Delta_0 = \frac{4}{9} \Delta_t$
150. The correct name of the compound  $[Cu(NH_3)_4](NO_3)_2$ , according to IUPAC system is:  
a) Cuprammonium nitrate  
b) Tetraamminecopper(II) dinitrate  
c) Tetraamminecopper(II) nitrate  
d) Tetraamminecopper(I) dinitrate
151. Which among the following will not show chain isomerism?  
a)  $C_3H_8$                                       b)  $C_4H_{10}$                                       c)  $C_5H_{12}O$                                       d)  $C_5H_{10}O$
152. Phenol (1 mole) reacts with bromine to give s-tribromophenol. How much bromine is needed?  
a) 1.5 mole                                      b) 3.0 mole                                      c) 4.5 mole                                      d) 6.0 mole
153. Dimethyl glyoxime forms a coloured complex with  
a) Ag                                      b) Ni                                      c) Cr                                      d) Zn
154. Which has regular tetrahedral geometry?  
a)  $[Ni(CN)_4]^{2+}$                                       b)  $SF_4$                                       c)  $[BF_4]^-$                                       d)  $XeF_4$
155. In haemoglobin the iron shows oxidation state:  
a) +2                                      b) +3                                      c) +1                                      d) +4
156. For the given complex  $[CoCl_2(en)(NH_3)_2]^+$ , the number of geometrical isomers, the number of optical

- isomers and total number of isomers of all type possible respectively are  
a) 2, 2 and 4                      b) 2, 2 and 3                      c) 2, 0 and 2                      d) 0, 2 and 2
157. Which can show aromatic character?  
a) Furan                      b) Pyrrol                      c) Benzene                      d) All of these
158. Of the following complexes, the one with the largest value of the crystal field splitting is:  
a)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$                       b)  $[\text{Ru}(\text{CN})_6]^{3-}$                       c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$                       d)  $[\text{Fe}(\text{NH}_3)_6]^{3+}$
159. The specific rotation of a pure enantiomer is  $+16^\circ$ . Its observed rotation if it is isolated from a reaction with 25% racemisation and 75% retention is  
a)  $-12^\circ$                       b)  $+12^\circ$                       c)  $+16^\circ$                       d)  $-16^\circ$
160. Lithium tetrahydridoaluminate is correctly represented as:  
a)  $\text{Al}[\text{LiH}_4]$                       b)  $\text{Al}_2[\text{LiH}_4]_3$                       c)  $\text{Li}[\text{AlH}_4]$                       d)  $\text{Li}[\text{AlH}_4]_2$
161. Which of the following compounds is generally used for hydrogenation of alkenes?  
a)  $\text{Ni}(\text{CO})_4$                       b)  $[(\text{C}_6\text{H}_5)_3\text{P}]_3\text{RhCl}$                       c)  $(\text{CH}_3)_3\text{Al}$                       d)  $(\text{C}_5\text{H}_5)_2\text{Fe}$
162. The end product of the reaction,  
 $\text{C}_6\text{H}_6 + \text{Cl}_2 \xrightarrow{\text{Sunlight}}$  is:  
a)  $\text{C}_6\text{H}_5\text{Cl}$                       b)  $o\text{-C}_6\text{H}_4\text{Cl}_2$                       c)  $\text{C}_6\text{H}_6\text{Cl}_6$                       d)  $p\text{-C}_6\text{H}_4\text{Cl}_2$
163.  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$  complex gives  
a) 4 ions                      b) 3 ions                      c) 2 ions                      d) 5 ions
164. Which does not obey EAN rule?  
a)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$                       b)  $[\text{Zn}(\text{OH})_4]^{2-}$                       c)  $[\text{HgI}_4]^{2-}$                       d)  $\text{Fe}(\text{CO})_5$
165. Oxidation number of Fe in  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is:  
a) +3                      b) +2                      c) +10                      d) 1
166. Which of the following is not an organometallic compound?  
a)  $\text{NaOC}_2\text{H}_5$                       b)  $(\text{CH}_3)_3\text{Al}$                       c)  $(\text{C}_2\text{H}_5)_4\text{Pb}$                       d)  $\text{RMgX}$
167. Considering  $\text{H}_2\text{O}$  as weak field ligand, the number of unpaired electrons in  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$  will be (Atomic no. of Mn=25)  
a) Three                      b) Five                      c) Two                      d) Four
168. The value of 'spin only' magnetic moment for one of the following configuration is 2.84 BM the correct one  
a)  $d^4$  (in weak ligand field)                      b)  $d^4$  (in strong ligand field)  
c)  $d^3$  (in weak as well as in strong field)                      d)  $d^5$  (in weak ligand field)
169. Fluorobenzene ( $\text{C}_6\text{H}_5\text{F}$ ) can be synthesized in the laboratory:  
a) By heating phenol with HF and KF  
b) From aniline by diazotisation followed by heating the diazonium salt with  $\text{HBF}_4$   
c) By direct fluorination of benzene with  $\text{F}_2$  gas  
d) By reacting bromobenzene with NaF solution
170. Which compound burns with a sooty flame?  
a)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$   
b)  $\text{C}_6\text{H}_5\text{COOH}$   
c)  $\text{CH}_3\text{OH}$   
d)  $\text{CH}_3\text{COC}_2\text{H}_5$
171. How many EDTA (ethylenediaminetetraacetic acid) molecules are required to make an octahedral complex with a  $\text{Ca}^{2+}$  ion?  
a) Six                      b) Three                      c) One                      d) Two
172. Intramolecular rearrangement of phenyl esters to give *o*- and *p*-derivatives in presence of  $\text{AlCl}_3$  is known as:  
a) Friedel-Craft's reaction  
b) Fries rearrangement  
c) Esterification  
d) Coupling
173. Which reaction can produce  $\text{R}-\text{CO}-\text{Ar}$  species?







174. Acidic character of phenol is due to:

- a) Resonance of phenoxide ion
- b) Tautomerism occurring in phenol
- c) The fact that the electronegativity of oxygen is more than that of hydrogen
- d) None of the above

175. In triethylenediamine cobalt(III) chloride the coordination number of cobalt is:

- a) 3
- b) 4
- c) 6
- d) 7

176. Mark the unidentical compound



177. A complex compound in which the oxidation number of a metal is zero, is

- a)  $\text{K}_4[\text{Fe}(\text{CN})_6]$
- b)  $\text{K}_3[\text{Fe}(\text{CN})_6]$
- c)  $[\text{Ni}(\text{CO})_4]$
- d)  $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_2$

178. In the halogenation of aromatic nucleus, the halogen carrier, used to generate the species is:

- a)  $\text{Cl}$
- b)  $\text{Cl}^+$
- c)  $\text{Cl}^-$
- d)  $\text{Cl}$

179. Among  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CO})_4]$ :

- a)  $[\text{Ni}(\text{CN})_4]^{2-}$  is square planar and  $[\text{NiCl}_4]^{2-}$ ,  $\text{Ni}(\text{CO})_4$  are tetrahedral
- b)  $[\text{NiCl}_4]^{2-}$  is square planar and  $[\text{NiCN}_4]^{2-}$ ,  $\text{Ni}(\text{CO})_4$  are tetrahedral
- c)  $\text{Ni}(\text{CO})_4$  is square planar and  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{NiCl}_4]^{2-}$  are tetrahedral
- d) None of the above

180. Benzene is obtained by:

- a) Condensation of three  $\text{C}_2\text{H}_2$  molecules
- b) Polymerization of three  $\text{C}_2\text{H}_2$  molecules
- c) Addition of three  $\text{C}_2\text{H}_2$  molecules
- d) Substitution of three acetylene molecules

181. IUPAC name of *t*-butyl chloride is

- a) 2-chloro butane
- b) 1-chloro-2-methylpropane
- c) 2-chloro-2-methylpropane
- d) None of the above

182. The *d*-electronic configuration of  $\text{Cr}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Ni}^{2+}$  are  $3d^4$ ,  $3d^5$ ,  $3d^6$  and  $3d^8$  respectively. Which of the following complex will show minimum paramagnetic behaviour?

- a)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$
- b)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$
- c)  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$
- d)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$

183. Phenol is more acidic than cyclohexanol because:

- a) Benzene ring exists in resonance
- b) Cyclohexane ring shows resonance
- c) Phenol is poor in hydrogen
- d) Cyclohexanol is rich in hydrogen

184. Total possible structural isomers (not stereo) of  $\text{C}_4\text{H}_6$  are

- a) 4
- b) 6
- c) 9
- d) 12

185. In the reaction of *p*-chlorotoluene with  $\text{KNH}_2$  in liq.  $\text{NH}_3$  the major product is:

- a) *o*-toluidine
- b) *m*-toluidine
- c) *p*-toluidine
- d) *p*-chloroaniline

186. The type of isomerism in the molecule of compounds  $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$  and  $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$  is referred as:

- a) Metamerism
- b) Chain isomerism
- c) Functional isomerism
- d) Tautomerism

187. Phenol is less soluble in water. It is due to:

- a) Non-polar nature of phenol

- b) Acidic nature of—OH group  
c) Non-polar hydrocarbons part in it  
d) None of the above

188. When phenol is treated with excess bromine water, it gives:


- a) *m*-bromophenol  
b) *o*-and *p*-bromophenol  
c) 2,4-dibromophenol  
d) 2,4,6-tribromophenol

189. Which have octahedral shape ( $d^2sp^3$ ) hybridization of central atom?

- a)  $[\text{Cr}(\text{NH}_3)_6]^{2+}$       b)  $[\text{Fe}(\text{CN})_6]^{3-}$       c)  $[\text{Cu}(\text{NH}_3)_6]^+$


d) All are correct

190. Which of the following molecules/species are aromatic in character?



- a) 2      b) 3      c) 4      d) 1


191. Among the following compounds ;



the order of basicity is :

- a) IV > III > II > I      b) II > I > III > IV      c) III > IV > II > I      d) I > III > IV > II

192.



- a) Tri- $\mu$ -carbonyl *bis*-(tricarbonyl)iron (0)  
b) Hexacarbonyl iron (III)  $\mu$ -tricarbonyl ferrate(0)  
c) Tricarbonyl iron(0)  $\mu$ -tricarbonyl iron(0) tricarbonyl  
d) Nonacarbonyl iron

193. Which is high spin complex?

- a)  $[\text{CoCl}_6]^{3-}$       b)  $[\text{FeF}_6]^{3-}$       c)  $[\text{Co}(\text{NH}_3)_6]^{2+}$       d) All are correct

194. The correct IUPAC name of tartaric acid is

- a) 1, 4-dicarboxy-2, 3-dihydroxy ethane      b)  $\alpha, \alpha'$ -dihydroxy butane-1,4-dioic acid  
c) 1, 4-dihydroxybutane-2, 3-dioic acid      d) 2, 3-dihydroxybutane-1, 4-dioic acid


195. What is the overall formation equilibrium constant for the ion  $[\text{ML}_4]^{2-}$  ion, given that  $\beta_4$  for this complex is  $2.5 \times 10^{13}$ ?

- a)  $2.5 \times 10^{13}$       b)  $5 \times 10^{-13}$       c)  $2.5 \times 10^{-14}$       d)  $4.0 \times 10^{-13}$

196. The oxidation state of Cr in  $[\text{Cr}(\text{NH}_3)_4 \text{Cl}_2]^+$  is

- a) 0      b) +1      c) +2      d) +3

197. Which of the following compounds has the most acidic nature?



198. The oxidation state of Mo in its oxo-complex species  $[\text{Mo}_2\text{O}_4(\text{C}_2\text{H}_4)_2(\text{H}_2\text{O})_2]^{2-}$  is:

- a) +2      b) +3      c) +4      d) +5


199.  $\text{CH}_3\text{MgI}$  is an organometallic compound due to

- a) Mg—I bond      b) C—I bond      c) C—Mg bond      d) C—H bond


200. The effective atomic number of Cr (At. No.=24) in  $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$  is




- a) 35                      b) 27                      c) 33                      d) 36
201. When aniline is heated with benzaldehyde, the product is:  
 a) Benzoin                      b) Schiff's base                      c) Unsaturated acid                      d) Azoxy benzene
202. Slow heating of salicylic acid gives:  
 a) Benzoic acid                      b) Phenol                      c) Benzaldehyde                      d) None of these
203. According to Hückel, monocyclic compounds will show aromaticity when:  
 a) It has  $4\pi$ -electrons  
 b) It has no  $\pi$ -electron  
 c) It has  $4\pi+2$  electrons  
 d) It has  $(4n+2)\pi$ -electrons
204. When phenol is distilled with zinc dust, it gives:  
 a) Benzene                      b) Toluene                      c)  $C_6H_5CHO$                       d) None of these
205. The IUPAC name of the given structure



- a) N-chloro-N-bromoethanamide                      b) N-bromo-N-chloroethanamide  
 c) N-bromo-N-chloroacetamide                      d) N-chloro-N-bromoacetamide
206. Acetophenone when reacted with a base  $C_2H_5ONa$ , yields a stable compound which has the structure:







207. Which of the following has maximum resonance energy?  
 a) Diphenyl                      b) Benzene                      c) Naphthalene                      d) Phenanthrene
208. Benzene sulphonic acid on treating with  $P_2O_5$  gives:  
 a) Salicylic acid                      b) Benzoic acid                      c) Acid anhydride                      d) Sodium benzoate
209. Compounds with following formula will show



- a) Position and functional isomerism  
 b) Chain and positional isomerism  
 c) Chain and functional isomerism  
 d) None of the above combinations
210. Which of the following statements is correct?  
 a) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferric ion  
 b) In  $(Cu(NH_3)_4)SO_4$ , the ligand has satisfied only the secondary valency of copper

- c) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied only the secondary valency of ferric ion  
 d) Both (b) and (c)
211. Which statement is not correct?  
 a)  $Fe(CO)_5$  reacts with  $Br_2Cl_4$   
 b) Carbonyl complexes are usually formed with transition metals  
 c) All transition metals form monometallic carbonyls  
 d) The decomposition of  $Ni(CO)_4$  to give Ni is used in the extraction of Ni by Mond's process
212. The complex showing a spin-only magnetic moment of 2.82 BM is  
 a)  $Ni(CO)_4$                       b)  $[NiCl_4]^{2-}$                       c)  $Ni(PPh_3)_4$                       d)  $[Ni(CN)_4]^{2-}$
213. The IUPAC name of  $[CoCl(NO_2)(en)_2]Cl$  is:  
 a) Chloronitro-*bis*(ethylenediamine) cobaltic(III) chloride  
 b) Chloronitro-*bis*(ethylenediamine)cobalt(II) chloride  
 c) Chloro-*bis*(ethylenediamine)nitrocobalt(III) chloride  
 d) *Bis*-(ethylenediamine)chloronitrocobalt(III) chloride
214. The product of acid catalysed hydration of 2-phenyl propene is:  
 a) 3-phenyl-2-propanol  
 b) 1-phenyl-2-propanol  
 c) 2-phenyl-2-propanol  
 d) 2-phenyl-1-propanol
215. Carbolic acid is the name used for:  
 a) Opium                      b) Phenol                      c) Chloroform                      d)  $H_2CO_3$
216. The major product of the following reaction





217. The oxidation number of cobalt in  $K[Co(CO)_4]$  is

- a) -1                      b) +3                      c) +1                      d) -3

218. Formaldehyde-phenol resin is:

- a) Orlon                      b) Nylon                      c) Teflon                      d) Bakelite

219. Among the ligands  $NH_3$ , en,  $CN^-$  and CO, the correct order of their increasing field strength, is

- a)  $CO < NH_3 < en < CN^-$                       b)  $NH_3 < en < CN^- < CO$   
c)  $CN^- < NH_3 < CO < en$                       d)  $en < CN^- < NH_3 < CO$

220. Cyclopentadienyl anion is aromatic due to the presence of:

- a)  $6\pi$ -electrons                      b)  $10\pi$ -electrons                      c)  $4\pi$ -electrons                      d)  $12\pi$ -electrons


221. The IUPAC name of  $K_4[Fe(CN)_6]$  is

- a) Potassium ferrocyanide                      b) Potassium hexa cyanoferrate (I)  
c) Tetra potassium hexa cyanoferrate (II)                      d) Potassium hexa cyanoferrate (II)

222. Which xylene is most easily sulphonated?

- a) *Ortho*                      b) *Para*                      c) *Meta*                      d) All at the same rate

223. The IUPAC name of following polyfunctional compound is



- a) 2,4-dioxo cyclohexanoic acid                      b) 2,4-dioxo cycloheptanoic acid  
c) 4-formyl-2-oxo cyclohexane-1-carboxylic acid                      d) 2,4-dioxo cyclohexane-1-carboxylic acid


224. Alkyl groups are *o*- and *p*-directing because of:



- a) Resonance effect  
b) Inductive effect  
c) Resonance effect through hyperconjugation  
d) All of the above

225. Racemic modification can be resolved by

- a) The use of enzymes                      b) Fractional crystallisation  
c) Fractional distillation                      d) None of the above

226. Which of the following structure contain 1 primary and 7 secondary hydrogen atoms?





227. Which of the following compounds does not dissolve in conc.  $\text{H}_2\text{SO}_4$  even on warming?

- a) Ethylene                      b) Benzene                      c) Hexane                      d) Aniline

228. In the complex  $\text{Fe}(\text{CO})_x$ , the value of  $x$  is and it is:

- a) 3, octahedral                      b) 4, tetrahedral                      c) 5, trigonal pyramidal                      d) 6, square pyramidal

229. The empirical formula of naphthalene is:

- a)  $\text{CH}_2$                       b)  $\text{C}_5\text{H}_4$                       c)  $\text{C}_2\text{H}$                       d)  $\text{C}_n\text{H}_{2n}$

230. The chemical formula of diammine silver (I) chloride is

- a)  $[\text{Ag}(\text{NH}_3)\text{Cl}]$                       b)  $[\text{Ag}(\text{NH}_3)_3]\text{Cl}$                       c)  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$                       d)  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$

231. For the square planar complex  $[\text{M}(a)(b)(c)(d)]$  (where,  $\text{M}$ =central metal and  $a, b, c$ , and  $d$  are monodentate ligands), the number of possible geometrical isomers are

- a) 1                      b) 2                      c) 3                      d) 4

232. Which group is *meta* directing?

- a)  $-\text{CCl}_3$                       b)  $-\text{OH}$                       c)  $-\text{NH}_2$                       d)  $-\text{CH}_3$

233. The IUPAC name of the compound  $[\text{Cu}(\text{NH}_3)_4(\text{NO}_3)_2]$  is:

- a) Cuprammonium nitrate  
b) Dinitratotetraamminecopper(II)  
c) Tetraamminecopper(II) dinitrite  
d) Tetraamminecopper(III) dinitrite

234. Coordination number of Fe in the complexes  $[\text{Fe}(\text{CN})_6]^{4-}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$  and  $[\text{FeCl}_4]^-$  would be respectively

- a) 6, 4, 6                      b) 6, 6, 4                      c) 6, 3, 3                      d) 2, 3, 3

235. Which statement is true for cyclohexane?

- a) It has two possible isomers  
b) It has three conformations  
c) Boat conformation is most stable  
d) Chair and boat conformations differ in energy by 44 kJ/mol

236. Ligands in a complex salt are:

- a) Anions linked by coordinate bonds to a central metal atom or ion  
b) Cations linked by coordinate bonds to a central metal atom or ion  
c) Molecules linked by coordinate bonds to a central metal atom or ion  
d) Ions or molecules linked by coordinate bonds to a central metal atom or ion

237. The IUPAC name of  $\text{CH}_3\text{COCH}_2\text{COOC}_2\text{H}_5$  is


- a) Ethyl butanoate                      b) Ethyl-(3-oxo)butanoate  
c) Ethyl butan-1-oate-2-one                      d) Ethyl butan-4-oate-2-one

238. When benzene is treated with CO and HCl in presence of anhydrous aluminium chloride, benzaldehyde is formed. This reaction is known as:

- a) Friedel-Craft's reaction  
b) Rosenmund's reaction  
c) Stephen's reaction  
d) Gattermann-Koch's reaction

239.  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  ion is:

- a) Paramagnetic                      b) Diamagnetic                      c) Square planar                      d) None of these
240. The following compound can exhibit
- $$\begin{array}{c}
 \text{CH}_3 \quad \quad \text{H} \\
 \diagdown \quad \diagup \\
 \text{C} = \text{C} \\
 \diagup \quad \diagdown \\
 \text{CH}_3 \quad \quad \text{CH}_3 \quad \quad \text{COOH}
 \end{array}$$
- a) Tautomerism                      b) Optical isomerism  
c) Geometrical isomerism                      d) Geometrical and optical isomerism
241. Which complex is diamagnetic?
- a)  $[\text{Fe}(\text{CN})_6]^{4-}$                       b)  $[\text{Cu}(\text{NH}_3)_4]^{3+}$                       c)  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$                       d) None of these
242. *Meso*-tartaric acid is optically inactive due to the presence of
- a) Molecular symmetry                      b) Molecular asymmetry  
c) External compensation                      d) Two asymmetric C-atoms
243. Complex forming tendency increases with:
- a) Increase in size of cation  
b) Decrease in size of cation  
c) Increase in size of anion  
d) None of the above
244. Ziegler-Natta catalyst is
- a)  $(\text{Ph}_3\text{P})_3\text{RhCl}$                       b)  $\text{Al}_2(\text{C}_2\text{H}_5)_6 + \text{TiCl}_4$   
c)  $\text{Fe}(\text{C}_2\text{H}_5)_2$                       d)  $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$
245. Among the following compounds the one that is most reactive towards electrophilic nitration is:
- a) Toluene                      b) Benzene                      c) Benzoic acid                      d) Nitrobenzene
246. Phenol on oxidation gives chloranil. The oxidant used is:
- a)  $\text{K}_2\text{S}_2\text{O}_8$                       b)  $\text{KMnO}_4$                       c)  $\text{KClO}_3 + \text{HCl}$                       d) None of these
247. The IUPAC name of the compound
- $$\begin{array}{c}
 \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{CH}_3 \\
 | \\
 \text{CH}_3 - \text{CH}_2 - \text{CH} - \text{CH} - \text{CH} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3 \\
 | \quad \quad | \\
 \text{CH}_3 \quad \quad \text{CH}_2 - \text{CH}_3
 \end{array}$$
- a) 3-sec-butyl-5-ethyl-3-methyloctane                      b) 4-sec-butyl-5-ethyl-3-methyloctane  
c) 5-sec-butyl-4-ethyl-3-methyloctane                      d) 4-sec-butyl-3-ethyl-5-methyloctane
248. All the common *m*-directing groups.....the benzene ring towards electrophilic substitution reactions.
- a) Deactivate                      b) Activate                      c) Both (a) and (b)                      d) None of these
249. Among the following, the coloured compound is :
- a)  $\text{CuCl}$                       b)  $\text{K}_3\text{C}_4(\text{CN})_4$                       c)  $\text{CuF}_2$                       d)  $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{BF}_3$
250. The existence of two different coloured complexes with the composition of  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$  is due to:
- a) Linkage isomerism  
b) Geometrical isomerism  
c) Coordination isomerism  
d) Ionisation isomersim
251.  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]$  possesses:
- a) Square planar geometry  
b) Tetrahedral geometry  
c) Tetrahedral nature  
d) Octahedral geometry
252. Which one does not belong to ligand?
- a)  $\text{PH}_3$                       b)  $\text{NO}^+$                       c)  $\text{BF}_3$                       d)  $\text{Cl}^-$
253. Product formed in the reaction;
- $$\text{Phenol} \xrightarrow[\text{Pyridine} - \text{SO}_3/(\text{CH}_3\text{CO})_2\text{O}]{(\text{CH}_3)\text{SO}} \text{Product; is:}$$





254. Which one of the following has square planar structure?

- a)  $[\text{Ni}(\text{CN})_4]^{2-}$       b)  $[\text{Ni}(\text{CO})_4]$       c)  $[\text{NiCl}_4]^{2-}$

d) All of these

255. 4-methyl benzene sulphonic acid reacts with sodium acetate to give:



256. Phthalein test is characteristics of ....and is given by it.

- a) Alcohols      b) Phenols      c) Aldehydes


d) Ketones

257. Which of the following compounds would exhibit coordination isomerism?


- a)  $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_2$       b)  $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$       c)  $[\text{Cr}(\text{en})_2]\text{NO}_2$

d)  $[\text{Ni}(\text{NH}_3)_6][\text{BF}_4]_2$

258. In a reaction of aniline a coloured product C was obtained.



The structure of C would be:



259. The carboxyl functional group ( $-\text{COOH}$ ) is present in:

- a) Picric acid  
b) Barbituric acid  
c) Ascorbic acid  
d) Aspirin

260. Which of the following is an example of electrophilic substitution reaction?

- a) Acylation      b) Alkylation      c) Benzoylation

d) All of these

261. The number of ions given by  $[\text{Co}(\text{NH}_3)_4]\text{Cl}_3$  in aqueous solution is:





- a) 2      b) 3      c) 1

d) 4

262. Which of the following is an organometallic compound?

- a)  $\text{Ti}(\text{OC}_6\text{H}_5)_4$       b)  $\text{Ti}(\text{OCOCH}_3)_4$       c)  $\text{Ti}(\text{OC}_2\text{H}_5)_4$       d)  $\text{Ti}(\text{C}_2\text{H}_5)_4$
263. A solution of  $\text{CuCl}$  in  $\text{NH}_4\text{OH}$  is used to measure the amount of which gas is a sample by simply measuring change in volume?
- a)  $\text{CO}_2$       b)  $\text{H}_2$       c)  $\text{CO}$       d) All of these
264. On passing benzene vapour through a tube at  $700\text{--}800^\circ\text{C}$  or through molten lead we get:
- a) Diphenyl      b) Phenol      c) Toluene      d) Benzaldehyde
265. Picric acid is a yellow coloured compound. Its chemical name is:
- a) *m*-nitrobenzoic acid      b) 2,4,6-trinitrophenol      c) Trinitrotoluene      d) Trinitroaniline
266. The ideal starting material for the synthesis of *m*-chloronitro benzene is:
- a) Benzene      b) Chlorobenzene      c) Toluene      d) Nitrobenzene
267. In a reaction involving ring substitution of  $\text{C}_6\text{H}_5\text{Y}$ , the major product is *meta*-isomer. The group *Y* can be:
- a)  $-\text{NH}_2$       b)  $-\text{COOH}$       c)  $-\text{CH}_3$       d)  $-\text{Cl}$
268. When ammonia is added to green aqueous solution of nickel(II) sulphate, the colour of the solution changes to blue violet. This is caused by:
- a) Nickel undergoing a change in oxidation state  
b) Ammonia molecules replacing water molecules surrounding nickel  
c) Change in coordination number of nickel  
d) Change in pH value of the solution
269. The compound, whose stereo chemical formula is written below, exhibits *x*-geometrical isomers and *y*-optical isomers. The value of *x* and *y* are
- 
- a) 4 and 4      b) 2 and 2      c) 2 and 4      d) 4 and 2
270. Among the following-phenol, benzoic acid, nitrobenzene and toluene, the compound that undergoes nitration readily is:
- a) Benzoic acid      b) Toluene      c) Phenol      d) Nitrobenzene
271. Which one is organometallic compound?
- a) Lithium acetate      b) Lithium methoxide  
c) Lithium dimethyl amide      d) Methyl lithium
272. What are the products formed when an equimolar mixture of benzaldehyde and formaldehyde is heated with concentrated  $\text{NaOH}$ ?
- a)  $\text{C}_6\text{H}_5-\text{CH}_2-\text{OH}$  and  $\text{H}-\text{COONa}$   
b)  $\text{C}_6\text{H}_5-\text{COONa}$  and  $\text{CH}_3-\text{OH}$   
c)  $\text{C}_6\text{H}_5-\text{CH}_2-\text{COONa}$   
d)  $\text{C}_6\text{H}_5-\text{COOH}$  and  $\text{CH}_3-\text{ONa}$
273. Gammexane (a  $\gamma$ -isomer of) is:
- a) BHC  
b) Benzene hexachloride  
c) Lindane  
d) All of these
274. Number of electrons gained by  $\text{Pd}$  in  $[\text{PdCl}_4]^{2-}$ :
- a) 4      b) 8      c) 10      d) 0
275. Which of the following is considered to be an anticancer species?





276. For benzaldehyde which of the following is incorrect?

- a) It is an aromatic aldehyde
- b) It is used in perfumery
- c) On oxidation it yields benzoic acid
- d) On reduction it yields phenol


277. The main source of aromatic compounds is:

- a) Wood
- b) Petroleum
- c) Coal
- d) Both (b) and (c)

278. Phenol on hydrogenation in presence of a nickel catalyst at 160°C gives:

- a) Benzene
- b) Cyclohexane
- c) Cyclohexanol
- d) *n*-hexanol

279. The IUPAC name of compound



- a) Hexane-1, 2, 5-tricarbonitrile
- b) Hexane-1, 3, 6-tricarbonitrile
- c) Butane-1, 2, 4-tricarbonitrile
- d) Butane-1, 3, 4-tircarbonitrile

280. *Fac-mer* isomerism is associated with which one of the following complexes? (*M*=central metal)

- a)  $[M(AA)_2]$
- b)  $[MA_3B_3]$
- c)  $[M(AA)_3]$
- d)  $[MABCD]$

281. Which of the following is the correct order of stability of the following four distinct conformation of *n* butane?

- a) Staggered > Gauche > Partially eclipsed > Fully eclipsed
- b) Gauche > Staggered > partially eclipsed > Fully eclipsed
- c) Staggered > Partially eclipsed > Gauche > Fully eclipsed
- d) Fully eclipsed > Staggered > Partially eclipsed > Gauche

282. *o*-nitrophenol can form hydrogen bonds within the molecule. It thus, has:

- a) Very high m.p.
- b) Very high viscosity
- c) Low m.p.
- d) none of these

283. The element which does not form mononuclear carbonyl is:

- a) Fe
- b) Mn
- c) Ni
- d) W

284. Which of the following is hexadentate ligand?

- a) Ethylene diamine
- b) Ethylene diamine tetra acetic acid
- c) 1,10-phenanthroline
- d) Acetyl acetonato

285. The molecular formula of a saturated compound is  $C_2H_4Cl_2$ . The formula permits the existence of two

- a) Functional isomers
- b) Position isomers
- c) Optical isomers
- d) *cis* – *trans* isomers

286. An octahedral complex is formed when hybrid orbitals of the following type are involved





- a)  $sp^3$
- b)  $dsp^2$
- c)  $d^2sp^3$
- d)  $sp^2d^2$


287. The IUPAC name of the given compound  $CH_3 - CH = CH - COOC_2H_5$  is


- a) Ethyl propenoate
- b) Ethyl-2-butenate
- c) Ethyl-1-butenate
- d) Propene ethyl methanoate


288. Which product is not obtained by heating wood or coal in the absence of air?



- a) Coal-tar
- b) Naphthalene
- c) Benzene
- d) Wax

289. Dry distillation of calcium benzoate with calcium formate gives:  
 a) Acetaldehyde                      b) Benzoic acid                      c) Benzaldehyde                      d) Benzoic anhydride
290. Which will give  $\text{Fe}^{3+}$  ions in solution?  
 a)  $[\text{Fe}(\text{CN})_6]^{3-}$   
 b)  $\text{Fe}_2(\text{SO}_4)_3$   
 c)  $[\text{Fe}(\text{CN})_6]^{4-}$   
 d)  $\text{NH}_4(\text{SO}_4)_2 \cdot \text{FeSO}_4 \cdot 6\text{H}_2\text{O}$
291. Each metal possesses:  
 a) Primary valencies satisfied by anions only  
 b) Secondary valencies satisfied by donor molecules  
 c) Coordination number  
 d) All of the above
292. Aspirin is:  
 a) Antibiotic                      b) Antipyretic                      c) Sedative                      d) Psychedelic
293. Hybridisation, shape and magnetic moment of  $\text{K}_3[\text{Co}(\text{CO}_3)_3]$  is  
 a)  $d^2 sp^3$ , octahedral, 4.9 BM                      b)  $sp^3 d^2$ , octahedral, 4.9 BM  
 c)  $dsp^2$ , square planer, 4.9 BM                      d)  $sp^3$ , tetrahedral, 4.9 BM
294. Among the following complexes ( $K-P$ ),  
 $\text{K}_3[\text{Fe}(\text{CN})_6](K)$ ,  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3(L)$ ,  
 $\text{Na}_3[\text{Co}(\text{ox})_3](M)$   
 $[\text{Ni}(\text{H}_2\text{O})_6]\text{Cl}_2(N)$ , and  
 $[\text{Zn}(\text{H}_2\text{O})_6](\text{NO}_3)_2(P)$  the diamagnetic complexes  
 a)  $K, L, M, N$                       b)  $K, M, O, P$                       c)  $L, M, O, P$                       d)  $L, M, N, O$
295. Aniline when diazotised in cold and then treated with dimethyl aniline gives a coloured product. Its structure would be:  
 a)   
 b)   
 c)   
 d) 
296. Pyridine possesses:  
 a) Aromatic nature  
 b) Unsaturated aliphatic nature  
 c) Alicyclic nature  
 d) Aliphatic nature
297. A reagent used for identifying nickel ion is:  
 a) Potassium ferrocyanide  
 b) Phenolphthalein  
 c) Dimethyl glyoxime  
 d) EDTA
298. Aniline was diazotised and subsequently reduced with stannous chloride and hydrochloric acid to yield:  
 a) Phenyl aniline                      b) Phenyl hydrazine                      c)  $p$ -amino azobenzene                      d) Diazoamino benzene
299. The reaction of toluene with  $\text{Cl}_2$  in presence of  $\text{FeCl}_3$  gives predominantly:  
 a)  $m$ -chlorobenzene  
 b) Benzoylchloride

- c) Benzyl chloride  
d) *o*- and *p*-chlorobenzene
300. Which statement is not correct in the case of  $[\text{Co}(\text{NH}_3)_6]^{3+}$  complex?  
a) It is octahedral in shape  
b) It involves  $d^2sp^2$ -hybridization  
c) It has diamagnetic nature  
d) None of the above
301. Pick out the complex compound in which the central metal atom obeys EAN rule strictly  
a)  $\text{K}_4[\text{Fe}(\text{CN})_6]$       b)  $\text{K}_3[\text{Fe}(\text{CN})_6]$       c)  $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3$       d)  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$
302. Amongst the following, the compound that can be most readily sulphonated is:  
a) Benzene      b) Methoxy benzene      c) Toluene      d) Chlorobenzene
303. *p*-chloroaniline and anilium hydrochloride can be distinguished by:  
a)  $\text{P}_2\text{O}_5$       b)  $\text{AgNO}_3$       c) Carbylamine test      d) Sandmeyer's reaction
304. Pyrogallol is..... trihydroxy benzene.  
a) 1, 2, 4      b) 1, 2, 3      c) 1, 3, 5      d) None of these
305. Phenol is weakly acidic but does not react with  $\text{NaHCO}_3$  like carboxylic acids hence:  
a) Phenol is weaker than carbonic acid  
b) Phenol is stronger than acid  
c) Phenol is stronger than carboxylic acid  
d) None of the above
306. *p*-cresol reacts with chloroform in alkaline medium to give compound (A) which adds hydrogen cyanide to form compound (B). The latter on acidic hydrolysis gives chiral carboxylic acid. The acid is:
- a) 

b) 

c) 





d) 
307. The number of isomeric xylenes is:  
a) 2      b) 3      c) 4      d) 1
308. The IUPAC name of  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl}$  is:  
a) Tetrahydrodichlorochromium(III) chloride  
b) Tetraaquodichlorochromium(III) chloride  
c) Tetraaquodichlorochromium(I) chloride  
d) None of the above
309. Among the following metal carbonyls, C—O bond order is lowest in  
a)  $[\text{Mn}(\text{CO})_6]^+$       b)  $[\text{Fe}(\text{CO})_5]$       c)  $[\text{Cr}(\text{CO})_6]$       d)  $[\text{V}(\text{CO})_6]^-$
310.  is  
a) 3-propyl-1,3-pentadiene      b) 3,3-dipropyl-1,3-pentadiene  
c) 3,3-diethenyl penta-1,4-diene      d) 4,4-diethenyl penta,1,2-diene
311. Which of the following shell, form an outer octahedral complex?  
a)  $d^4$       b)  $d^8$       c)  $d^6$       d) None of these
312. Friedel-Craft's reaction of bromobenzene with methyl iodide gives:  
a) *o*-bromotoluene  
b) *p*-bromotoluene  
c) *o*- and *p*-bromotoluene

d) *m*-bromotoluene

313. An organic compound  $C_7H_8O$  is neither soluble in NaOH nor gives blue colour with  $FeCl_3$ , is:

- a)  $C_6H_5 \cdot CH_2OH$       b)  $C_6H_4 \begin{matrix} \nearrow CH_3 \\ \searrow OH \end{matrix}$       c)  $C_6H_5 \cdot OCH_3$       d) None of these

314. Which exist as a pair of mirror image isomers?

- a)       b)       c)       d) 


315. Benzene double bonds are not so reactive as those of hexatriene because:

- a) The three double bonds are caged in a ring  
b) Benzene is aromatic and has six  $\pi$ -resonating electrons  
c) Benzene has no double bond  
d) Benzene is non-polar

316. The most stable ion is

- a)  $[Fe(OH)_5]^{3-}$       b)  $[FeCl_6]^{3-}$       c)  $[Fe(CN)_6]^{3-}$       d)  $[Fe(H_2O)_6]^{3+}$

317. Which of the following is/are threo isomers?




- a) Only (i)      b) Only (ii)  
c) Only (iii)      d) All (i), (ii) and (iii)

318. In the coal-tar distillation of middle oil, the aromatic compounds present are:

- a) Benzene, naphthalene, anthracene  
b) Naphthalene, pyridine, phenol  
c) Naphthalene, pyridine  
d) None of the above

319. The correct order of increasing reactivity of  $C-X$  bond towards nucleophilic in the following compound is:



- a)  $I < II < IV < III$       b)  $II < III < I < IV$       c)  $IV < III < I < II$       d)  $III < II < I < IV$


320. Which of the following system is most stable for a chelate?

- a) Two fused cyclic system      b) Three fused cyclic system  
c) Four fused cyclic system      d) Five fused cyclic system

321. Which of the following reaction take place when a mixture of concentrated  $HNO_3$  and  $H_2SO_4$  reacts on benzene at 300 K?





- a) Sulphonation      b) Nitration      c) Hydrogenation      d) Dehydration



322. Consider the following reaction:








- a) Benzene      b) Toluene      c) Benzaldehyde      d) Benzoic acid

323. The shortest  $C-O$  bond order exists in:

- a)  $[\text{Mn}(\text{CO})_6]^+$       b)  $[\text{Fe}(\text{CO})_5]$       c)  $[\text{Cr}(\text{CO})_6]$       d)  $[\text{V}(\text{CO})_6]^-$
324. Between *p*-nitrophenol and salicylaldehyde, solubility in base is:  
 a) Almost nil in both cases  
 b) Higher in *p*-nitrophenol  
 c) Higher for salicylaldehyde  
 d) Equal in nature
325. (+) and (-) forms of optically active compounds are different in  
 a) Boiling points      b) Melting points      c) Specific gravity      d) Specific rotation
326. Benzene on treatment with dry HCN and HCl in presence of anhy.  $\text{AlCl}_3$  followed by hydrolysis forms:  
 a) Chlorobenzene      b) Benzoic acid      c) Benzaldehyde      d) Cyanobenzene
327. In which of the following compounds does the central atom obey EAN rule?  
 a)  $\text{K}_3\text{Fe}(\text{CN})_6$       b)  $\text{K}_4\text{Fe}(\text{CN})_6$       c)  $\text{Cu}(\text{NH}_3)_4\text{SO}_4$       d) All of these
328. Pick the correct name of  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$   
 a) Chloropentammine cobalt (III) chloride      b) Chloropentammine cobalt (III)  
 c) Chloropentammine cobalt (II) chloride      d) Pentammine chloro cobalt(III) chloride
329. The geometry of  $\text{Ni}(\text{CO})_4$  and  $\text{Ni}(\text{PPh}_3)_2\text{Cl}_2$  are  
 a) Square planar and tetrahedral respectively      b) Both tetrahedral  
 c) Tetrahedral and square planar respectively      d) Both square planar
330. Select pair of chain isomers from the following
- (I)       (II) 
- (III)       (IV) 
- a) I and II      b) II and III      c) I and IV      d) II and III
331. Which ligand produces a high crystal field splitting (a strong ligand field)?  
 a) CO      b)  $\text{NO}_2^-$       c)  $\text{CN}^-$       d) All are correct
332. Benzene reacts with *n*-propyl chloride in the presence of anhydrous  $\text{AlCl}_3$  to give predominantly:  
 a) Isopropyl benzene  
 b) No reaction  
 c) *n*-propylbenzene  
 d) 3-propyl-1-chlorobenzene
333. Which of the following coordination compounds would exhibit optical isomerism?  
 a) Pentaamminenitrocobalt (III) iodide      b) Diamminedinitroplatinum (II)  
 c) *trans*-dicyanobis (ethylenediamine)      d) Tris-(ethylenediamine) cobalt(III) bromide
334. What is the magnetic moment of  $\text{K}_3[\text{FeF}_6]$ ?  
 a) 3.87 BM      b) 4.89 BM      c) 5.91 BM      d) 6.92 BM
335. The EAN of Cr in  $[\text{Cr}(\text{SCN})_6]^{3-}$  is:  
 a) 35      b) 33      c) 34      d) 37
336. Which has maximum paramagnetic character?  
 a)  $[\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$       c)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$       d)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$
337. Phenol, when it first reacts with concentrated sulphuric acid and then with concentrated nitric acid, gives:  
 a) Nitrobenzene  
 b) 2, 4, 6-trinitrobenzene  
 c) *o*-nitrophenol  
 d) *p*-nitrophenol
338. Activation of benzene ring by  $-\text{NH}_2$  in aniline can be reduced by treating with:  
 a) Dil. HCl      b) Ethyl alcohol      c) Acetic acid      d) Acetyl chloride
339. Sulphonation of benzoic acid produces mainly:  
 a) *o*-sulphobenzoic acid

- b) *m*-sulphobenzoic acid  
 c) *p*-sulphobenzoic acid  
 d) *o-p*-disulphobenzoic acid
340. The IUPAC name for the complex  $[\text{Co}(\text{NO}_2)(\text{NH}_3)_5]\text{Cl}_2$  is  
 a) Nitrito -N- pentamminecobalt (III) chloride      b) Nitrito -N- pentamminecobalt (II) chloride  
 c) Pentammine nitrito-N- cobalt (II) chloride      d) Pentaammine nitrito-N- cobalt (III) chloride
341. The ionisation isomer of  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}(\text{NO}_2)\text{Cl}]$  is  
 a)  $[\text{Cr}(\text{H}_2\text{O})_4(\text{O}_2\text{N})]\text{Cl}_2$       b)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2](\text{NO}_2)$   
 c)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}(\text{ONO})]\text{Cl}$       d)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2(\text{NO}_2)] \cdot \text{H}_2\text{O}$
342. Salicylic acid, aspirin, nylon, plastics and picric acid have a common raw material, namely:  
 a) Methane      b) Formic acid      c) Phenol      d) Alcohol
343. Ulmann's reaction is used for the preparation of:  
 a) Diphenyl      b) Iodobenzene      c) Toluene      d) Naphthalene
344. Which of the following statements is/are incorrect for *D* – (+) –glyceradehyde?  
 a) The symbol *D* not indicates the dextrorotatory nature of the compound  
 b) The sign (+) indicates the dextrorotatory nature of the compound  
 c) The symbol *D* indicates that hydrogen atom lies left to the chiral centre in the Fischer projection diagram  
 d) The symbol *D* indicates that hydrogen atom lies right to the chiral centre in the Fischer projection diagram
345. Complexes with  $\text{CN}^-$  ligands are usually:  
 a) High spin complexes      b) Low spin complexes      c) Both (a) and (b)      d) None of these
346. The IUPAC of  is  
 a) 2-cyclopentyl propane      b) 1, 1-dimethyl-1-cyclopentyl methane  
 c) 1-(1-methyl) ethyl cyclopentane      d) None of the above
347. Which ion is paramagnetic?  
 a)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$       b)  $[\text{Fe}(\text{CN})_6]^{4-}$       c)  $[\text{Ni}(\text{CO})_4]$       d)  $[\text{Ni}(\text{CN})_4]^{2-}$
348. Configuration of  are  
 a) *R, R*      b) *R, S*      c) *S, S*      d) *S, R*
349. Dow process is used for the conversion of chlorobenzene to:  
 a) Benzene      b) Nitrobenzene      c) Phenol      d) Gammexane
350. Phenolphthalein is produced on heating phthalic anhydride and conc. sulphuric acid with:  
 a) Salicylic acid      b) Phenol      c) Phenacetin      d) Phenanthrene
351. Benzene is converted to toluene by:  
 a) Friedel-Crafts reaction  
 b) Grignard reaction  
 c) Wurtz reaction  
 d) Perkin's reaction
352. The number of ions formed when hexamine copper (II) sulphate is dissolved in water is?  
 a) 1      b) 2      c) 4      d) 6
353. In a set of reactions *m*-bromobenzoic acid gave a product *D*, Identify the product *D*:





354. In  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$ , the isomerism shown is:

- a) Ligand                                      b) Optical                                      c) Geometrical                                      d) Ionization

355. The hybridization of Fe in  $\text{K}_4[\text{Fe}(\text{CN})_6]$  complex is:

- a)  $d^2sp^2$                                       b)  $d^2sp^3$                                       c)  $dsp^2$                                       d)  $sp^3$

356. The correct name of  $\text{CH}_3-\text{CH}_2-\text{C}\equiv\text{C}-\text{CH}=\text{CH}_2$  is

- a) Hex-3-yn-5-ene                                      b) Hex-5-en-3-yne                                      c) Hex-3-yn-1-ene                                      d) Hex-1-en-3-yne

357. Nickel metal is in highest oxidation state in:

- a)  $\text{Ni}(\text{CO})_4$                                       b)  $\text{K}_2\text{NiF}_6$                                       c)  $[\text{Ni}(\text{NH}_3)_6](\text{BF}_4)_2$                                       d)  $\text{K}_4[\text{Ni}(\text{CN})_6]$

358. Which of the following complexes show six coordination number?

- a)  $[\text{Zn}(\text{CN})_4]^{2-}$                                       b)  $[\text{Ni}(\text{NH}_3)_4]^{2+}$                                       c)  $[\text{Cu}(\text{CN})_4]^{2-}$                                       d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$

359. Which of the following statements is wrong?

- a) The IUPAC name of alkenes ends with suffix-ene  
b) The IUPAC name of alkynes ends with suffix-yne  
c) The IUPAC name of acid amide is alkanamide  
d) The substituents get lower number in comparison to principal functional group

360. The possible number of isomers for the complex  $[\text{MCl}_2\text{Br}_2]\text{SO}_4$  is:

- a) 1                                      b) 2                                      c) 4                                      d) 5

361.  $\text{K}_3[(\text{Al})(\text{C}_2\text{O}_4)_3]$  is called

- a) Potassium aluminium (III) oxalate                                      b) Potassium alumino oxalate  
c) Potassium trioxalato aluminate (VI)                                      d) Potassium trioxalato aluminate (III)





362. In  $\text{Fe}(\text{CO})_5$ , the Fe — C bond possesses

- a)  $\pi$  —Character only                                      b) Both  $\sigma$  and  $\pi$  —characters  
c) Ionic characters                                      d)  $\sigma$  —Character only

363. The reaction,  $[\text{Fe}(\text{CNS})_6]^{3-} \rightarrow [\text{FeF}_6]^{3-}$  taken place with

- a) Decrease in magnetic moment                                      b) Increase in magnetic moment  
c) Decrease in coordination number                                      d) Increase in coordination number

364. Which chloro derivative of benzene among the following would undergo hydrolysis most readily with aqueous NaOH to furnish the corresponding hydroxyl derivative?



365. Some salts although containing two different metallic elements give test for only one of them in solution. Such salts are:

- a) Complex salts                                      b) Double salts                                      c) Normal salts                                      d) None of these

366. Mixture X = 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  and 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  was prepared in 2 litre of



solution.

1 litre of mixture  $X + \text{excess AgNO}_3 \rightarrow Y$ .

1 litre of mixture  $X + \text{excess BaCl}_2 \rightarrow Z$ .

No. of moles of  $Y$  and  $Z$  are.

- a) 0.01, 0.01                      b) 0.02, 0.01                      c) 0.01, 0.02                      d) 0.02, 0.02

367. The hybridization of central metal ion and shape of Wilkinson's catalyst is

- a)  $sp^3d$ , trigonal bipyramidal                      b)  $sp^3$ , tetrahedral  
c)  $dsp^2$ , square planar                      d)  $d^2sp^2$ , octahedral

368. The  $d$ -electron configurations of  $\text{Cr}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$  and  $\text{Co}^{2+}$  are  $d^4$ ,  $d^5$ ,  $d^6$  and  $d^7$  respectively. Which one of the following will exhibit minimum paramagnetic behaviour?

- a)  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$   
b)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$   
c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$   
d)  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$   
(At. Nos. Cr = 24, Mn = 25, Fe = 26, Co = 27)

369. An enantiomerically pure acid is treated with racemic mixture of an alcohol having one chiral carbon. The ester formed will be

- a) Optically active mixture                      b) Pure enantiomer  
c) *meso* compound                      d) Racemic mixture

370. Which of the following ring is most strained?

- a) Cyclohexane                      b) Cyclopentane                      c) Cyclobutane                      d) Cyclopropane

371. Formylchloride has not been prepared so far. Which can function as formylchloride in formylation?

- a)  $\text{HCHO} + \text{HCl}$                       b)  $\text{HCOOCH}_3 + \text{HCl}$                       c)  $\text{CO} + \text{HCl}$                       d)  $\text{HCONH}_2 + \text{HCl}$


372. In hexacyanomanganate (II) ion the Mn-atom assumes  $d^2sp^3$ -hybrid state. The number of unpaired electrons in the complex is:

- a) 1                      b) 2                      c) 3                      d) 0


373. Which one of the following does not give a white precipitate with silver nitrate solution?

- a)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$                       b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$                       c)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$                       d)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$

374. In a set of reactions, ethyl benzene yielded a product  $D$ .



$D$  would be:



375. The oxidation number of Pt in  $[\text{Pt}(\text{C}_2\text{H}_4)\text{Cl}_3]$  is

- a) +1                      b) +2                      c) +3                      d) +4


376. Among  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$ ,  $[\text{Fe}(\text{Cl})_6]^{3-}$  species, the hybridization state of the Fe atom are, respectively


- a)  $d^2sp^3$ ,  $d^2sp^3$ ,  $sp^3d^2$                       b)  $sp^3d^2$ ,  $d^2sp^3$ ,  $d^2sp^3$                       c)  $sp^3d^2$ ,  $d^2sp^3$ ,  $sp^3d^2$                       d) None of these

377. Of the following complex ions, which is diamagnetic in nature?


- a)  $[\text{CoF}_6]^{3-}$                       b)  $[\text{NiCl}_4]^{2-}$                       c)  $[\text{Ni}(\text{CN})_4]^{2-}$                       d)  $[\text{CuCl}_4]^{2-}$

378. The IUPAC name of compound



- a) 2-methoxycarbonylbenzoic acid  
c) 2-carboxy phenyl ethanoate
- b) Methyl-2-carboxy benzoate  
d) *o*-carboxyphenyl acetate
379. Which of the following are produced from coal-tar?  
a) Synthetic dyes      b) Drugs      c) Perfumes      d) All of these
380. Chlorine is least reactive in:  
a)  $\text{CH}_3\text{Cl}$       b)  $\text{CH}_2=\text{CHCl}$       c)  $\text{C}_6\text{H}_5\text{Cl}$       d)  $\text{C}_2\text{H}_5\text{Cl}$
381. Correct IUPAC name of compound  $(\text{CH}_3)_2\text{C}(\text{CH}_2\text{CH}_3)\text{CH}_2\text{CH}(\text{Cl})\text{CH}_3$  is  
a) 5-chloro-3,3-dimethylhexane      b) 3-chloro-2-ethyl-2-methylpentane  
c) 2-chloro-4-ethyl-4-methylpentane      d) None of the above
382. 
- In the above reaction 'X' stands for:  
a)  $\text{NH}_2$       b)  $\text{Cl}$       c)  $\text{SnCl}_2$       d)  $\text{NH}_3^+\text{Cl}^-$
383. Which follows EAN rule?  
a)  $\text{Fe}(\text{CO})_5$       b)  $\text{Ni}(\text{CO})_4$       c)  $\text{K}_4[\text{Fe}(\text{CN})_6]$       d) All are correct
384. Which one is bidentate ligand?  
a)  $\text{C}_2\text{O}_4^{2-}$       b)  $\text{NH}_2 \cdot \text{CH}_2 \cdot \text{CH}_2 \cdot \text{NH}_2$       c) Both (a) and (b)      d) None of these
385. The reagent used for conversion of benzene diazonium chloride to benzene is:  
a)  $\text{H}_3\text{PO}_2 + \text{H}_2\text{O}$       b)  $\text{Na}_2\text{SnO}_2 + \text{NaOH}$       c)  $\text{C}_2\text{H}_5\text{OH}$       d) All of these
386. Which will not give the usual test for iron?  
a)  $\text{K}_2\text{Fe}_2(\text{SO}_4)_4 \cdot 24\text{H}_2\text{O}$   
b)  $(\text{NH}_4)_2\text{Fe}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$   
c)  $\text{K}_3[\text{Fe}(\text{CN})_6]$   
d)  $\text{Fe}_2(\text{SO}_4)_3$
387.  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  and  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  are a pair of ..... isomers.  
a) Ionisation      b) Ligand      c) Coordination      d) Hydrate
388. The first organic compound prepared in the laboratory was  
a) Acetic acid      b) Acetylene      c) Urea      d) Methane
389. Aniline on heating with conc.  $\text{H}_2\text{SO}_4$  at 460 K gives:  
a) Aniline sulphate  
b) Benzene sulphonic acid  
c) Sulphanilic acid  
d) None of the above
390. Which of the following statements regarding phenols is not correct?  
a) Phenols are stronger acid than water and alcohols  
b) Phenols are weaker acids than carboxylic acids  
c) Phenols are soluble in both aqueous  $\text{NaOH}$  and aqueous  $\text{NaHCO}_3$   
d) Phenoxide ions are more stable than the corresponding phenols
391. Which would decolourise cold, *aq.* potassium permanganate solution?  
a) Benzoic acid      b) Cinnamic acid      c) *p*-toluic acid      d) *m*-toluic acid
392. The magnetic moment of  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is found to be 1.7 BM. How many unpaired electron (s) is/are present per molecule?  
a) 1      b) 2      c) 3      d) 4


393. The IUPAC name of the compound



- a) Bicyclo [2,5,0] nonane  
 c) Bicyclo [5,2,0] nonane

- b) Bicyclo [5,0,2] nonane  
 d) Bicyclo [0,2,5] nonane


394. The IUPAC name of the compound



- a) 2-oxocyclohexane-1-carboxylic acid  
 c) 6-oxocyclohexane-1-carboxylic acid

- b) Cyclohexane-2-oxo-1-carboxylic acid  
 d) None of the above

395. The IUPAC name of



- a) Spiro [3.2.1] octane    b) Bicyclo [3.2.2] octane    c) Bicyclo [3.2.1] octane    d) None of these


396. Which of the following deactivates benzene substitution?

- a) —NHR                      b) —OH                      c) —OR                      d) —COOR

397. Aniline, chloroform and alc. KOH on heating give:

- a) Phenyl isocyanide    b) Phenyl cyanide    c) Chlorobenzene    d) Phenol

398. In the chemical reactions,



- a) Nitrobenzene and chlorobenzene  
 b) Nitrobenzene and fluorobenzene  
 c) Phenol and benzene  
 d) Benzenediazonium chloride and fluorobenzene

399. The incorrect statement for IUPAC system of nomenclature is

- a) In an organic compound, the longest carbon chain is always selected for assigning the root word  
 b) There is no compound with the name 3-ethyl pentane  
 c) Out of —NH<sub>2</sub> and —OH groups present in an organic compound, —NH<sub>2</sub> is treated as substituent  
 d) Different alkyl groups are written alphabetically while, writing the IUPAC name

400. When sodium benzene sulphonate is fused with sodium hydroxide (solid), followed by hydrolysis the product formed is:

- a) Benzene                      b) Sod. phenoxide                      c) Benzene thiophenol                      d) Phenol

401. The correct order of stability of conformations of cyclohexane is

- a) Chair > twist boat > boat                      b) Twist boat > chair > boat  
 c) Boat > chair > twist boat                      d) Boat > twist boat > chair

402. Phenol with dilute HNO<sub>3</sub> gives:

- a) *meta* and *para* nitrophenol  
 b) *ortho* and *para* nitrophenol  
 c) Trinitrophenol  
 d) *ortho* and *meta* nitrophenol

403. The increasing order of boiling points of compounds given below is:

- (I) 1,2-dihydroxy benzene  
 (II) 1,3-dihydroxy benzene  
 (III) 1,4-dihydroxy benzene  
 (IV) Hydroxyl benzene

a)  $I < II < III < IV$       b)  $I < II < IV < III$       c)  $IV < I < II < III$       d)  $IV < II < I < III$

404. The pair of the compounds in which both the metals are in the higher possible oxidation state is

- a)  $\text{CrO}_2\text{Cl}_2, \text{MnO}_4^-$       b)  $[\text{Co}(\text{CN})_6]^{3-}, \text{MnO}_3$   
 c)  $\text{TiO}_3, \text{MnO}_2$       d)  $[\text{Fe}(\text{CN})_6]^{3-}, [\text{Co}(\text{CN})_6]^{3-}$

405. The number of ions given by  $\text{K}_2[\text{PtCl}_6]$  in aqueous solution is:

- a) 2      b) 3      c) 4      d) Zero

406. Which of the following are functional isomers?

- a)  $\text{CH}_3\text{CH}_2\text{Cl}$  and  $\text{CH}_3\text{CH}_2\text{Br}$       b)  $\text{CH}_3\text{CHBr}_2$  and  $\text{CH}_2\text{Br}_2 \cdot \text{CH}_2\text{Br}$   
 c)  $\text{C}_2\text{H}_5\text{OC}_2\text{H}_5$  and  $\text{CH}_3\text{OC}_3\text{H}_7$       d)  $\text{CH}_3\text{CH}_2\text{CHO}$  and  $\text{CH}_3-\text{CH}=\text{CH}_2$


407. Phenol is:

- a) Strongly acidic      b) Weakly acidic      c) Strongly basic      d) Weakly basic

408. The correct IUPAC name of  $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$  is:

- a) Aluminium potassium sulphate-12-water  
 b) Potassium aluminium(III) sulphate-12-water  
 c) Potassium aluminate(III) sulphatehydrate  
 d) Aluminium(III) potassium sulphate hydrate-12

409. A complex shown below can exhibit:






- a) Optical isomerism only  
 b) Geometrical isomerism only  
 c) Both optical and geometrical isomerism  
 d) None of the above

410. The IUPAC name of the complex  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$  is

- a) Dichloro tetraammine cobalt (III) chloride      b) Tetraammine dichloro cobalt(III) chloride  
 c) Tetraammine dichloro cobalt (II) chloride      d) Tetraammine dichloro cobalt (IV) chloride

411. The correct decreasing order of their reactivity towards hydrolysis is:

(i)  $\text{C}_6\text{H}_5\text{COCl}$



- a) (i) > (ii) > (iii) > (iv)      b) (iv) > (ii) > (i) > (iii)      c) (ii) > (iv) > (i) > (iii)      d) (ii) > (iv) > (iii) > (i)


412. Nitrobenzene is generally used for:

- a) Preparing shoe polish      b) Preparing floor polish      c) Preparing aniline      d) All of these

413. In the coordination compound,  $\text{K}_4[\text{Ni}(\text{CN})_4]$ , the oxidation state of nickel is

- a) -1      b) 0      c) +1      d) +2

414. Salicylic acid as compared to benzoic acid:  
 a) Is more acidic                      b) Has same acidity                      c) Has less acidity                      d) None of these
415. Which ligand is expected to be bidentate?  
 a)  $\text{C}_2\text{O}_4^{2-}$                       b)  $\text{CH}_3\text{C} \equiv \text{N}$                       c)  $\text{Br}^-$                       d)  $\text{CH}_3\text{NH}_2$
416. Which one of the following is most reactive towards aqueous NaOH?  
 a)  $\text{C}_6\text{H}_5\text{Cl}$                       b)  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$                       c)  $\text{C}_6\text{H}_5\text{Br}$                       d)  $\text{BrC}_6\text{H}_4\text{Br}$
417. Which is not an aromatic compound?  
 a) Pyridine                      b) Naphthalene                      c) Xylene                      d) Cyclohexane
418. Which one of the following is wrongly matched?  
 a)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$  – Square planar                      b)  $[\text{Ni}(\text{CO})_4]$  – Neutral ligand  
 c)  $[\text{Fe}(\text{CN})_6]^{3-}$  –  $sp^3 d^2$                       d)  $[\text{Co}(\text{en})_3]^{3+}$  – Follows EAN rule
419. Stereoisomers have different  
 a) Molecular formula                      b) Structural formula  
 c) Configuration                      d) Molecular mass
420. Which of the following will show optical isomerism?  
 a)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$   
 b)  $[\text{ZnCl}_4]^{2-}$   
 c)  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$   
 d)  $[\text{Co}(\text{CN})_6]^{3-}$
421. A complex of cobalt has five ammonia molecules, one nitro group and two chlorine atoms for each cobalt atom. One mole of this compound produces three mole ions in aqueous solution which on treating with excess of  $\text{AgNO}_3$  give two mole of  $\text{AgCl}$ . The formula of the compound is:  
 a)  $[\text{Co}(\text{NH}_3)_4\text{NO}_2\text{Cl}][(\text{NH}_3)_2\text{Cl}]$                       b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}][\text{ClNO}_2]$                       c)  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$                       d)  $[\text{Co}(\text{NH}_3)_5][(\text{NO}_2)_2\text{Cl}_2]$
422. Which one group is trivalent in nature?  
 a) Benzo                      b) Benzal                      c) Benzyl                      d) All of these
423. Benzene contains double bonds but does not give addition reactions because:  
 a) Double bonds in benzene are strong  
 b) Double bonds change their position rapidly  
 c) Resonance lowers the energy of benzene molecule and leads to greater stabilization  
 d) None of the above
424. Low spin complex of  $d^6$ -cation in an octahedral field will have the following energy:  
 a)  $\frac{-12}{5}\Delta_0 + P$   
 b)  $\frac{-12}{5}\Delta_0 + 3P$   
 c)  $\frac{-2}{5}\Delta_0 + 2P$   
 d)  $\frac{-2}{5}\Delta_0 + P$   
 ( $\Delta_0$  = Crystal field splitting energy in an octahedral field,  $P$  = Electron pairing energy)
425.  $\text{C}_7\text{H}_8\text{O}$  show how many isomers?  
 a) 2                      b) 3                      c) 4                      d) 5
- 426.
- The above structural formula refers to:  
 a) BHC                      b) DNA                      c) DDT                      d) RNA
427. The compound



Have its IUPAC name as

- a) Octa dec-9-enoic acid  
b) Oleic acid  
c) Ethyl hexadic-9-enoic acid  
d) All of these
428. The type of isomerism present in nitropentaammine-chromium (III) chloride is :  
a) Optical  
b) Linkage  
c) Ionization  
d) polymerization
429. Which complex compound possesses  $sp^3d^2$  hybridisation?  
a)  $[\text{Fe}(\text{NH}_3)_6]^{3+}$   
b)  $[\text{Fe}(\text{CN})_6]^{4-}$   
c)  $[\text{Fe}(\text{CN})_6]^{3-}$   
d)  $[\text{Fe}(\text{Cl})_6]^{3-}$
430. Amongst the following carboxylic acids the strongest acid is:  
a) Benzoic acid  
b) *o*-methoxybenzoic acid  
c) *m*-nitrobenzoic acid  
d) *p*-nitrobenzoic acid
431. When EDTA solution is added to  $\text{Mg}^{2+}$  ion solution, then which of the following statements is not true?  
a) Four coordinate sites of  $\text{Mg}^{2+}$  are occupied by EDTA and remaining two sites are occupied by water molecules.  
b) All six coordinate sites of  $\text{Mg}^{2+}$  are occupied.  
c)  $\text{pH}$  of the solution is decreased.  
d) Colourless  $[\text{Mg} - \text{EDTA}]^{2-}$  chelate is formed.
432. The energy difference between chair and the boat conformation of cyclohexane is  
a) 29.7 kJ  
b) 44 kJ  
c) 151 kJ  
d) 36 kJ
433. Compounds having the same molecular formula but different properties are called  
a) Isotopes  
b) Isobars  
c) Isomers  
d) Isomorphs
434.  $\text{CH}_3\text{OCH}_2\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$  is  
a) Ethylmethylpropyl diether  
b) Ethylmethoxypropyl ether  
c) 3-ethoxy-1-methoxy propane  
d) 1-ethoxy-3-methoxy propane
435. The benzene molecule contains:  
a) Six  $sp^2$ -hybridized carbons  
b) Three  $sp^2$ -hybridized carbons  
c) Six  $sp^3$ -hybridized carbons  
d) Three  $sp^3$ -hybridized carbons
436. The correct order of stability of conformations of  $\text{NH}_2 - \text{CH}_2 - \text{CH}_2 - \text{OH}$  is  
a) Gauche > eclipsed > anti  
b) Gauche > anti > eclipsed  
c) Eclipsed > gauche > anti  
d) Anti > eclipsed > gauche
437. The solubility of  $\text{AgCN}$  increases by the addition of  $\text{KCN}$  because of:  
a) Complex formation  
b) Redox change  
c) Salt formation  
d) None of these
438. Alicyclic compounds are  
a) Aromatic cyclic compounds  
b) Aliphatic cyclic compounds  
c) Both (a) and (b)  
d) None of the above
439. Which of the following compounds reacts slower than benzene in electrophilic bromination?  
a)  $\text{C}_6\text{H}_5 - \text{NO}_2$   
b)  $\text{C}_6\text{H}_5 - \text{NH}_2$   
c)  $\text{C}_6\text{H}_5 - \text{OH}$   
d)  $\text{C}_6\text{H}_6 - \text{CH}_3$
440. The fraction of chlorine precipitated by  $\text{AgNO}_3$  solution from  $[\text{Cu}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$  is:  
a)  $1/2$   
b)  $2/3$   
c)  $1/3$   
d)  $1/4$
441. Number of possible optical isomers in  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$  is  
a) 2  
b) 3  
c) 4  
d) 6
442. Dimethyl glyoxime gives a red precipitate with  $\text{Ni}^{2+}$  which is used for its detection. To get this precipitate readily, the best pH range is






a) &lt; 1

b) 3 – 4

c) 9 – 11

d) 2 – 3

443. Predict the product:



444. Replacement of Cl of chlorobenzene to give phenol requires drastic conditions but chlorine of 2, 4-dinitrochlorobenzene is readily replaced because:

- a)  $\text{NO}_2$  makes the electron rich ring at *ortho* and *para* positions  
 b)  $\text{NO}_2$  withdraws electrons at *meta* position  
 c)  $\text{NO}_2$  donate electrons at *meta*-position  
 d)  $\text{NO}_2$  withdraws electrons at *ortho* and *para* positions

445. Salicylic acid on heating with soda lime forms:

- a) Phenol                      b) Benzyl alcohol                      c) Benzene                      d) Benzoic acid


446. Which of the following is an organometallic compound?

- a)  $\text{Ti}(\text{C}_2\text{H}_5)_4$                       b)  $\text{Ti}(\text{OC}_2\text{H}_5)_4$                       c)  $\text{Ti}(\text{OCOCH}_3)_4$                       d)  $\text{Ti}(\text{OC}_6\text{H}_5)_4$


447. Which kind of isomerism is exhibited by octahedral  $\text{Co}(\text{NH}_3)_4\text{Br}_2\text{Cl}$ ?

- a) Geometrical and ionisation                      b) Geometrical and optical  
 c) Optical and ionisation                      d) Geometrical only

448. Which of the following is the strongest base?



449. Which of the following will be aromatic?







450. The correct symbol relating the two Kekule structure of benzene is:  
 a)  $\rightarrow$                       b)  $\rightleftharpoons$                       c)  $\leftrightarrow$                       d)  $\rightleftarrows$
451. Benzaldehyde can be obtained by the hydrolysis of:  
 a) Benzyl chloride              b) Benzal chloride              c) Benzonitrile              d) Benzoic acid
452. Which of the following has an optical isomer?  
 a)  $[\text{Co}(\text{en})(\text{NH}_3)_2]^{2+}$       b)  $[\text{Co}(\text{H}_2\text{O})_4(\text{en})]^{3+}$       c)  $[\text{Co}(\text{en})_2(\text{NH}_3)_2]^{3+}$       d)  $[\text{Co}(\text{NH}_3)_3\text{Cl}]^+$
453. Chromium carbonyl is:  
 a)  $\text{Cr}(\text{CO})_4$                       b)  $\text{Cr}(\text{CO})_5$                       c)  $\text{Cr}(\text{CO})_6$                       d) None of these
454. Which of the following reagents may be used to distinguish between phenol and benzoic acid?  
 a) Aqueous NaOH              b) Tollen's reagent              c) Molisch reagent              d) Neutral  $\text{FeCl}_3$
455. Which of the following complex species do not involve  $d^2sp^3$ -hybridization?  
 a)  $[\text{CoF}_6]^{3-}$                       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                       c)  $[\text{Fe}(\text{CN})_6]^{3-}$                       d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
456. Which one of the following shows maximum value of paramagnetic behaviour?  
 a)  $[\text{Sc}(\text{CN})_6]^{3-}$                       b)  $[\text{Co}(\text{CN})_6]^{3-}$                       c)  $[\text{Ni}(\text{CN})_4]^{2-}$                       d)  $[\text{Cr}(\text{CN})_6]^{3-}$
457. The IUPAC name of  

$$\begin{array}{c} \text{HOOC} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH} - \text{CH}_2 - \text{COOH} \\ | \\ \text{CH}_2\text{COOH} \end{array}$$
 is  
 a) 3-(carboxymethyl) heptane-1,7-dioic acid  
 b) 5-(carboxymethyl) heptane-1,7-dioic acid  
 c) 2-(carboxymethyl) pentane dicarboxylic acid  
 d) 4-(carboxymethyl) pentane dicarboxylic acid
458. Which of the following species will be diamagnetic?  
 a)  $[\text{Fe}(\text{CN})_6]^{4-}$                       b)  $[\text{FeF}_6]^{3-}$                       c)  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$                       d)  $[\text{CoF}_6]^{3-}$
459. Which one of the following is an outer orbital complex and exhibits paramagnetic behaviour?  
 a)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$                       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                       c)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$                       d)  $[\text{Zn}(\text{NH}_3)_6]^{2+}$
460. Moth balls contain:  
 a) Camphor                      b) Benzoic acid                      c) Naphthalene                      d) Cinnamic acid
461. The number of unidentate ligands in the complex ion is called  
 a) Oxidation number                      b) Primary valency  
 c) Coordination number                      d) EAN
462. According to Hückel rule, the number of  $\pi$ -electrons in anthracene is:  
 a) 12                      b) 14                      c) 10                      d) 20
463. In ethane and cyclohexane which one of the following pairs of conformations are more stable?  
 a) Eclipsed and chair conformations                      b) Staggered and chair conformations  
 c) Staggered and boat conformations                      d) Eclipsed and boat conformations
464. Among the following which is not  $\pi$ -bonded organometallic compound?  
 a)  $\text{K}[\text{PtCl}_3(\eta^2 - \text{C}_2\text{H}_4)]$       b)  $\text{Fe}(\eta^5 - \text{C}_5\text{H}_5)_2$                       c)  $\text{Cr}(\eta^6 - \text{C}_6\text{H}_6)_2$                       d)  $(\text{CH}_3)_4\text{Sn}$
465. *o*, *p*-directing groups are generally:  
 a) Activating groups              b) Deactivating groups              c) Neutral groups              d) None of these
466. Aryl halides are less reactive towards nucleophilic substitution reaction as compared to alkyl halides to:  
 a) The formation of less stable carbonium ion  
 b) Resonance stabilization  
 c) Longer carbon-halogen bond  
 d) The inductive effect
467. Which would be least reactive towards bromine?  
 a) Nitrobenzene                      b) Anisole                      c) Phenol                      d) Chlorobenzene
468. Which has a smell of oil of winter green?



- a) Benzaldehyde      b) Benzoic acid      c) Ethyl salicylate      d) Methyl salicylate
469. The coordination number of Pt in  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$  ion is  
 a) 2      b) 4      c) 6      d) 8
470.  $\text{C}_6\text{H}_5\text{Cl}$  on treating with  $\text{NaOH}$  at  $300^\circ\text{C}$  gives phenol. However the yield is poor because of side reaction producing:  
 a)  $\text{C}_6\text{H}_5\text{Na}$       b)  $\text{C}_6\text{H}_5\text{OCH}_3$       c)  $\text{C}_6\text{H}_5\text{OC}_6\text{H}_5$       d) None of these
471. In  $\text{Cr}(\text{NH}_3)_4\text{Cl}_2\text{Cl}$  the ligands are:  
 a)  $\text{NH}_3$  only      b)  $\text{Cl}^-$  only      c) Both  $\text{NH}_3$  and  $\text{Cl}^-$       d)  $\text{Cr}$ ,  $\text{NH}_3$ ,  $\text{Cl}^-$
472. Which statement is not correct regarding aniline?  
 a) It is less basic than ethyl amine  
 b) It can be steam distilled  
 c) It reacts with sodium to give hydrogen  
 d) It is soluble in water
473. Among the following, identify the species with an atom of +6 oxidation state:  
 a)  $[\text{MnO}_4]^-$       b)  $[\text{Cr}(\text{CN})_6]^{3-}$       c)  $[\text{NiF}_6]^{2-}$       d)  $\text{CrO}_2\text{Cl}_2$
474. Which of the following alkanes contain primary, secondary, tertiary and quaternary carbon atom together?  
 a)  $(\text{CH}_3)_3\text{CH}$       b)  $(\text{C}_2\text{H}_5)_3\text{CH}$       c)  $(\text{CH}_3)_3\text{CCH}_2\text{CH}(\text{CH}_3)_2$       d)  $(\text{CH}_3)_4\text{C}$
475. The hardness of water is estimated by:  
 a) Conductivity method      b) EDTA method      c) Titrimetric method      d) Distillation method
476.  $\text{I}_2$  is stirred in between two liquids,  $\text{C}_6\text{H}_6$  and water. It:  
 a) Dissolves more in  $\text{C}_6\text{H}_6$   
 b) Dissolves more in  $\text{H}_2\text{O}$   
 c) Dissolve equally  
 d) Dissolves in neither  $\text{C}_6\text{H}_6$  nor water
477. The number of tertiary C-atoms in 2,2,4,4-tetra methyl pentane is  
 a) 1      b) 2      c) 3      d) 4
478. Hydrogenation of benzoyl chloride in the presence of Pd on  $\text{BaSO}_4$  gives:  
 a) Benzyl alcohol      b) Benzaldehyde      c) Benzoic acid      d) Phenol
479. The Clemmensen reduction of benzaldehyde gives:  
 a)  $\text{C}_6\text{H}_5\text{NH}_2$       b)  $\text{C}_6\text{H}_5\text{OH}$       c)  $\text{C}_6\text{H}_5\text{CH}_3$       d)  $\text{C}_6\text{H}_5\text{COOH}$
480. Which of the following ligand has lowest  $\Delta_o$  value?  
 a)  $\text{CN}^-$       b)  $\text{CO}$       c)  $\text{F}^-$       d)  $\text{NH}_3$
481. Which one of the following has an optical isomer?  
 (en=ethylenediamine)  
 a)  $[\text{Zn}(\text{en})(\text{NH}_3)_2]^{2+}$       b)  $[\text{Co}(\text{en})_3]^{3+}$       c)  $[\text{Co}(\text{H}_2\text{O})_4(\text{en})]^{3+}$       d)  $[\text{Zn}(\text{en})_2]^{2+}$
482. Trichloroacetaldehyde,  $\text{CCl}_3\text{CHO}$  reacts with chlorobenzene in presence of sulphuric acid and produces:





483. Which fraction of coal-tar is rich in arene?

- a) Light oil                      b) Heavy oil                      c) Green oil                      d) Middle oil

484. The coordination number and oxidation number of  $X$  in the following compound  $[X(SO_4)(NH_3)_5]Cl$  will be

- a) 10 and 3                      b) 2 and 6                      c) 6 and 3                      d) 6 and 4

485. Benzyl chloride is formed by treating toluene with  $Cl_2$  in:

- a) Presence of light  
b) Absence of light  
c) Treating benzene with anhy.  $AlCl_3$   
d) Treating benzene with  $As_2S_3$

486. Which complex cannot ionize in solution?

- a)  $[CoCl_3(NH_3)_3]$                       b)  $K_4(Fe(CN)_6)$                       c)  $K_2[Pt(F_6)]$                       d)  $[Pt(NH_3)_6]Cl_4$

487.  $[Ni(CN)_4]^{2-}$ ,  $[MnBr_4]^{2-}$  and  $[CoF_6]^{3-}$ , geometry, hybridisation and magnetic moment of the ions respectively, are

- a) Tetrahedral, square planar, octahedral :  
 $sp^3, dsp^2, sp^3d^2$ : 5.9, 0, 4.9  
b) Tetrahedral, square planar, octahedral :  
 $dsp^2, sp^3, sp^3d^2$ : 0, 5.9, 4.9  
c) Square planar, tetrahedral, octahedral :  
 $dsp^2, sp^3, d^2sp^3$ : 5.9, 4.9, 0  
d) Square planar, tetrahedral, octahedral :  
 $dsp^2, sp^3, sp^3d^2$ : 0, 5.9, 4.9


488. Ozonolysis of benzene gives:

- a) 1 molecule of glyoxal  
b) 2 molecules of glyoxal  
c) 3 molecules of glyoxal  
d) None of these

489. In benzene, C—C bond length is 1.39 Å; the C—H bond length is:

- a) 1.39                      b) 1.08                      c) 1.54                      d) 1.46

490. The IUPAC name of following compound is




- a) N,N-dimethyl, 3-methyl pentan-3-amine                      b) 3-N,N-dimethyl, 3-methyl pentanamine  
c) 3-methyl-3-N, N-dimethyl pentane                      d) 3-methyl-3-N, N-dimethyl butane

491. Which of the following may be used as food preservative?

- a) Benzene  
b) Ethylene  
c) Sodium benzoate  
d) Sodium metaaluminate


492. Which compound is formed when sodium phenoxide is heated with ethyl iodide?  
 a) Phenetole                      b) Ethyl phenyl alcohol    c) Phenol                      d) None of these
493. In metal carbonyl (organometallic) complexes, the  $M-C$  bond is  
 a) Ionic                                      b) Covalent with ionic character  
 c) Covalent                                  d) Coordinate covalent

494. Octahedral complex



- a) *cis*                                      b) *trans*                                      c) *mer*                                      d) *fac*
495. The correct order of magnetic moments (spin only values in BM) among the following is  
 (Atomic no. Mn=25, Fe=26, Co=27)  
 a)  $[MnCl_4]^{2-} > [CoCl_4]^{2-} > [Fe(CN)_6]^{4-}$                       b)  $[MnCl_4]^{2-} > [Fe(CN)_6]^{4-} > [CoCl_4]^{2-}$   
 c)  $[Fe(CN)_6]^{4-} > [MnCl_4]^{2-} > [CoCl_4]^{2-}$                       d)  $[Fe(CN)_6]^{4-} > [CoCl_4]^{2-} > [MnCl_4]^{2-}$
496. Aniline and methyl amine can be differentiated by:  
 a) Diazotisation followed by coupling with phenol  
 b) Reaction with chloroform and aqueous solution of KOH  
 c) Reaction with  $HNO_2$   
 d) None of the above
497. The functional group present in cresols is:  
 a) Alcoholic ( $-OH$ )                      b) Aldehydic ( $-CHO$ )                      c) Phenolic ( $-OH$ )                      d) Carboxylic ( $-COOH$ )





498. In the reaction;



the structure of the product T is:

- a)
- b)
- c)
- d)

499. Which one of the following compounds is most acidic?



500. The most unstable configuration of cyclohexane is

- a) Boat                                      b) Chair

c) Twist boat

d) Half chair


501. In which compound synergic effect is present?

- a)  $[\text{Ni}(\text{CO})_4]$                                       b)  $[\text{NiCl}_4]^{2-}$

c)  $[\text{CuCl}_4]^{2-}$

d)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$

502. The IUPAC name of the compound



a) 4-amino-2-ethyl pent-1-ene

c) Amino-4-pentene

b) 2-ethyl pentan-4-amine

d) 4-ethyl pent-4-en-2-amine

503. Aqua regia reacts with Pt to yield:

- a)  $\text{Pt}(\text{NO}_3)_4$                                       b)  $\text{H}_2[\text{PtCl}_6]$


c)  $\text{PtCl}_4$

d)  $\text{PtCl}_2$

504.  $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$  is called:

- a) Potassium aluminooxalate  
b) Potassium alumino(III) oxalate  
c) Potassium trioxalatoaluminate  
d) Potassium trioxalatoaluminate(III)

505. The IUPAC name of



a) 6-oxo-1,2,2-tri methyl bicycle [2.2.1] heptane

c) 1,5,5-trimethyl bicyclo [2.1.1] hexane-2-one

b) 1,7,7-trimethyl bicyclo [2.2.1] heptan-2-one

d) 1,7,7-trimethyl bicyclo [2.1.2] heptan-2-one

506. Nitration of toluene takes place at:

- a) *ortho* position  
b) *meta* position  
c) *para* position  
d) Both *ortho* and *para* position

507. Estimation of calcium and magnesium is done by

- a) EDTA                                      b) Oxalate                                      c) Phosphate                                      d) None of these


508. How many enantiomer pairs are obtained by monochlorination of 2, 3-dimethyl butane?

- a) Four                                      b) Two                                      c) Three                                      d) One

509. Common reactions of benzene and its derivatives are:

- a) Electrophilic addition reactions  
b) Electrophilic substitution reactions  
c) Nucleophilic substitution reactions  
d) Nucleophilic addition reactions

510. The IUPAC name of the compound




a) 1, 3, 5-triheptene






c) 2, 4, 6-heptatriene

b) 2, 4, 6-triheptene


d) Hepta-1, 3, 5-triene


511. Name of compound






- a) 1, 2, 3-triformylpentane  
c) 3-formylpentane-1, 5-dial
- b) Propane-1, 2, 3-tricarbaldehyde  
d) Propane-1, 2, 3-trial
512. The attacking species in aromatic sulphonation is:  
a)  $\text{SO}_3$   
b)  $\text{H}_3\text{SO}_4^+$   
c)  $\text{HSO}_4$   
d)  $\text{SO}_2^+$
513. Which one of the following compound does not react with bromine?  
a) Ethyl amine  
b) Propene  
c) Phenol  
d) Chloroform
514. The magnetic moment (spin only) of  $[\text{NiCl}_4]^{2-}$  is  
a) 1.82 BM  
b) 5.46 BM  
c) 2.82 BM  
d) 1.41 BM
515.  undergoes electrophilic substitution reaction preferentially :  
a) At position-2  
b) At position-3  
c) At position-4  
d) At positions-2 and 4
516. Ionization of  $\text{K}[\text{Ag}(\text{CN})_2]$  will give:  
a)  $\text{K}^+$  and  $[\text{Ag}(\text{CN})_2]^-$  ion  
b) KCN and AgCN  
c)  $\text{K}^+$ ,  $\text{Ag}^+$ ,  $\text{CN}^-$   
d) None of the above
517. The coordination number and oxidation state of Cr in  $\text{K}_3[\text{Cr}(\text{C}_2\text{O}_4)_3]$  are respectively  
a) +6 and +3  
b) 3 and 0  
c) 4 and +2  
d) 3 and +3
518. A complex of platinum, ammonia and chlorine produces four ions per molecule in the solution. The structure consistent with the observation is:  
a)  $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_4$   
b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_4]$   
c)  $[\text{Pt}(\text{NH}_3)_5\text{Cl}]\text{Cl}_3$   
d)  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}_2$
519. The type of magnetism exhibited by  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$  ion is  
a) Paramagnetism  
b) Diamagnetism  
c) Both (a) and (b)  
d) None of these
520. According to effective atomic number rule the central metal acquires:  
a) Inert gas configuration  
b) Duplet  
c) Octet  
d) Quartet
521.  $\text{K}_3\text{CoF}_6$  is high spin complex. What is the hybrid state of Co-atom in this complex?  
a)  $sp^3d$   
b)  $sp^3d^2$   
c)  $d^2sp^3$   
d)  $dsp^2$
522. The correct structure of ethylenediaminetetraacetic acid (EDTA) is  
a)   
b)   
c)   
d) 
523.  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  are examples of which type of isomerism?  
a) Linkage  
b) Optical  
c) Geometrical  
d) Ionisation
524. The coordination number of a central metal atom in a complex is determined by  
a) The number of ligands around a metal ion bonded by  $\sigma$  – bonds  
b) The number of ligands around a metal ion bonded by  $\pi$  –bonds  
c) The number of ligands around a metal ion bonded by  $\sigma$  –and  $\pi$  – bonds both

- d) The number of only anionic ligands bonded to the metal ion
525. Action of benzoic acid with hydrazoic acid in presence of  $\text{N}_3\text{H}$  gives:  
 a) Aniline                      b) Benzamide                      c) Phenyl cyanide                      d) All of these
526. Which ion shows usually the coordination number 6?  
 a)  $\text{Cr}^{3+}$                       b)  $\text{Fe}^{3+}$                       c)  $\text{Fe}^{2+}$                       d) All of these
527. Which of the following represents hexadentate ligand?  
 a) 2, 2-bipyridyl                      b) DMG                      c) Ethylenediamine                      d) None of these
528. Nitrobenzene can be prepared from benzene by using a mixture of conc.  $\text{HNO}_3$  and conc.  $\text{H}_2\text{SO}_4$ . In the mixture, nitric acid acts as a/an:  
 a) Catalyst                      b) Reducing agent                      c) Acid                      d) Base
529. The value of the 'spin only' magnetic moment for one of the following configurations is 2.84 BM. The correct one is  
 a)  $d^5$  (in strong ligand field)                      b)  $d^3$  (in weak as well as strong ligand fields)  
 c)  $d^4$  (in weak ligand field)                      d)  $d^4$  (in strong ligand field)
530. The IUPAC name of the compound




- a) 6-(3-oxobutyl) cyclohexan-1-one                      b) 6-(2-oxobutyl) cyclohexan-1-one  
 c) 2-(3-oxobutyl) cyclohexan-1-one                      d) 2-(2-oxobutyl) cyclohexan-1-one
531. Hybridisation, shape and magnetic moment of  $[\text{Ni}(\text{CN})_4]^{2-}$  ion  
 a)  $dsp^2$ , square planar, zero                      b)  $dsp^2$ , square planar, 1.73  
 c)  $sp^2d^2$ , octahedral, zero                      d)  $d^2sp^3$ , octahedral, 1.73
532. Choose the IUPAC name of   
 a) Dicyclobutane                      b) Bicyclo [2.2.0] hexane  
 c) Bicyclo [2.2.1] hexane                      d) None of these
533. Which of the following is a heterocyclic compound?  
 a) Phenanthrene                      b) Thiophene                      c) Phenol                      d) Aniline
534.  $[\text{Sc}(\text{H}_2\text{O})_6]^{3+}$  ion is  
 a) Colourless and diamagnetic                      b) Coloured and octahedral  
 c) Colourless and paramagnetic                      d) Coloured and paramagnetic
535. Benzene reacts with  $\text{CH}_3\text{Cl}$  in the presence of anhydrous  $\text{AlCl}_3$  to form:  
 a) Xylene                      b) Toluene                      c) Chlorobenzene                      d) Benzylchloride
536. The magnetic moment of  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is  
 a) 1.73                      b) 2.83                      c) 6.6                      d) Zero
537. The correct order of reactivity towards electrophilic substitution is:  
 a) Phenol > Benzene > Chlorobenzene > Benzoic acid  
 b) Benzoic acid > Chlorobenzene > Benzene > Phenol  
 c) Phenol > Chlorobenzene > Benzene > Benzoic acid  
 d) Benzoic acid > Phenol > Benzene > Chlorobenzene
538. The product formed by the reaction of  $\text{C}_6\text{H}_5\text{CN}$  and  $\text{CH}_2\text{N}_2$  is:




d) None of these




539. Increasing order of expected enol content  
 a)  $\text{CH}_3\text{COCH}_2\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_2\text{COCH}_3$   
 b)  $\text{CH}_3\text{COCH}_2\text{COCH}_3 > \text{CH}_3\text{COCH}_2\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{CHO}$   
 c)  $\text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{COCH}_2\text{CHO} > \text{CH}_3\text{COCH}_2\text{COCH}_3$

- d)  $\text{CH}_3\text{COCH}_3 > \text{CH}_3\text{COCH}_2\text{COCH}_3 > \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_2\text{CHO}$
540. Out of the following the metal which forms polynuclear carbonyl is:  
 a) Na                                      b) Mg                                      c) Mn                                      d) All of these
541. Picric acid and benzoic acid can be distinguished by:  
 a) Aqueous  $\text{NaHCO}_3$                       b) Aqueous NaOH                      c) Aqueous  $\text{FeCl}_3$                       d) Aqueous  $\text{Na}_2\text{CO}_3$
542. The compound having the lowest oxidation state of iron is  
 a)  $\text{K}_4\text{Fe}(\text{CN})_6$                       b)  $\text{K}_2\text{FeO}_4$                       c)  $\text{Fe}_2\text{O}_3$                       d)  $\text{Fe}(\text{CO})_5$
543. The name of  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ ,  $[\text{PtCl}_4]^{2-}$  is  
 a) Tetramminedichloroplatinum(IV) tetrachloro platinate(II)  
 b) Dichloroplatinum (IV) tetrachloroplatinate  
 c) Tetrachloroplatinum (II) tetrammineplatinate  
 d) Tetrachloroplatinum (II) dichlorotetraammine platinate
544. *m*-dihydroxybenzene is also called:  
 a) Catechol                                      b) Resorcinol                                      c) Quinol                                      d) Pyrogallol
545. The ion which exhibits green colour  
 a)  $\text{Cu}^{2+}$                                       b)  $\text{Mn}^{2+}$                                       c)  $\text{Co}^{2+}$                                       d)  $\text{Ni}^{2+}$
546.  $\text{X} \xrightarrow{\text{Cl}_2} \text{Benzotrichloride} \xrightarrow{\text{Hydrolysis}} \text{Y}$   
 X and Y respectively are:  
 a) Benzene, benzaldehyde  
 b) Toluene, benzaldehyde  
 c) Toluene, benzoic acid  
 d) Benzene, benzoic acid
547. Geometrical isomerism is found in coordination compounds having coordination number:  
 a) 2                                      b) 3                                      c) 4 (tetrahedral)                      d) 6
548. Which one of the following complexes is not expected to exhibit isomerism?  
 a)  $[\text{Ni}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$                       b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$                       c)  $[\text{Ni}(\text{NH}_3)_2\text{Cl}_2]$                       d)  $[\text{Ni}(\text{en})_3]^{2+}$
549. The correct acidity order of the following is:



- a) (III) > (IV) > (II) > (I)  
 b) (IV) > (III) > (I) > (II)  
 c) (III) > (II) > (I) > (IV)  
 d) (II) > (III) > (IV) > (I)
550. Identify 'Z' in the reaction;





551. Pure aniline is a:

- a) Brown coloured liquid
- b) Colourless liquid
- c) Brown coloured solid
- d) Colourless solid

552. Aromatic compounds undergo most easily:

- a) Nucleophilic substitution
- b) Electrophilic substitution
- c) Nucleophilic addition
- d) Electrophilic addition

553. The colour of  $\text{CoCl}_3 \cdot 5\text{NH}_3 \cdot \text{H}_2\text{O}$  is:

- a) Orange yellow
- b) Orange
- c) Green
- d) Pink

554. The value of  $x$  on the  $[\text{Ni}(\text{CN})_4]^x$  is:


- a) +2
- b) -2
- c) Zero
- d) +4

555. Complexes with halide ligands are generally:

- a) High spin complexes
- b) Low spin complexes
- c) Both (a) and (b)
- d) None of these

556. The hybridization involved in  $[\text{CoF}_6]^{3-}$  is:

- a)  $d^2 sp^3$
- b)  $d^3 sp^2$
- c)  $dsp^3$
- d)  $sp^3 d^2$



Will have the name

- a) N-ethyl-N-methylethanamine
- b) N,N-diethylmethanamine
- c) N,N-diethylethanamide
- d) None of the above

558. The oxidation state of Fe in the brown ring complex  $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]\text{SO}_4$  is

- a) +3
- b) 0
- c) +2
- d) +1

559. The metal ion in complex A has EAN identical to the atomic number of krypton. A is

(At. no. of Cr=24, Fe=26, Pd=46)

- a)  $[\text{Pd}(\text{NH}_3)_6]\text{Cl}_4$
- b)  $[\text{Cr}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$
- c)  $\text{Na}_4[\text{Fe}(\text{CN})_6]$
- d)  $\text{K}_3[\text{Fe}(\text{CN})_6]$

560. Which one of the following is expected to exhibit optical isomerism [en = ethylenediamine]?



- a) *trans* -  $[\text{Co}(\text{en})_2\text{Cl}_2]$
- b) *cis* -  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$
- c) *cis* -  $[\text{Co}(\text{en})_2\text{Cl}_2]$
- d) *Trans* -  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

561. What is the magnetic moment of  $\text{K}_3[\text{FeF}_6]$ ?

- a) 5.91 BM
- b) 4.89 BM
- c) 3.87 BM
- d) 6.92 BM

562. Identify 'Y' in the change;





563. Among the following statements on the nitration of aromatic compounds, the false one is:

- The rate of nitration of benzene is almost the same as that of hexadeuterobenzene
- The rate of nitration of toluene is greater than that of benzene
- The rate of nitration of benzene is greater than that of hexadeuterobenzene
- Nitration is an electrophilic substitution reaction


564. The bond length of C—O bond in carbon monoxide is 1.128 Å. The C—O bond in  $\text{Fe}(\text{CO})_5$  is:

- 1.15 Å
- 1.128 Å
- 1.72 Å
- 1.118 Å

565. Which one is not correct for homologous series?

- All members are represented by same general formula
- All members have same chemical properties
- All members have same physical properties
- All members have same functional group

566.



is named in IUPAC as

- 2, 3-dimethyl bicyclo [2.2.1] hept-5-ene
- 1, 2-dimethyl bicyclo [2.2.1] hept-4-ene
- 5, 6-dimethyl bicyclo [2.2.1] hept-2-ene
- 4, 5-dimethyl bicyclo [2.2.1] hept-1-ene

567. Ferric ion forms a prussian blue coloured solution due to the formation of:

- $\text{K}_4[\text{Fe}(\text{CN})_6]$
- $\text{Fe}(\text{CNS})_3$
- $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$
- $\text{K}_3[\text{Fe}(\text{CN})_6]$

568. What is the magnetic moment of  $[\text{FeF}_6]^{3-}$ ?

- 5.92
- 5.49
- 2.34
- 4

569. Which of the following can exhibit geometrical isomerism?

- $[\text{MnBr}_4]^{2-}$
- $[\text{Pt}(\text{NH}_3)_3\text{Cl}]^+$
- $[\text{PtCl}_2, \text{P}(\text{C}_2\text{H}_5)_3]_2$
- $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+}$

570. A compound contains 2 dissimilar asymmetric C-atoms. The number of optical isomers are

- 2
- 3
- 4
- 5

571. Coordination number of Ni in  $[\text{Ni}(\text{C}_2\text{O}_4)_3]^{4-}$  is:

- 3
- 6
- 4
- 5

572. Which compound exhibits optical isomerism?

- Pentaamminenitrocobalt (III) iodide
- Diamminedichloroplatinum (II)
- Trans*-dicyano-bis-(ethylenediamine) chromium (III) chloride
- Tris-(ethylenediamine)cobalt (III) bromide

573. Ruthenium carbonyl is:

- $\text{Ru}(\text{CO})_4$
- $\text{Ru}(\text{CO})_5$
- $\text{Ru}(\text{CO})_8$
- $\text{Ru}(\text{CO})_6$

574. Oxidation state of nitrogen is incorrectly given for


- | Compound                                            | Oxidation state |
|-----------------------------------------------------|-----------------|
| a) $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ | 0               |
| b) $\text{NH}_2\text{OH}$                           | -1              |
| c) $(\text{N}_2\text{H}_5)_2\text{SO}_4$            | +2              |
| d) $\text{Mg}_3\text{N}_2$                          | -3              |

575. Which of the following can participate in linkage isomerism?


- $\text{NH}_3$
- $\text{H}_2\text{O}$
- $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$
- $\text{NO}_2^-$

576. *Ortho*-nitrophenol is less soluble in water than *p*- and *m*-nitrophenols because:


- o*-nitrophenol shows intramolecular H-bonding

- b) *o*-nitrophenol shows intermolecular H-bonding  
 c) Melting point of *o*-nitrophenol is lower than those of *m*- and *p*-isomers  
 d) *o*-nitrophenol is more volatile in steam than those of *m*- and *p*-isomers
577. Among the following most basic compound is:  
 a) Benzyl amine                      b) Aniline                      c) Acetanilide                      d) *p*-nitro aniline
578. The EAN of platinum in potassium hexachloroplatinate (IV) is:  
 a) 46                      b) 86                      c) 36                      d) 84
579. The number of ions formed when copper ammonium sulphate is dissolved in water is:  
 a) 1                      b) 2                      c) 4                      d) Zero
580. Which of the following cannot show linkage isomerism?  
 a)  $\text{NO}_2^-$                       b)  $\text{NH}_3$                       c)  $\text{CN}^-$                       d)  $\text{SCN}^-$
581. Xylenes on oxidation with acidic  $\text{KMnO}_4$  gives:  
 a) Phthalic acid                      b) Isophthalic acid                      c) Terephthalic acid                      d) All of these
582. The ratio of  $\sigma$ - and  $\pi$ -bonds in benzene is:  
 a) 2                      b) 4                      c) 6                      d) 8
583. The order of decreasing reactivity towards  $S_E$  reaction for the given compound is:  
 (i)  $\text{C}_6\text{H}_6$   
 (ii)  $\text{C}_6\text{H}_5\text{CH}_3$   
 (iii)  $\text{C}_6\text{H}_5\text{Cl}$   
 (iv)  $\text{C}_6\text{H}_5\text{OH}$   
 a) (ii) > (iv) > (i) > (iii)                      b) (iv) > (iii) > (ii) > (i)                      c) (iv) > (ii) > (i) > (iii)                      d) (i) > (ii) > (iii) > (iv)
584. Which of the following compounds is not optically active?
- 


a)



b)



c)



d)
585. The number of geometrical isomers of  $[\text{Co}(\text{NH}_3)_3(\text{NO}_2)_3]$  are:  
 a) Zero                      b) 2                      c) 3                      d) 4
586. Phenol is less acidic than:  
 a) Water                      b) *p*-methoxyphenol                      c) *p*-nitrophenol                      d) Ethanol
587. In the reaction,  

$$\text{C}_6\text{H}_5\text{CH}_3 \xrightarrow{\text{Oxidation}} \text{A} \xrightarrow{\text{NaOH}} \text{B} \xrightarrow[\Delta]{\text{Soda lime}} \text{C}_s$$
  
 a)  $\text{C}_6\text{H}_5\text{OH}$                       b)  $\text{C}_6\text{H}_6$                       c)  $\text{C}_6\text{H}_5\text{COONa}$                       d)  $\text{C}_6\text{H}_5\text{ONa}$
588. Incorrect statement is  
 a) Ethane can have an infinite number of conformations  
 b) Cyclopropane molecule has considerable angle strain  
 c) Eclipsed form of ethane is less stable than staggered conformation  
 d) Staggered conformation possess maximum energy
589. The complex  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  will give white ppt. with:  
 a)  $\text{PbCl}_2$                       b)  $\text{AgNO}_3$                       c) KI                      d) None of these
590. Which of the following complexes exhibits the highest paramagnetic behaviour?  
 a)  $[\text{Fe}(\text{en})(\text{bpy})(\text{NH}_3)_2]^{2+}$   
 b)  $[\text{Co}(\text{OX})_2(\text{OH})_2]^-$   
 c)  $[\text{Ti}(\text{NH}_3)_6]^{3+}$   
 $[\text{V}(\text{gly})_2(\text{OH})_2(\text{NH}_3)_2]^+$   
 d) Where, gly = glycine, en = ethylenediamine and bpy = bipyridylmoities  
 (At. No. Ti=22, V=23, Fe=26, Co=27)
591. The coordination number in a/an ..... complex may increase to 8.  
 a) Cobalt                      b) Osmium                      c) Nickel                      d) Iron

592. Compound used for covering wounds caused by bite of mad dog is:  
 a) Benzoic acid                      b) Aniline                      c) Phenol                      d) Salicylic acid
593. Cinnamic acid on decarboxylation gives:  
 a) Benzene                      b) Toluene                      c) Styrene                      d) Benzaldehyde
594. In which of the following pairs both the complex show optical isomerism?  
 a)  $Cis-[Cr(C_2O_4)_2Cl_2]^{3-}$ ;  $cis-[Co(NH_3)_4Cl_2]$                       b)  $[PtCl(dien)]Cl$ ,  $[NiCl_2Br_2]^{2-}$   
 c)  $[Co(NO_3)_3(NH_3)_3]$ ,  $cis-[Pt(en)_2Cl_2]$                       d)  $[Co(en)_3]Cl_3$ ,  $cis-[Co(en)_2Cl_2]Cl$
595. The name of the ring structure complex compound formed between metal ion and polydentate ligand is  
 a) Simple complex                      b) Chelate complex                      c) Polynuclear complex                      d) None of the above
596. IUPAC name of  

$$\begin{array}{c} Cl_2CH - CH - CH - CCl_3 \\ | \quad \quad | \\ C_2H_5 \quad C_2H_5 \end{array}$$
  
 a) 1,1,1,4,4-pentachloro-2,3-diethyl-butane  
 b) 3-(dichloromethyl)-4-(trichloromethyl)-hexane  
 c) 3-(trichloromethyl)-4-(dichloromethyl)-hexane  
 d) 1,1,4,4,4-pentachloro-2,3-diethyl butane
597. Which statement is wrong with regard to acetaldehyde and benzaldehyde?  
 a) Both react with hydroxylamine to form oximes  
 b) Both react with HCN to form cyanohydrin  
 c) Both react with NaOH to form polymers  
 d) Both react with hydrazine to form hydrazones
598. The coordination number of Cu in complex  $[Cu(H_2O)_4]^{2+}$  is  
 a) 4                      b) 3                      c) 2                      d) 1
599. Which reaction sequence would be best to prepare 3-chloroaniline from benzene?  
 a) Chlorination, nitration, reduction  
 b) Nitration, chlorination, reduction  
 c) Nitration, reduction, chlorination  
 d) Nitration, reduction, acetylation, chlorination, hydrolysis
600. The complexes  $(Co(NH_3)_6)[Cr(C_2O_4)_3]$  and  $[Cr(NH_3)_6][Co(C_2O_4)_3]$   
 a) Geometrical isomerism                      b) Ionization energy  
 c) Coordination isomerism                      d) Linkage isomerism
601. The reaction,  

$$C_6H_5NHCOCH_3 \xrightarrow{B_2/Fe} BrC_6H_4NHCOCH_3$$
  
 is an example of:  
 a) Substitution reaction  
 b) Addition reaction  
 c) Condensation reaction  
 d) Elimination reaction
602. Given the molecular formula of the hexa coordinated complexes is  
 (A)  $CoCl_3 \cdot 6NH_3$   
 (B)  $CoCl_3 \cdot 5NH_3$   
 (C)  $CoCl_3 \cdot 4NH_3$   
 If the number of coordinated  $NH_3$  molecules in A, B and C respectively are 6, 5 and 4 the primary valency in (A), (B) and (C) are  
 a) 6, 5, 4                      b) 3, 2, 1                      c) 0, 1, 2                      d) 3, 3, 3
603.  $C_6H_{14}$  has two tertiary carbons. The IUPAC name is  
 a) n-hexane                      b) 2-methylpentane                      c) 3-methylpentane                      d) 2,3-dimethylbutane
604. The compound  $[Co(NO_2)(NH_3)_5]Cl_2$  and  $[Co(ONO)(NH_3)_5]Cl_2$  are examples of:  
 a) Geometrical isomers                      b) Linkage isomers                      c) Ligand isomers                      d) Ionization isomers


605. Which is not a  $\pi$ -bonded complex?

- a) Zeise salt                      b) Ferrocene                      c) Dibenzene chromium                      d) Tetraethyl lead


606. When phenol is treated with  $\text{PCl}_5$ , the yield of chlorobenzene is generally poor because of the formation of:

- a) Benzoyl chloride                      b) *p*-chlorophenol                      c) *o*-chlorophenol                      d) Tertiary phosphate

607. Which will show tautomerism?




608. The IUPAC name of compound



- a) N-phenylaminoethanone                      b) N-phenylethanamide  
c) N-phenylmethanamide                      d) N-phenylaminomethane

609. Which one of the following is most reactive towards electrophilic reagent?



610. Which of the following shows  $dsp^2$  hybridisation?

- a)  $\text{NiCl}_4^{2-}$                       b)  $\text{SCl}_4$                       c)  $\text{NH}_4^+$                       d)  $\text{PtCl}_4^{2-}$

611. Which one of the following is not an explosive?

- a) Trinitroglycerine                      b) *o*-aminotoluene                      c) Dynamite                      d) TNT

612. Spin only magnetic moment of the compound  $\text{Hg}[\text{Co}(\text{SCN})_4]$  is

- a)  $\sqrt{3}$                       b)  $\sqrt{15}$                       c)  $\sqrt{24}$                       d)  $\sqrt{8}$

613. When phenol is treated with  $\text{NH}_3$  and  $\text{ZnCl}_2$ , it changes to:

- a) Aniline                      b) Salicylic acid                      c) Cyclohexanol                      d) None of these

614. In which complex is the transition metal in zero oxidation state?

- a)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_2$                       b)  $[\text{Fe}(\text{H}_2\text{O})_6]\text{SO}_4$                       c)  $[\text{Ni}(\text{CO})_4]$                       d)  $[\text{Fe}(\text{H}_2\text{O})_3](\text{OH})_2$

615. The species having tetrahedral shape is

- a)  $[\text{NiCl}_4]^{2-}$                       b)  $[\text{Ni}(\text{CN})_4]^{2-}$                       c)  $[\text{PdCl}_4]^{2-}$                       d)  $[\text{Pd}(\text{CN})_4]^{2-}$

616. An imperfect complex of a complex compound is 100% ionized; the compound is called:

- a) Double salt                      b) Complex salt                      c) Acid salt                      d) Normal salt

617. For which transition metal ions are low spin complexes impossible?

- a)  $\text{Zn}^{2+}$                       b)  $\text{Zr}^{2+}$                       c)  $\text{Ag}^+$                       d) All are correct

618. (A)  $\text{K}_4[\text{Fe}(\text{CN})_6]$

(B)  $\text{K}_3[\text{Cr}(\text{CN})_6]$

(C)  $\text{K}_3[\text{Co}(\text{CN})_6]$

(D)  $\text{K}_2[\text{Ni}(\text{CN})_6]$

Select the complexes which are diamagnetic.





- a) (A), (B) and (C)                      b) (B), (C) and (D)                      c) (A), (C) and (D)                      d) (A), (B) and (D)

619. Wilkinson's catalyst,  $(\text{Ph}_3\text{P})_3\text{RhCl}$  is used for  
 a) Hydrogenation of oils  
 b) Hydrogenation of alkynes  
 c) Hydrogenation of alkenes  
 d) Polymerization of alkenes
620. Among the following compounds, the most acidic is:  
 a) *p*-nitrophenol  
 b) *p*-hydroxybenzoic acid  
 c) *o*-hydroxybenzoic acid  
 d) *p*-toluic acid
621. An aromatic primary amine with cold nitrous acid leads to the formation of:  
 a) Alcohol  
 b) Nitrite  
 c) Diazonium salt  
 d) Benzene
622. Chlorobenzene gives DDT when it reacts with:  
 a) Phenol  
 b) Naphthalene  
 c) Chloral  
 d) Acetaldehyde
623. Under suitable conditions  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$  (A),  $\text{C}_6\text{H}_5\text{OH}$  (B) and  $\text{C}_6\text{H}_5\text{COOH}$  (C) can act as acids. The increasing order of their acidic strengths is:  
 a)  $A < B < C$   
 b)  $A < C < B$   
 c)  $B < A < C$   
 d)  $C < B < A$
624. Which is considered to be an anticancer species?
- a)

b)

c)

d)
625. The compound required for the formation of thermosetting polymer with methanal is:  
 a) Phenol  
 b) Benzene  
 c) Benzaldehyde  
 d) All of these
626. Which one of the following has highest number of isomers?  
 a)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$   
 b)  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$   
 c)  $[\text{Ru}(\text{NH}_3)_4\text{Cl}]^-$   
 d)  $[\text{In}(\text{PP}_3)_2\text{H}(\text{CO})]^{2+}$
627. Which group is *o*- and *p*-directing?  
 a)  $-\text{NO}_2$   
 b)  $-\text{SO}_3\text{H}$   
 c)  $-\text{COOH}$   
 d)  $-\text{NHCOCH}_3$
628. When benzyl chloride is boiled with aqueous solution of lead nitrate in current of carbon dioxide, the main product is:  
 a) Benzoic acid  
 b) Benzyl alcohol  
 c) Benzaldehyde  
 d) Nitrobenzene
629. Ligands in complex compounds  
 a) Donates electron pair  
 b) Accept electron pair  
 c) Neither accept electron pair nor donate  
 d) All of the above
630. Aniline is separated by:  
 a) Fractional crystallisation  
 b) Fractional distillation  
 c) Steam distillation  
 d) Vacuum distillation
631. In which of the following octahedral complexes of Co (at. No. 27), will be magnitude of  $\Delta_0$  be the highest?  
 a)  $[\text{Co}(\text{CN})_6]^{3-}$   
 b)  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$   
 c)  $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$   
 d)  $[\text{Co}(\text{NH}_3)_6]^{3+}$
632. The IUPAC name of  $\text{K}_2[\text{PtCl}_6]$  is  
 a) Hexachloroplatinate potassium  
 b) Potassium hexachloroplatinate (IV)  
 c) Potassium hexachloroplatinate  
 d) Potassium hexachloroplatinum(IV)
633. Aqueous solution of nickel sulphate on treating with pyridine and then adding a solution of sodium nitrite gives dark blue crystals of:  
 a)  $[\text{Ni}(\text{py})_4]\text{SO}_4$   
 b)  $[\text{Ni}(\text{py})_2(\text{NO}_2)_2]$   
 c)  $[\text{Ni}(\text{py})_4(\text{NO}_2)_2]$   
 d)  $[\text{Ni}(\text{py})_3(\text{NO}_2)]_2\text{SO}_4$
634. Benzyl alcohol is obtained from benzaldehyde by:  
 a) Fittig's reaction  
 b) Cannizzaro's reaction  
 c) Kolbe's reaction  
 d) Wurtz's reaction
635. The structure of the compound that gives a tribromo derivative on treatment with bromine water is:



636. The coordination number and the oxidation state of the element 'E' in the complex  $[E(en)_2(C_2O_4)]NO_2$  (where (en) is ethylene diamine) are, respectively :

- a) 6 and 3                      b) 6 and 2                      c) 4 and 2                      d) 4 and 3

637. Benzaldehyde reacts with  $PCl_5$  to give:

- a) Benzyl chloride              b) Benzo trichloride              c) Benzal chloride              d) Chlorobenzene

638. Which one of the following complex ions has geometrical isomers?

- a)  $[Co(en)_3]^{3+}$                       b)  $[Ni(NH_3)_5Br]^+$                       c)  $[Co(NH_3)_2(en)_2]^{3+}$                       d)  $[Cr(NH_3)_4(en)]^{3+}$

639. The strongest acid among the following aromatic compounds is:

- a) *Ortho*-nitrophenol              b) *para*-chlorophenol              c) *para*-nitrophenol              d) *meta*-nitrophenol

640. The isomers observed in alkanes is

- a) Metamerism                                              b) Chain isomerism  
c) Position isomerism                                      d) Geometrical isomerism





641. The two compounds pentaamminesulphatocobalt (III) bromide and pentaamminesulphatocobalt(III) chloride represent:

- a) Linkage isomerism  
b) Ionization isomerism  
c) Coordination isomerism  
d) No isomerism

642. Both  $[Ni(CO)_4]$  and  $[Ni(CN)_4]^{2-}$  are diamagnetic. The hybridisation of nickel in the compounds respectively are :

- a)  $sp^3, sp^3$                       b)  $sp^3, dsp^2$                       c)  $dsp^2, sp^3$                       d)  $dsp^3, dsp^2$

643. The following compounds on hydrolysis in aqueous acetone will give:



- a) Mixture of (K) and (L)    b) Mixture of (K) and (M)    c) Only (M)                      d) Only (K)

644. The number of  $\pi$ -electrons in cyclo hepta trienyl anion is:

- a) 2                                      b) 3                                      c) 8                                      d) 5

645. In the Grignard reaction, which metal forms an organometallic bond?

- a) Sodium                              b) Titanium                              c) Magnesium                      d) Palladium

646. Aromatic hydrocarbons are the derivatives of:


- a) Benzene  
b) Methane  
c) Normal series of paraffins  
d) None of the above

647. Benzene easily shows:

- a) Ring fission reactions since it is unstable  
b) Addition reactions since it is unsaturated  
c) Electrophilic substitution reactions due to stable ring and high  $\pi$ -electron density

d) Nucleophilic substitution reactions due to stable ring and minimum electron density


648. The IUPAC name of the compound



is

- a) Tetra phenyl methane  
b) 1,1,1,1-tetraphenyl methane  
c) 1,1,1,1-tetracyclohexyl methane  
d) Methyno-1,1,1-1-tetracyclohexane

649.



having the IUPAC name as

- a) 2,4,4-trimethyl pentanal  
b) 4,4,2-trimethyl pentanal  
c) 1,3,3-trimethyl butanal  
d) 3,3,1-trimethyl butanal

650. When benzoic acid is heated with soda lime, we get:

- a) Phenol  
b) Benzyl alcohol  
c) Benzene  
d) Benzaldehyde

651. If a compound absorbs violet colour from the sunlight, then the observed colour is:

- a) Yellow  
b) Orange  
c) Blue  
d) Green

652. Sulphonic acid is used in the manufacture of:

- a) Antipyretics  
b) Antitoxine  
c) Antibiotics  
d) Dyes

653. In the silver plating of Cu,  $K[Ag(CN)_2]$  is used instead of  $AgNO_3$ . The reason is:

- a) A thin layer of Ag is formed on Cu  
b) More heat is required  
c)  $Ag^+$  ions are completely removed from solution  
d) Less availability of  $Ag^+$  ion as Cu cannot displace Ag from  $Ag(CN)_2$

654. The strongest *o*-, *p*-directing group among the following is:

- a)  $-OH$   
b)  $-Cl$   
c)  $-C_6H_5$   
d)  $-Br$

655. Out of  $TiF_6^{2-}$ ,  $CoF_6^{3-}$ ,  $Cu_2Cl_2$  and  $NiCl_4^{2-}$  ( $Z$  of Ti = 22, Co = 27, Cu = 29, Ni = 28) the colourless species are:

- a)  $CoF_6^{3-}$  and  $NiCl_4^{2-}$   
b)  $TiF_6^{2-}$  and  $CoF_6^{3-}$   
c)  $Cu_2Cl_2$  and  $NiCl_4^{2-}$   
d)  $TiF_6^{2-}$  and  $Cu_2Cl_2$

656. Which is true in the case of  $[Fe(CN)_6]^{3-}$  complex?

- a)  $d^2sp^3$ -hybridization of Fe  
b) Paramagnetic  
c) One unpaired electron  
d) All of the above are correct

657. The IUPAC name of  $[Ni(PPh_3)_2Cl_2]^{2+}$  is

- a) Bis-dichloro (triphenylphosphine)nickel(II)  
b) Dichloro bis (triphenylphosphine)nickel(II)  
c) Dichloro triphenylphosphine nickel(II)  
d) Triphenyl phosphine nickel (II) dichloride

658. The complex  $[Co(NH_3)_3Cl_3]$  is:

- a) Neutral  
b) Cationic  
c) Anionic  
d) None of these

659. From the stability constant (hypothetical values) given below, predict which is the strongest ligand?

- a)  $Cu^{2+} + 4NH_3 \rightleftharpoons [Cu(NH_3)_4]^{2+}$  ; ( $K = 4.5 \times 10^{11}$ )  
b)  $Cu^{2+} + 4CN \rightleftharpoons [Cu(CN)_4]^{2-}$  ; ( $K = 2.0 \times 10^{27}$ )  
c)  $Cu^{2+} + 2en \rightleftharpoons [Cu(en)_2]^{2+}$  ; ( $K = 3.0 \times 10^{15}$ )  
d)  $Cu^{2+} + 4H_2O \rightleftharpoons [Cu(H_2O)_4]^{2+}$  ; ( $K = 9.5 \times 10^8$ )

660. Which has highest m.p.?

- a) *o*-bromophenol  
b) *m*-bromophenol  
c) *p*-bromophenol  
d) *m*-chlorophenol

661. Hexafluorocobaltate(III) ion is found to be high spin complex, the probable hybrid state of cobalt in it is:

- a)  $d^2sp^3$   
b)  $sp^3$   
c)  $sp^3d$   
d)  $sp^3d^2$



662. Which isomeric dibromotoluene is most difficult to make from toluene?  
 a) 2,3                                      b) 2,4                                      c) 3,5                                      d) 2,6
663. Which one of the following forms with an excess of  $\text{CN}^-$  (cyanide) a complex?  
 a)  $\text{Cu}^+$                                       b)  $\text{Ag}^+$                                       c)  $\text{Ni}^{2+}$                                       d)  $\text{Fe}^{2+}$
664. Nitration of salicylic acid gives:  
 a) 2,4,6-trinitrosalicylic acid  
 b) 2,4,6-trinitrophenol  
 c) 2,4,6-trinitrobenzoic acid  
 d) None of the above
665. The IUPAC name of the compound  

$$\begin{array}{c} \text{CH}_3-\text{CH}_2-\text{C}-\text{CH}_2-\text{CH}_3 \\ \parallel \\ \text{N}-\text{OH} \end{array}$$
 is  
 a) N-hydroxy-3-amino pentane                                      b) N-hydroxyamino pentane  
 c) N-hydroxy-3-imino pentane                                      d) None of the above
666. Which is not true of the coordination compound  $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$ ?  
 a) Exhibits geometrical isomerism                                      b) Exhibits optical isomerism  
 c) Exhibits ionisation isomerism                                      d) Is an octahedral complex
667. The IUPAC name of  

$$\begin{array}{c} \text{CH}_3 \qquad \qquad \text{O} \\ | \qquad \qquad \parallel \\ \text{CH}_3-\text{CH}-\text{CH}-\text{C}-\text{Cl} \\ | \\ \text{CH}_2\text{Br} \end{array}$$
 is  
 a) 3-(bromomethyl)-2-methyl butanoyl chloride                                      b) 3-(bromomethyl)-2-methyl propanoyl chloride  
 c) 2-(bromomethyl)-3-methyl butanoyl chloride                                      d) None of the above
668. Aniline is reacted with bromine water and the resulting product is treated with an aqueous solution of sodium nitrite in the presence of dilute HCl. The compound so formed is treated with fluoroboric acid which is subsequently heated dry. The final product is:  
 a) *p*-bromofluorobenzene  
 b) *p*-bromoaniline  
 c) 2,4,6-tribromofluorobenzene  
 d) 1,3,5-tribromobenzene
669. Which of the following is a common donor atom in ligands?  
 a) Nitrogen                                      b) Oxygen                                      c) Arsenic                                      d) Both (b) and (c)
670. The reaction of aniline with acetyl chloride in presence of NaOH gives:  
 a) Acetanilide                                      b) Aniline hydrochloride                                      c) *p*-chloroaniline                                      d) A red dye
671. In the reaction, the compound "X" is:  

$$\text{Me}-\text{C}_6\text{H}_4-\text{CHO} + \text{X} \xrightarrow[\text{H}_2\text{O}]{\text{CH}_3\text{COONa}}$$

$$\text{Me}-\text{C}_6\text{H}_4-\text{CH}=\text{CHCOOH}$$
  
 a)  $\text{CH}_3\text{COOH}$   
 b)  $\text{Br} \cdot \text{CH}_2\text{COOH}$   
 c)  $(\text{CH}_3\text{CO})_2\text{O}$   
 d)  $\text{CHO} \cdot \text{COOH}$
672. Which of the following will exhibit maximum ionic conductivity?  
 a)  $\text{K}_4[\text{Fe}(\text{CN})_6]$                                       b)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$                                       c)  $[\text{Cu}(\text{NH}_3)_4]\text{Cl}_2$                                       d)  $[\text{Ni}(\text{CO})_4]$
673. Dipole moment of *p*-nitroaniline, when compared to nitrobenzene (X) and aniline (Y) will be:  
 a) Greater than (X) and (Y)  
 b) Smaller than (X) and (Y)  
 c) Greater than (X) but smaller than (Y)



d) Equal to zero


674. The structure of iron pentacarbonyl is:

- a) Square planar      b) Trigonal bipyramidal      c) Triangular      d) None of these

675. Turnbull's blue is:

- a) Ferricyanide      b) Ferrous ferricyanide      c) Ferrous cyanide      d) Ferri ferrocyanide

676. The correct IUPAC name of



- a) 2-hydroxypropane-1, 2, 3-tricarboxylic acid      b) 3-carboxy-3-hydroxy-pentane-1, 5-dioic acid  
c) 2 carboxy-4 hydroxy-pentane-1, 5-dioic acid      d) 3-carboxy-3-hydroxy-hexane-1, 6-dioic acid

677. The trivial name among the following is

- a) Acetone      b) Acetylene      c) Uric acid      d) None of these

678. The IUPAC name of  $[\text{Pt}(\text{NH}_3)_4(\text{NO}_2)\text{Cl}]\text{SO}_4$  is

- a) Chloronitro tetrammine platinum (IV) sulphate  
b) Tetrammine chloronitro platinum (II) sulphate  
c) Tetrammine chloronitro platinum (IV) sulphate  
d) Chlorotetrammine nitroplatinum (IV) sulphate

679. The overlapping in benzene is in carbon-carbon orbitals of the type:

- a)  $p - p$       b)  $sp - sp$       c)  $sp^2 - sp^2$       d)  $sp^3 - sp^3$


680. Change in composition of coordination sphere yields which type of isomer?





- a) Geometrical      b) Ionization      c) Optical      d) None of these

681. The IUPAC name of  $\text{K}_2[\text{Ni}(\text{CN})_4]$  is

- a) Potassium tetracyanonickelate (II)      b) Potassium tetracyanonickelate (III)  
c) Potassium tetracyanonickel (II)      d) Potassium tetracyanonickel (III)

682. Aniline in a set of the following reactions yielded a coloured compound Y:



- a) 
- b) 
- c) 
- d) 

683. The effective atomic number rule is less likely to apply if the metal-ligand bond:

- a) Is extremely weak  
b) Has a covalent character  
c) Has a large amount of ionic character  
d) None is correct

684. Potassium ferrocyanide is an example of

- a) Tetrahedral      b) Octahedral      c) Square planar      d) Linear


685. 1-phenyl, 2-chloropropane on treatment with aqueous KOH gives mainly:

- a) 1-phenylpropane      b) 3-phenylpropane      c) 1-phenylpropan-2-ol      d) 1-phenylpropan-3-ol
686. Which class of compounds can exhibit geometrical isomerism?
- a)  $C_6H_5CH=NOH$       b)  $CH_3CH=CHCH_3$   
 c)  $HOOCCH=CH_2-CHCOOH$       d) All of the above

687. The product of oxidation of aniline with  $K_2Cr_2O_7$  and conc.  $H_2SO_4$  will be:

- a) p-amino phenol  
 b) p-benzoquinone  
 c) Aniline black dye  
 d) Phenyl hydroxylamine

688. Among the following the Newmann projections of *meso*-2, 3-butanediol are




- a) P, Q      b) P, R      c) R, S      d) Q, S

689. A new carbon-carbon bond is formed in:

- a) Cannizzaro's reaction  
 b) Friedel-Crafts reaction  
 c) Clemmensen reduction  
 d) None of the above

690. Which of the following compounds can exhibit tautomerism?



691. The most basic compound among the following is:

- a) Benzylamine      b) Aniline      c) Acetanilide      d) p-nitroaniline

692. Which of the following has least oxidation state of Fe?

- a)  $K_3[Fe(OH)_6]$       b)  $K_2[FeO_4]$   
 c)  $FeSO_4(NH_4)_2SO_4 \cdot 6H_2O$       d)  $[Fe(CN)_6]^{3-}$

693. The spin only magnetic moment value (in Bohr magneton units) of  $Cr(CO)_6$  is

- a) 0      b) 2.84      c) 4.90      d) 5.92

694. Which is an excellent antiseptic?

- a) Phenol      b) Benzyl alcohol      c) Benzaldehyde      d) Acetic acid

695. Scientist who explained the structures and isomerism in the complex compound was:

- a) Sidgwick      b) Pauling      c) Powell      d) Werner

696. The cation that does not form an ammine complex with excess of ammonia is:

- a)  $Al^{3+}$       b)  $Ag^+$       c)  $Cu^{2+}$       d)  $Cd^{2+}$

697. The complex ion which has the highest magnetic moment among the following is

- a)  $[CoF_6]^{3-}$       b)  $[Co(NH_3)_6]^{3+}$       c)  $[Ni(NH_3)_4]^{2+}$       d)  $[Ni(CN)_4]^{2-}$

698. For square planar complex of platinum (II),  $[Pt(NH_3)(Br)(Cl)Py]^{2+}$ , how many isomeric forms are possible?

- a) Two      b) Three      c) Four      d) Six

699. Which of the following has highest boiling point?

- a) Benzene                      b) Phenol                      c) Toluene                      d) Ethyl benzene

700. A nitrogen containing organic compound on heating with chloroform and alcoholic KOH evolved very unpleasant smelling vapours. The compound could be:

- a) Nitrobenzene                      b) Benzamide                      c) *N,N*-dimethyl amine                      d) Aniline


701. Which of the following 0.1 M complex compound solutions will have the minimum electrical conductivity?

- a) Hexammine platinum (IV) chloride                      b) Chloropenta ammine platinum (IV) chloride  
c) Dichloro tetrammine platinum (IV) chloride                      d) Trichloro triammine platinum (IV) chloride

702. False statement is

- a) Aprotic solvents increase the enol content in tautomerism  
b) Any deviation from the normal bond angles introduces angle strain in molecule  
c) Diastereomers have identical physical properties  
d) Chain isomers can also be position isomers

703. The correct IUPAC name of the compound is







- a) 4-bromo-5-chloro-3-iodo hexane                      b) 3-bromo-2-chloro-4-iodo hexane  
c) 3-bromo-4-iodo -2-chloro hexane                      d) 2-bromo-3-bromo-4-iodo hexane


704. Benzyl chloride ( $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ ) can be prepared from toluene by chlorination with:

- a)  $\text{SO}_2\text{Cl}_2$                       b)  $\text{SOCl}_2$                       c)  $\text{S}_2\text{Cl}_2$                       d)  $\text{NaOCl}$

705. The compound 2,2'-bipyridine has the structure




706. The IUPAC name of



- a) 4-formyl-6-oxocyclohexane-1-carboxylic acid                      b) 2-oxo-4-formyl cyclohexane-1-carboxylic acid  
c) 6-oxo-4-formyl cyclohexane-1-carboxylic acid                      d) 4-formyl-2-oxo cyclohexane-1-carboxylic acid

707. The diazonium salt  $\text{H}_3\text{C}-\text{C}_6\text{H}_3(\text{Br})-\text{N}_2\text{Cl}$ , gives,



- a)  $\text{HCl}/\text{CuCl}$                       b)  $\text{HNO}_2/\text{Cu}$                       c)  $\text{C}_2\text{H}_5\text{OH}/\text{Cu}$                       d)  $\text{SnCl}_2/\text{HCl}$

708. Diethylenetriamine is:

- a) Chelating agent                      b) Polydentate ligand                      c) Tridentate ligand                      d) All of these


709. The no. of ions given by  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}_2$  in aqueous solution is:

- a) 2                      b) 3                      c) 4                      d) 5



710. Aniline reacts with excess of bromine to give:

- a) Benzyl bromide and hydrobromic acid  
b) 2,4,6-tribromoaniline  
c) 2-bromotoluene and hydrobromic acid

- d) 2-bromophenol and hydrobromic acid
711. The coordination compounds,  
 $[\text{Co}(\text{NH}_3)_6]^{3+}[\text{Cr}(\text{CN})_6]^{3-}$   
 and  $[\text{Cr}(\text{NH}_3)_6]^{3+}[\text{Co}(\text{CN})_6]^{3-}$  are example of  
 a) Linkage isomerism  
 b) Coordination isomerism  
 c) Ionisation isomerism  
 d) Geometrical isomerism
712. Both  $\text{Co}^{3+}$  and  $\text{Pt}^{4+}$  have a coordination number of six. Which of the following pairs of complexes will show approximately the same electrical conductance for their 0.001 M, aqueous solutions?  
 a)  $\text{CoCl}_3 \cdot 4\text{NH}_3$  and  $\text{PtCl}_4 \cdot 4\text{NH}_3$   
 b)  $\text{CoCl}_3 \cdot 3\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$   
 c)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$   
 d)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 3\text{NH}_3$
713. In SCN ligand if N is attached to central atom, the name of ligand is:  
 a) Thiocyanato-N  
 b) Cyanato-N  
 c) Thiocyanato-S  
 d) Cyanato-S
714. The product formed on heating



715. Oxidation of ethyl benzene by  $\text{KMnO}_4$  gives:  
 a) Benzyl alcohol  
 b) Benzophenone  
 c) Acetophenone  
 d) Benzoic acid
716. One of the following statements regarding Reimer-Tiemann reaction is false:  
 a) Reaction of phenol with  $\text{CHCl}_3$  and  $\text{KOH}$   
 b)  $\text{CCl}_2$  acts as a nucleophile  
 c) Reaction of phenol with  $\text{CCl}_4$  and  $\text{NaOH}$   
 d) Reaction of phenol with formaldehyde to form bakelite
717. The structure representing a heterocyclic compound is



718. Phenol reacts with  $\text{Br}_2$  in  $\text{CCl}_4$  at low temperature to give:

- a) *m*-bromophenol
- b) *o*- and *p*-bromophenol
- c) *p*-bromophenol
- d) 2,4,6-tribromophenol


719. The correct name of the compound  $[\text{Cu}(\text{NH}_3)_4](\text{NO}_3)_2$ , according to IUPAC system is

- a) Cuprammonium nitrate
- b) Tetrammine copper (II) dinitrate
- c) Tetrammine copper (II) nitrate
- d) Tetrammine copper (II) dinitrite

720. Nitroethane can exhibit one of the following kind of isomerism

- a) Metamerism
- b) Optical activity
- c) Tautomerism
- d) Position isomerism

721. What would be the correct IUPAC name of



- a) 3,3-dimethyl-3-cyclopentyl propanal
- b) 3-methyl-3-cyclopentyl butan-1-al
- c) 1-(1-methyl-1-formyl) methylethyl cyclopropane
- d) None of above

722. The number of unpaired electrons in the square planar  $[\text{Pt}(\text{CN})_4]^{2-}$  ion is

- a) 2
- b) 1
- c) 0
- d) 3

723. The oxidation number of cobalt in  $\text{K}[\text{Co}(\text{CO})_4]$  is

- a) +1
- b) +3
- c) -1
- d) -3

724. IUPAC name of  $\text{Na}_3[\text{Co}(\text{NO}_2)_6]$  is

- a) Sodium hexanitrito cobaltate (II)
- b) Sodium hexanitro cobaltate (III)
- c) Sodium hexanitrito cobaltate (III)
- d) Sodium cobaltinitrite(II)

725. The total number of possible isomers for the complex compound  $[\text{Cu}(\text{NH}_3)_4][\text{PtCl}_4]$

- a) 6
- b) 5
- c) 4
- d) 3


726. Benzaldehyde reacts with excess of anhydrous ethyl alcohol in the presence of HCl, gives:

- a)  $\text{C}_6\text{H}_5\text{COCl}$
- b)  $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$
- c)  $\text{C}_6\text{H}_5\text{CH}(\text{OC}_2\text{H}_5)_2$
- d)  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$

727. Which pair of isomerism is not possible together?




- a) Chain and position
- b) Functional and position
- c) Tautomerism and functional
- d) All of the above

728. Which type of conformation is shown by I and II?



- a) I is eclipsed, II is staggered
- b) II is eclipsed, I is staggered
- c) Both are eclipsed
- d) Both are staggered


729. Which will give chiral molecule?




730. The neutral ligand is:

- a) Chloro                      b) Hydroxo                      c) Ammine                      d) Oxalato
731. The effective atomic number of cobalt in the complex  $[\text{Co}(\text{NH}_3)_6]^{3+}$  is  
 a) 36                      b) 24                      c) 33                      d) 30
732.  $\text{K}_4[\text{Fe}(\text{CN})_6]$  is used to detect the presence of:  
 a) Metallic ion                      b) Ferrous ion                      c) Ferric ion                      d) None of these
733. *p*-nitro benzaldehyde reacts with concentrated NaOH solution at room temperature to give:  
 a) *p*-nitrobenzamide  
 b) *p*-nitro benzyl alcohol and sod. *p*-nitrobenzoate  
 c) Benzaldehyde  
 d) *p*-nitrotoluene
734. But-1-ene and cyclobutane exhibit  
 a) Ring chain isomerism  
 b) Position isomerism  
 c) Tautomerism  
 d) Functional isomerism
735. The groups satisfying the secondary valencies of a cation in a complex are called:  
 a) Ligands                      b) Radicals                      c) Primary valencies                      d) None of these
736. Benzene was discovered by:  
 a) Cavendish                      b) Faraday                      c) Berzelius                      d) Wöhler
737. The number of structural and configurational isomers of a bromo compound  $\text{C}_5\text{H}_9\text{Br}$  obtained by the addition of HBr on 2-pentyne respectively are  
 a) 1, 2                      b) 2, 4                      c) 4, 2                      d) 2, 1
738. The primary valency of Fe in  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is:  
 a) 3                      b) 2                      c) 1                      d) Zero
739. Which complex compound obeys 18-electron rule?  
 a)  $[\text{V}(\text{CO})_5]$                       b)  $[\text{Fe}(\text{NH}_3)_6]^{2+}$                       c)  $[\text{Ni}(\text{CO})_6]$                       d)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$
740. Two isomers *X* and *Y* with the formula  $\text{Cr}(\text{H}_2\text{O})_5\text{ClBr}_2$  were taken for experiment on depression in freezing point. It was found that one mole of *X* gave depression corresponding to 2 moles of particles and one mole of *Y* gave depression due to 3 moles of particles. The structural formula of *X* and *Y* respectively, are  
 a)  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Br}_2$ ;  $[\text{Cr}(\text{H}_2\text{O})_4\text{Br}_2]\text{Cl} \cdot \text{H}_2\text{O}$                       b)  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Br}_2$ ;  $[\text{Cr}(\text{H}_2\text{O})_3\text{ClBr}_2] \cdot 2\text{H}_2\text{O}$   
 c)  $[\text{Cr}(\text{H}_2\text{O})_5\text{Br}]\text{BrCl}$ ;  $[\text{Cr}(\text{H}_2\text{O})_4\text{ClBr}]\text{Br} \cdot \text{H}_2\text{O}$                       d)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Br}_2]\text{ClH}_2\text{O}$ ;  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Br}_2$
741. The IUPAC name of  

$$\text{OHC}-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\text{COOH}$$
 is  
 a) 1-formyl-3-oxo-pentanoic acid                      b) 5-formyl-3-oxo pentanoic acid  
 c) 3-oxo-5-formyl pentanoic acid                      d) 3-oxo-1-formyl pentanoic acid
742. The two complexes given below are:



and



- a) Geometrical isomers    b) Position isomers    c) Optical isomers    d) Identical

743. Which of the following statements is not correct?

- a) In oxyhaemoglobin  $\text{Fe}^{2+}$  is paramagnetic  
 b) During respiration the size of  $\text{Fe}^{2+}$  increases when it changes from diamagnetic to paramagnetic state  
 c) Four haeme groups are present in haemoglobin  
 d) Haeme is the prosthetic group and it is non-protein part

744. Chlorination of toluene in the presence of light and heat followed by treatment with aqueous NaOH gives:

- a) *o*-cresol    b) *p*-cresol    c) 2,4-dihydroxytoluene    d) Benzoic acid


745. Which of the following has maximum probability of showing tautomerism?



746. The halide which undergoes nucleophilic substitution most readily is:

- a) *p*- $\text{H}_3\text{CC}_6\text{H}_4\text{Cl}$     b) *o*- $\text{H}_3\text{COC}_6\text{H}_4\text{Cl}$     c) *p*- $\text{ClC}_6\text{H}_4\text{Cl}$     d) *p*- $\text{O}_2\text{NC}_6\text{H}_4\text{Cl}$

747. The major product (70% to 80%) of the reaction between *m*-dinitrobenzene with  $(\text{NH}_4)_2\text{S}_x$  is:



748. The '*E*'-isomer is



749. The Baeyer angle strain is minimum in

- a) Cyclopropane    b) Cyclobutane    c) Cyclopentane    d) Cyclohexane


750. Among the following ions, which one has the highest unpaired electrons?

- a)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$     b)  $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$     c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$     d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$

751. Which will give a white precipitate with  $\text{AgNO}_3$  in aqueous solution?

- a)  $[\text{Co}(\text{NH}_3)_5\text{Cl}](\text{NO}_2)_2$     b)  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$     c)  $[\text{Pt}(\text{en})\text{Cl}_2]$     d)  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$

752. The organic product formed in the reaction;



- a)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$  and  $\text{CH}_3\text{OH}$   
 b)  $\text{C}_6\text{H}_5\text{COOH}$  and  $\text{CH}_4$   
 c)  $\text{C}_6\text{H}_5\text{CH}_3$  and  $\text{CH}_3\text{OH}$   
 d)  $\text{C}_6\text{H}_5\text{CH}_3$  and  $\text{CH}_4$


753. Complexes with bidentate ligands are called:

- a) Ligands    b) Chelates    c) Complexes    d) None of these

754. Excited state configuration of  $\text{Mn}^{2+}$  is

- a)  $t_{2g}^4$     b)  $t_{2g}^3 e_g^2$     c)  $t_{2g}^4 e_g^2$     d)  $t_{2g}^5 e_g^0$

755. The IUPAC name of






- $$\text{C}_6\text{H}_5\text{NH}_2 \xrightarrow[\text{HCl}]{\text{NaNO}_2} \text{A} \xrightarrow{\text{CuCN}} \text{B} \xrightarrow[\text{Ni}]{\text{H}_2} \text{C} \xrightarrow{\text{HNO}_2} \text{D}$$

a)  $\text{C}_6\text{H}_5\text{NHCH}_2\text{CH}_3$       b)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$       c)  $\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$       d)  $\text{C}_6\text{H}_5\text{NHOH}$

- PHONE NO: 8583042324 Page | 62



769. Tautomerism is not exhibited by :

- a)  $C_6H_5 - CH = CH - OH$     b)     c)     d) 

770. Benzaldehyde reacts with  $NH_3$  to give:

- a) Aniline    b) Benzamide    c) Phenylcyanide    d) Hydrobenzamide

771. In coal-tar fraction of heavy oil, the aromatic compound present is:

- a) Cresol  
b) Pyridine  
c) Benzene  
d) Anthracene

772. Optical isomerism is shown by octahedral complexes

- a) Having all monodentate ligands    b) Having all the three bidentate ligands  
c) Having two *trans* bidentate ligands    d) Having two *trans* monodentate ligands

773. Which can be hydrolysed most easily?

- a)  $(C_6H_5)_3CCl$     b)  $C_6H_5CH_2Cl$     c)  $(C_6H_5)_2CHCl$     d)  $C_6H_5Cl$


774. The most stable configuration of *n* butane will be

- a) Skew boat    b) Eclipsed    c) Gauche    d) Staggered-anti

775. Anhydrous aluminium chloride is used in Friedel-Craft's reaction because it is:

- a) Electron rich  
b) Soluble in ether  
c) Ionizable to chloride and aluminium ions  
d) Electron deficient molecule

776. The two isomers given below are



- a) Enantiomers    b) Diastereomers    c) Measomers    d) Position isomers

777. Which of the following has lowest boiling point?

- a) Phenol    b) *o*-nitrophenol    c) *m*-nitrophenol    d) *p*-nitrophenol

778. The IUPAC name of  $[Ni(NH_3)_4][NiCl_4]$  is


- a) Tetrachloro nickel (II) – tetraammine nickel (II)  
b) Tetraammine nickel (II) –tetrachloro nickel(II)  
c) Tetraammine nickel (II) –tetrachloro nickelate(II)  
d) Tetrachloro nickel (II) –tetraammine nickelate(0)

779. All ligands are:


- a) Lewis acid    b) Lewis base    c) Neutral    d) None of these

780. Aspirin is known as:

- a) Phenyl salicylate      b) Acetyl salicylate      c) Methyl salicylic acid      d) Acetyl salicylic acid
781. Which of the following has on optical isomer?  
 a)  $[\text{Co}(\text{NH}_3)_3\text{Cl}]^+$       b)  $[\text{Co}(\text{en})(\text{NH}_3)_2]^{2+}$       c)  $[\text{Co}(\text{H}_2\text{O})_4(\text{en})]^{3+}$       d)  $[\text{Co}(\text{en})_2(\text{NH}_3)_2]^{3+}$
782. The IUPAC name of the compound



- a) 1,1,1-trichloro-2,2-diphenyl ethane      b) 2,4,5-trichloro hexanol  
 c) 2,2,2-trichloro bicyclo [4.4.0] nenone      d) 2,2,2-trichloro-1,1-diphenyl ethane
783. The property by virtue of which a compound can rotate the plane of polarised light is known as  
 a) Polarisability      b) Phosphorescence      c) Optical activity      d) Polarization
784. The molecules represented by the following two structures are



- a) Epimers      b) Diastereomers      c) Enantiomers      d) Identical
785. The IUPAC name of the coordination compound  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is  
 a) Tripotassium hexacyanoiron (II)      b) Potassium hexacyanoiron(II)  
 c) Potassium hexacyanoferrate (III)      d) Potassium hexacyanoferrate (II)
786. Which one of the following is an inner orbital complex as well as diamagnetic in nature?  
 a)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       c)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$       d)  $[\text{Zn}(\text{NH}_3)_6]^{2+}$
787. How many unpaired electrons are present in the central metal ion of  $[\text{CoCl}_4]^{2-}$ ?  
 a) 3      b) 4      c) 5      d) 2
788. Show the coordination number of the metal ion, its oxidation number, the number of electrons in  $d$ -orbitals and the number of unpaired electrons  $d$ -orbitals respectively in complex  $[\text{Co}(\text{H}_2\text{O})_4\text{SO}_3]\text{Cl}$ .  
 a) 6, 3, 6, 4      b) 6, 3, 6, 0      c) 5, 3, 6, 4      d) 5, 3, 6, 0
789. Benzene reacts with.....to give acetophenone.  
 a) Acetyl chloride  
 b) Acetyl chloride in presence of anhy.  $\text{AlCl}_3$   
 c) Anhy.  $\text{AlCl}_3$   
 d) None of the above
790. Which group would you introduce into a drug or a dye to make it water soluble?  
 a)  $-\text{NO}_2$       b)  $-\text{Cl}$       c)  $-\text{SO}_3\text{H}$       d)  $-\text{OH}$
791. In the coordination compound,  $\text{K}_4[\text{Ni}(\text{CN})_4]$ , oxidation state of nickel is  
 a) -1      b) +1      c) 0      d) +2
792. The IUPAC name of  $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{NO}_3$  is:  
 a) Tetraaminodichlorochromium (I) nitrate  
 b) Tetraaminodichlorochromium (III) nitrate  
 c) Dichlorotetraamminechromium (III) nitrate  
 d) Tetraaminodichlorochromium (II) nitrate
793. Vanillin, used as a flavouring agent is:  
 a) An aliphatic alcohol      b) An aromatic aldehyde      c) A hydrocarbon      d) A carbohydrate
794. Which of the following will exhibit optical isomerism?  
 a)  $[\text{Cr}(\text{en})(\text{H}_2\text{O})_4]^{3+}$       b)  $[\text{Cr}(\text{en})_3]^{3+}$       c)  $\text{trans}-[\text{Cr}(\text{en})_2\text{Cl}_2]^+$       d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
795. Which one is a mixed ketone?  
 a) Benzophenone      b) Benzenone      c) Acetophenone      d) Dibenzyl ketone

796. Transition metals can form complexes in:  
 a) Zero oxidation state    b) Cation form    c) Anion form    d) All of these
797. Toluene on oxidation with air in presence of  $V_2O_5$  yields:  
 a) Phenol    b) Benzoic acid    c) Benzaldehyde    d) Benzyl alcohol
798.  $[Pt((NH_3)_4)Cl_2]$  is  
 a) Pyramidal    b) Pentagonal    c) Tetrahedral    d) Square planar
799. In  $Fe(CO)_5$ , the  $Fe-C$  bond possess:  
 a)  $\pi$ -character only  
 b) Both  $\sigma$  and  $\pi$ -characters  
 c) Ionic character  
 d)  $\sigma$ -character only
800. Which molecule has tetrahedral geometry?  
 a)  $[Co(NH_3)_6]^{3+}$     b)  $[Ni(CN)_4]^{2-}$     c)  $Fe(CO)_5$     d)  $[NiCl_4]^{2-}$
801.  $[Co(NH_3)_5Br]SO_4$  and  $[Co(NH_3)_5SO_4]Br$  are the examples of:  
 a) Linkage isomerism  
 b) Geometrical isomerism  
 c) Ionization isomerism  
 d) Optical isomerism

802. The compounds  $R-NO_2$  and  $R-ONO$  are

- a) Geometrical isomers    b) Functional isomers  
 c) Metamers    d) Optical isomers


803. Which of the following Fischer projection formula is same as D-glyceraldehyde?



804.  $[Fe(NO_2)_3Cl_3]$  and  $[Fe(O-NO)_3Cl_3]$  shows

- a) Linkage isomerism    b) Geometrical isomerism  
 c) Optical isomerism    d) None of the above

805.



The IUPAC name of the compound is

- a) 2-ethenyl-3-methyl cyclohexa-1, 3-diene    b) 2, 5-dimethyl hepta-2, 6-dienoic acid  
 c) 2, 6-dimethyl hepta-2, 5dienoic acid    d) 2, 3-dimethyl epoxyethane

806. When benzene sulphonic acid and *p*-nitrophenol are treated with  $NaHCO_3$ , the gases released respectively are:

- a)  $SO_2, NO_2$     b)  $SO_2, NO$     c)  $SO_2, CO_2$     d)  $NO_2, CO_2$

807. Which of the following is non-ionizable?


- a)  $[Co(NH_3)_3Cl_3]$     b)  $[Co(NH_3)_4Cl_2]Cl$     c)  $[Co(NH_3)_5Cl]Cl_2$     d)  $[Co(NH_3)_6]Cl_2$

808. Increasing order of expected keto content






- a)  $CH_3COC_2H_5 > CH_3CHO > CH_3COCH_3 > CH_3COCH_2COCH_3$   
 b)  $CH_3COCH_3 > CH_3CHO > CH_3COC_2H_5 > CH_3COCH_2COCH_3$   
 c)  $CH_3CHO > CH_3COC_2H_5 > CH_3COCH_3 > CH_3COCH_2COCH_3$   
 d)  $CH_3COCH_2COCH_3 > CH_3CHO > CH_3COCH_3 > CH_3COC_2H_5$

809. Which is colourless complex?

- a)  $Cu_2(CH_3COO)_4 \cdot H_2O$   
 b)  $Cu_2Cl_2$   
 c)  $CuSO_4 \cdot 5H_2O$



810. Which is not a reasonable structure for dimethyl benzene?



The IUPAC name of the compound is

a) Propionic anhydride

b) Dipropanoic anhydride

c) Ethoxy propanoic acid

d) Propanoic anhydride

812. A mixture of benzene and aniline can be separated by:

a) Alcohol

b) Dil. HCl

c) Dil. NaOH

d) Hot water


813. The correct IUPAC name of the complex  $\text{Fe}(\text{C}_5\text{H}_5)_2$  is

a) Cyclopentadienyl iron (II)

b) Bis (Cyclopentadienyl)iron (II)

c) Dicyclo pentadienyl ferrate (II)

d) Ferrocane



The electrophile involved in the above reaction is:

a) dichloromethyl cation ( $\text{CHCl}_2^+$ )

b) Dichlorocarbene ( $:\text{CCl}_2$ )

c) Trichloromethyl anion ( $\text{CCl}_3^-$ )

d) Formyl cation ( $\text{CHO}^+$ )

815. Benzoyl Chloride is prepared from benzoic acid by:

a)  $\text{Cl}_2, h\nu$

b)  $\text{SO}_2\text{Cl}_2$

c)  $\text{SOCl}_2$

d)  $\text{Cl}_2, \text{H}_2\text{O}$

816. Which of the following ions forms most stable complex compound?

a)  $\text{Fe}^{3+}$

b)  $\text{Mn}^{2+}$

c)  $\text{Ni}^{2+}$

d)  $\text{Cu}^{2+}$

817. Which one of the following cyano complexes would exhibit the lowest value of paramagnetic behaviour?

(Atomic no. Cr=24, Mn=25, Fe=26, Co=27)

a)  $[\text{Co}(\text{CN})_6]^{3-}$

b)  $[\text{Fe}(\text{CN})_6]^{3-}$

c)  $[\text{Mn}(\text{CN})_6]^{3-}$

d)  $[\text{Cr}(\text{CN})_6]^{3-}$

818. Which of the following statements is not correct?

a) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in the state of hybridisation of nickel.

b) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in the magnetic properties.

c) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in geometry.

d) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in primary valencies of nickel.

819. In the complexes  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$ ,  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$  and  $[\text{FeCl}_6]^{3-}$ , more stability is shown by:


a)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$

b)  $[\text{Fe}(\text{CN})_6]^{3-}$

c)  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$

d)  $[\text{FeCl}_6]^{3-}$

820. In the reaction,



the intermediate 'X' is:

a) Phthalic anhydride

b) Phthalic acid

c) o-xylene

d) Benzoic acid

821. Which of the following is  $\pi$  complex?

a) Trimethyl aluminium

b) Ferrocene

c) Diethyl zinc

d) Nickel carbonyl

822. When phenol is reacted with chloroform and an alkali like NaOH, the compound formed is salicylaldehyde. If we use pyrene in place of chloroform the product obtained is:  
 a) Salicylaldehyde      b) Phenolphthalein      c) Salicylic acid      d) Cyclohexanol
823. Among the properties (a) reducing (b) oxidizing (c) complexing, the set of properties shown by  $\text{CN}^-$  ion towards metal species is  
 a) B, c      b) A, b, c      c) C, a      d) A, b
824. Which of the following is most powerful *meta* directing group?  
 a)  $-\text{NO}_2$       b)  $-\text{SO}_3\text{H}$       c)  $-\text{CHO}$       d)  $-\text{COOH}$
825. Which among the following compounds will show metamerism?  
 a)  $\text{CH}_3\text{COC}_3\text{H}_7$       b)  $\text{CH}_3\text{OC}_2\text{H}_5$       c)  $\text{CH}_3\text{SC}_2\text{H}_5$       d)  $\text{CH}_3\text{OCH}_3$
826. The hybridization of  $[\text{PtCl}_6]^{2-}$  ion is:  
 a)  $d^2sp^3$       b)  $sp^2d^3$       c)  $sp^3d$       d)  $sp^3d^2$
827. The correct name of  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2][\text{PtCl}_4]$  is  
 a) Tetrachloro platinum (II) dichloro tetrammine platinate  
 b) Dichloro tetrammine platinum (IV) tetrachloro platinate(II)  
 c) Tetrammine dichloro platinum (IV) tetrachloro platinate (II)  
 d) Tetrachloro platinum (II) tetrammine platinate (IV)
828. The oxidation state of iron in  $\text{K}_4[\text{Fe}(\text{CN})_6]$  is  
 a) 1      b) 4      c) 3      d) 2
829. Formation of complex compound can be detected by:  
 a) Change in colour      b) Change in solubility      c) Change in pH      d) All are correct
830. The complex that violates the EAN:  
 a) Potassium ferrocyanide  
 b) Potassium ferricyanide  
 c) Nickel carbonyl  
 d) Cobalt(III) hexaammine chloride
831. Chlorobenzene on heating with aqueous  $\text{NH}_3$  under pressure in presence of  $\text{Cu}_2\text{Cl}_2$  gives:  
 a) Aniline      b) Benzamide      c) *o*-dichlorobenzene      d) Chloroaminobenzene
832. The complex,  $[\text{Pt}(\text{Py})(\text{NH}_3)\text{BrCl}]$  will have how many geometrical isomers?  
 a) 2  
 b) 3  
 c) 4  
 d) 0
833. Which one doesn't have  $\pi$  -bond?  
 a) Grignard reagent      b) Dibenzene chromium  
 c) Zeise's salt      d) Ferrocene
834. The IUPAC name of the compound  $\text{CH}_2 - \text{CH} - \text{COOH}$  is  

$$\begin{array}{cc} | & | \\ \text{NH}_2 & \text{OH} \end{array}$$
  
 a) 1-hydroxy-2-aminopropanoic acid      b) 2-hydroxy-3-aminopropanoic acid  
 c) 3-amino-2-hydroxypropanoic acid      d) 2-hydroxy-1-aminopropanoic acid
835. EDTA is a.....ligand.  
 a) Monodentate  
 b) Hexadentate  
 c) Bidentate  
 d) Tridentate
836. Thymol, a phenol derivative is mainly used as:  
 a) Germicide  
 b) Insecticide  
 c) Antibiotic

d) Fragrance compound and antiseptic


837. Which of the following complex has zero magnetic moment (spin only)?

- a)  $[\text{Ni}(\text{NH}_3)_6]\text{Cl}_2$       b)  $\text{Na}_3[\text{FeF}_6]$       c)  $[\text{Cr}(\text{H}_2\text{O})_6]\text{SO}_4$       d)  $\text{K}_4[\text{Fe}(\text{CN})_6]$

838. Which compound is zero valent metal complex?

- a)  $[\text{Ni}(\text{CO})_4]$       b)  $\text{K}_3[\text{Fe}(\text{CN})_6]$       c)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$       d)  $[\text{Cu}(\text{NH}_3)_4\text{SO}_4]$

839.

The IUPAC name of  is

- a) Bicyclo [5.5.0] nonane      b) Biphenyl  
c) Cyclopropyl cyclohexane      d) Spiro [3.5] nonane

840. The tetrahedral crystal field splitting is only .....of the octahedral splitting.

- a) 1/9      b) 2/9      c) 4/9      d) 5/9


841. IUPAC name of  $[\text{Co}(\text{ONO})(\text{NH}_3)_5]\text{Cl}_2$  is

- a) Pentammine nitrocobalt (II) chloride      b) Pentammine nitrosocobalt (III) chloride  
c) Pentammine nitritocobalt (III) chloride      d) Pentammine oxo-nitrocobalt (III) chloride

842. Point out the central ion ligand in the complex  $\text{K}_2[\text{Cd}(\text{CN})_4]$  ;

- a)  $\text{Cd}^+$ ,  $\text{CN}^{1-}$       b)  $\text{Cd}^{2+}$ ,  $\text{CN}^{1-}$       c)  $\text{Cd}^{2+}$ ,  $\text{CN}^{4-}$       d)  $\text{Cd}^{2+}$ ,  $\text{CN}^{2-}$

843. Number of chiral centres in



- a) 1      b) 2      c) 3      d) 4

844. From the equation,  $3\text{C}_2\text{H}_2 \rightarrow \text{C}_6\text{H}_6$ , find the volume of acetylene (NTP) for the manufacture of 3 mole of benzene:

- a) 67.2 litre      b) 134.4 litre      c) 201.6 litre      d) 33.8 litre

845. According to IUPAC nomenclature sodium nitroprusside is named as

- a) Sodium pentacyanonitrosyl ferrate(II)      b) Sodium pentacyanonitrosyl ferrate(III)  
c) Sodium nitroferrocyanide      d) Sodium nitroferrocyanide

846. Among  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  and  $[\text{NiCl}_4]^{2-}$  species the hybridisation states of Ni atom are respectively:

- a)  $sp^3$ ,  $dsp^2$ ,  $dsp^2$       b)  $sp^3$ ,  $dsp^2$ ,  $sp^3$       c)  $sp^3$ ,  $sp^3$ ,  $dsp^2$       d)  $dsp^2$ ,  $sp^3$ ,  $sp^3$

847. The chemical name of DDT is:

- a) Dichloro dinitro toluene  
b) Dichloro dimethyl toluene  
c) *p, p'*-dichloro diphenyl trichloroethane  
d) None of the above

848. The stability of complexes of  $\text{Cu}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Co}^{2+}$  and  $\text{Fe}^{2+}$  varies in the order

- a)  $\text{Cu}^{2+} > \text{Ni}^{2+} > \text{Co}^{2+} > \text{Fe}^{2+}$       b)  $\text{Cu}^{2+} > \text{Fe}^{2+} > \text{Ni}^{2+} > \text{Co}^{2+}$   
c)  $\text{Ni}^{2+} > \text{Co}^{2+} > \text{Fe}^{2+} > \text{Cu}^{2+}$       d)  $\text{Cu}^{2+} < \text{Ni}^{2+} < \text{Co}^{2+} < \text{Fe}^{2+}$


849. The number of unpaired electrons in  $\text{Ni}(\text{CO})_4$  is

- a) 0      b) 1      c) 3      d) 4

850. In sodium tetrafluorooxochromate(....),  $\text{Na}_3[\text{Cr}(\text{O})\text{F}_4]$  the left out place should be filled with which of the following roman numerals?

- a) VI      b) III      c) IV      d) None of these


851. The IUPAC name of compound



- a) 3-methoxy-4-cyano methyl butanoate      b) Methyl-4-cyano-3-methoxy butanoate  
c) 4-cyano-3-methoxy methyl butanoate      d) Methyl-3-methoxy-4-cyano butanoate


852. Cumene is:

- a) *o*-methyl phenol      b) *p*-cresol      c) Isopropyl benzene      d) Phenyl *n*-propane
853. In Etard's reaction toluene is oxidised to benzaldehyde using:  
 a)  $\text{H}_2\text{O}_2$   
 b)  $\text{Cl}_2$   
 c) Chromium trioxide or  $\text{CrO}_2\text{Cl}_2$   
 d)  $\text{KMnO}_4$
854. Which of the following will exhibit geometrical isomerism?  
 a) Propene      b) Butene-2  
 c) Butene-1      d) 1, 1-dichloro butane
855. Ferrocene is:  
 a)  $\text{Fe}(\eta^5 - \text{C}_5\text{H}_5)_2$       b)  $\text{Fe}(\eta^2 - \text{C}_5\text{H}_5)_2$       c)  $\text{Cr}(\eta^5 - \text{C}_5\text{H}_5)_5$       d)  $\text{Os}(\eta^5 - \text{C}_5\text{H}_5)_2$
856. Which one is an outer orbital complex?  
 a)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$       b)  $[\text{Mn}(\text{CN})_6]^{4-}$       c)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       d)  $[\text{Fe}(\text{CN})_6]^{4-}$
857. The pair of  $[\text{Co}(\text{SO}_4)(\text{NH}_3)_5]\text{Cl}$  and  $[\text{CoCl}(\text{NH}_3)_5]\text{SO}_4$  constitutes  
 a) Optical isomers      b) Linkage isomers      c) Coordination isomers      d) Ionisation isomers
858. The IUPAC name of  $\text{K}_2[\text{Cr}(\text{CN})_2\text{O}_2(\text{O})_2(\text{NH}_3)]$  is  
 a) Potassiumammine dicyanodioxoperoxochromate (VI)  
 b) Potassiumammine cyanoperoxodioxochrometic (IV)  
 c) Potassiumammine dicyanodioxoperoxochromium (IV)  
 d) Potassiumammine dicyanodioxoperoxochromium (IV)
859. In spectrochemical series chlorine is above than water *i.e.*,  $\text{Cl} > \text{H}_2\text{O}$ , this is due to  
 a) Good  $\pi$ -acceptor properties of Cl  
 b) Strong  $\sigma$  -donor and good  $\pi$ -acceptor properties of Cl  
 c) Good  $\pi$  -donor properties of Cl  
 d) Larger size of Cl than  $\text{H}_2\text{O}$
860. The type of isomerism shown by  $[\text{Co}(\text{en})_2(\text{NCS})_2]\text{Cl}$  and  $[\text{Co}(\text{en})_2(\text{NCS})\text{Cl}]\text{NCS}$  is:  
 a) Coordination      b) Ionization      c) Linkage      d) All of these
861. Which ion shows only the coordination number 4 in complexes?  
 a)  $\text{Pt}^{2+}$       b)  $\text{Cr}^{3+}$       c)  $\text{Fe}^{3+}$       d)  $\text{Pt}^{4+}$
862. The spin magnetic moment of cobalt in  $\text{Hg}[\text{Co}(\text{SCN})_4]$  is :  
 a)  $\sqrt{3}$       b)  $\sqrt{8}$       c)  $\sqrt{15}$       d)  $\sqrt{24}$
863. Which of the following is not an isomer of but-1-yne?  
 a) But-2-yne  
 b) Buta-1-3-diene  
 c) Methyl cyclopropene  
 d) But-2-ene
864. How many unpaired electrons are present in the central metal ion of  $[\text{CoCl}_4]^{2-}$ ?  
 a) 2      b) 3      c) 4      d) 5
865. The brown ring complex compound is formulated as  $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]\text{SO}_4$ . The oxidation state of Fe is:  
 a) +1      b) +2      c) +3      d) Zero
866. Correct IUPAC name of



- a) Gammexane      b) Dichloro diphenyl trichloroethane  
 c) Diparachlorophenyl trichloroethane      d) 1,1,1-trichloro-2,2-bis (4-chlorophenyl) ethane
867. IUPAC name of





a) Cumene

c) Phenyl propane

b) 2-phenyl propane

d) 1-(2-propyl) benzene

868. Which of the following gives violet colour with an alcoholic solution of  $\text{FeCl}_3$ ?

a) Benzoic acid

b) Toluene

c) Salicylic acid

d) Nitrobenzene

869. Which of the following is wrong statements?

a)  $\text{Ni}(\text{CO})_4$ , has zero oxidation number for Nib)  $\text{Ni}(\text{CO})_4$ , has oxidation number +4 for Ni


c) Ni is metal

d) CO is gas

870. Which of the following represents a chelating ligand?

a)  $\text{H}_2\text{O}$ b)  $\text{Cl}^-$ c)  $\text{OH}^-$ 

d) DMG

871. The correct order of reactivity of  $\text{PhMgBr}$  with;a)  $\text{I} > \text{II} > \text{III}$ b)  $\text{III} > \text{I} > \text{II}$ c)  $\text{II} > \text{III} > \text{I}$ d)  $\text{II} > \text{I} > \text{III}$ 

872. Which of the following will give maximum number of isomers?

a)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]$ b)  $[\text{Ni}(\text{en})(\text{NH}_3)_4]^{2+}$ c)  $[\text{Ni}(\text{C}_2\text{O}_4)(\text{en})_2]$ d)  $[\text{Cr}(\text{SCN})_2(\text{NH}_3)_4]^+$ 873.  $\text{CuCl}$  reacts with  $\text{KCN}$  solution forming a complex. Coordination number of copper in the complex is:

a) 2

b) 3

c) 4

d) 6

874. The terms stereoisomers, enantiomers and diastereomers will refer

a) Only to configurational isomers including geometric isomers

b) Only to configurational isomers

c) To both configurational as well as conformational isomers

d) To neither configuration nor conformational isomers

875. Aniline was acetylated. The product on nitration followed by alkaline hydrolysis gave:

a) *o*-nitroacetanilideb) *o*- and *p*-nitroanilinec) *m*-nitroaniline

d) Acetanilide

876. The IUPAC name of the compound  $[\text{CuCl}_2(\text{CH}_3\text{NH}_2)_2]$  is





a) Dichloro bis (dimethyl amine) copper(II)

b) Dichloro bis (methyl amine) copper(II)

c) Dimethyl amine copper (II) chloride

d) Bis (dimethyl amine ) copper (II) chloride

877. Which is the structure of compound 2-(1-cyclobutenyl)-1-hexene?



878. On explosion TNT gives:

a)  $\text{CO} + \text{N}_2 + \text{H}_2 + \text{CH}_4 + \text{CO}_2$ b)  $\text{CO} + \text{N}_2 + \text{H}_2$ c)  $\text{CO}_2 + \text{N}_2 + \text{H}_2\text{O}$





- b) Electrophilic substitution
- c) Electrophilic addition
- d) Nucleophilic addition


895. Reimer-Tiemann reaction involves a:

- a) Carbonium ion intermediate
- b) Carbene intermediate
- c) Carbanion intermediate
- d) Free radical intermediate

896. Which does not have a carboxyl group?

- a) Picric acid
- b) Ethanoic acid
- c) Aspirin
- d) Benzoic acid

897. In Cannizzaro's reaction given below:



the slowest step is:

- a) The transfer of hydride to the carbonyl group
- b) The abstraction of proton from the carboxylic group
- c) The deprotonation of  $\text{PhCH}_2\text{OH}$
- d) The attack of  $:\ddot{\text{O}}\text{H}^-$  at the carboxyl group

898. The oxidation state of Ag in Tollens' reagent is:

- a) Zero
- b) +1
- c) +2
- d) +1.5


899. Hybridization of Fe in  $[\text{K}_3\text{Fe}(\text{CN})_6]$  is

- a)  $sp^3$
- b)  $d^2sp^3$
- c)  $sp^3d^2$
- d)  $dsp^3$

900. Which of the following is not isomeric with diethyl ether?

- a) Methyl *n*-propyl ether
- b) Butan-1-ol
- c) 2-methyl propan-2-ol
- d) Butan-2-one

901. In the given conformation  $\text{C}_2$  is rotated about  $\text{C}_2 - \text{C}_3$  bond anticlockwise by an angle of  $120^\circ$  then the conformation obtained is



- a) Fully eclipsed conformation
- b) Partially eclipsed conformation
- c) Gauche conformation
- d) Staggered conformation


902. Crystal field stabilization energy for high spin  $d^4$  octahedral complex is:

- a)  $-1.8 \Delta_0$
- b)  $-1.6 \Delta_0 + P$
- c)  $-1.2 \Delta_0$
- d)  $-0.6 \Delta_0$

903. Which kind of isomerism is exhibited by octahedral  $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}$ ?

- a) Geometrical and ionisation
- b) Geometrical and optical
- c) Optical and ionisation
- d) Geometrical only

904. The IUPAC name of the following compound is



- a) 5-cyclopropyl pent-2-en-1-oic acid
- b) 6-cyclopropyl pent-2-en-1-oic acid
- c) 5-cyclopropyl pent-1-en carboxylic acid
- d) 6-cyclopropyl pent-1-en carboxylic acid

905. Which of the following compounds will show a negative test with phenyl hydrazine?

- a) Glucose
- b) Ethyl alcohol
- c) A cetaldehyde
- d) Benzophenone

906. Friedel-Craft's reaction is not possible in:  
 a)  $C_6H_5OH$                       b)  $C_6H_5C_2H_5$                       c)  $C_6H_5NO_2$                       d)  $C_6H_5CH_3$
907. The geometry of  $Ni(CO)_4$  and  $Ni(PPh_3)_2Cl_2$  are  
 a) Both square planar                      b) Tetrahedral and square planar respectively  
 c) Both tetrahedral                      d) Square planar and tetrahedral respectively
908. The number of isomers possible for square planar complex  $K_2[PdClBr_2SCN]$  is:  
 a) 2                      b) 3                      c) 4                      d) 6
909. The correct order for the wavelength of absorption in the visible region is  
 a)  $[Ni(NO_2)_6]^{4-} < [Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+}$                       b)  $[Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+} < [Ni(NO_2)_6]^{4-}$   
 c)  $[Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+} < [Ni(NO_2)_6]^{4-}$                       d)  $[Ni(NO_2)_6]^{4-} < [Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+}$
910. The IUPAC name of  $CCl_3CH_2CHO$  is  
 a) Chloral                      b) 1,1,1-trichloropropanol  
 c) 2,2,2-trichloropropanol                      d) 3,3,3-trichloropropanol
911. The coordination number of Cu in  $[Cu(H_2O)_4]^{2+}$  complex is  
 a) 2                      b) 1                      c) 3                      d) 4
912. Among the following, the correct statement is  
 a) Prefixes are written before the name of compound  
 b) Suffixes are written after the name of compound  
 c) The IUPAC name is always written as a single word  
 d) All of the above
913. In which of the following *p*-electrons of the halogens are not involved in delocalisation?  
 a) Chlorobenzene                      b) Bromobenzene                      c) Allyl chloride                      d) Vinyl chloride
914. Which of the following does not have optical isomer?  
 a)  $[Co(en)(NH_3)_2Cl_2]Cl$                       b)  $[Co(en)_2Cl_2]Cl$                       c)  $[Co(NH_3)_3Cl]$                       d)  $[Co(en)_3]Cl_3$
915. Ethylene diamine is an example of  
 a) Monodentate ligand                      b) Bidentate ligand                      c) Tridentate ligand                      d) Polydentate ligand
916. In chlorobenzene, the  $-Cl$  group:  
 a) Activates the benzene ring more via resonance effect than deactivating it via inductive effect  
 b) Deactivates the benzene ring more via inductive effect than activating it via resonance effect  
 c) Activates the benzene ring via resonance effect and deactivates it via inductive effect. Both these effects are more evenly matched  
 d) None of the above
917. The *R*-isomer among the following are
- $$\begin{array}{c} \text{CHO} \\ | \\ \text{H} - \text{C} - \text{OH} \\ | \\ \text{CH}_2\text{OH} \end{array}$$

(i)

$$\begin{array}{c} \text{H} \\ | \\ \text{D} - \text{C} - \text{OH} \\ | \\ \text{CH}_3 \end{array}$$

(ii)
- $$\begin{array}{c} \text{CH}_3 \\ | \\ \text{H} - \text{C} - \text{OH} \\ | \\ \text{CH}_2\text{CH}_3 \end{array}$$

(iii)

$$\begin{array}{c} \text{COOH} \\ | \\ \text{H}_3\text{C} - \text{C} - \text{NH}_2 \\ | \\ \text{H} \end{array}$$

(iv)
- a) (i) and (ii)                      b) (ii) and (iii)                      c) (iii) and (iv)                      d) (i) and (iii)
918. Which possesses tetrahedral shape ( $sp^3$ -hybridization of central atom)?  
 a)  $[Zn(NH_3)_4]^{2+}$                       b)  $[Ni(CO)_4]$                       c)  $[Cd(NH_3)_4]^{2+}$                       d) All are correct
919. The reaction,  

$$C_6H_5CHO + CH_3CHO \xrightarrow{Dil. NaOH} C_6H_5CH=CHCHO$$
 is called:  
 a) Benzoin condensation  
 b) Claisen condensation

- c) Perkin's reaction  
d) Cannizaro's reaction

920. Complexation is shown by:

- a) Ag                                      b) Au                                      c) Cu                                      d) All of these


921. AgO in Ag(II) complex which is:

- a) Diamagnetic                                      b) Paramagnetic                                      c) Ferromagnetic                                      d) Neutral

922. Acylation of benzene to produce aliphatic aromatic ketones is called:

- a) Benzoin condensation  
b) Hydroformylation  
c) Friedel-Crafts reaction  
d) None of these

923. The structure of the major product formed in the given reaction



- a)
- b)
- c)
- d)

924. Chlorobenzene is prepared commercially by:

- a) Grignard reaction                                      b) Raschig process                                      c) Wurtz - Fittig reaction                                      d) Friedel-Crafts reaction

925. An aqueous solution of  $\text{CoCl}_2$  on addition of excess of concentrated  $\text{HCl}$  turns blue to formation of

- a)  $[\text{CoCl}_4]^{2-}$                                       b)  $[\text{Co}(\text{H}_2\text{O})_2\text{Cl}_4]^{2-}$                                       c)  $[\text{Co}(\text{H}_2\text{O})_2\text{Cl}_4]^{2-}$                                       d)  $[\text{Co}(\text{H}_2\text{O})_4\text{Cl}_2]$


926. Which one of the following will not show geometrical isomerism?

- a)  $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$                                       b)  $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$                                       c)  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$                                       d)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

927. When ethyl benzoate is hydrolysed with aqueous alkali, the products present in the medium are:


- a)  $\text{C}_6\text{H}_5\text{COOH}$ ,  $\text{C}_2\text{H}_5\text{O}^-$                                       b)  $\text{C}_6\text{H}_5\text{COO}^-$ ,  $\text{C}_6\text{H}_5\text{OH}$                                       c)  $\text{C}_2\text{H}_5\text{OH}$ ,  $\text{C}_6\text{H}_5\text{COOH}$                                       d)  $\text{C}_6\text{H}_5\text{COO}^-$ ,  $\text{C}_2\text{H}_5\text{O}^-$

928. The IUPAC name of



- a) 2-carbamoyl hexanal  
c) 6-keto-2-methylhexanamide
929. Which of the following is more basic than aniline?  
a) *p*-Nitroaniline      b) Benzylamine      c) Diphenylamine      d) Triphenylamine
930. Name of some compounds are given below. Which one is not in IUPAC system?
- a)  $\text{CH}_3 - \text{CH} - \text{CH} - \text{CH}_3$   
          |        |  
          OH     CH<sub>3</sub>  
          4-methyl-2-butanol
- b)  $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}(\text{CH}_3)_2$   
          4 methyl-2-pentyne
- c)  $\text{CH}_3\text{CH}_2 - \text{C} - \text{CH} - \text{CH}_3$   
                  ||     |  
                  CH<sub>2</sub> CH<sub>3</sub>  
          2 - ethyl-3- methyl - but -1- ene
- d)  $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH} - \text{CH} - \text{CH}_2\text{CH}_3$   
                                  |        |  
                                  CH<sub>2</sub>CH<sub>3</sub> CH<sub>3</sub>  
                                  3-methyl-4-ethyl heptane
931. For which transition metal ions are low spin complexes possible?  
a) Rh<sup>3+</sup>      b) Mn<sup>3+</sup>      c) Ru<sup>2+</sup>      d) All are correct
932. Which one is monodentate ligand?  
a) F<sup>-</sup>      b) NO<sub>2</sub><sup>-</sup>      c) H<sub>2</sub>O      d) All are correct
933. Cyclic hydrocarbon molecule A has all the carbons and hydrogens in a single plane. All the carbon-carbon bonds are of same length and less than 1.54 Å and more than 1.34 Å. The C—C—C bond angle will be:  
a) 120°      b) 180°      c) 100°      d) 109°28'
934. Chlorine reacts with benzaldehyde to give:  
a) Benzyl chloride      b) Benzal chloride      c) Benzoyl chloride      d) Chlorobenzene
935. Phenol is:  
a) A base weaker than NH<sub>3</sub>  
b) An acid stronger than carbonic acid  
c) An acid weaker than carbonic acid  
d) Neutral
936. Which one is example of octahedral complex?  
a) Cu(NH<sub>3</sub>)<sub>4</sub><sup>2+</sup>      b) FeF<sub>6</sub><sup>3-</sup>      c) Zn(NH<sub>3</sub>)<sub>4</sub><sup>2+</sup>      d) Ni(CN)<sub>4</sub><sup>2-</sup>
937. Which one of the following statement is correct?  
a) Ferric ions give a deep green precipitate on adding potassium ferrocyanide solution.  
b) On boiling a solution having K<sup>+</sup>, Ca<sup>2+</sup> and HCO<sub>3</sub><sup>-</sup> ions, we get a precipitate of K<sub>2</sub>Ca(CO<sub>3</sub>)<sub>2</sub>  
c) Manganese salt give a violet vortex test in reducing flame  
d) From a mixed precipitate of AgCl and AgI, ammonia solution dissolves only AgCl
938. Which of the following fractions obtained in fractional distillation of coal-tar contains benzene and toluene?  
a) Light oil  
b) Heavy oil  
c) Middle oil  
d) Green oil
939. The tetrahedral complexes have coordination number  
a) 3      b) 6      c) 4      d) 8
940. The C—C bond length in benzene is .....than C—C bond length in alkenes.  
a) Less      b) More      c) Equal      d) None of these
941. Which are generally used for preparing derivative of aldehydes and ketones?  
a) Hydroxylamine hydrochloride  
b) 2,4-dinitrophenylhydrazine  
c) Phenylhydrazinehydrochloride





946. In an octahedral structure, the pair of *d*-orbitals involved in  $d^2 sp^3$  hybridisation is

- a)  $d_{x^2-y^2}, d_{z^2}$       b)  $d_{xz}, d_{x^2-y^2}$       c)  $d_{z^2}, d_{xz}$       d)  $d_{xy}, d_{yz}$

947. In which of the following ions has the metal atom EAN as 36?

- a)  $[\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{Fe}(\text{CN})_6]^{3-}$       c)  $[\text{PbCl}_4]^{2-}$       d)  $[\text{Pd}(\text{CN})_6]^{2-}$

948. The number of ions given by  $\text{K}[\text{Pt}(\text{NH}_3)_5\text{Cl}_5]$  in aqueous solution is:

- a) 2      b) 3      c) 4      d) 1


949.  $\text{CuCl}$  is sparingly soluble in  $\text{H}_2\text{O}$  but it dissolves in  $\text{KCl}$  solution due to the formation of:

- a)  $\text{K}_2(\text{CuCl}_4)$       b)  $\text{K}_3(\text{CuCl}_4)$       c)  $\text{K}(\text{CuCl}_2)$       d) None of these

950. A characteristics group test for phenolic gp. is:


- a) Libermann's nitroso reaction  
b) Coupling with diazonium salt  
c) aq.  $\text{FeCl}_3$   
d) All of the above

951. Write the IUPAC name of the compound



- a) 5, 6-dimethyl bicyclo [2,2,1] heptane      b) 2, 3-dimethyl bicyclo [2,2,1] heptane  
c) 2, 3-dimethyl bicyclo [1,2,2] heptane      d) 3, 4-dimethyl bicyclo [2,1,2] heptane

952. Choose the correct statement from the ones given below for two anilium in:



- a) II is not an acceptable canonical structure because carbonium ions are less stable than ammonium ions  
b) II is not an acceptable canonical structure because it is non-aromatic  
c) II is not an acceptable canonical structure because the nitrogen has 10 valence electrons  
d) II is an acceptable canonical structure


953. Which of the following statements is/are incorrect?

- a) Metamerism belongs to the category of structural isomerism  
b) Tautomeric structures are the resonating structures of a molecule  
c) The violet colouration produce by a molecule with neutral ferric chloride solution indicates the presence of enolic group in the molecule  
d) Geometrical isomerism is not shown by alkenes

954. Gives are (i) cyclohexanol; (ii) acetic acid; (iii) 2, 4, 6-trinitrophenol; and (iv) phenol. In these the order of decreasing acidic character will be:


- a) (iii) > (ii) > (iv) > (i)      b) (ii) > (iii) > (i) > (iv)      c) (ii) > (iii) > (iv) > (i)      d) (iii) > (iv) > (ii) > (i)

955. Phenol and benzoic acid can be distinguished by:  
 a) Aqueous  $\text{NaHCO}_3$       b) Aqueous  $\text{NaNO}_3$       c) Aqueous  $\text{NaOH}$       d) Conc.  $\text{H}_2\text{SO}_4$
956. The functional groups  $-\text{OH}$ ,  $-\text{COOH}$ ,  $-\text{CHO}$ ,  $-\text{OCH}_3$  attached to a chiral carbon is in the preference order  
 a)  $\text{OH} > \text{COOH} > \text{CHO} > \text{OCH}_3$       b)  $\text{OCH}_3 > \text{OH} > \text{CHO} > \text{COOH}$   
 c)  $\text{OCH}_3 > \text{OH} > \text{COOH} > \text{CHO}$       d)  $\text{OCH}_3 > \text{COOH} > \text{CHO} > \text{OH}$
957. The hypothetical complex chloro diaquatrammine cobalt(II) chloride can be represented as:  
 a)  $[\text{CoCl}(\text{NH}_3)_3(\text{H}_2\text{O})_2]\text{Cl}_2$  b)  $[\text{Co}(\text{NH}_3)_3(\text{H}_2\text{O})\text{Cl}_3]$  c)  $[\text{Co}(\text{NH}_3)_3(\text{H}_2\text{O})_2\text{Cl}]$  d)  $[\text{Co}(\text{NH}_3)_3(\text{H}_2\text{O})_3]\text{Cl}_3$
958. Which is expected to be paramagnetic?  
 a)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$       b)  $[\text{Ni}(\text{CO})_4]$       c)  $[\text{Zn}(\text{NH}_3)_4]^{2+}$       d)  $[\text{Co}(\text{NH}_3)_6]^{3+}$
959. The molecular formula of diphenyl methane



How many structural isomers are possible when one of the hydrogen is replaced by a chlorine atom?

- a) 8      b) 7      c) 6      d) 4
960. Among the properties (A) reducing, (B) oxidising (C) complexing, the set of properties shown by  $\text{CN}^-$  ion towards metal species is  
 a) A, B      b) B, C      c) C, A      d) A, B, C



In the double bonds are

- a) *cis, cis*      b) *cis, trans*  
 c) *trans, cis*      d) *trans, trans*
962. The reaction of toluene with  $\text{Cl}_2$  in presence of  $\text{FeCl}_3$  gives 'X' and the reaction in presence of light gives 'Y'. Thus, 'X' and 'Y' are:  
 a) X = benzal chloride; Y = *o*-chlorotoluene  
 b) X = *m*-chlorotoluene; Y = *p*-chlorotoluene  
 c) X = *o*- and *p*-chlorotoluene; Y = trichloro methyl benzene  
 d) X = benzal chloride; Y = *m*-chlorotoluene

963. Among the following four compounds:  
 a) Phenol      b) Methyl phenol      c) *meta*-nitrophenol      d) *para*-nitrophenol
964. Which gives phthalic anhydride on reaction with hot, conc.  $\text{H}_2\text{SO}_4$  in presence of Hg?  
 a) Naphthalene      b) Phenol      c) *p*-xylene      d) *m*-xylene
965. *Cis-trans*-isomerism is found in square planar complexes of the molecular formula: (a and b are monodentate ligands)  
 a)  $\text{Ma}_4$       b)  $\text{Ma}_3\text{b}$       c)  $\text{Ma}_2\text{b}_2$       d)  $\text{Mab}_3$
966. Which ion produces a small crystal field splitting (a weak ligand field)?  
 a)  $\text{I}^-$       b)  $\text{Cl}^-$       c)  $\text{F}^-$       d) All of these


967. Benzene undergoes substitution reaction more easily than addition because:  
 a) It has a cyclic structure  
 b) It has three double bonds  
 c) It has six hydrogen atoms  
 d) Of resonance

968. Isomers have essentially identical  
 a) Structural formula      b) Chemical properties  
 c) Physical properties      d) Molecular formula


969. Which of the following pair is not correctly matched?  
 a) Absorption peak for  $[\text{Cr}^{\text{III}}(\text{NH}_3)_6]^{3+} = 21680 \text{ cm}^{-1}$   
 b) Effective atomic no. of Pt in  $[\text{PtCl}_6]^{2-} = 84$   
 c) Crystal field stabilization energy of  $d^2$  in weak ligand field =  $(-)\text{0.8 } \Delta_0$   
 d) Example of weak ligand field for  $d^5$  configuration =  $[\text{Mn}^{\text{II}}\text{F}_6]^{4-}$



970. Aspirin (or acetyl salicylic acid) is obtained by action of  $\text{CH}_3\text{COCl}$  with:  
 a) Salicylic acid                      b) Phenol                      c) Benzaldehyde                      d) Aniline
971.  $\text{CuCl}$  dissolves in ammonia forming a complex. The coordination number of copper in the complex is:  
 a) 1                      b) 2                      c) 4                      d) 6
972. IUPAC name of the following cycloalkane is




- a) 8-methyl bicyclo [4,3,0] nonane                      b) 1-methyl bicyclo [4,3,0] nonane  
 c) 3-methyl bicyclo [4,3,0] nonane                      d) 4-methyl bicyclo [4,3,0] nonane
973. Schiff's bases are formed when aniline is condensed with:  
 a) Phenols                      b) Aromatic aldehydes                      c) Aryl chlorides                      d) Aliphatic alcohols
974. Which of the following is not an organometallic compound?  
 a) Zeise's salt                      b) TEL                      c) Sodium ethoxide                      d) Ferrocene
975. Molecular formula  $\text{C}_5\text{H}_{12}\text{O}$  will show  
 a) Position  
 b) Optical isomerism  
 c) Functional isomerism  
 d) All of these
976. Both  $\text{Co}^{3+}$  and  $\text{Pt}^{4+}$  have a coordination number of six. Which of the following pairs of complexes will show approximately the same electrical conductance for their 0.001 M aqueous solutions?  
 a)  $\text{CoCl}_2 \cdot 4\text{NH}_3$  and  $\text{PtCl}_4 \cdot 4\text{NH}_3$                       b)  $\text{CoCl}_3 \cdot 3\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$   
 c)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$                       d)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 3\text{NH}_3$
977. Which of the following is not an organometallic compound?  
 a) Sodium ethoxide                      b) Trimethyl aluminium  
 c) Tetraethyl lead                      d) Ethyl magnesium bromide
978. The number of water molecule(s) directly bonded to the metal centre in  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  is  
 a) 1                      b) 2                      c) 3                      d) 4
979. The formula of sodium nitroprusside is:  
 a)  $\text{Na}_4[\text{Fe}(\text{CN})_5\text{NO}_2]$                       b)  $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]$                       c)  $\text{NaFe}[\text{Fe}(\text{CN})_6]$                       d)  $\text{Na}_2[\text{Fe}(\text{CN})_6\text{NO}_2]$
980. The IUPAC name of the compound  
 $\text{H}_2\text{N} - \text{CH} - \text{CH}_2\text{OH}$   
 $\quad \quad |$   
 $\quad \quad \text{COOH}$   
 a) 2-amino-2-carboxy pentanol                      b) 1-amino-2-hydroxy propanoic acid  
 c) 1-hydroxy-2-amino-3-propanoic acid                      d) 2-amino-3-hydroxy propanoic acid
981. Which of the following complex species does not involve inner orbital hybridisation?  
 a)  $[\text{CoF}_6]^{3-}$                       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                       c)  $[\text{Fe}(\text{CN})_6]^{3-}$                       d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
982. The EAN of nickel in  $\text{K}_2[\text{Ni}(\text{CN})_4]$  is:  
 a) 35                      b) 34                      c) 36                      d) 38
983. The type of isomerism shown by, 6,6'-disitrodiphenic acid is  
 a) Conformational                      b) Optical                      c) Geometrical                      d) Functional
984. Which one of the following compounds forms benzoic acid on oxidation?  
 a) Chlorophenol                      b) Benzylchloride                      c) Chlorobenzene                      d) Chlorotoluene
985. Glycinato ligand is:



- b) Bidentate ligand  
 c) Two donor sites N and  $\text{O}^-$



- d) All of the above
986. Which one is the most likely structure of  $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ , if 1/3 of total chlorine of the compound is precipitated by adding  $\text{AgNO}_3$  to its aqueous solution?
- $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$
  - $[\text{Cr}(\text{H}_2\text{O})_3\text{Cl}_3] \cdot (\text{H}_2\text{O})_3$
  - $[\text{CrCl}_2(\text{H}_2\text{O})_4] \cdot \text{Cl} \cdot 2\text{H}_2\text{O}$
  - $[\text{CrCl}(\text{H}_2\text{O})_5]\text{Cl}_2 \cdot \text{H}_2\text{O}$
987. Carbon in benzene undergoes  $sp^2$ -hybridization and the bond angle is  $120^\circ$ . The shape of benzene molecule is:
- Linear
  - Planar
  - Pyramidal
  - Planar hexagonal
988. The example of coordination isomerism is
- $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$  and  $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$
  - $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$
  - $\text{Co}(\text{NH}_3)_5\text{NO}_3]\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{NO}_3$
  - $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Br}_2$  and  $[\text{Pt}(\text{NH}_3)_4\text{Br}_2]\text{Cl}_2$
989. Coordination compounds have great importance in biological systems. In this context which of the following statement is incorrect?
- Haemoglobin is the red pigment of blood and contains iron
  - Cyanocobalamin is  $\text{B}_{12}$  and contains cobalt
  - Chlorophylls are green pigments in plants and contains calcium
  - Carbocypeptidase-A an enzyme and contains zinc
- 990.




The IUPAC name of the given structure






H is

- Diisohexane
  - Isohexane
  - 2, 2-dimethylbutane
  - 2, 3-dimethylbutane
991. Aniline on treating with phosgene gives:
- Phenyl isocyanate
  - A secondary base
  - A neutral substance
  - A tertiary base
992. On boiling with conc. hydrobromic acid, phenylethylether will yield:
- Phenol and ethyl bromide
  - Bromobenzene and ethanol
  - Phenol and ethane
  - Bromobenzene and ethane
993. Ammonia gas does not evolve from the complex  $\text{FeCl}_3 \cdot 4\text{NH}_3$  but it gives white precipitate with aqueous solution of  $\text{AgNO}_3$ . Coordination number of central metal ion in above complex is six. Give IUPAC name of the complex.
- Ammonium trichloro triammine ferrum(III)
  - Tetra ammine ferrum (III) chloride
  - Dichloro tetraammine ferrate (II) chloride
  - Dichloro tetraammine ferrum (III) chloride
994. Nickel ( $Z=28$ ) combines with a uninegative monodentate ligand  $X^-$  to form a paramagnetic complex  $[\text{NiX}_4]^{2-}$ . The number of unpaired electron ( $s$ ) in the nickel and geometry of this complex ion are respectively
- One, tetrahedral
  - Two, tetrahedral
  - One, square planar
  - Two, square planar
995. Amongst the compounds given, the one that would form a brilliant coloured dye on treatment with  $\text{NaNO}_2$  in dil.  $\text{HCl}$  followed by addition to an alkaline solution of  $\beta$ -naphthol is:








996. Identify 'Z' in the reaction given below;



997. A solution containing 2.675 g of  $\text{CoCl}_3 \cdot 6\text{NH}_3$  (molar mass = 267.5 g mol<sup>-1</sup>) is passed through a cation exchanger. The chloride ions obtained in solution were treated with excess of  $\text{AgNO}_3$  to give 4.78 g of  $\text{AgCl}$  (molar mass = 143.5 g mol<sup>-1</sup>). The formula of the complex is (Atomic mass of Ag = 108 u)



998.  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  ion has d-electrons equal to:





a) 2

b) 3

c) 4

d) 5

999. Enol form is more stable in



100 The coordination number of cobalt in  $[\text{Co}(\text{en})_2\text{Br}_2]\text{Cl}_2$  is:

0.

a) 2


b) 4

c) 6

d) 8





100 Which one readily accepts a proton?


1.



100 Identify 'Z' in the reaction;

2.






100 The number of  $\sigma$  and  $\pi$ -bonds in a molecule of benzene is:

3.

- a)  $6\sigma$  and  $9\pi$                       b)  $9\sigma$  and  $3\pi$                       c)  $12\sigma$  and  $3\pi$                       d)  $6\sigma$  and  $6\pi$


100 The phenomenon of optical activity will be shown by:

4.



100

5.



The correct name of the compound

- a) 1,3,4-trimethyldecaline                      b) 1,3,9-trimethyldecaline  
c) 1,8,10-trimethyldecaline                      d) 1,3,10-trimethyldecaline

100 If  $\text{NH}_4\text{OH}$  is added to the  $(\text{PtCl}_4)^{2-}$  ion, the complex formed represents:

6.

- a) Zero dipole                      b) Finite dipole                      c) Infinite dipole                      d) All of these

100 Which one of the following will be able to show *cis-trans*-isomerism?

7.

- a)  $M_{A_3B}$   
b)  $M_{(AA')_2}$   
c)  $M_{A_2BCD}$   
d)  $M_{A_4}$   
( $AA'$  is unsymmetrical bidentate ligand,  $ABCD$  are unidentate ligands.)

100 The coordination number of a metal in coordination compound is

8.

- a) Same as primary valency                      b) Sum of primary and secondary valencies  
c) Same as secondary valency                      d) None of the above

100 The IUPAC name of  $\text{K}_4[\text{Ni}(\text{CN})_4]$  is

9.

- a) Tetrapotassium tetracyanonickelate (II)                      b) Potassium tetracyanonickel (II)  
c) Potassium tetracyanonickelate (0)                      d) Potassium tetracyanonickelate (II)

101 Which of the following compounds shows optical isomerism?

0.

- a)  $[\text{Co}(\text{CN})_6]^{3-}$                       b)  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$                       c)  $[\text{ZnCl}_4]^{2-}$                       d)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$

101  $[\text{C}_6\text{H}_5]_2\text{Pd}(\text{SCN})_2$  and  $[(\text{C}_6\text{H}_5)_2\text{Pd}(\text{NCS})_2]$  are:

1.

- a) Linkage isomers                      b) Coordination isomers                      c) Ionization isomers                      d) Geometrical isomers


101 Mark the correct statement

2.




- Ethane has two conformations of which staggered conformation is more stable than the eclipsed conformation
- Ethane has an infinite number of conformations of which eclipsed conformation is more stable than the staggered conformation
- Ethane has an infinite number of conformation of which staggered conformation has the maximum energy
- Ethane has an infinite number of conformation of which the staggered conformation is possessed by majority of the molecules at room temperature

101

3.



Of the following on heating with alkali followed with acid hydrolysis?



101 Which among the following will be named as dibromidobis (ethylene diamine) chromium (III) bromide?

4.

- $[\text{Cr}(\text{en})_2\text{Br}_2]\text{Br}$
- $[\text{Cr}(\text{en})\text{Br}_4]^-$
- $[\text{Cr}(\text{en})\text{Br}_2]\text{Br}$
- $[\text{Cr}(\text{en})_3]\text{Br}_3$

101 Which one of the following complex is an outer orbital complex?

5. (Atomic no. Mn=25, Fe=24, Co=27, Ni=28)

- $[\text{Fe}(\text{CN})_6]^{4-}$
- $[\text{Mn}(\text{CN})_6]^{4-}$
- $[\text{Co}(\text{NH}_3)_6]^{3+}$
- $[\text{Ni}(\text{NH}_3)_6]^{2+}$

101 Benzene can be directly obtained from:

6.

- $\text{CH}\equiv\text{CH}$
- $\text{CH}_2=\text{CH}_2$  and butadiene
- Chlorobenzene
- All of the above





101 Chlorobenzene on treatment with Raney nickel or Al in presence of alkali gives:

7.

- Benzene
- Chlorophenol
- Phenol
- None of these




101 The compound that undergoes decarboxylation most readily under mild condition is:

8.



101 Which ion is paramagnetic?

9.



102 Which kind of isomerism is exhibited by octahedral  $[\text{Co}(\text{NH}_3)_4\text{Br}_2\text{Cl}]$ ?

0.

a) Geometrical and ionization

b) Geometrical only

c) Geometrical and optical

d) Optical and ionisation

102 Resorcinol and conc.  $\text{H}_2\text{SO}_4$  in presence of phthalic anhydride produce a compound which is:

1.

a) A dye




b) An antiseptic

c) An indicator

d) A detergent

102 Which of the following compounds shows optical isomerism?

2.



102 The IUPAC name of  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is

3.

a) Hexamine cobalt (II) chloride

b) Triammine cobalt (III) trichloride

c) Hexamine cobalt (III) chloride

d) None of the above

102 In the following compounds, the order of acidity is:


4.



(I)



(II)



(III)



(IV)

a)  $\text{III} > \text{IV} > \text{I} > \text{II}$


b)  $\text{I} > \text{IV} > \text{III} > \text{II}$

c)  $\text{II} > \text{I} > \text{III} > \text{IV}$


d)  $\text{IV} > \text{III} > \text{I} > \text{II}$

102 Consider the following structure and choose the correct statements


5.



I



II



III

a) I and II have *R*-configuration



b) I and III have *R*-configuration

c) Only III has *S*-configuration

d) Both (a) and (c) are correct


102 Benzaldehyde, when heated with concentrated KOH solution, gives:

6.



102 Write the IUPAC name of the compound

7.



a) Bicyclo-[2.2.2] octane-2,6-dione

b) Bicyclo-[2.2.2] octane-3,5-dione

c) Bicyclo-[2.2] octane 2,6-dione

d) Bicyclo [2,2] octane-3,5-dione

102 3-chloro-4-methyl benzene sulphonic acid on steam distillation gives:

8.

- a) Toluene                      b) *m*-chloro benzene sulphonic acid                      c) *p*-methyl benzene sulphonic acid                      d) *o*-chloro toluene


102 The oxidation number of platinum in  $[\text{Pt}(\text{NH}_3)_5\text{Cl}]\text{Cl}_3$  is 9.

- a) 2                      b) 3                      c) 4                      d) 6

103 Which of the following is not an organometallic compound? 0.

- a)  $\text{C}_2\text{H}_5\text{ONa}$                       b)  $\text{CH}_3\text{MgI}$                       c) Tetraethyl tin                      d)  $\text{KC}_4\text{H}_9$

103 Which of the following pairs of compounds are enantiomers? 1.



103 Which complex has square planar shape  $dsp^2$ -hybridization? 2.

- a)  $[\text{Ni}(\text{CN})_4]^{2-}$                       b)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$                       c)  $[\text{PtCl}_4]^{2-}$                       d) All of these

103 The complex used as an anticancer agent is 3.

- a) *cis*- $[\text{PtCl}_2(\text{NH}_3)_2]$                       b)  $\text{Na}_2\text{CO}_3$   
c) *trans*- $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$                       d) *cis*- $\text{K}_2[\text{PtCl}_2\text{Br}_2]$

103 Dyes are formed when diazonium salts react with: 4.

- a) Phenols  
b) Aldehydes  
c) Ketones  
d) Alcohols

103 Potassium ferrocyanide is a 5.

- a) Complex salt                      b) Double salt                      c) Normal salt                      d) Mixed salt

103 The primary and secondary valencies of chromium in the complex ion, dichlorodioxalatochromium (III), are respectively. 6.

- a) 3,4                      b) 4,3                      c) 3,6                      d) 6,3

103 The reaction,  $\text{C}_6\text{H}_6 + \text{CH}_3\text{Cl} \xrightarrow[\text{AlCl}_3]{\text{Anhydrous}} \text{C}_6\text{H}_5\text{CH}_3 + \text{HCl}$  is an example of: 7.

- a) Friedel-Craft's reaction  
b) Kolbe's synthesis  
c) Wurtz's reaction  
d) Grignard synthesis

103 The correct statement related to IUPAC nomenclature is 8.

- a) If 2 or more chains of equal length are seen in the compound then the chain with minimum number of side chains will be preferred

- b) If double and triple bonds are at symmetrical positions in a compound then triple bond gets lower preference  
 c) Correct IUPAC name of  $\text{CH}_3\text{COC}_2\text{H}_5$  is ethyl methyl ketone  
 d) As far as possible, the IUPAC name of a compound is written as a single word

103 Which of the following isomerism is shown by ethyl acetoacetate?

9.

- a) Geometrical isomerism  
 b) Keto-enol tautomerism  
 c) Enantiomerism  
 d) Diastereoisomerism


104 The number of moles of ions given on complete ionisation of one mole of  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is/are





0.

- a) 4  
 b) 3  
 c) 2  
 d) 1

104 The major products ( $P$ ,  $Q$ ) in the given reaction are:

1.



- a)  and  $\text{CH}_3\text{CHO}$   
 b)  and  $\text{CH}_3\text{COCH}_3$   
 c)  and  $\text{CH}_3\text{COCH}_3$   
 d)  and  $\text{CH}_3\text{CH}_2\text{CHO}$

104  $p$ -nitrophenol is stronger acid than phenol because nitro group is:

2.

- a) Electron withdrawing  
 b) Electron donating  
 c) Basic  
 d) Acidic

104 Among the following group, which deactivates benzene ring for electrophilic substitution:

3.

- a) Methyl  
 b) Amino  
 c) Hydroxyl  
 d) Chlorine

104 Iron has lowest oxidation state in:

4.

- a)  $\text{Fe}(\text{CO})_5$   
 b)  $\text{Fe}_2\text{O}_3$   
 c)  $\text{K}_2\text{FeO}_4$   
 d)  $\text{FeSO}_4 \cdot (\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$

104 The dihedral angle between the two methyl groups in Gauch conformation of  $n$  butane is






5.

- a)  $120^\circ$   
 b)  $180^\circ$   
 c)  $45^\circ$   
 d)  $60^\circ$

104 Increasing order of acid strength among  $p$ -methoxyphenol,  $p$ -methylphenol and  $p$ -nitrophenol is:

6.







- a) *p*-methylphenol < *p*-methoxyphenol < *p*-nitrophenol  
 b) *p*-methoxyphenol < *p*-methylphenol < *p*-nitrophenol  
 c) *p*-nitrophenol < *p*-methoxyphenol < *p*-methylphenol  
 d) *p*-nitrophenol < *p*-methylphenol < *p*-methoxyphenol
- 104 Total number of isomeric alcohols with formula  $C_4H_{10}O$  are
7. a) 1 b) 2 c) 3 d) 4
- 104 What is the IUPAC name of  $Na_2[Fe(CN)_5NO]$ ?
8. a) Pentacyanonitroso sodium ferrate b) Pentacyanonitroso sodium ferrate(II)  
 c) Sodium pentacyanonitroso ferrate(II) d) Sodium pentacyanonitroso ferrate
- 104 Which of the following cations does not form an ammine complex with excess of ammonia?
9. a)  $Ag^+$  b)  $Cu^{2+}$  c)  $Cd^{2+}$  d)  $Na^+$
- 105 In the complex  $K_2[Fe(CN)_6]$
0. a) The complex is high spin complex b) Both Fe atoms are in the same oxidation state  
 c) The coordination number of iron is 4 d) Both Fe atoms are in different oxidation state
- 105 The number of chiral carbon atoms present in the molecule
1.  is
- a) 3 b) 4 c) 2 d) 1
- 105 The complex that doesn't give a precipitate with  $AgNO_3$  solution
2. a)  $[Co(NH_3)_3Cl_3]$  b)  $[Co(NH_3)_6]Cl_3$  c)  $[Ag(NH_3)_2]Cl$  d)  $[Cr(NH_3)_4Cl_2]Cl$
- 105 The IUPAC name of the given compound  $[Co(NH_3)_5Cl]Cl_2$  is
3. a) Penta amino cobalt chloride chlorate b) Cobalt penta ammine chloro chloride  
 c) Pentamine chloro cobalt (III) chloride. d) Penta amino cobalt (III) chlorate
- 105 Amongst  $Ni(CO)_4$ ,  $[Ni(CN)_4]^{2-}$  and  $[NiCl_4]^{2-}$
4. a)  $Ni(CO)_4$  is diamagnetic,  $[NiCl_4]^{2-}$  and  $[Ni(CN)_4]^{2-}$  are paramagnetic  
 b)  $Ni(CO)_4$  and  $[NiCl_4]^{2-}$  are diamagnetic and  $[Ni(CN)_4]^{2-}$  is paramagnetic  
 c)  $Ni(CO)_4$  and  $[Ni(CN)_4]^{2-}$  are diamagnetic and  $[NiCl_4]^{2-}$  is paramagnetic  
 d)  $[NiCl_4]^{2-}$  and  $[Ni(CN)_4]^{2-}$  are diamagnetic and  $Ni(CO)_4$  is paramagnetic
- 105 Which aromatic acid among the following is weaker than simple benzoic acid?
5. a)  b)  c)  d) 
- 105 Which statement is incorrect?
6. a)  $Ni(CO)_4$ -tetrahedral, paramagnetic  
 b)  $[Ni(CN)_4]^{2-}$ -square planar, diamagnetic  
 c)  $Ni(CO)_4$ -tetrahedral, diamagnetic

d)  $[\text{NiCl}_4]^{2-}$  -tetrahedral, paramagnetic

105 Which of the following has asymmetric C-atom?

7.



105 The IUPAC name of

8.  $\text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{OH}$



is

a) 1-phenyl-3-propanol

b) 3-phenyl-1-propanol

c) 1-hydroxy-3-phenyl-propane

d) None of the above

105 The complexes  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$  and  $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$  are the examples of which type of

9. isomerism?

a) Geometrical isomerism

b) Linkage isomerism

c) Ionization isomerism

d) Coordination isomerism

106 Racemic tartaric acid is optically inactive due to

0.

a) External compensation

b) Internal compensation

c) Presence of plane of symmetry

d) All of the above

106 Nitration of aniline is done in:

1.

a) Acidic medium

b) Alkaline medium

c) Neutral medium

d) In acidic medium by first converting it into acetanilide before nitration

106 A bridging ligand possesses:

2.

a) Polydentate or monodentate nature

b) Two or more donor centres

c) The tendency to get itself attached to two metal ions

d) All of the above

106 What is the neutralization equivalent of benzoic acid?

3.

a) 122

b) 61

c) 244

d) 488

106 *m*-chlorobenzaldehyde on reaction with conc. KOH at room temperature gives:

4.

a) Potassium *m*-chlorobenzoate and *m*-hydroxy benzaldehyde

b) *m*-hydroxybenzaldehyde and *m*-chlorobenzylalcohol

c) *m*-chlorobenzylalcohol and *m*-hydroxy benzylalcohol

d) Potassium *m*-chlorobenzoate and *m*-chlorobenzyl alcohol

106 The oxidation number of Fe in brown ring  $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+}$  is

5.

a) 0


b) +1

c) +2

d) +3

106  $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3$  (at. No. of Cr=24) has a magnetic moment of 3.83 BM. The correct distribution of 3d-

6. electrons in the chromium of the complex:  
 a)  $3d_{xy}^1, 3d_{yz}^1, 3d_{xz}^1$       b)  $3d_{xy}^1, 3d_{yz}^1, 3d_{z^2}^1$       c)  $(3d_{x^2-y^2}^1, 3d_{z^2}^1, 3d_{xz}^1)$       d)  $3d_{xy}^1, (3d_{x^2-y^2}^1, 3d_{yz}^1)$
- 106 Excess of silver nitrate solution is added to 100 mL of 0.01 M pentaqua chloro chromium (III) chloride solution. The mass of silver chloride obtained in grams is [Atomic mass of silver is 108].  
 a)  $287 \times 10^{-3}$       b)  $143.5 \times 10^{-3}$       c)  $143.5 \times 10^{-2}$       d)  $287 \times 10^{-2}$
- 106 The total number of possible structural isomers of the compound  $[\text{Cu}^{\text{II}}(\text{NH}_3)_4][\text{Pt}^{\text{II}}\text{Cl}_4]$  are:  
 8. a) 3      b) 5      c) 4      d) 6
- 106 A similarity between optical and geometrical isomerism is that :  
 9. a) Each gives equal number of isomers for a given compound  
 b) If in a compound one is present then so is the other  
 c) Both are included in stereoisomerism  
 d) They have no similarity
- 107 In  $[\text{Ni}(\text{NH}_3)_4]\text{SO}_4$ , the valency and coordination number of Ni will be respectively  
 0. a) 3 and 6      b) 4 and 4      c) 4 and 2      d) 2 and 4
- 107  $\text{C}_6\text{H}_5\text{CHO}$  is different from aliphatic aldehyde in its reaction towards:  
 1. a) Tollen's reagent      b) Schiff's reagent      c)  $\text{NaHSO}_3$       d) Fehling's solution
- 107 Oxidation of naphthalene by acidic  $\text{KMnO}_4$  gives:  
 2. a) Toluene      b) Benzaldehyde      c) Phthalic acid      d) Benzoic acid
- 107 The number of possible theoretical conformations of *n*-butane are  
 3. a) Two      b) Three      c) Four      d) Infinite
- 107 Which is correct order for acidic nature of following acids?  
 4. (I)  $\text{PhCOOH}$       (II)  $o\text{-NO}_2\text{C}_6\text{H}_4\text{COOH}$   
 (III)  $p\text{-NO}_2\text{C}_6\text{H}_4\text{COOH}$       (IV)  $m\text{-NO}_2\text{C}_6\text{H}_4\text{COOH}$   
 a)  $\text{II} > \text{III} > \text{IV} > \text{I}$       b)  $\text{II} > \text{IV} > \text{III} > \text{I}$       c)  $\text{II} > \text{IV} > \text{I} > \text{III}$       d)  $\text{I} > \text{II} > \text{III} > \text{IV}$
- 107 Salicylic acid when treated with zinc dust gives:  
 5. a) Phenol      b) Salicylaldehyde      c) Benzene      d) Benzoic acid
- 107 Action of  $\text{PCl}_5$  on salicylic acid produces:  
 6. a) *o*-chlorobenzoyl  
 b) *o*-hydroxybenzoyl chloride  
 c) *o*-chlorobenzoic acid  
 d) None of the above
- 107 Which of the following species is most stable?  
 7. a)  $p\text{-O}_2\text{N}-\text{C}_6\text{H}_4-\text{CH}_2^+$   
 b)  $\text{C}_6\text{H}_5-\text{CH}_2^+$   
 c)  $p\text{-Cl}-\text{C}_6\text{H}_4-\text{CH}_2^+$   
 d)  $p\text{-CH}_3\text{O}-\text{C}_6\text{H}_4-\text{CH}_2^+$
- 107 Give the IUPAC name of the following



- a) 5-ethyl-4, 4-dimethyloctane  
b) 4-ethyl-5, 5-dimetyloctane  
c) 3-ethyl-2-methyl-2-propyl hexane  
d) 4-ethyl-5-methyl, 5-propyl hexane

107 Which of the following reacts with KCN to form benzoin?

9.

- a)  $C_6H_5CHO$                       b)  $C_6H_5Cl$                       c)  $C_2H_5Cl$                       d)  $C_6H_5CH_3$





108 Which one is an organometallic compound in the following?

0.

- a)  $C_2H_5ONa$                       b)  $C_2H_5-S-S-C_2H_5$   
c)  $Al_2(CH_3)_6$                       d)  $Al(C_6H_5S)_3$

108 The formula of picramide is:

1.





108 An alkane forms isomers if minimum number of C-atom is:

2.

- a) 1                      b) 2                      c) 3                      d) 4

108 Which will form geometrical isomers?

3.



d) All of these

108 Choose the option which show correct preferential order of groups among the following

4.

- a)  $-COOH, -CHO, -OH, -NH_2$                       b)  $-NH_2, -OH, CHO, -COOH$   
c)  $-COOH, -OH, -NH_2, -CHO$                       d)  $-COOH, -NH_2, -CHO, -OH$

108 The number of precipitable halide ions in  $[Pt(NH_3)Cl_2Br]Cl$  is:

5.

- a) 2                      b) 3                      c) 4                      d) 1

108 Which of the following is polycyclic compound?

6.

- a) Xylene                      b) Cumene                      c) Styrene                      d) Naphthalene

108 Among acetic acid, phenol and *n*-hexanol, which of the compound(s) will react with  $NaHCO_3$  solution to


7. give sodium salt and  $CO_2$ ?

- a) Acetic acid and phenol  
b) Acetic acid  
c) Phenol  
d) *n*-hexanol

108 Nitrosobenzene can be isolated from nitrobenzene under:

8.

- a) Metal and acid  
b) Zn dust and  $NH_4Cl$

- c) Alkaline sodium arsenite  
d) None of the above
- 108 Which of the following complexes is an outer orbital complex?  
9.  
a)  $[\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       c)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$       d) None of these
- 109 In which of the following complex ion, the central metal ion is in a state of  $sp^3 d^2$  hybridisation?  
0.  
a)  $[\text{Co}(\text{F}_6)]^{3-}$       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       c)  $[\text{Fe}(\text{CN})_6]^{3-}$       d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
- 109 Give name of the complex, name should specify the position of ligands  
1.  
  
a) Bistransphosphinecarbonylchloroiridium [II]      b) Carbonylchlorobistransphosphineiridium[III]  
c) Carbonylchlorobistransphosphineiridium[I]      d) Chlorocarbonylbistransphosphineiridium [I]
- 109 The function of anhydrous aluminium chloride in the Friedel-Crafts reaction is:  
2.  
a) To absorb water  
b) To absorb hydrochloric acid  
c) To produce an electrophile  
d) To produce nucleophile
- 109 Coordination isomerism is caused by interchange of ligands between the  
3.  
a) Complex cation and complex anion      b) Inner sphere and outer sphere  
c) Low oxidation and higher oxidation states      d) *cis* and *trans* structure
- 109 Which aldehyde is used in the manufacture of perfumes?  
4.  
a) Cinnamaldehyde      b) Salicylaldehyde      c) Benzaldehyde      d) None of these
- 109 Which of the following statements is not correct?  
5.  
a) A *meso* compound has chiral centres but exhibits no optical activity  
b) A *meso* compound has no chiral centres and thus are optically inactive  
c) A *meso* compound has molecules in which one half of molecule is superimposable on the other even through chiral centre is present in them  
d) A *meso* compound is optically inactive because the rotation caused by one half of molecule is cancelled by the rotation produced by another half
- 109 The volume (in mL) of 0.1 M  $\text{AgNO}_3$  required for complete precipitation of chloride ions present in 30 mL of 0.01 M solution of  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2$ , as silver chloride is close to  
6.  
a) 3      b) 4      c) 5      d) 6
- 109 Benzene is a resonance hybrid mainly of two Kekule structures. Hence:  
7.  
a) Half of the molecules correspond to one structure, and half to the second structure  
b) At low temperatures benzene can be separated into two structures  
c) Two structures make equal contribution to resonance hybrid  
d) An individual benzene molecule changes back and forth between two structures
- 109 Keto form is more stable in  
8.  
a)  $\text{CH}_3\text{COCH}_2\text{COOC}_2\text{H}_5$       b)  $\text{CH}_3\text{COCH}_2\text{COCH}_3$       c)  $\text{CH}_3\text{COCH}_3$       d)  $\text{CH}_3\text{COCH}_2\text{COC}_2\text{H}_5$
- 109 The oxidation state and effective atomic number(EAN) of cobalt ( $\text{CoF}_6$ ) $^{2-}$  are respectively  
9.  
a) 3 and 36      b) 4 and 35      c) 4 and 37      d) 2 and 35





110 Benzamide on reaction with  $\text{POCl}_3$  gives:

0.

- a) Aniline                      b) Chlorobenzene                      c) Benzylamine                      d) Benzonitrile

110 Which pair of carbon skeleton is an example of isomerism?

1.



110 Electrolytic reduction of nitrobenzene in weak acidic medium gives:

2.

- a) Aniline  
b) *p*-hydroxy aniline  
c) Nitrobenzene  
d) *N*-phenyl hydroxyl amine

110 In complexes, metal atom acts as:

3.

- a) Lewis base                      b) Bronsted acid                      c) Bronsted base                      d) Lewis acid

110 When benzene is treated with concentrated  $\text{HNO}_3$  at room temperature it will give:

4.

- a)  $\text{CO}_2$  and  $\text{H}_2\text{O}$                       b) Nitrochlorobenzene                      c) Dark red colour                      d) Dinitrobenzene


110 Which of the following compounds exhibit linkage isomerism?

5.

- a)  $[\text{Co}(\text{en})_3]\text{Cl}_3$                       b)  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$                       c)  $[\text{Co}(\text{en})_2\text{NO}_2\text{Cl}]\text{Br}$                       d)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Br}_2$

110 The compound

6.



have IUPAC name as

- a) Tricyclopropyl                      b) Tricyclopropane  
c) 1,1', 2', 1''-tercyclo propane                      d) None of the above

110 The most stable conformation of chlorohydrin at room temperature is

7.

- a) Fully eclipsed                      b) Partially eclipsed                      c) Gauche                      d) Staggered

110 Among  $\text{Ni}(\text{CO})_4$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  and  $[\text{Ni}(\text{Cl})_4]^{2-}$  :

8.

- a)  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{NiCl}_4]^{2-}$  are diamagnetic and  $[\text{Ni}(\text{CN})_4]^{2-}$  is paramagnetic
- b)  $[\text{NiCl}_4]^{2-}$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic and  $[\text{Ni}(\text{CO})_4]$  is paramagnetic
- c)  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic and  $[\text{NiCl}_4]^{2-}$  is paramagnetic
- d)  $[\text{Ni}(\text{CO})_4]$  is diamagnetic and  $[\text{NiCl}_4]^{2-}$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  are paramagnetic

110 The complex  $\text{Hg}[\text{Co}(\text{CNS})_4]$  is correctly named as:

9.

- a) Mercury tetrathiocyanatocobaltate(II)
- b) Mercury cobalt tetrasulphocyno(II)
- c) Mercury tetrasulphocyanidecobaltate(II)
- d) Mercury sulphocyanatocobalt(II)

111 Which of the following compounds is not coloured?

0.

- a)  $\text{Na}_2[\text{CuCl}_4]$
- b)  $\text{Na}_2[\text{CdCl}_4]$
- c)  $\text{K}_4[\text{Fe}(\text{CN})_6]$
- d)  $\text{K}_3[\text{Fe}(\text{CN})_6]$

111 Which one has square planar geometry?

1.

- a)  $[\text{CoCl}_4]^{2-}$
- b)  $[\text{FeCl}_4]^{2-}$
- c)  $[\text{NiCl}_4]^{2-}$
- d)  $[\text{PtCl}_4]^{2-}$

111 Which exhibits highest molar conductivity?

2.

- a)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
- b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
- c)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
- d)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$

111 Coordination compounds have great importance in biological systems. In this context which of the

3. following statement is incorrect?

- a) Chlorophyll is green pigment in plants and contain calcium.
- b) Haemoglobin is the red pigment of blood and contains iron.
- c) Cyanocobalamin is vitamin  $\text{B}_{12}$  and contains cobalt.
- d) Carboxypeptidase-A is an enzyme and contains zinc.

111 Complex salt can be made by the combination of  $[\text{Co}^{\text{III}}(\text{NH}_3)_5\text{Cl}]^x$  with

4.

- a)  $\text{Cl}^-$
- b)  $2\text{Cl}^-$
- c)  $\text{PO}_4^{3-}$
- d)  $2\text{K}^+$

111 Which of the following pairs represents linkage isomers?

5.

- a)  $[\text{Cu}(\text{NH}_3)_4][\text{PtCl}_4]$  and  $[\text{Pt}(\text{NH}_3)_4][\text{CuCl}_4]$
- b)  $[\text{Pd}(\text{PPh}_3)_2(\text{NCS})_2]$  and  $[\text{Pd}(\text{PPh}_3)_2(\text{SCN})_2]$
- c)  $[\text{Co}(\text{NH}_3)_5]\text{NO}_3\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{NO}_3$
- d)  $[\text{PtCl}_2(\text{NH}_3)_4]\text{Br}_2$  and  $[\text{PtBr}_2(\text{NH}_3)_4]\text{Cl}_2$

111 The reaction products of  $\text{C}_6\text{H}_5\text{OCH}_3 + \text{HI} \xrightarrow{\Delta}$  is:

6.

- a)  $\text{C}_6\text{H}_5\text{OH} + \text{CH}_3\text{I}$
- b)  $\text{C}_6\text{H}_5\text{I} + \text{CH}_3\text{OH}$
- c)  $\text{C}_6\text{H}_5\text{CH}_3 + \text{HOI}$
- d)  $\text{C}_6\text{H}_6 + \text{CH}_3\text{OI}$

111 An aromatic amine (A) was treated with alcoholic potash and another compound (Y) when a foul smelling

7. gas was formed with formula  $\text{C}_6\text{H}_5\text{NC}$ . (Y) was formed by reacting a compound (Z) with  $\text{Cl}_2$  in the presence of slaked lime. Compound (Z) is:

- a)  $\text{C}_6\text{H}_5\text{NH}_2$
- b)  $\text{CH}_3\text{OH}$
- c)  $\text{CH}_3\text{COCH}_3$
- d)  $\text{CHCl}_3$

111 Chlorine is most reactive in:

8.

- a)  $\text{CH}_3\text{Cl}$
- b)  $\text{CH}_2=\text{CHCl}$
- c)  $\text{C}_6\text{H}_5\text{Cl}$
- d)  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$

111 The C—C bond order in benzene is close to:

9.

- a) 1.5
- b) 2.5
- c) 3.0
- d) 6.0

112 Mixture X = 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  and 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  was prepared in 2 L of

0. solution

1 L of mixture X + excess  $\text{AgNO}_3 \rightarrow \text{Y}$

1 L of mixture  $X + \text{excess BaCl}_2 \rightarrow Z$

Number of moles of  $Y$  and  $Z$  are

a) 0.01, 0.01

b) 0.01, 0.02

c) 0.02, 0.01

d) 0.02, 0.02

112 Phenol can be converted into salicylic acid by:

1.

a) Etard's reaction

b) Kolbe's reaction

c) Reimer-Tiemann reaction

d) Both (b) and (c)

112  $\text{Fe}_2(\text{CO})_9$  is diamagnetic. Which of the following reasons is correct?

2.

a) Presence of one CO as bridge group

b) Presence of monodentate ligand

c) Metal-metal (Fe-Fe) bond in molecule

d) Resonance hybridization of CO

112 The formula of dichlorobis (urea) copper(II) is:

3.

a)  $[\text{CuO} = \text{C}(\text{NH}_2)_2]\text{Cl}_2$

b)  $[\text{CuCl}_2\{\text{O} = \text{C}(\text{NH}_2)\}]$

c)  $[\text{Cu}\{\text{O} = \text{C}(\text{NH}_2)_2\text{Cl}\}]\text{Cl}$

d)  $[\text{CuCl}_2][\text{O} = \text{C}(\text{NH}_2)_2]\text{H}_2$

112 Which of the following facts about the complex  $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$  is wrong?

4.


a) The complex involves  $d^2sp^3$  hybridisation and is b) The complex is paramagnetic.  
octahedral in shape.

c) The complex is an outer orbital complex.

d) The complex gives white precipitate with silver nitrate solution.

112 The compounds P, Q and S were separately subjected to nitration using  $\text{HNO}_3/\text{H}_2\text{SO}_4$  mixture. The major product formed in each case respectively, is:



5.




(P)

(Q)




(S)




a)



b)








112 Aromaticity of benzene is due to:

6.

- a) Ring
- b) Three double bonds
- c) Delocalisation of  $\pi$ -electrons
- d) None of the above

112

7.

The IUPAC name of  is

- a) 2, 2, 4, 4-tetramethyl pentane
- b) 2, 2-dimethyl propane
- c) 4-ethyl-3-methyl hex-3-ene
- d) Ethyl isopropyl ethene

112 Phenol is heated with a solution of mixture of KBr and  $\text{KBrO}_3$ . The major product obtained in the above

8. reaction is:

- a) 2-bromophenol
- b) 3-bromophenol
- c) 4-bromophenol
- d) 2,4,6-tribromophenol

112 The coordination number of a central metal atom in a complex is determined by

9.

- a) The number around a metal ion bonded by pi-bonds
- b) The number of only anionic ligands bonded to the metal ion
- c) The number of ligands around a metal ion bounded by sigma and pi-bonds both
- d) The number of ligands around a metal ion bonded by sigma bonds

113 The true statement about benzene is:

0.

- a) Because of unsaturation benzene easily undergoes additions
- b) There are two types of C—C bonds in benzene molecule
- c) There is a cyclic delocalisation of  $\pi$ -electrons in benzene
- d) Monosubstitution of benzene gives three isomeric products

113 Which reagent can convert  $\text{>CO}$  group to  $\text{>C(C}_6\text{H}_5\text{)OH}$ ?

1.

- a)  $\text{C}_6\text{H}_5\text{OH}$
- b)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$
- c)  $\text{C}_6\text{H}_5\text{MgBr}$
- d)  $\text{C}_6\text{H}_5\text{Cl}$

113 Which has highest paramagnetism?

2.

- a)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
- b)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$
- c)  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$
- d)  $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$

113 Which is not true ligands metal complex?

3.

- a) Larger the ligand, the more stable is the metal-ligand complex
- b) Highly charged ligand forms stronger bonds

- c) Larger the permanent dipole moment of ligand, the more stable is the bond  
d) Greater the ionization potential of central metal, the stronger the bond
- 113  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{NO}_2$  and  $[\text{Co}(\text{NH}_3)_4\text{Cl} \cdot \text{NO}_2]\text{Cl}$  are  
4. a) Optical isomers      b) Geometrical isomers      c) Ionization isomers      d) Linkage isomers
- 113 Acetophenone on oxidation by perbenzoic acid gives phenyl acetate. The reaction is named as:  
5. a) Baeyer-Villiger oxidation  
b) Perkin's reaction  
c) Claisen condensation  
d) Reformatsky reaction
- 113 Friedel-Craft's reaction does not occur in case of:  
6. a) Toluene      b) Benzene      c) Naphthalene      d) pyridine
- 113 One mole of a complex compound  $\text{Co}(\text{NH}_3)_5\text{Cl}_3$  gives three moles of ions on dissolution in water. One of the same complex reacts with two moles of  $\text{AgNO}_3$  solution to yield two moles of  $\text{AgCl}(s)$ . The structure of the complex is  
7. a)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3] \cdot 2\text{NH}_3$       b)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2] \cdot \text{Cl} \cdot \text{NH}_3$   
c)  $[\text{Co}(\text{NH}_3)_4\text{Cl}]\text{Cl}_2 \cdot \text{NH}_3$       d)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
- 113  $\text{C}_6\text{H}_6$  is a very good industrial solvent for:  
8. a) Oil      b) Fat      c) Rubber      d) All of these
- 113 Salol is used as:  
9. a) Antiseptic      b) Antipyretic      c) Both (a) and (b)      d) None of these
- 114 Presence of nitro gp. in benzene ring:  
0. a) Deactivates the ring for  $\text{S}_\text{E}$  reaction  
b) Activates the ring for  $\text{S}_\text{E}$  reactions  
c) Renders the ring basic  
d) Deactivates the ring for  $\text{S}_\text{N}$  reaction
- 114 Which of the following complexes will show geometrical as well as optical isomerism? (en = ethylene diamine)  
1. a)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$       b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_4]$       c)  $[\text{Pt}(\text{en})_3]^{4+}$       d)  $[\text{Pt}(\text{en})_2\text{Cl}_2]$
- 114 The huge number of organic compounds is due to the fact that  
2. a) Tetravalency of carbon      b) Carbon possesses property of catenation  
c) Carbon compounds exhibits polymerisation      d) Both (b) and (c)
- 114 When nitrobenzene is treated with  $\text{Br}_2$  in presence of  $\text{FeBr}_3$ , the major product formed is *m*-bromonitrobenzene. Statements which are related to obtain the *m*-isomer are:  
3. a) The relative electron density on *meta* carbon is more than that of *ortho* and *para* positions  
b) Loss of aromaticity when  $\text{Br}^+$  attacks at the *ortho* and *para* positions and not at *meta* position  
c) Easier loss of  $\text{H}^+$  to regain aromaticity from the *meta* position than from *ortho* and *para* positions  
d) None of the above
- 114 Which one of the following compounds when dissolved in water, gives a solution with pH more than 7?  
4. a)  $\text{C}_6\text{H}_5\text{NH}_2$       b)  $\text{C}_6\text{H}_5\text{OH}$       c)  $\text{C}_2\text{H}_5\text{OH}$       d)  $\text{CH}_3\text{COCH}_3$
- 114 Formula of ferrocene is:  
5. a)  $[\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{Fe}(\text{CN})_6]^{3+}$       c)  $[\text{Fe}(\text{CO})_5]$       d)  $[(\text{C}_6\text{H}_5)_2\text{Fe}]$

114 What is the EAN of nickel in  $\text{Ni}(\text{CO})_4$ ?

6.

- a) 38                                      b) 30                                      c) 36                                      d) 32

114 One mole of the complex compound  $\text{Co}(\text{NH}_3)_5\text{Cl}_3$ , gives 3 moles of ions on dissolution in water. One mole

7. of the same complex reacts with two moles of  $\text{AgNO}_3$  solution to yield two moles of  $\text{AgCl}(s)$ . The structure of the complex is

- a)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$                                       b)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_2] \cdot 2\text{NH}_3$   
c)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl} \cdot \text{NH}_3$                                       d)  $[\text{Co}(\text{NH}_3)_4\text{Cl}]\text{Cl}_2 \cdot \text{NH}_3$

114 Which one of the following has largest number of isomers?

8. ( $R$ =alkyl group, en=ethylenediamine)

- a)  $[\text{Ru}(\text{NH}_3)_4\text{Cl}_2]^+$                       b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$                       c)  $[\text{Ir}(\text{PR}_3)_2\text{H}(\text{CO})]^{2+}$                       d)  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$

114 Which complex is likely to show optical activity?

9.

- a) *Trans*- $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$ .  
b)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$   
c) *Cis*- $[\text{Co}(\text{NH}_3)_2(\text{en})_2]^{3+}$   
d) *Trans*- $[\text{Co}(\text{NH}_3)_2(\text{en})_2]^{3+}$


115 A square planar complex is formed by hybridization of which atomic orbitals?

10.

- a)  $s, p_x, p_y, d_{yz}$                       b)  $s, p_x, p_y, d_{x^2-y^2}$                       c)  $s, p_x, p_y, d_{z^2}$                       d)  $s, p_x, p_y, d_{xy}$

115 The IUPAC name of the compound

1.  $\text{CH}_2 - \text{CH} - \text{CH} - \text{CH}_2\text{COCl}$  is



- a) 1, 2, 3, 4-butanetetrachlorocarbonyl                      b) 1, 2, 3, 4-butanetetrachloroformyl  
c) 1, 2, 4-butanetricarboxylic acid                      d) None of the above

115 Nitrobenzene can be prepared from benzene by using a mixture of conc.  $\text{HNO}_3$  and conc.  $\text{H}_2\text{SO}_4$ . In the

2. nitrating mixture  $\text{HNO}_3$  acts as a:

- a) Base                      b) Acid                      c) Reducing agent                      d) Catalyst

115 In the compound lithium tetrahydroaluminate, the ligand is

3.

- a)  $\text{H}$                                       b)  $\text{H}^+$                                       c)  $\text{H}^-$                                       d) None of these