


Date :
Time :
Marks :

PHYSICS

MAGNETISM AND MATTER

Single Correct Answer Type

- Magnetic intensity for an axial point due to a short bar magnet of magnetic moment M is given by
 a) $\frac{\mu_0}{4\pi} \times \frac{M}{d^3}$ b) $\frac{\mu_0}{4\pi} \times \frac{M}{d^2}$ c) $\frac{\mu_0}{2\pi} \times \frac{M}{d^3}$ d) $\frac{\mu_0}{2\pi} \times \frac{M}{d^2}$
- To measure which of the following, is a tangent galvanometer used
 a) Charge b) Angle c) Current d) Magnetic intensity
- A long magnet is placed vertically with its S –pole resting on the table. A neutral point is obtained 10 cm from the pole the geographic north of it. If $H = 3.2 \times 10^{-5} \text{ T}$, then the pole strength of magnet is
 a) 8 ab-A-cm $^{-1}$ b) 16 ab-A-cm $^{-1}$ c) 32 ab-A-cm $^{-1}$ d) 64 ab-A-cm $^{-1}$
- The magnetic moment of a magnet of length 10 cm and pole strength 4.0 Am will be
 a) 0.4 Am 2 b) 1.6 Am 2 c) 20 Am 2 d) 8.0 Am 2
- Two magnets held together in earth's magnetic field with same polarity together make 12 vib – min $^{-1}$ and when opposite poles together make 4 vib – min $^{-1}$. The ratio of magnetic moments is
 a) 9 : 1 b) 1 : 3 c) 1 : 9 d) 10 : 8
- In which direction, the magnetic field on the axis at a distance z from the centre of the bar magnet would be?
 a) In the perpendicular direction of the magnetic moment (\mathbf{M}) of the magnet
 b) In the direction of the magnetic dipole moment (\mathbf{M}) of the magnet
 c) Its direction depends on the magnitude of the magnetic moment (\mathbf{M}) of the magnet
 d) In the opposite direction of the magnetic dipole moment (\mathbf{M}) of the magnet
- Two short magnets have equal pole strengths but one is twice as long as other. The shorter magnet is placed 20 cm in tan A position from the compass needle. The longer magnet must be placed on the other side of the magnetometer for no deflection at a distance equal to
 a) 20 cm b) $20 \times (2)^{1/3}$ cm c) $20 \times (2)^{2/3}$ cm d) $20 \times (2)$ cm
- Electromagnets are made of soft iron because soft iron has
 a) Low susceptibility and low retentivity b) Low susceptibility and high retentivity
 c) High permeability and low retentivity d) High permeability and high coercivity
- A long magnet is cut into two equal parts, such that the length of each half is same as that of original magnet. If the period of original magnetic is T , the period of new magnet is
 a) T b) $\frac{T}{2}$ c) $\frac{T}{4}$ d) $2T$
- If the $B - H$ curves of two samples of P and Q of iron are as shown below, then which one of the following statements is correct?



- Both P and Q are suitable for making permanent magnet
- P is suitable for making permanent magnet and Q for making electromagnet


- c) P is suitable for making electromagnet and Q is suitable for permanent magnet
d) Both P and Q are suitable for making electromagnets
11. A short magnetic needle is pivoted in a uniform magnetic field of strength 1 T . When another magnetic field of strength $\sqrt{3}\text{ T}$ is applied to the needle in a perpendicular direction, the needle deflects through an angle θ , where θ is
a) 30° b) 45° c) 90° d) 60°
 12. The magnetic susceptibility does not depend upon the temperature in
a) Ferrite substances b) Ferromagnetic substances
c) Diamagnetic substances d) Paramagnetic substances
 13. The area of hysteresis loop of a material is equivalent to 250 joule . When 10 kg material is magnetized by an alternating field of 50 Hz then energy lost in one hour will be (density of material is 7.5 gm/cm^3)
a) $6 \times 10^4\text{ J}$ b) $6 \times 10^4\text{ erg}$ c) $3 \times 10^2\text{ J}$ d) $3 \times 10^2\text{ erg}$
 14. A magnetic dipole is placed in two perpendicular magnetic fields \vec{B} and \vec{H} and is in equilibrium taking angle θ with \vec{B} . Then
a) $B = H$ b) $B \cos\theta = H \sin\theta$ c) $B \sin\theta = H \cos\theta$ d) $B = H \tan\theta$
 15. Keeping dissimilar poles of two magnets of equal pole strength and length same side, their time period will be
a) Zero b) One second c) Infinity d) Any value
 16. When two magnetic moments are compared using equal distance method the deflections produced are 45° and 30° . If the length of magnets are in the ratio $1 : 2$, the ratio of their pole strength is
a) $3 : 1$ b) $3 : 2$ c) $\sqrt{3} : 1$ d) $2\sqrt{3} : 1$
 17. If a magnetic dipole of dipole moment M rotated through an angle θ with respect to the direction of the field H , then the work done is
a) $MH \sin\theta$ b) $MH(1 - \sin\theta)$ c) $MH \cos\theta$ d) $MH(1 - \cos\theta)$
 18. The radius of the coil of a tangent galvanometer which has 10 turns is 0.1 m . The current required to produce a deflection of 60° ($B_H = 4 \times 10^{-5}\text{ T}$) is
a) 3 A b) 1.1 A c) 2.1 A d) 1.5 A
 19. At a temperature of 30° C , the susceptibility of a ferromagnetic material is found to be X . Its susceptibility at 333° C is
a) X b) $0.5 X$ c) $2 X$ d) $0.09 X$
 20. Magnetic moment of bar magnet is M . The work done to turn the magnet by 90° of magnet in direction of magnetic field B will be
a) Zero b) $\frac{1}{2} MB$ c) $2 MB$ d) MB
 21. Two identical short bar magnets, each having magnetic moment of 10 Am^2 , are arranged such that their axial lines are perpendicular to each other and their centres be along the same straight line in a horizontal plane. If the distance between their centres is 0.2 m , the resultant magnetic induction at a point midway between them is
($\mu_0 = 4\pi \times 10^{-7}\text{ Hm}^{-1}$)
a) $\sqrt{2} \times 10^{-7}\text{ tesla}$ b) $\sqrt{5} \times 10^{-7}\text{ tesla}$ c) $\sqrt{2} \times 10^{-3}\text{ tesla}$ d) $\sqrt{5} \times 10^{-3}\text{ tesla}$
 22. Relative permeability of iron is 5500, then its magnetic susceptibility will be
a) 5500×10^7 b) 5500×10^{-7} c) 5501 d) 5499
 23. A very small magnet is placed in the magnetic meridian with its south pole pointing north. The null point is obtained 20 cm away from the centre of the magnet. If the earth's magnetic field (horizontal component) at this point is 0.3 gauss , the magnetic moment of the magnet is
a) $8.0 \times 10^2\text{ e.m.u}$ b) $1.2 \times 10^3\text{ e.m.u}$ c) $2.4 \times 10^3\text{ e.m.u}$ d) $3.6 \times 10^3\text{ e.m.u}$
 24. Hysteresis loss is minimized by using
a) Alloy of steel b) Shell type of core
c) Thick wire which has low resistance d) Mu metal

25. A magnet of length 0.1 m and pole strength 10^{-4} A-m is kept in a magnetic field of 30 Wb m^{-2} at an angle 30° . The couple acting on it is $\times 10^{-4}$ Nm.
 a) 7.5 b) 3.0 c) 4.5 d) 1.5
26. If the angular momentum of an electron is \vec{J} then the magnitude of the magnetic moment will be
 a) $\frac{eJ}{m}$ b) $\frac{eJ}{2m}$ c) $eJ 2m$ d) $\frac{2m}{eJ}$
27. A deflection magnetometer is adjusted in the usual way. When a magnet is introduced, the deflection observed is θ , and the period of oscillation of the needle in the magnetometer is T . When the magnet is removed, the period of oscillation is T_0 . Find the relation between T and T_0 is
 a) $T^2 = T_0^2 \cos \theta$ b) $T = T_0 \cos \theta$ c) $T = \frac{T_0}{\cos \theta}$ d) $T^2 = \frac{T_0^2}{\cos \theta}$
28. A long magnetic needle of length $2L$, magnetic moment M and pole strength m units is broken into two pieces at the middle. The magnetic moment and pole strength of each piece will be
 a) $\frac{M}{2}, \frac{m}{2}$ b) $M, \frac{m}{2}$ c) $\frac{M}{2}, m$ d) M, m
29. A solenoid has core of a material with relative permeability 500 and its windings carry a current of 1A. The number of turns of the solenoid is 500 per metre. The magnetization of the material is nearly
 a) $2.5 \times 10^3 \text{ Am}^{-1}$ b) $2.5 \times 10^5 \text{ Am}^{-1}$ c) $2.0 \times 10^3 \text{ Am}^{-1}$ d) $2.0 \times 10^5 \text{ Am}^{-1}$
30. Two short magnets with their axes horizontal and perpendicular to the magnetic meridian are placed with their centres 40 cm east and 50 cm west of magnetic needle. If the needle remains undeflected, the ratio of their magnetic moments $M_1 : M_2$ is
 a) 4 : 5 b) 16 : 25 c) 64 : 125 d) 2 : $\sqrt{5}$
31. The lines of force due to earth's horizontal component of magnetic field are
 a) Parallel straight lines b) Concentric circles c) Elliptical d) Parabolic
32. The coil in a tangent galvanometer is 16 cm in radius. If a current of 20 mA is to produce a deflection of 45° then the number of turns wound on it, is (Take horizontal component of earth's magnetic field = $0.36 \times 10^{-4} \text{ T}$ and $\mu_0 = 4\pi \times 10^{-7} \text{ Wb A}^{-1} \text{ m}^{-1}$)
 a) 229 b) 458 c) 689 d) 916
33. A compass needle which is allowed to move in a horizontal plane is taken to a geomagnetic pole. It
 a) Will become rigid showing no movement
 b) Will stay in any position
 c) Will stay in north-south direction only
 d) Will stay in east-west direction only
34. The magnetic moment of a diamagnetic atom is
 a) Much greater than one b) 1
 c) Between zero and one d) Equal to zero
35. A tangent galvanometer has a coil with 50 turns and radius equal to 4 cm. A current of 0.1 A is passing through it. The plane of the coil is set parallel to the earth's magnetic meridian. If the value of the earth's horizontal component of the magnetic field is $7 \times 10^{-5} \text{ tesla}$ and $\mu_0 = 4\pi \times 10^{-7} \text{ weber/amp} \times \text{m}$, then the deflection in the galvanometer needle will be
 a) 45° b) 48.2° c) 50.7° d) 52.7°
36. With a standard rectangular bar magnet of length (l), breadth (b ; $b \ll l$) and magnetic moment M , the time period of the magnet in vibration magnetometer is 4 s. If the magnet is cut normal to its length into four equal pieces, the time period (in second) with one of the pieces
 a) 16 b) 2 c) 1 d) $\frac{1}{4}$
37. A wire of length L metre carrying current i , ampere is bent in the form of a circle. What is the magnitude of magnetic of magnetic dipole moment?
 a) $iL^2/4\pi$ b) $i^2L^2/4\pi$ c) $i^2L/8\pi$ d) $iL^2/8\pi$

38. Two tangent galvanometer having coils of the same radius are connected in series. A current flowing in them produces of 60° and 45° respectively. The ratio of the number of turns in the coil is

a) $4/3$ b) $(\sqrt{3} + 1)/1$ c) $\frac{\sqrt{3} + 1}{\sqrt{3} - 1}$ d) $\frac{\sqrt{3}}{1}$

39. Two identical bar magnets with a length 10 cm and weight 50 g-weight are arranged freely with their poles facing in a inverted vertical glass tube. The upper magnet hangs in the air above the lower one so that the distance between the nearest pole of the magnet is 3 mm . Pole strength of the poles of each magnet will be




- a) $6.64\text{ amp} \times \text{m}$ b) $2\text{ amp} \times \text{m}$ c) $10.25\text{ amp} \times \text{m}$ d) None of these
40. The magnetized wire of moment M and length l is bent in the form of semicircle of radius r . Then its magnetic moment is
- a) $\frac{2M}{\pi}$ b) $2M$ c) $\frac{M}{\pi}$ d) Zero
41. Among the following properties describing diamagnetism identify the property that is wrongly stated
- a) Diamagnetic material do not have permanent magnetic moment
b) Diamagnetism is explained in terms of electromagnetic induction
c) Diamagnetic materials have a small positive susceptibility
d) The magnetic moment of individual electrons neutralize each other
42. A bar magnet of magnetic moment M and moment of inertia I is freely suspended such that the magnetic axial line is in the direction of magnetic meridian. If the magnet is displaced by a very small angle (θ), the angular acceleration is (Magnetic induction of earth's horizontal field = B_H)
- a) $\frac{MB_H \theta}{I}$ b) $\frac{IB_H \theta}{M}$ c) $\frac{M\theta}{IB_H}$ d) $\frac{I\theta}{MB_H}$
43. A tangent galvanometer has a coil of 25 turns and a radius of 15 cm . The horizontal component of the earth's magnetic field is $3 \times 10^{-5}\text{ T}$. The current required to produce a deflection of 45° in it is
- a) 0.29 A b) 0.14 A c) 1.2 A d) $3.6 \times 10^{-5}\text{ A}$
44. A magnet makes 40 oscillations per minute at a place having magnetic field intensity of $0.1 \times 10^{-5}\text{ T}$. At another place, it takes 2.5 sec to complete one vibration. The value of earth's horizontal field at that place is+
- a) $0.25 \times 10^{-6}\text{ T}$ b) $0.36 \times 10^{-6}\text{ T}$ c) $0.66 \times 10^{-8}\text{ T}$ d) $1.2 \times 10^{-6}\text{ T}$
45. Two identical magnetic dipoles of magnetic moment 2 Am^2 are placed at a separation of 2 m with their axis perpendicular to each other in air. The resultant magnetic field at a mid-point between the dipoles is
- a) $4\sqrt{5} \times 10^{-5}\text{ T}$ b) $2\sqrt{5} \times 10^{-5}\text{ T}$ c) $4\sqrt{5} \times 10^{-7}\text{ T}$ d) $2\sqrt{5} \times 10^{-7}\text{ T}$
46. Two short magnets with pole strengths of 900 ab amp-cm and 100 ab-amp-cm are placed with their axes in the same vertical line, with similar poles facing each other. Each magnet has a length of 1 cm . When separation between the nearer poles is 1 cm , the weight of upper magnet is supported by the repulsive force between the magnets. If $g = 1000\text{ cms}^{-2}$, then the mass of upper magnet is
- a) 100 g b) 55 g c) 45 g d) 77.5 g
47. Nickel shows ferromagnetic property at room temperature. If the temperature is increased beyond Curie temperature, then it will show
- a) Paramagnetism b) Anti-ferromagnetism
c) No magnetic property d) Diamagnetism
48. To shield an instrument from external magnetic field, it is placed inside a cabin made of
- a) Wood b) Ebonite

c) Iron


d) Diamagnetic substance

49. The basic magnetization curve for a ferromagnetic material is shown in figure. Then, the value of relative permeability is highest for the point




- a) P b) Q c) R d) S
50. A small bar magnet A oscillates in a horizontal plane with a period T at a place where the angle of dip is 60° . When the same needle is made to oscillate in a vertical plane coinciding with the magnetic meridian, its period will be
- a) $\frac{T}{\sqrt{2}}$ b) T c) $\sqrt{2}T$ d) $2T$
51. A dip circle is at right angles to the magnetic meridian.
- a) 0° b) 90° c) 45° d) $4 : 1$
52. A magnet of magnetic moment $50 \hat{i} \text{ A-m}^2$ is placed along the x -axis in a magnetic field $\vec{B} = (0.5 \hat{i} + 3.0 \hat{j})T$. The torque acting on the magnet is
- a) $175 \hat{k} \text{ N-m}$ b) $150 \hat{k} \text{ N-m}$ c) $75 \hat{k} \text{ N-m}$ d) $25\sqrt{37} \hat{k} \text{ N-m}$
53. When a metallic plate swings between the poles of a magnet
- a) No effect on the plate
b) Eddy currents are set inside the plate and the direction of the current is along the motion of the plate
c) Eddy currents are set up inside the plate and the direction of the current oppose the motion of the plate
d) Eddy currents are set up inside the plate
54. Susceptibility of Mg at 300 K is 1.2×10^{-5} . The temperature at which susceptibility will be 1.8×10^{-5} is
- a) 450 K b) 200 K c) 375 K d) None of these
55. The magnetic susceptibility of paramagnetic materials is
- a) Positive, but very high b) Negative, but very small
c) Negative, but very high d) Positive, but small
56. Which of the following is represented by the area enclosed by a hysteresis loop (B - H curve)?
- a) Permeability b) Retentivity
c) Heat energy lost per unit volume in the sample d) Susceptibility
57. A straight wire carrying current i is turned into a circular loop. If the magnitude of magnetic moment associated with it in M.K.S. unit is M , the length of wire will be
- a) $4\pi iM$ b) $\sqrt{\frac{4\pi M}{i}}$ c) $\sqrt{\frac{4\pi i}{M}}$ d) $\frac{M\pi}{4i}$
58. A magnet performs 10 oscillations per minute in a horizontal plane at a place where the angle of dip is 45° and the total intensity is 0.707 units. The number of oscillations per minute at a place where dip angle is 60° and total intensity is 0.5 CGS units will be
- a) 5 b) 7 c) 9 d) 11
59. The relative permeability of a substance X is slightly less than unity and that of substance Y is slightly more than unity, then
- a) X is paramagnetic and Y is ferromagnetic b) X is diamagnetic and Y is ferromagnetic
c) X and Y both are paramagnetic d) X is diamagnetic and Y is paramagnetic
60. A bar magnet of magnetic moment 3.0 A-m^2 is placed in a uniform magnetic induction field of $2 \times 10^{-5} \text{ T}$. If each pole of the magnet experiences a force of $6 \times 10^{-4} \text{ N}$, the length of the magnet is


- a) 0.5 m b) 0.3 m c) 0.2 m d) 0.1 m
61. A magnet freely suspended in a vibration magnetometer makes 40 oscillations per minute at a place A and 20 oscillations per minute at a place B . If the horizontal component of earth's magnetic field at A is $36 \times 10^{-6} \text{T}$, then its value at B is
- a) $36 \times 10^{-6} \text{T}$ b) $9 \times 10^{-6} \text{T}$ c) $144 \times 10^{-6} \text{T}$ d) $228 \times 10^{-6} \text{T}$
62. Tangent galvanometer measures
- a) Capacitance b) Current c) Resistance d) Potential difference
63. At a certain place the horizontal component of the earth's magnetic field is B_0 and the angle of dip is 45° . The total intensity of the field at that place will be
- a) B_0 b) $\sqrt{2} B_0$ c) $2 B_0$ d) B_0^2
64. If a bar magnet of length l and cross-sectional area A is cut into two equal parts as shown in figure, then the pole strength of each pole becomes



- a) Half b) Double c) One-fourth d) Four time
65. At a point on the right bisector of a magnetic dipole magnetic
- a) Potential varies as $\frac{1}{r^2}$
- b) Potential is zero at all points on the right bisector
- c) Field varies as r^2
- d) Field is perpendicular to the axis of dipole
66. Two short magnets placed along the same axis with their like poles facing each other repel each other with a force which varies inversely as
- a) Square of the distance b) Cube of the distance
- c) Distance d) Fourth power of the distance
67. A bar magnet of magnetic moment \vec{M} is placed in a magnetic field of induction \vec{B} . The torque exerted on it is
- a) $\vec{M} \cdot \vec{B}$ b) $-\vec{M} \cdot \vec{B}$ c) $\vec{M} \times \vec{B}$ d) $\vec{B} \times \vec{M}$
68. A dip needle which is free to move in a vertical plane perpendicular to magnetic meridian will remain
- a) Horizontal b) Vertical
- c) Neither horizontal nor vertical d) Inclined
69. Relative permittivity and permeability of a material are ϵ_r and μ_r , respectively. Which of the following values of these quantities are allowed for a diamagnetic material?
- a) $\epsilon_r = 0.5, \mu_r = 1.5$ b) $\epsilon_r = 1.5, \mu_r = 0.5$ c) $\epsilon_r = 0.5, \mu_r = 0.5$ d) $\epsilon_r = 1.5, \mu_r = 1.5$
70. At a place the angle of dip is 30° . If the horizontal component of earth's magnetic field is B_H , then the total field intensity is
- a) $\frac{B_H}{2}$ b) $\frac{2B_H}{\sqrt{3}}$ c) $B_H \sqrt{2}$ d) $B_H \sqrt{3}$
71. Two magnets of equal mass are joined at 90° each other as shown in figure. Magnet $N_1 S_1$ has a magnetic moment $\sqrt{3}$ times that of $N_2 S_2$. The arrangement is pivoted so that it is free to rotate in horizontal plane. When in equilibrium, what angle should $N_1 S_1$ make with magnetic meridian?



- a) 75° b) 60° c) 30° d) 45°
72. Two normal uniform magnetic fields contain a magnetic needle making an angle 60° with F . Then the ratio of $\frac{F}{H}$ is
 a) $1 : 2$ b) $2 : 1$ c) $\sqrt{3} : 1$ d) $1 : \sqrt{3}$
73. A short bar magnet experiences a torque of magnitude 0.64 J. When it is placed in a uniform magnetic field of 0.32 T, taking an angle of 30° with the direction of the field. The magnetic moment of the magnet is
 a) 1 Am^2 b) 4 Am^2 c) 6 Am^2 d) None of these
74. A circular loop of radius 0.0157 m carries a current of 2.0 A. The magnetic field at the center of the loop is $[\mu_0 = 4\pi \times 10^{-7} \text{ Wb} - \text{A}^{-1}\text{m}^{-1}]$
 a) $1.57 \times 10^{-5} \text{ Wb} - \text{m}^2$
 b) $8.0 \times 10^{-5} \text{ Wb} - \text{m}^2$
 c) $2.0 \times 10^{-5} \text{ Wb} - \text{m}^2$
 d) $3.14 \times 10^{-5} \text{ Wb} - \text{m}^2$
75. A magnet is suspended in such a way that it oscillates in the horizontal plane. It makes 20 oscillations per minute at a place where dip angle is 30° and 15 oscillations per minute at a place where dip angle 60° . The ratio of total earth's magnetic field at the two places is
 a) $3\sqrt{3} : 8$ b) $16 : 9\sqrt{3}$ c) $4 : 9$ d) $2\sqrt{3} : 9$
76. A certain amount of current when flowing in a properly set tangent galvanometer, produces a deflection of 45° . If the current be reduced by a factor of $\sqrt{3}$, the deflection would
 a) Decrease by 30° b) Decrease by 15° c) Increase by 15° d) Increase by 30°
77. The magnetic field of earth is due to
 a) Motion and distribution of some material in and outside the earth
 b) Interaction of cosmic rays with the current of earth
 c) A magnetic dipole buried at the centre of the earth
 d) Induction effect of the sun
78. A bar magnet is placed north-south with its north pole due north. The points of zero magnetic field will be in which direction from center of magnet
 a) North and south b) East and west
 c) North-east and south-west d) North-east and south-east
79. A deflection magnetometer is adjusted in the usual way. When a magnet is introduced, the deflection observed is θ , and the period of oscillation of the needle in the magnetometer is T . When the magnet is removed, the period of oscillation is T_0 . The relation between T and T_0 is
 a) $T^2 = T_0^2 \cos \theta$ b) $T = T_0 \cos \theta$ c) $T = \frac{T_0}{\cos \theta}$ d) $T^2 = \frac{T_0^2}{\cos \theta}$
80. A compass needle placed at a distance r from a short magnet in a tan A position shows a deflection of 60° . If the distance is increased to $r(2)^{1/3}$, the deflection of compass needle is
 a) 30° b) 60° c) 45° d) 0°
81. At the magnetic poles of the earth, a compass needle will be
 a) Vertical b) Bent slightly
 c) Horizontal d) Inclined at 45° to the horizontal
82. The length of magnet is large compared to its width and breadth. The time period of its oscillation in a vibration magnetometer is 2 s. The magnet is cut along its length into three equal parts and these parts are then placed on each other with their like poles together. The time period of this combination will be

- a) $\frac{2}{\sqrt{3}}s$ b) $\frac{2}{3}s$ c) $2\sqrt{3}s$ d) $2s$
83. Water is
a) Diamagnetic b) Paramagnetic c) Ferromagnetic d) None of these
84. The ratio of magnetic moments of two bar magnets is 13 : 5. These magnets held together in a vibration magnetometer oscillate with 15 oscillations per minute in earth's magnetic field with like poles together. What will be the frequency of oscillations of system if unlike poles are together
a) 10 oscillations/min b) 15 oscillations/min c) 12 oscillations/min d) $\frac{75}{13}$ oscillations/min
85. Curie temperature is the one above which
a) Paramagnetic substance changes of ferromagnetic
b) Paramagnetic changes to diamagnetic
c) Diamagnetic changes to paramagnetic
d) Ferromagnetic changes to paramagnetic
86. Time period for a magnet is T . If it is divided in four equal parts along its axis and perpendicular to its axis as shown then time period for each part will be
- 
- a) $4T$ b) $T/4$ c) $T/2$ d) T
87. A tangent galvanometer is connected directly to an ideal battery. If the number of turns in the coil is doubled, the deflection will
a) Increase b) Decrease
c) Remain unchanged d) Either increase or decrease
88. Curie-Weiss law is obeyed by iron
a) At Curie temperature only b) At all temperatures
c) Below Curie temperature d) Above Curie temperature
89. The susceptibility of a paramagnetic material is K at 27°C . At what temperature will its susceptibility be $K/2$?
a) 600°C b) 287°C c) 54°C d) 327°C
90. Two bar magnets with magnetic moments $2M$ and M are fastened together at right angles to each other at their centres to form a crossed system, which can rotate freely about a vertical axis through the centre. The crossed system sets in earth's magnetic field with magnet having magnetic moment $2M$ making an angle θ with the magnetic meridian such that
a) $\theta = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$ b) $\theta = \tan^{-1}(\sqrt{3})$ c) $\theta = \tan^{-1}\left(\frac{1}{2}\right)$ d) $\theta = \tan^{-1}\left(\frac{3}{4}\right)$
91. When a magnet is placed vertically on horizontal board, number of neutral points obtained on the board is
a) Four b) Three c) Two d) One
92. Two similar bar magnets P and Q , each of magnetic moment M , are taken, If P is cut along its axial line and Q is cut along its equatorial line, all the four pieces obtained have
a) Equal pole strength b) Magnetic moment $\frac{M}{4}$ c) Magnetic moment $\frac{M}{2}$ d) Magnetic moment M
93. When a piece of a ferromagnetic substance is put in a uniform magnetic field, the flux density inside it is four times the flux density away from the piece. The magnetic permeability of the material (in N/A^2) is
a) 1 b) 2 c) 3 d) 4
94. At a certain place, a magnet makes 30 oscillations per min. At another place where the magnetic field is double, its time period will be
a) $4s$ b) $2s$ c) $1/2s$ d) $\sqrt{2}s$
95. Substances in which the magnetic moment of a single atom is not zero, is known as

- a) Diamagnetism b) Ferromagnetism
c) Paramagnetism d) Ferrimagnetism

96. A bar magnet has a magnetic moment equal to $5 \times 10^{-5} \text{ Wb-m}$. It is suspended in a magnetic field which has a magnetic induction B equal to $8\pi \times 10^{-4} \text{ T}$. The magnet vibrates with a period of vibration equal to 15 s. The moment of inertia of magnet is
a) $4.54 \times 10^4 \text{ kg - m}^2$ b) $4.54 \times 10^{-5} \text{ kg - m}^2$ c) $4.54 \times 10^{-4} \text{ kg - m}^2$ d) $4.54 \times 10^5 \text{ kg - m}^2$

97. What happens to the force between magnetic poles when their pole strength and the distance between them are both doubled
a) Force increases to two times the previous value
b) No change
c) Force decreases to half the previous value
d) Force increases to four times the previous value

98. The vertical component of earth's magnetic field is zero at or The earth's magnetic field always has a vertical component except at the
a) Magnetic poles b) Geographical poles c) Every place d) Magnetic equator

99. A bar magnet is held at right angles to a uniform magnetic field. The couple acting on the magnet is to be halved by rotation it from this position. The angle of rotation is
a) 60° b) 45° c) 30° d) 75°

100. The angle of dip at a place on the earth gives
a) The horizontal component of the earth's magnetic field
b) The location of the geographic meridian
c) The vertical component of the earth's field
d) The direction of the earth's magnetic field

101. The magnetic lines of force inside a bar magnet
a) Are from north-pole to south-pole of the magnet
b) Do not exist
c) Depend upon the area of cross-section of the bar magnet
d) Are from south-pole to north-pole of the magnet

102. A current loop placed in a magnetic field behaves like a
a) Magnetic dipole b) Magnetic substance c) Magnetic pole d) All are true

103. Two small bar magnets are placed in a line with like poles facing each other at a certain distance d apart. If the length of each magnet is negligible as compared to d , the force between them will be inversely proportional to
a) d b) d^2 c) $\frac{1}{d^2}$ d) d^4

104. Magnetic dipole moment is a
a) Scalar quantity b) Vector quantity c) Constant quantity d) None of these

105. The materials suitable for making electromagnets should have
a) High retentivity and high coercivity b) Low retentivity and low coercivity
c) High retentivity and low coercivity d) Low retentivity and high coercivity

106. Isogonic lines on magnetic map will have
a) Zero angle of dip b) Zero angle of declination
c) Same angle of declination d) Same angle of dip


107. The time period of a vibration magnetometer is T_0 . Its magnet is replaced by another magnet whose moment of inertia is 3 times and magnetic moment is $1/3$ of the initial magnet. The time period now will be
a) $3T_0$ b) T_0 c) $T_0/\sqrt{3}$ d) $T_0/3$

108. If magnetic lines of force are drawn by keeping magnet vertical, then number of neutral points will be
a) One b) Two c) Four d) Five

109. The magnetic moment of a magnet is $0.1 \text{ amp} \times \text{m}^2$. It is suspended in a magnetic field of intensity $3 \times 10^{-4} \text{ Wbm}^{-2}$. The couple acting upon it when deflected by 30° from the magnetic field is

- a) $1 \times 10^{-5} \text{ N m}$ b) $1.5 \times 10^{-5} \text{ N m}$ c) $2 \times 10^{-5} \text{ N m}$ d) $2.5 \times 10^{-5} \text{ N m}$

110. The variation of magnetic susceptibility (χ) with magnetising field for a paramagnetic substance is



111. Due to the earth's magnetic field, charged cosmic ray particles

- a) Require greater kinetic energy to reach the equator than the poles
b) Require less kinetic energy to reach the equator than the poles
c) Can never reach the equator
d) Can never reach the poles


112. Magnetic lines of force due to a bar magnet do not intersect because

- a) A point always has a single net magnetic field
b) The lines have similar charges and so repel each other
c) The lines always diverge from a single point
d) The lines need magnetic lenses to be made to intersect

113. Tangent galvanometer is used to measure

- a) Steady currents b) Current impulses
c) Magnetic moments of bar magnets d) Earth's magnetic field

114. The figure illustrates how B , the flux density inside a sample of unmagnetised ferromagnetic material, varies with B_0 , the magnetic flux density in which the sample is kept. For the sample to be suitable for making a permanent magnet



- a) OQ should be large, OR should be small
b) OQ and OR should both be large
c) OQ should be small and OR should be large
d) OQ and OR should both be small

115. A bar magnet of magnetic moment 10^4 J/T is free to rotate in a horizontal plane. The work done in rotating the magnet slowly from a direction parallel to a horizontal magnetic field of $4 \times 10^{-5} \text{ T}$ to a direction 60° from the field will be

- a) 0.2 J b) 2.0 J c) 4.18 J d) $2 \times 10^2 \text{ J}$

116. Two like magnetic poles of strength 10 and 45 SI units are separated by a distance 30 cm. The intensity of magnetic field is zero on the line joining them

- a) At a point 10 cm from the stronger pole b) At a point 20 cm from the stronger pole
c) At the mid-point d) At infinity


117. Domain formation is the necessary feature of

- a) Ferromagnetism b) Paramagnetism c) Diamagnetism d) All of these

118. A short bar magnet placed with its axis at 30° with a uniform external magnetic field of 0.16 tesla experiences a torque of magnitude 0.032 J. The magnetic moment of bar magnet will be

- a) 0.23 J/T b) 0.40 J/T c) 0.80 J/T d) Zero

119. Some equipotential surfaces of the magnetic scalar potential are shown in the figure. Magnetic field at a point in the region is



- a) $10^{-4}T$ b) $2 \times 10^{-4}T$ c) $0.5 \times 10^{-4}T$ d) None of these
120. A bar magnet when placed at an angle of 30° to the direction of magnetic field induction of $5 \times 10^{-2} T$, experiences a moment of couple $25 \times 10^{-6} N - m$. If the length of the magnet is 5 cm, its pole strength is
a) $2 \times 10^{-2} A-m$ b) $5 \times 10^{-2} A-m$ c) 2 A-m d) 5 A-m
121. A vibration magnetometer consists of two identical bar magnets placed one over the other such that they are perpendicular and bisect each other. The time period of oscillation in a horizontal magnetic field is $2^{2/5}s$. One of the magnets is removed and if the other magnet oscillates in the same field, then the time period in second is
a) $2^{1/4}$ b) $2^{1/2}$ c) 2 d) 4
122. If a magnet is hanged with its magnetic axis then it stops in
a) Magnetic meridian b) Geometric meridian c) Angle of dip d) None of these
123. Needles N_1, N_2 and N_3 are made of a ferromagnetic, a paramagnetic and a diamagnetic substance respectively. A magnet when brought close to them will
a) Attract N_1 and N_2 strongly but repel N_3
b) Attract N_1 strongly, N_2 weakly and repel N_3 weakly
c) Attract N_1 strongly, but repel N_2 and N_3 weakly
d) Attract all three of them
124. Two lines of force due to a bar magnet
a) Intersect at the neutral point
b) Intersect near the poles of the magnet
c) Intersect on the equatorial axis of the magnet
d) Do not intersect at all
125. At a certain place, the angle of dip is 30° and the horizontal component of earth's magnetic field is 0.50 oersted. The earth's total magnetic field (in oersted) is
a) $\sqrt{3}$ b) 1 c) $\frac{1}{\sqrt{3}}$ d) $\frac{1}{2}$
126. A vibration magnetometer is placed at south pole, then the time period will be
a) Zero b) Infinity
c) Same as at magnetic equator d) Same as at any other place on earth
127. Iron would become paramagnetic at about
a) $200^\circ C$ b) $400^\circ C$ c) $600^\circ C$ d) $800^\circ C$
128. The earth's magnetic field at a certain place has a horizontal component 0.3 gauss and the total strength 0.5 gauss. The angle of dip is
a) $\tan^{-1} \frac{3}{4}$ b) $\sin^{-1} \frac{3}{4}$ c) $\tan^{-1} \frac{4}{3}$ d) $\sin^{-1} \frac{3}{5}$
129. A coil in the shape of an equilateral triangle of side 0.02 m is suspended from its vertex such that it is hanging in a vertical plane between the pole pieces of permanent magnet producing a uniform field of $5 \times 10^{-2} T$. If a current of 0.1 A is passed through the coil, what is the couple acting
a) $5\sqrt{3} \times 10^{-7} N - m$ b) $5\sqrt{3} \times 10^{-10} N - m$ c) $\frac{\sqrt{3}}{5} \times 10^{-7} N - m$ d) None of these
130. Curie's law can be written as
a) $\chi \propto (T - T_c)$ b) $\chi \propto \frac{1}{T - T_c}$ c) $\chi \propto \frac{1}{T}$ d) $\chi \propto T$
131. Two bar magnets of the same mass, same length and breadth but having magnetic moments M and $3M$ are joined together pole for pole and suspended by a string.


The time period of assembly in a magnetic field of strength H is 3 s. If now the polarity of one of the magnets is reversed and the combination is again made to oscillate in the same field, the time of oscillation is

- a) 3 s b) $3\sqrt{3}$ s c) $3/\sqrt{3}$ s d) 6 s
132. A magnet of magnetic moment M is rotated through 360° in a magnetic field H . The work done will be
a) MH b) $2MH$ c) $2\pi MH$ d) Zero
133. A coil of 50 turns and area $1.25 \times 10^{-3} \text{ m}^2$ is pivoted about a vertical diameter in a uniform horizontal magnetic field and carries a current of 2 A. When the coil is held with its plane is $N - S$ of 2A. When the coil is held with its plane in $N - S$ direction, it experience a couple of 0.04 N-m; and when its plane is $E - W$, the corresponding couple is 0.03 N-m. The magnetic induction is
a) 0.2 T b) 0.3 T c) 0.4 T d) 0.5 T
134. A short bar magnet has a length $2l$ and a magnetic moment 10 Am^2 . Find the magnetic field at a distance of $z = 0.1 \text{ m}$ from its centre on the axial line. Here, l is negligible as compared to z .
a) $2 \times 10^{-3} \text{ T}$ b) $3 \times 10^{-3} \text{ T}$ c) $1 \times 10^{-3} \text{ T}$ d) $4 \times 10^{-3} \text{ T}$
135. Lines which represent places of constant angle of dip are called
a) Isobaric lines b) Isogonic lines c) Isoclinic lines d) Isodynamic lines
136. Two identical short bar magnets, each having magnetic moment M , are placed a distance of $2d$ apart with axes perpendicular to each other in a horizontal plane. The magnetic induction at a point midway between them is
a) $\frac{\mu_0}{4\pi}(\sqrt{2})\frac{M}{d^3}$ b) $\frac{\mu_0}{4\pi}(\sqrt{3})\frac{M}{d^3}$ c) $\left(\frac{2\mu_0}{\pi}\right)\frac{M}{d^3}$ d) $\frac{\mu_0}{4\pi}(\sqrt{5})\frac{M}{d^3}$
137. The distance of two points on the axis of a magnet from its centre is 10 cm and 20 cm respectively. The ratio of magnetic intensity at these points is 12.5 : 1. The length of the magnet will be
a) 5 cm b) 25 cm c) 10 cm d) 20 cm
138. The time period of oscillation of a freely suspended bar magnet with usual notations is given by
a) $T = 2\pi\sqrt{\frac{I}{MB_H}}$ b) $T = 2\pi\sqrt{\frac{MB_H}{I}}$ c) $T = \sqrt{\frac{I}{MB_H}}$ d) $T = 2\pi\sqrt{\frac{B_H}{MI}}$
139. The force between two magnetic poles is F . If the distance between the poles and pole strengths of each pole are doubled, then the force experienced is
a) $2F$ b) $\frac{F}{2}$ c) $\frac{F}{4}$ d) F
140. When a ferromagnetic material is heated to temperature above its curie point, the material
a) Is permanently magnetized b) Remains ferromagnetic
c) Behaves like a diamagnetic material d) Behaves like a paramagnetic material
141. A magnetic needle is kept in a non-uniform magnetic field. It experiences force and torque both due to unequal forces acting on poles.
a) A torque but not a force b) Neither a force nor a torque
c) A force and a torque d) A force but not a torque
142. The value of the horizontal component of the earth's magnetic field and angle of dip are $1.8 \times 10^{-5} \text{ weber/m}^2$ and 30° respectively at some place. The total intensity of earth's magnetic field at that place will be
a) $2.08 \times 10^{-5} \text{ weber/m}^2$ b) $3.67 \times 10^{-5} \text{ weber/m}^2$
c) $3.18 \times 10^{-5} \text{ weber/m}^2$ d) $5.0 \times 10^{-5} \text{ weber/m}^2$
143. The torque on a bar magnet due to the earth's magnetic field is maximum when the axis of the magnet is
a) Perpendicular to the field of the earth
b) Parallel of the vertical component of the earth's field
c) At an angle of 33° with respect $N-S$ direction
d) Along the North-South ($N-S$) direction

144. If the magnetic is cut into four equal parts such that their lengths and breadths are equal. Pole strength of each part is

- a) m b) $m/2$ c) $m/4$ d) $m/8$

145. Two equal bar magnets are kept as shown in the figure. The direction of resultant magnetic field, indicated by arrow head at the point P is (approximately)



- a) \longrightarrow b) \nearrow c) \searrow d) \uparrow

146. A bar magnet of length 10 cm and having pole strength equal to 10^{-3} Wb is kept in a magnetic field having magnetic induction B equal to $4\pi \times 10^{-3}$ T. It makes an angle of 30° with the direction of magnetic induction. The value of the torque acting on the magnet is

- a) 0.5 Nm b) $2\pi \times 10^{-5}$ Nm c) $\pi \times 10^{-5}$ Nm d) 0.5×10^{-5} Nm

147. The magnetising field required to be applied in opposite direction to reduce residual magnetism to zero is called

- a) Coercivity b) Retentivity c) Hysteresis d) None of these


148. A paramagnetic liquid is taken in a U-tube and arranged so that one of its limbs is kept between pole pieces of the magnet. The liquid level in the limb

- a) Goes down b) Rises up
c) Remains same d) First goes down and then rises

149. A bar magnet is oscillating in the Earth's magnetic field with a period T . What happens to its period of motion if its mass is quadrupled?

- a) Motion remains SHM with time period $= T/2$ b) Motion remains SHM and period remains nearly constant
c) Motion remains SHM with time period $= 2T$ d) Motion remains SHM with time period $= 4T$

150. For substances hysteresis (B - H) curves are given as shown in figure. For making temporary magnet which of the following is best



151. A magnet makes 5 oscillations per min in $B = 0.3 \times 10^{-4}$ T. By what amount should the field be increased so that number of oscillations is 10 in the same time?

- a) 0.3×10^{-4} T b) 0.6×10^{-4} T c) 0.9×10^{-4} T d) 1.2×10^{-4} T

152. The length of a magnet is large compared to its width and breadth. The time period of its oscillation in a vibration magnetometer is 2 s. The magnet is cut along its length into three equal parts and three parts are then placed on each other with their like poles together. The time period of this combination will be

- a) 2 s b) $2/3$ s c) $2\sqrt{3}$ s d) $2/\sqrt{3}$ s

153. The mass of specimen of a ferromagnetic material is 0.6 kg and the density is $7.8 \times 10^3 \text{ kg m}^{-3}$. If the area of hysteresis loop of alternating magnetizing field of frequency 50 Hz is 0.722 MKS units, then hysteresis loss per second will be

- a) 27.77×10^{-5} J b) 2.777×10^{-5} J c) 27.27×10^{-4} J d) 27.77×10^{-6} J

154. A small bar magnet has a magnetic moment 1.2 A-m^2 . The magnetic field at a distance 0.1 m on its axis will be: ($\mu_0 = 4\pi \times 10^{-7} \text{ T-m/A}$)

- a) $1.2 \times 10^{-4} \text{ T}$ b) $2.4 \times 10^{-4} \text{ T}$ c) $2.4 \times 10^4 \text{ T}$ d) $1.2 \times 10^4 \text{ T}$

155. Unit of magnetic flux density (or magnetic induction) is

- PHONE NO: 8583042324 Page | 14

169. The magnetic field due to a small magnetic dipole of magnetic moment M , at distance r from the centre on the equatorial line is given by (in M.K.S system)

- a) $\frac{\mu_0}{4\pi} \times \frac{M}{r^2}$ b) $\frac{\mu_0}{4\pi} \times \frac{M}{r^3}$ c) $\frac{\mu_0}{4\pi} \times \frac{2M}{r^2}$ d) $\frac{\mu_0}{4\pi} \times \frac{2M}{r^3}$


170. A uniform magnetic needle is suspended from its centre by a thread. Its upper end is now loaded with a mass 50 mg, and the needle becomes horizontal. If the strength of each pole is 98.1 ab-amp-cm and $g=981 \text{ cms}^{-2}$, then the vertical component of earth's magnetic induction is

- a) 0.50 G b) 0.25 G c) 0.005 G d) 0.05 G

171. The magnetism of a magnet is due to

- a) The earth b) Cosmic rays
c) The spin motion of electrons d) Pressure of big magnet inside the earth

172. Which curve may best represent the current deflection in a tangent galvanometer



- a) A b) B c) C d) D

173. The period of oscillation of a vibration magnetometer depends on which of the following factors

Where I is the moment of inertia of the magnet about the axis of suspension, M is the magnetic moment of the magnet and H is the external magnetic field

- a) I and M only b) M and H only c) I and H only d) I , M and H only

174. Two short magnets AB and CD are in the X - Y plane and are parallel to X -axis and co-ordinates of their centers respectively are $(0, 2)$ and $(2, 0)$. Line joining the north-south poles of CD is opposite to that of AB and lies along the positive X -axis. The resultant field induction due to AB and CD at a point $P(2, 2)$ is $100 \times 10^{-7} \text{ T}$. When the poles of the magnet CD are reversed, the resultant field induction is $50 \times 10^{-7} \text{ T}$. The value of magnetic moments of AB and CD (in Am^2) are

- a) 300; 200 b) 600; 400 c) 200; 100 d) 300; 150

175. The time period of a bar magnet suspended horizontally in the earth's magnetic field and allowed to oscillate

- a) Is directly proportional to the square root of its mass
b) Is directly proportional to its pole strength
c) Is inversely proportional to its magnetic moment
d) Decrease if the length increases but pole strength remains same

176. The needle of a deflection galvanometer shows a deflection of 60° due to a short bar magnet at a certain distance in tan A position. If the distance is double the deflection is

- a) $\sin^{-1} \left[\frac{\sqrt{3}}{8} \right]$ b) $\cos^{-1} \left[\frac{\sqrt{3}}{8} \right]$ c) $\tan^{-1} \left[\frac{\sqrt{3}}{8} \right]$ d) $\cot^{-1} \left[\frac{\sqrt{3}}{8} \right]$

177. Aurora Borealis is a luminous electrical discharge in the upper layers of the atmosphere, which is visible more frequently in


- a) Polar regions b) Equator
c) Lunar eclipse d) Regions of earth's magnetic poles

178. The direction of the null points is on the equatorial line of a bar magnet, when the north pole of the magnet is pointing

- a) North b) South c) East d) West

179. Each atom of an iron bar ($5\text{cm} \times 1\text{cm} \times 1\text{cm}$) has a magnetic moment $1.8 \times 10^{-23} \text{ Am}^2$. Knowing that the density of iron is $7.78 \times 10^3 \text{ kg m}^{-3}$, atomic weight is 56 and Avogadro's number of 6.02×10^{23} the magnetic moment of bar in the state of magnetic saturation will be

- a) 4.75 Am^2 b) 5.74 Am^2 c) 7.54 Am^2 d) 75.4 Am^2
180. Magnetic meridian is a
a) Point b) Horizontal plane c) Vertical plane d) Line along N-S
181. A very long magnet is placed vertically with one pole on the table. A neutral point was found at 20 cm from the pole. What is the pole strength if the vertical component of earth's field is $0.4 \times 10^{-4} \text{ Wbm}^{-2}$?
a) 16 A-m b) 8 A-m c) 4 A-m d) None of these
182. The only property possessed by ferromagnetic substance is
a) Hysteresis b) Susceptibility
c) Directional property d) Attracting magnetic substances
183. The mathematical equation for magnetic field lines of force is
a) $\vec{\nabla} \cdot \vec{B} = 0$ b) $\vec{\nabla} \cdot \vec{B} \neq 0$ c) $\vec{\nabla} \cdot \vec{B} > 0$ d) $\vec{\nabla} \cdot \vec{B} < 0$
184. The hysteresis cycle for the material of permanent magnet is
a) Short and wide b) Tall and narrow c) Tall and wide d) Short and narrow
185. An electron moving around the nucleus with an angular momentum l has a magnetic moment
a) $\frac{e}{m} l$ b) $\frac{e}{2m} l$ c) $\frac{2e}{m} l$ d) $\frac{e}{2\pi m} l$
186. The magnetic susceptibility of a material of a rod is 499. The absolute permeability of vacuum is $4\pi \times 10^{-7} \text{ HM}^{-1}$. The absolute permeability of the material of a rod is
a) $\pi \times 10^{-4} \text{ HM}^{-1}$ b) $2\pi \times 10^{-4} \text{ HM}^{-1}$ c) $3\pi \times 10^{-4} \text{ HM}^{-1}$ d) $4\pi \times 10^{-4} \text{ HM}^{-1}$
187. The work done in turning a magnet of magnetic moment ' M ' by an angle of 90° from the meridian is ' n ' times the corresponding work done to turn it through an angle at 60° , when ' n ' is given by
a) $1/2$ b) 2 c) $1/4$ d) 1
188. A magnet of magnetic moment 20 CGS units is freely suspended in a uniform magnetic field of intensity 0.3 CGS units. The amount of work done in deflecting it by an angle of 30° in CGS units is
a) 6 b) $3\sqrt{3}$ c) $3(2 - \sqrt{3})$ d) 3
189. The magnetic force required to demagnetize the material is
a) Retainivity b) Coercivity c) Energy loss d) Hysteresis
190. The angle between magnetic meridian and geographical meridian is known as
a) Magnetic dip b) Magnetic latitude
c) Magnetic Declination d) Magnetic longitude
191. An inductor of 10 mH shows 50 mH when operate with a core made of ferrite. The susceptibility of ferrite is
a) 5 b) 4 c) 3 d) None of these
192. A magnet 20 cm long with its poles concentrated at its ends is placed vertically with its north pole on the table. At a point due 20 cm south (magnetic) of the pole, a neutral point is obtained. If $H = 0.3 \text{ G}$, then the pole strength of the magnet is approximately
a) 185 ab-amp-cm b) 185 amp-m c) 18.5 ab-amp-cm d) 18.5 amp-cm
193. The time period of a freely suspended magnet is 2 sec. If it is broken in length into two equal parts and one parts is suspended in the same way, then its time period will be
a) 4 sec b) 2 sec c) $\sqrt{2} \text{ sec}$ d) 1 sec
194. A short bar magnet of magnetic moment 255 JT^{-1} is placed with its axis perpendicular to earth's field direction. At what distance from the center of the magnet, the resultant field is inclined at 45° with earth's field, $H = 0.4 \times 10^{-4} \text{ T}$?
a) 5 m b) 0.5m c) 2.5 m d) 0.25 m
195. A bar-magnet of moment of inertia $49 \times 10^{-2} \text{ kg} - \text{m}^2$ vibrate in a magnetic field of induction $0.5 \times 10^{-4} \text{ T}$. The time period of vibration is 8.8 s. The magnetic moment of the bar magnet is
a) $350 \text{ A} - \text{m}^2$ b) $490 \text{ A} - \text{m}^2$ c) $3300 \text{ A} - \text{m}^2$ d) $5000 \text{ A} - \text{m}^2$


196. A magnet performs 10 oscillations per minute in a horizontal plane at a place where the angle of dip is 45° and the total intensity is 0.707 CGS units. The number of oscillations per minute at a place where dip angle is 60° and total intensity is 0.5 CGS units will be
 a) 5 b) 7 c) 9 d) 11
197. There is no couple acting when two bar magnets are placed coaxially separated by a distance because
 a) There are no forces on the poles
 b) The force are parallel and their lines of action do not coincide
 c) The forces are perpendicular to each other
 d) The forces act along the same line
198. The points A and B are situated perpendicular to the axis of 2 cm long bar magnet at large distances x and $3x$ from the centre on opposite sides. The ratio of magnetic fields at A and B will be approximately equal to
 a) 27 : 1 b) 1 : 27 c) 9 : 1 d) 1 : 9
199. The correct relation is
 [Where B_H = Horizontal component of earth's magnetic field; B_V = Vertical component of earth's magnetic field and B = Total intensity of earth's magnetic field]
 a) $B = \frac{B_V}{B_H}$ b) $B = B_V \times B_H$ c) $|B| = \sqrt{B_H^2 + B_V^2}$ d) $B = B_H + B_V$
200. A short bar magnet with the north pole facing north forms a neutral point P in the horizontal plane. If the magnet is rotated by 90° in the horizontal plane, the net magnetic induction at P is (Horizontal component of earth's magnetic field = B_H)
 a) Zero b) $2 B_H$ c) $\frac{\sqrt{5}}{2} B_H$ d) $\sqrt{5} B_H$
201. At which place, earth's magnetism becomes horizontal
 a) Magnetic pole b) Geographical pole c) Magnetic meridian d) Magnetic equator
202. The angle of dip at a certain place on earth is 60° and the magnitude of earth's horizontal component of magnetic field is 0.26 G. The magnetic field at the place on earth is
 a) 0.13 G b) 0.26 G c) 0.52 G d) 0.65 G
203. The variation of magnetic susceptibility (χ) with absolute temperature T for a ferromagnetic is given in figure, by

204. At a certain place, horizontal component is $\sqrt{3}$ times the vertical component. The angle of dip at this place is
 a) Zero b) $\pi/3$ c) $\pi/6$ d) None of these
205. The resultant magnetic moment of neon atom will be
 a) Infinity b) μ_B c) Zero d) $\frac{\mu_B}{2}$
206. The horizontal component of the earth's magnetic field is 0.22 gauss and total magnetic field is 0.4 gauss. The angle of dip is
 a) $\tan^{-1}(1)$ b) $\tan^{-1}(\infty)$ c) $\tan^{-1}(1.518)$ d) $\tan^{-1}(\pi)$
207. A short magnet oscillates with a time period 0.1 s at a place where horizontal magnetic field is $24 \mu\text{T}$. A downward current of 18 A is established in a vertical wire 20 cm east of the magnet. The new time period of oscillator
 a) 0.1 s b) 0.089 s c) 0.076 s d) 0.057 s
208. The relative magnetic permeability of ferromagnetic materials is of the order of
 a) 10 b) 100 c) 1000 d) 10000

209. If the magnetic dipole moment of an atom of diamagnetic material, paramagnetic material and ferromagnetic material is denoted by μ_d, μ_p, μ_f respectively then
 a) $\mu_d \neq 0$ and $\mu_f \neq 0$ b) $\mu_p \neq 0$ and $\mu_f \neq 0$ c) $\mu_d \neq 0$ and $\mu_p \neq 0$ d) $\mu_d \neq 0$ and $\mu_p \neq 0$
210. As magnetising field on a ferromagnetic material is increased, its permeability
 a) Increases b) Decreases c) Remains constant d) Cannot say
211. A short bar magnet of magnetic moment 0.4 J T^{-1} is placed in a uniform magnetic field of 0.16 T . The magnet is in stable equilibrium when the potential energy is
 a) -0.082 J b) 0.064 J c) -0.064 J d) Zero

212. The material of permanent magnet has
 a) High retentivity, low coercivity b) Low retentivity, high coercivity
 c) Low retentivity, low coercivity d) High retentivity, high coercivity
213. The variation of magnetic susceptibility (χ) with temperature for a diamagnetic substance is best represented by figure



214. The period of oscillations of a magnet is 2 s . When it is magnetized that the pole strength is 4 times, its period will be
 a) 4 s b) 1 s c) 2 s d) $\frac{1}{2} \text{ s}$
215. A magnet of distance moment 2 J T^{-1} is aligned in the direction of magnetic field of 0.1 T . What is the net work done to bring, the magnet normal to the magnetic field?
 a) 0.1 J b) 0.2 J c) 1 J d) 2 J
216. The variation of intensity of magnetization (I) with respect to the magnetizing field (H) in a diamagnetic substance is described by the graph in figure.




- a) OD b) OC c) OB d) OA
217. A bar magnet having centre O has a length of 4 cm . Point P_1 is in the broad side-on and P_2 is in the end side-on position with $OP_1 = OP_2 = 10 \text{ metres}$. The ratio of magnetic intensities H at P_1 and P_2 is
 a) $H_1 : H_2 = 16 : 100$ b) $H_1 : H_2 = 1 : 2$ c) $H_1 : H_2 = 2 : 1$ d) $H_1 : H_2 = 100 : 16$
218. At a place, if one earth's horizontal and vertical components of magnetic fields are equal, then the angle of dip will be
 a) 30° b) 90° c) 45° d) 0°
219. Two magnets are held together in a vibration magnetometer and are allowed to oscillate in the earth's magnetic field. With like poles together, 12 oscillations per minute are made but for unlike poles together only 4 oscillations per minute are executed. The ratio of their magnetic moment is
 a) $3 : 1$ b) $1 : 3$ c) $3 : 5$ d) $5 : 4$
220. The relation between B, H and I in SI unit is
 a) $B = \mu_0(H + I)$ b) $B = H + 4\mu I$ c) $H = \mu_0(B + I)$ d) None of these
221. Two short magnets having magnetic moments in the ratio $27 : 8$, when placed on opposite sides of a deflection magnetometer, produce no deflection. If the distance of the weaker magnet is 0.12 m from the centre of deflection magnetometer, the distance of the stronger magnet from the centre is
 a) 0.06 m b) 0.08 m c) 0.12 m d) 0.18 m
222. Ferromagnetic materials used in a transformer must have
 a) Low permeability and high hysteresis loss b) High permeability and low hysteresis loss

- c) High permeability and high hysteresis loss d) Low permeability and low hysteresis loss
223. A current carrying small loop behaves like a small magnet. If A be its area and M its magnetic moment, the current in the loop will be
 a) M/A b) A/M c) MA d) AM^2
224. A bar magnet 20 cm in length is placed with its south pole towards geographic north. The neutral points are situated at a distance of 40 cm from centre of the magnet.
 If horizontal component of earth's field = 3.2×10^{-5} T, then pole strength of magnet is
 a) 5 AM b) 10 AM c) 45 AM d) 20 AM
225. A compass needle whose magnetic moment is $60 \text{ amp} \times \text{m}^2$ pointing geographical north at a certain place, where the horizontal component of earth's magnetic field is $40 \mu \text{ Wb/m}^2$, experiences a torque $1.2 \times 10^{-3} \text{ N} \times \text{m}$. What is the declination at this place
 a) 30° b) 45° c) 60° d) 25°
226. Two small magnets each of magnetic moment 10 A-m^2 are placed in end-on position 0.1 m apart from their centres. The force acting between them is
 a) $0.6 \times 10^7 \text{ N}$ b) $0.06 \times 10^7 \text{ N}$ c) 0.6 N d) 0.06 N
227. In a vibration magnetometer, the time period of a bar magnet oscillating in horizontal component of earth's magnetic field is 2 s. When a magnet is brought near and parallel to it, the time period reduces to 1 s. The ratio $\frac{F}{H}$ of the fields, F due to magnet and H , the horizontal component will be
 a) $\sqrt{3}$ b) $\frac{1}{\sqrt{3}}$ c) $\frac{1}{3}$ d) 3
228. Which of the following is the most suitable material for making permanent magnet
 a) Steel b) Soft iron c) Copper d) Nickel
229. The length of a magnet is large compared to its width and breadth. The time period of its oscillation in a vibration magnetometer is T . The magnet is cut along its length into six parts and these parts are then placed together as shown in the figure. The time period of this combination will be




- a) T b) $\frac{T}{\sqrt{3}}$ c) $\frac{T}{2\sqrt{3}}$ d) Zero
230. In sum and difference method vibration magnetometer, the time period is more if
 a) Similar poles of both magnets are on same sides
 b) Opposite poles of both magnets are on same sides
 c) Both magnets are perpendicular to each other
 d) Nothing can be said
231. At two places A and B using vibration magnetometer, a magnet vibrates in a horizontal plane and its respective periodic time are 2 sec and 3 sec and at these places the earth's horizontal components are H_A and H_B respectively. Then the ratio between H_A and H_B will be
 a) 9 : 4 b) 3 : 2 c) 4 : 9 d) 2 : 3
232. For ferromagnetic material, the relative permeability (μ_r) versus magnetic intensity (H) has the following shape



233. The period of oscillations of a magnetic needle in a magnetic field is 1.0 sec. If the length of the needle is halved by cutting it, the time period will be

- a) 1.0 sec b) 0.5 sec c) 0.25 sec d) 2.0 sec


234. A curve between magnetic moment and temperature of magnet is



235. A magnetic dipole is placed at right angles to the direction of lines of force of magnetic induction B . If it is rotated through an angle of 180° , then the work done is

- a) MB b) $2MB$ c) $-2MB$ d) Zero

236. The given figure represents a material which is



- a) Paramagnetic b) Diamagnetic c) Ferromagnetic d) None of these

237. Which of the following statements is not true

- a) While taking reading of tangent galvanometer, the plane of the coil must be set at right angles to the earth's magnetic meridian
b) A short magnet is used in a tangent galvanometer since a long magnet would be heavy and may not easily move
c) Measurement with the tangent galvanometer will be more accurate when the deflection is around 45°
d) A tangent galvanometer can not be used in the polar region

238. Two tangent galvanometers A and B have coils of radii 8 cm and 16 cm respectively and resistance $8\ \Omega$ each. They are connected in parallel with a cell of emf 4 V and negligible internal resistance. The deflections produced in the tangent galvanometers A and B are 30° and 60° respectively. If A has 2 turns, then B must have

- a) 18 turns b) 12 turns c) 6 turns d) 2 turns

239. A magnetic needle lying parallel to a magnetic field requires W units of work to turn it through 60° . The torque required to keep the needle in this position will be

- a) $2W$ b) W c) $\frac{W}{\sqrt{2}}$ d) $\sqrt{3}W$

240. The time period of a freely suspended magnet does not depend upon





- a) Length of magnet b) Pole strength of magnet
c) Horizontal component of earth's field d) Length of the suspension


241. Magnetic field intensity is defined as

- a) Magnetic moment per unit volume
b) Magnetic induction force acting on a unit magnetic pole
c) Number of lines of force crossing per unit area
d) Number of lines of force crossing per unit volume


242. A magnetic needle is placed on a cork floating in a still lake in the northern hemisphere. Does the needle together with the cork move towards the north of the lake

- a) Yes
b) No
c) May be or may not be move
d) Nothing can be said
243. The field due to a magnet at a distance R from the centre of the magnet is proportional to
a) R^2 b) R^3 c) $1/R^2$ d) $1/R^3$
244. If the angles of dip at two places are 30° and 45° respectively, then the ratio of horizontal components of earth's magnetic field at the two places will be
a) $\sqrt{3}:\sqrt{2}$ b) $1:\sqrt{2}$ c) $1:\sqrt{3}$ d) $1:2$
245. Diamagnetic substance are
a) Feebly attracted by magnets b) Strongly attracted by magnets
c) Feebly repelled by magnets d) Strongly repelled by magnets
246. Magnets A and B are geometrically similar but the magnetic moment of A is twice that of B . If T_1 and T_2 be the time periods of the oscillation when their like poles and unlike poles are kept together respectively, then $\frac{T_1}{T_2}$ will be
a) $\frac{1}{3}$ b) $\frac{1}{2}$ c) $\frac{1}{\sqrt{3}}$ d) $\sqrt{3}$
247. In the case of bar magnet, lines of magnetic induction
a) Start from the north pole and end at the south pole
b) Run continuously through the bar and outside
c) Emerge in circular paths from the middle of the bar
d) Are produced only at the north pole like rays of light from a bulb
248. If the total magnetic field due to earth is 28 Am^{-1} ; then the total magnetic induction due to earth is
a) 28 T b) $280 \text{ ab} - \text{Acm}^{-1}$ c) 0.352 G d) 0.325 T
249. Intensity of magnetic field due to earth at a point inside a hollow steel box is
a) Less than outside b) More than outside c) Same d) Zero
250. Two magnets A and B are identical in mass, length and breadth but have different magnetic moments. In a vibration magnetometer, if the time period of B is twice the time period of A . The ratio of the magnetic moments M_A/M_B of the magnets will be
a) $1/2$ b) 2 c) 4 d) $1/4$
251. Which one of the following is a non-magnetic substance
a) Iron b) Nickel c) Cobalt d) Brass
252. The relative permeability is represented by μ and the susceptibility is denoted by χ for a magnetic substance. Then for a paramagnetic substance
a) $\mu_r < 1, \chi < 0$ b) $\mu_r < 1, \chi > 0$ c) $\mu_r > 1, \chi < 0$ d) $\mu_r > 1, \chi > 0$
253. The time period of a freely suspended bar magnet in a field is 2 s . It is cut into two equal parts along its axis, then the time period is
a) 4 s b) 0.5 s c) 2 s d) 0.25 s
254. Due to a small magnet, intensity at a distance x in the end on position is 9 gauss . What will be the intensity at a distance $\frac{x}{2}$ on broad side on position
a) 9 gauss b) 4 gauss c) 36 gauss d) 4.5 gauss
255. For an isotropic medium B, μ, H and M are related as (where B, μ_0, H and M have their usual meanings in the context of magnetic material)
a) $(B - M) = \mu_0 H$ b) $M = \mu_0 (H + M)$ c) $H = \mu_0 (H + M)$ d) $B = \mu_0 (H + M)$
256. Which one of the following is not a characteristics of diamagnetism?
a) The diamagnetic materials are repelled by a bar magnet
b) The magnetic susceptibility of the materials is small and negative
c) The origin of dia magnetism is the spin of electrons

- d) The material move from a region of strong magnetic field to weak magnetic field
257. A superconductor exhibits perfect
- a) Ferrimagnetism b) Ferromagnetism c) Paramagnetism d) Diamagnetism
258. On applying an external magnetic field, to a ferromagnetic substance domains
- a) Align in the direction of magnetic field b) Align in the direction opposite to magnetic field
- c) Remain unaffected d) None of the above
259. The magnetic field of a small bar magnet varies in the following manner by the influence of a magnet placed at a large distance d .
- a) $\frac{1}{d}$ b) $\frac{1}{d^2}$ c) $\frac{1}{d^3}$ d) $\frac{1}{d^4}$
260. The correct $I - H$ curve for a paramagnetic material is represented by, figure.
- a)  b)  c)  d) 
261. The distance between the poles of a horse shoe magnet is 0.1 m and its pole strength is 0.01 amp-m . The induction of magnetic field at a point midway between the poles will be




- a) $2 \times 10^{-5} \text{ T}$ b) $4 \times 10^{-6} \text{ T}$ c) $8 \times 10^{-7} \text{ T}$ d) Zero
262. The magnetic field due to a short magnet at a point on its axis at distance $X \text{ cm}$ from the middle point of the magnet is 200 gauss . The magnetic field at a point on the neutral axis at a distance $X \text{ cm}$ from the middle of the magnet is
- a) 100 gauss b) 400 gauss c) 50 gauss d) 200 gauss
263. Two short bar magnets of equal dipole moment M are fastened perpendicularly at their centers, figure. The magnitude of resultant of two magnetic field at a distance d from the center on the bisector of the right angle is



- a) $\frac{\mu_0}{4\pi} \frac{2\sqrt{2}M}{d^3}$ b) $\frac{\mu_0}{4\pi} \frac{2M}{d^3}$ c) $\frac{\mu_0}{4\pi} \frac{M}{d^3}$ d) $\frac{\mu_0}{4\pi} \frac{2\sqrt{2}M}{d^3}$
264. The unit for molar susceptibility is
- a) m^3 b) kg-m^{-3} c) kg^{-1}m^3 d) No units
265. A bar magnet A of magnetic moment M_A is found to oscillate at a frequency twice that of magnet B of magnetic moment M_B when placed in a vibrating *magneto-meter*. We may say that
- a) $M_A = 2M_B$ b) $M_A = 8M_B$ c) $M_A = 4M_B$ d) $M_B = 8M_A$
266. A thin rectangular magnet suspended freely has a period of oscillation equal to T . Now it is broken into two equal halves (each having half of the original length) and one piece is made to oscillate freely in the same field. If its period of oscillation is T' , the ratio T'/T

- a) $\frac{1}{2\sqrt{2}}$ b) $\frac{1}{2}$ c) 2 d) $\frac{1}{4}$
267. The error in measuring the current with a tangent galvanometer is minimum when the deflection is about
a) 0° b) 30° c) 45° d) 60°
268. A magnet is suspended in the magnetic meridian with an untwisted wire. The upper end of wire is rotated through 180° to deflect the magnet by 30° from magnetic meridian. When this magnet is replaced by another magnet, the upper end of wire is rotated through 270° to deflect the magnet 30° from magnetic meridian. The ratio of magnetic moments of magnets is
a) 1 : 5 b) 1 : 8 c) 5 : 8 d) 8 : 5
269. When 2 amperes current is passed through a tangent galvanometer, it gives a deflection of 30° . For 60° deflection, the current must be
a) 1 amp b) $2\sqrt{3}$ amp c) 4 amp d) 6 amp
270. A bar magnet of magnetic moment 200 A-m^2 is suspended in a magnetic field of intensity 0.25 N/A-m . The couple required to deflect it through 30° is
a) 50 N-m b) 25 N-m c) 20 N-m d) 15 N-m
271. The angle of dip at a place is 37° and the vertical component of the earth's magnetic field is $6 \times 10^{-5} \text{ T}$. The earth's magnetic field at this place is ($\tan 37^\circ = 3/4$)
a) $7 \times 10^{-5} \text{ T}$ b) $6 \times 10^{-5} \text{ T}$ c) $5 \times 10^{-5} \text{ T}$ d) 10^{-4} T
272. The effect due to uniform magnetic field on a freely suspended magnetic needle is as follows
a) Both torque and net force are present b) Torque is present but no net force
c) Both torque and net force are absent d) Net force is present but not torque
273. If a diamagnetic substance is brought near north or south pole of a bar magnet, it is
a) Attracted by the poles
b) Repelled by the poles
c) Repelled by the north pole and attracted by the south pole
d) Attracted by the north pole and repelled by the south pole
274. When $\sqrt{3}$ ampere current is passed in a tangent galvanometer, there is a deflection of 30° in it. The deflection obtained when 3 amperes current is passed, is
a) 30° b) 45° c) 60° d) 75°
275. A small rod of bismuth is suspended freely between the poles of a strong electromagnet. It is found to arrange itself at right angles to the magnetic field. This observation establishes that bismuth is
a) Diamagnetic b) Paramagnetic c) Ferri-magnetic d) Antiferro-magnetic
276. Two identical thin bar magnets each of length l and pole strength m are placed at right angle to each other with north pole of one touching south pole of the other. Magnetic moment of the system is
a) ml b) $2ml$ c) $\sqrt{2}ml$ d) $\frac{1}{2}ml$
277. If a magnet of pole strength m is divided into four parts such that the length and width of each part is half that of initial one, then the pole strength of each part will be
a) $m/4$ b) $m/2$ c) $m/8$ d) $4m$
278. An example of a diamagnetic substance is
a) Aluminium b) Copper c) Iron d) Nickel
279. Demagnetisation of magnets can be done by
a) Rough handling b) Heating
c) Magnetising in the opposite direction d) All the above
280. Using a bar magnet P , a vibration magnetometer has time period 2seconds. When a bar Q (identical to P in mass and size) is placed on top of P , the time period is unchanged. Which of the following statements is true
a) Q is of non-magnetic material
b) Q is a bar magnet identical to P , and its north pole is placed on top of P 's north pole

- c) Q is of unmagnetized ferromagnetic material
 d) Nothing can be said about Q 's properties
281. The intensity of magnetic field is H and moment of magnet is M . The maximum potential energy is
 a) MH b) $2MH$ c) $3MH$ d) $4MH$
282. A frog can be levitated in magnetic field produced by a current in a vertical solenoid placed below the frog. This is possible because the body of the frog behaves as
 a) Paramagnetic b) Diamagnetic c) Ferromagnetic d) Anti-ferromagnetic
283. The period of oscillation of a bar magnet in a vibration magnetometer is 2 s. The period of oscillation of another bar magnet whose moment is 4 times that of 1st magnet is
 a) 4 s b) 1 s c) 2 s d) 0.5 s
284. A paramagnetic substance of susceptibility 3×10^{-4} is placed in a magnetic field of $4 \times 10^{-4} \text{ Am}^{-1}$. Then the intensity of magnetization in the units of Am^{-1} is
 a) 1.33×10^8 b) 0.75×10^{-8} c) 12×10^{-8} d) 14×10^{-8}
285. A vibration magnetometer consists of two identical bar magnets placed one over the other such that they are mutually perpendicular and bisect each other. The time period of combination is 4 s. If one of the magnets is removed, find the period of other
 a) 5 s b) 3.36 s c) 4.36 s d) 5.36 s
286. The vertical component of the earth's magnetic field is zero at a place where the angle of dip is
 a) 0° b) 45° c) 60° d) 90°
287. A magnet is suspended horizontally in the earth's magnetic field. When it is displaced and then released it oscillates in a horizontal plane with a period T . If a piece of wood of the same moment of inertia (about the axis of rotation) as the magnet is attached to the magnet, what would be the new period of oscillation of the system




- a) $\frac{T}{3}$ b) $\frac{T}{2}$ c) $\frac{T}{\sqrt{2}}$ d) $T\sqrt{2}$
288. The magnet can be completely demagnetized by
 a) Breaking the magnet into small pieces b) Heating it slightly
 c) Dropping it into ice cold water d) A reverse field of appropriate strength
289. The magnetic susceptibility is negative for
 a) Paramagnetic materials b) Diamagnetic materials
 c) Ferromagnetic materials d) Paramagnetic and ferromagnetic materials
290. The angle of dip at a certain place where the horizontal and vertical components of the earth's magnetic field are equal is
 a) 30° b) 90° c) 60° d) 45°
291. Let B_V and B_H be the vertical and horizontal components of earth's magnetic field at any point on earth. Near the north pole
 a) $B_V \gg B_H$ b) $B_V \ll B_H$ c) $B_V = B_H$ d) $B_V = B_H = 0$
292. In an experiment with vibration magnetometer, the value of $4\pi^2 I/T^2$ for a short bar magnet is observed as 36×10^{-4} . In the experiment with deflection magnetometer with the same magnet, the value of $4\pi d^3/2\mu_0$ is observed as $10^8/36$. The magnetic moment of the magnet used is
 a) 50 A-m b) 100 A-m c) 200 A-m d) 1000 A-m

293. A dip needle vibrates in the vertical plane perpendicular to magnetic meridian. The time period of vibration is found to be 2 s. The same needle is then allowed to vibrate in the horizontal plane and time period is again found to be 2s. Then the angle of dip is
 a) 0° b) 30° c) 45° d) 90°
294. Vibration magnetometer works on the principle of
 a) Torque acting on the bar magnet
 b) Force acting on the bar magnet
 c) Both the force and the torque acting on the bar magnet
 d) None of these
295. A magnetic dipole is placed in a uniform magnetic field. The net magnetic force on the dipole
 a) Is always zero b) Depends on the orientation of the dipole
 c) Can never be zero d) Depends on the strength of the dipole
296. The time period of a thin bar magnet in earth's magnetic field is T . If the magnet is cut into equal parts perpendicular to its length, the time period of each part in the same field will be
 a) $T/2$ b) $T/4$ c) $\sqrt{2} T$ d) $2 T$
297. If two identical bar magnets, each of length l , pole strength m and magnet moment M , are placed perpendicular to each other with their unlike poles in contact, the magnetic moment of the combination is
 a) $\frac{M}{\sqrt{2}}$ b) $lm(\sqrt{2})$ c) $2lm(\sqrt{2})$ d) $2M$
298. For a paramagnetic material, the dependence of the magnetic susceptibility X on the absolute temperature is given as
 a) $X \propto T$ b) $X \propto 1/T^2$ c) $X \propto 1/T$ d) Independent
299. The deflection magnetometer is most sensitive when deflection θ is
 a) Nearly zero b) Nearly 30° c) Nearly 45° d) Nearly 90°
300. There are four light-weight-rod samples, A, B, C, D separately suspended by threads. A bar magnet is slowly brought near each sample and the following observations are noted
 (i) A is feebly repelled
 (ii) B is feebly attracted
 (iii) C is strongly attracted
 (iv) D remains unaffected
 Which one of the following is true
 a) A is of a non-magnetic material
 b) B is of a paramagnetic material
 c) C is of a diamagnetic material
 d) D is of a ferromagnetic material
301. A superconducting material is
 a) Ferromagnetic b) Ferroelectric c) Diamagnetic d) Paramagnetic
302. The magnetic susceptibility of a paramagnetic substance at -73°C is 0.0060, then its value at -173°C will be
 a) 0.0030 b) 0.0120 c) 0.0180 d) 0.0045
303. A vibration magnetometer placed in magnetic meridian has a small bar magnet. The magnet executes oscillations with a time period of 2 sec in earth's horizontal magnetic field of 24 microtesla. When a horizontal field of 18 microtesla is produced opposite to the earth's field by placing a current carrying wire, the new time period of magnet will be
 a) 4s b) 1s c) 2s d) 3s
304. Which of the following is diamagnetic
 a) Aluminium b) Quartz c) Nickel d) Bismuth
305. The period of oscillation of a freely suspended bar magnet is 4 s. If it is cut into two equal parts in length, then the time period of each part will be

- a) 4 s b) 2 s c) 0.5 s d) 0.25 s
306. A magnet oscillating in a horizontal plane has a time period of 2 *second* at a place where the angle of dip is 30° and 3 *seconds* at another place where the angle of dip is 60° . The ratio of resultant magnetic fields at the two places is
- a) $\frac{4\sqrt{3}}{7}$ b) $\frac{4}{9\sqrt{3}}$ c) $\frac{9}{4\sqrt{3}}$ d) $\frac{9}{\sqrt{3}}$
307. Ratio of magnetic intensities for an axial point and a point on broad side-on position at equal distance d from the centre of magnet will be or The magnetic field at a distance d from a short bar magnet in longitudinal and transverse positions are in the ratio
- a) 1 : 1 b) 2 : 3 c) 2 : 1 d) 3 : 2
308. A magnet is parallel to a uniform magnetic field. If it is rotated by 60° , the work done is 0.8 J. How much work is done in moving it 30° further
- a) $0.8 \times 10^7 \text{ erg}$ b) 0.4 J c) 8 J d) 0.8 erg
309. A rigid circular loop of radius r and mass m lies in the $x - y$ plane of a flat table and has a current i flowing in it. At this particular place the earth's magnetic field is $\mathbf{B} = B_x \hat{i} + B_z \hat{k}$. The value of i so that the loop start tilting is
- a) $\frac{mg}{\pi r \sqrt{B_x^2 + B_z^2}}$ b) $\frac{mg}{\pi r B_x}$ c) $\frac{mg}{\pi r B_z}$ d) $\frac{mg}{\pi r \sqrt{B_x B_z}}$
310. The magnetic susceptibility is
- a) $\chi = \frac{I}{H}$ b) $\chi = \frac{B}{H}$ c) $\chi = \frac{M}{V}$ d) $\chi = \frac{M}{H}$
311. A small bar magnet of moment M is placed in a uniform field H . If magnet makes an angle of 30° with field, the torque acting on the magnet is
- a) MH b) $\frac{MH}{2}$ c) $\frac{MH}{3}$ d) $\frac{MH}{4}$
312. Two magnets, each of magnetic moment ' M ' are placed so as to form a cross at right angles to each other. The magnetic moment of the system will be
- a) 2 M b) $\sqrt{2} M$ c) 0.5 M d) M
313. A magnet of magnetic moment M oscillating freely in earth's horizontal magnetic field makes n oscillations per minute. If the magnetic moment is quadrupled and the earth's field is doubled, the number of oscillations made per minute would be
- a) $\frac{n}{2\sqrt{2}}$ b) $\frac{n}{\sqrt{2}}$ c) $2\sqrt{2}n$ d) $\sqrt{2}n$
314. Let ϕ_1 and ϕ_2 be the angles of dip observed in two vertical planes at right angles to each other and ϕ be the true angle of dip, then
- a) $\cos^2 \phi = \cos^2 \phi_1 + \cos^2 \phi_2$ b) $\sec^2 \phi = \sec^2 \phi_1 + \sec^2 \phi_2$
c) $\tan^2 \phi = \tan^2 \phi_1 + \tan^2 \phi_2$ d) $\cot^2 \phi = \cot^2 \phi_1 + \cot^2 \phi_2$
315. Which of the following is true
- a) Diamagnetism is temperature dependent
b) Paramagnetism is temperature dependent
c) Paramagnetism is temperature dependent
d) None of these
316. A bar magnet is situated on a table along east-west direction in the magnetic field of earth. The number of neutral points, where the magnetic field is zero, are
- a) 2 b) 0 c) 1 d) 4
317. A steel wire of length l has a magnetic moment M . It is bent at its middle point at an angle of 60° . Then the magnetic moment of new shape of wire will be
- a) $M/\sqrt{2}$ b) $M/2$ c) M d) $\sqrt{2}M$

318. Two bar magnets having same geometry with magnetic moments M and $2M$ are firstly placed in such a way that their poles are same side. Time period of oscillations is T_1 . Now the polarity of one of the magnets is reversed, and time period of oscillations is T_1 . Now the polarity of one of the magnets is reversed, and time period of oscillations is T_2 .
- a) $T_1 < T_2$ b) $T_1 = T_2$ c) $T_1 > T_2$ d) $T_2 = \infty$
319. Permanent magnet has properties retentivity and coercivity respectively
- a) High-high b) Low-low c) Low-high d) High-low
320. When the N -pole of a bar magnet points towards the south and S -pole towards the north, the null points are at the
- a) Magnetic axis b) Magnetic centre
c) Perpendicular divider of magnetic axis d) N and S poles
321. A bar magnet is placed in the position of stable equilibrium in a uniform magnetic field of induction B . If it is rotated through an angle 180° , then the work done is (M = Magnetic dipole moment of bar magnet)
- a) MB b) $2MB$ c) $\frac{MB}{2}$ d) Zero
322. The incorrect statement regarding the lines of force of the magnetic field B is
- a) Magnetic intensity is a measure of lines of force passing through unit area held normal to it
b) Magnetic lines of force form a closes curve
c) Inside a magnet, its magnetic lines of force move from north pole of a magnet towards its south pole
d) Due to a magnet magnetic lines of force never cut each other
323. Magnetic permeability is maximum for
- a) Diamagnetic substance b) Paramagnetic substance
c) Ferromagnetic substance d) All of these
324. The magnetic needle of a vibration magnetometer makes 12 oscillations per minute in the horizontal component of earth's magnetic field. When an external short bar magnet is placed at some distance along the axis of the needle in the same line, it makes 15 oscillations per minute. If the poles of the bar magnet are interchanged, the number of oscillations it makes per minute is
- a) $\sqrt{61}$ b) $\sqrt{63}$ c) $\sqrt{65}$ d) $\sqrt{67}$
325. The true value of angle of dip at a place is 60° , the apparent dip in a plane inclined at an angle of 30° with magnetic meridian is
- a) $\tan^{-1} \frac{1}{2}$ b) $\tan^{-1}(2)$ c) $\tan^{-1} \left(\frac{2}{3} \right)$ d) None of these
326. When a magnetic substance is heated, then it
- a) Becomes a strong magnet b) Losses its magnetism
c) Does not effect the magnetism d) Either (a) or (c)
327. A bar magnet has coercivity $4 \times 10^3 \text{ Am}^{-1}$. It is desired to demagnetise it by inserting it inside a solenoid 12 cm long and having 60 turns. The current that should be sent through the solenoid is
- a) 2 A b) 4 A c) 6 A d) 8 A
328. The figure below shows the north and south poles of a permanent magnet in which n turn coil of area of cross-section A is resting, such that for a current i passed through the coil, the plane of the coil makes an angle θ with respect to the direction of magnetic field B . If the plane of the magnetic field and the coil are horizontal and vertical respectively, the torque on the coil will be



a) $\tau = niAB \cos \theta$

b) $\tau = niAB \sin \theta$

c) $\tau = niAB$

d) None of the above, since the magnetic field is radial


329. Weber/m² is equal to

- a) Volt b) Henry c) Tesla d) All of these

330. The horizontal component of flux density of earth's magnetic field is $1.7 \times 10^{-5} \text{ T}$. The value of horizontal component of intensity of earth's magnetic field will be?

- a) 24.5 Am^{-1} b) 13.5 Am^{-1} c) 1.53 Am^{-1} d) 0.35 Am^{-1}


331. A uniform magnetic field parallel to the plane of paper, existed in space initially directed from left to right. When a bar of soft iron is placed I the field parallel to it, the lines of force passing through it will be represented by figure.



332. A magnet of length 14 cm and magnetic moment M is broken into two parts of length 6 cm and 8 cm. They are put at a right angle to each other with opposite poles together. The magnetic moment of the combination is

- a) $M/10$ b) M c) $M/1.4$ d) $2.8 M$

333. The $\chi - 1/T$ graph for an alloy of paramagnetic nature is shown in fig. the curie constant is




- a) 57 K b) $2.8 \times 10^{-3} \text{ K}$ c) 570 K d) $17.5 \times 10^{-3} \text{ K}$

334. A short bar magnet, placed with its axis at 30° with an external magnetic field of 0.16 T, experiences a torque of magnitude 0.032 J. The magnetic moment of the bar magnet is (in units of J/T)


- a) 4 b) 0.2 c) 0.5 d) 0.4

335. Two magnets A and B are identical and these are arranged as shown in the figure. Their length is negligible in comparison to the separation between them. A magnetic needle is placed between the magnets at point P which gets deflected through an angle θ under the influence of magnets. The ratio of distance d_1 and d_2 will be



- a) $(2 \tan \theta)^{1/3}$ b) $(2 \tan \theta)^{-1/3}$ c) $(2 \cot \theta)^{1/3}$ d) $(2 \cot \theta)^{-1/3}$

336. The most appropriate magnetization M versus magnetizing field H curve for a paramagnetic substance is




- a) A b) B c) C d) D

337. Resultant force acting on a diamagnetic material in a magnetic field is in direction

- a) From stronger to the weaker part of the magnetic field
b) From weaker to the stronger part of the magnetic field

- c) Perpendicular to the magnetic field
d) In the direction making 60° to the magnetic field
338. A magnet is placed on a paper in a horizontal plane for locating neutral points. A dip needle placed at the neutral point will be horizontal at the
a) Magnetic poles b) Magnetic equator c) Latitude angle 45° d) Latitude angle of 60°
339. The dimensions of magnetic permeability are
a) $[MLT^{-2}A^{-2}]$ b) $[ML^2T^{-2}A^{-2}]$ c) $[ML^2T^{-2}A^{-1}]$ d) $[M^{-1}LT^{-2}A^{-2}]$
340. The tangent galvanometers having coils of the same radius are connected in series. Same current flowing in them produces deflections of 60° and 45° respectively. The ratio of the number of turns in the coil is
a) $\frac{4}{\sqrt{3}}$ b) $\frac{\sqrt{3} + 1}{1}$ c) $\frac{\sqrt{3} + 1}{\sqrt{3} - 1}$ d) $\frac{\sqrt{3}}{1}$
341. At a place, the magnitudes of the horizontal component and total intensity of the magnetic field of the earth are 0.3 and 0.6 oersted respectively. The value of the angle of dip at this place will be
a) 60° b) 45° c) 30° d) 0°
342. The magnetic induction in air at a distance d from an isolated point pole of strength m unit will be
a) $\frac{m}{d}$ b) $\frac{m}{d^2}$ c) md d) md^2
343. At a certain place the horizontal component of the earth's magnetic field is B_0 and the angle of dip is 45° then total intensity of field at that place will be
a) B_0^2 b) $2B_0$ c) $\sqrt{2}B_0$ d) B_0
344. Rate of change of torque τ with deflection θ is maximum for a magnet suspended freely in a uniform magnetic field of induction B , when
a) $\theta = 0^\circ$ b) $\theta = 45^\circ$ c) $\theta = 60^\circ$ d) $\theta = 90^\circ$
345. Two magnets of equal mass are joined at right angles to each other as shown. Magnet 1 has a magnetic moment 3 times that of magnet 2. This arrangement is pivoted so that it is free to rotate in the horizontal plane. In equilibrium what angle will the magnet 1 subtend with the magnetic meridian



- a) $\tan^{-1}\left(\frac{1}{2}\right)$ b) $\tan^{-1}\left(\frac{1}{3}\right)$ c) $\tan^{-1}(1)$ d) 0°
346. The direction of lines of magnetic field of bar magnet is
a) From south pole to north pole
b) From north pole to south pole
c) Across the bar magnet
d) From south pole to north pole inside the magnet and from north pole to south pole outside the magnet
347. A bar magnet having a magnetic moment of $2 \times 10^4 \text{ J T}^{-1}$ is free to rotate in a horizontal plane. A horizontal magnetic field $B = 6 \times 10^{-4} \text{ T}$ exists in the space. The work done in taking the magnet slowly from a direction parallel to the field to a direction 60° from the field is
a) 0.6 J b) 12 J c) 6 J d) 2 J
348. The magnetic moment of atomic neon is equal to
a) Zero b) $\frac{1}{2} \mu_B$ c) $\frac{3}{2} \mu_B$ d) $2 \mu_B$
349. A vibration magnetometer consists of two identical bar magnets placed one over the other such that they are perpendicular and bisect each other. The time period of oscillation in a horizontal magnetic field is

$2^{5/4}$ second. One of the magnets is removed and if the other magnet oscillates in the same field, then the time period in second is

- a) $2^{1/4}$ b) $2^{1/2}$ c) 2 d) $2^{5/4}$

350. The correct measure of magnetic hardness of a material is

- a) Remanant magnetism b) Hysteresis loss
c) Coercivity d) Curie temperature

351. A dip circle is so that its needle moves freely in the magnetic meridian. In this position, the angle of dip is 40° . Now the dip circle is rotated so that the plane in which the needle moves makes an angle of 30° with the magnetic meridian. In this position, the needle will dip by an angle

- a) 40° b) 30° c) More than 40° d) Less than 40°

352. The value of angle of dip is zero at the magnetic equator because on it

- a) V and H are equal b) The value of V and H is zero
c) The value of V is zero d) The value of H is zero

353. A magnet when placed perpendicular to a uniform field of strength 10^{-4} Wb/m^2 experiences a maximum couple of moment $4 \times 10^{-5} \text{ N/m}$. What is its magnetic moment

- a) $0.4A \times m^2$ b) $0.2A \times m^2$ c) $0.16A \times m^2$ d) $0.04A \times m^2$


354. Identify the paramagnetic substance

- a) Iron b) Aluminium c) Nickel d) Hydrogen

355. At the magnetic north pole of the earth, the value of the horizontal component of earth's magnetic field and angle of dip are respectively

- a) Zero, maximum b) Maximum, minimum c) Maximum, maximum d) Minimum, minimum

356. The relative permeability (μ_r) of a ferromagnetic substance varies with temperature (T) according to the curve



- a) A b) B c) C d) D

357. Which of the following is most suitable for the core of electromagnets

- a) Soft iron b) Steel c) Copper-nickel alloy d) Air

358. Direction of magnetic field at equatorial point is

- a) Parallel to \mathbf{M} b) Perpendicular to \mathbf{M}
c) Making an angle of angle 45° with \mathbf{M} d) Antiparallel to \mathbf{M}

359. Choose the correct statement

- a) A paramagnetic material tends to move from a strong magnetic field to weak magnetic field
b) A magnetic material is in the paramagnetic phase below its Curie temperature
c) The resultant magnetic moment in an atom of a diamagnetic substance is zero
d) Typical domain size of a ferromagnetic material is 1 nm

360. The space inside a toroid is filled with tungsten whose susceptibility is 6.8×10^{-5} . The percentage increase in the magnetic field will be

- a) 0.0068% b) 0.068% c) 0.68% d) None of these

361. An electron of charge e moves in a circular orbit of radius r around the nucleus at a frequency ν . The magnetic moment associated with the orbital motion of the electron is.

- a) $\pi \nu e r^2$ b) $\frac{\pi \nu r^2}{e}$ c) $\frac{\pi \nu e}{r}$ d) $\frac{\pi e r^2}{\nu}$


362. The magnetic field due to short bar magnet of magnetic dipole moment M and length $2l$, on the axis at a distance z (where $z \gg l$) from the center of the magnet is given by formula

- a) $\frac{\mu_0 M}{4\pi z^3} \hat{M}$ b) $\frac{2\mu_0 M}{4\pi z^3} \hat{M}$ c) $\frac{4\pi M}{\mu_0 z^2} \hat{M}$ d) $\frac{\mu_0 M}{2\pi z^3} \hat{M}$

363. Magnetic moment of two bar magnets may be compared with the help of

- a) Deflection magnetometer b) Vibration magnetometer
c) Both of the above d) None of the above

364. A bar magnet of length 3 cm has points A and B along its axis at distances of 24 cm and 48 cm on the opposite sides. Ratio of magnetic fields at these points will be




- a) 8 b) $1/2\sqrt{2}$ c) 3 d) 4

365. If the magnetic flux is expressed in *weber*, then magnetic induction can be expressed in


- a) *Weber/m²* b) *Weber/m* c) *Weber-m* d) *Weber-m²*

366. Two short magnets of magnetic moment 1000 Am^2 are placed as shown at the corners of a square of side 10 cm . The net magnetic induction at P at



- a) 0.1 T b) 0.2 T c) 0.3 T d) 0.4 T

367. The figure shows the various positions (labelled by subscripts) of small magnetised needles P and Q . The arrows show the direction of their magnetic moment. Which configuration corresponds to the lowest potential energy among all the configurations shown



- a) PQ_3 b) PQ_4 c) PQ_5 d) PQ_6

368. Before using the tangent galvanometer, its coil is set in

- a) Magnetic meridian (or vertically north south)
b) Perpendicular to magnetic meridian
c) At angle of 45° to magnetic meridian
d) It does not require any setting

369. In the hysteresis cycle, the value of H needed to make the intensity of magnetization zero is called

- a) Retentivity b) Coercive force c) Lorentz force d) None of the above

370. A current carrying coil is placed with its axis parallel to N-S direction. Let horizontal component of earth's magnetic field be H_0 and magnetic field inside the loop is H . If a magnet is suspended inside the loop, it makes angle θ with H . Then θ is equal to

- a) $\tan^{-1}\left(\frac{H_0}{H}\right)$ b) $\tan^{-1}\left(\frac{H}{H_0}\right)$ c) $\operatorname{cosec}^{-1}\left(\frac{H}{H_0}\right)$ d) $\cot^{-1}\left(\frac{H_0}{H}\right)$

371. Ferromagnetic show their properties due to

- a) Filled inner subshells b) Vacant inner subshells
c) Partially filled inner subshells d) All the subshells equally filled

a) Diamagnetic b) Paramagnetic c) Ferromagnetic d) Antiferromagnetic

388. The magnetic needle of a tangent galvanometer is deflected at angle of 30° due to a current in its coil. The horizontal component of earth's magnetic field is 0.34×10^{-4} T, then magnetic field at the center of the coil due to current

- a) 1.96×10^{-5} T b) 1.96×10^{-4} T c) 1.96×10^4 T d) 1.96×10^5 T

389. A copper rod is suspended in a non homogeneous magnetic field region. The rod when in equilibrium will align itself

- a) In the region where magnetic field is strongest b) In the region where magnetic field is weakest and parallel to direction of magnetic field there
- c) In the direction in which it was originally suspended d) In the region where magnetic field is weakest and perpendicular to the direction of magnetic field there


390. The north pole of the earth's magnet is near the geographical

- a) South b) East c) West d) North

391. If a diamagnetic solution is poured into a U-tube and one arm of this U-tube is placed between the poles of a strong magnet, with the meniscus in line with the field, then the level of solution will

- a) Rise b) Fall c) Oscillate slowly d) Remain as such

392. The magnetic field lines due to a bar magnet are correctly shown in



393. The magnetic field at a point x on the axis of a small bar magnet is equal to the field at a point y on the equator of the same magnet. The ratio of the distances of x and y from the centre of the magnet is

- a) 2^{-3} b) $2^{-1/3}$ c) 2^3 d) $2^{1/3}$

394. A bar magnet is oscillating in the earth's magnetic field with time period T . If its mass is increased four times, then its time period will be

- a) $4T$ b) $2T$ c) T d) $\frac{T}{2}$

395. The magnetic needle of a tangent galvanometer is deflected at an angle 30° due to a magnet. The horizontal component of earth's magnetic field 0.34×10^{-4} T is along the plane of the coil. The magnetic intensity is

- a) 1.96×10^{-4} T b) 1.96×10^4 T c) 1.96×10^{-5} T d) 1.96×10^5 T

396. The angle of dip at the magnetic equator is

- a) 0° b) 45° c) 30° d) 90°

397. The hysteresis cycle for the material of a transformer core is

- a) Short and wide b) Tall and narrow c) Tall and wide d) Short and narrow





398. If a piece of metal was thought to be magnet, which one of the following observations would offer conclusive evidence

- a) It attracts a known magnet b) It repels a known magnet
- c) Neither (a) nor (b) d) It attracts a steel screw driver


399. A tangent galvanometer has a reduction factor of 1 A and it is placed with the plane of its coil perpendicular to the magnetic meridian. The deflection produced when a current of 1 A is passed through it is

- a) 60° b) 45° c) 30° d) None of these

400. Two uniform magnetic fields B and H are perpendicular to each other at a place. When a magnetic needle is placed in the field, it rest making angle 60° and 30° with B and H respectively. The value of $B:H$ is


- a) 1 : 2 b) 2 : 1 c) $\sqrt{3} : 1$ d) $1 : \sqrt{3}$
401. The effective length of a magnet is 31.4 cm and its pole strength is 0.5 Am. Calculate its magnetic moment. If it is bent in form of semicircle, then magnetic moment will be
 a) $0.157 \text{ Am}^2, 0.01 \text{ Am}^2$ b) $0.357 \text{ Am}^2, 0.01 \text{ Am}^2$
 c) $1.157 \text{ Am}^2, 1.01 \text{ Am}^2$ d) None of these
402. The hysteresis curve is studied generally for
 a) Ferromagnetic materials b) Paramagnetic materials
 c) Diamagnetic materials d) All of the above
403. A rod of ferromagnetic material with dimensions $10 \text{ cm} \times 0.5 \text{ cm} \times 0.2 \text{ cm}$ is placed in a magnetic field of strength $0.5 \times 10^4 \text{ A} - \text{m}^{-1}$ as a result of which of which a magnetic moment of $0.5 \text{ A} - \text{m}^{-2}$ is produced in rod. The value of magnetic induction will be
 a) 0.54 T b) 6.28 T c) 0.358 T d) 2.591 T
404. A magnetic needle is made to vibrate in uniform field H , then its time period is T . If it vibrates in the field of intensity $4H$, its time period will be
 a) $2T$ b) $T/2$ c) $2/T$ d) T
405. The bob of a simple pendulum is replaced by a magnet. The oscillation are set along the length of the magnet. A copper coil is added so that one pole of the magnet passes in and out of the coil. The coil is short-circuited. Then which one of the following happens
 a) Period decreases b) Period does not change
 c) Oscillations are damped d) Amplitude increases
406. A current carrying loop is placed in a uniform magnetic field in four different orientations, I, II, III and IV, arrange them in the decreasing order of potential energy
- I.  II. 
- III.  IV. 
- a) $I > III > II > IV$ b) $I > II > III > IV$ c) $I > IV > II > III$ d) $III > IV > I > II$
407. The line on the earth surface joining the point where the field is horizontal, is called
 a) Magnetic equator b) Magnetic Line c) Magnetic axis d) Magnetic inertia
408. The time of vibration of a dip needle vibrating in the vertical plane is 3s. When magnetic needle is made to vibrate in the horizontal plane, the time of vibration is $3\sqrt{2}$ s. Then the angle of dip is
 a) 30° b) 45° c) 60° d) 90°
409. Magnetic lines of force
 a) Always intersect b) Are always closed
 c) Tend to crowd far away from the poles of magnet d) Do not pass through vacuum

410. A bar magnet is 10 cm long, and is kept with its north (N)- pole pointing north. A neutral point is formed at a distance of 15 cm from each pole. Given the horizontal component of earth's field to be 0.4 Gauss, the pole strength of the magnet is
 a) 9 A-m b) 6.75 A-m c) 27 A-m d) 1.35 A-m
411. With a standard rectangular bar magnet the time period of a vibration magnetometer is 4 s. The bar magnet is cut parallel to its length into four equal pieces. The time period of vibration magnetometer when one piece is used (in second) (bar magnet breadth is small) is
 a) 16 b) 8 c) 4 d) 2
412. A short magnet of moment 6.75 Am^2 produces a neutral point on its axis. If horizontal component of earth's magnetic field is $5 \times 10^{-5} \text{ Wb/m}^2$, then the distance of the neutral point should be
 a) 10 cm b) 20 cm c) 30 cm d) 40 cm
413. A domain in a ferromagnetic substance is in the form of a cube of side length $1 \mu\text{m}$. If it contains 8×10^{10} atoms and each atomic dipole has a dipole moment of $9 \times 10^{-24} \text{ Am}^2$, then magnetization of the domain is
 a) $7.2 \times 10^5 \text{ Am}^{-1}$ b) $7.2 \times 10^3 \text{ Am}^{-1}$ c) $7.2 \times 10^9 \text{ Am}^{-1}$ d) $7.2 \times 10^{12} \text{ Am}^{-1}$
414. The earth's magnetic induction at a certain point is $7 \times 10^{-5} \text{ Wbm}^{-2}$. This is to be annulled by the magnetic induction at the center of a circular conducting loop of radius 15 cm. The required current in the loop is
 a) 0.56 A b) 5.6 A c) 0.28 A d) 2.8 A
415. The time period of oscillation of a bar magnet suspended horizontally along the magnetic meridian is T_0 . If this magnet is replaced by another magnet of the same size and pole strength but with double the mass, the new time period will be
 a) $\frac{T_0}{2}$ b) $\frac{T_0}{\sqrt{2}}$ c) $\sqrt{2}T_0$ d) $2T_0$
416. If a ferromagnetic material is inserted in a current carrying solenoid, the magnetic field of solenoid
 a) Large increases b) Slightly increases c) Largely decreases d) Slightly decreases
417. Which one of the following characteristics is not associated with a ferromagnetic material?
 a) It is strongly attracted by a magnet
 b) It tends to move from a region of strong magnetic field to a region of low magnetic field
 c) Its origin is the spin of electrons
 d) Above the Curie temperature, it exhibits paramagnetic properties
418. A small coil C with $N = 200$ turns is mounted on one end of a balance beam and introduced between the poles of an electromagnet as shown in figure. The cross sectional area of coil is $A = 1.0 \text{ cm}^2$, length of arm OA of the balance beam is $l = 30 \text{ cm}$. When there is no current in the coil the balance is in equilibrium. On passing a current $I = 22 \text{ mA}$ through the coil the equilibrium is restored by putting the additional counter weight of mass $\Delta m = 60 \text{ mg}$ on the balance pan. Find the magnetic induction at the spot where coil is located



- a) 0.4 T b) 0.3 T c) 0.2 T d) 0.1 T
419. If a magnetic substance is kept in a magnetic field then which of the following substance is thrown out?
 a) Paramagnetic b) Ferromagnetic c) Diamagnetic d) Antiferromagnetic
420. A loop of area 0.5 m^2 is placed in a magnetic field of strength 2 T in direction making an angle of 30° with the field. The magnetic flux linked with the loop will be
 a) $\frac{1}{2} \text{ Wb}$ b) $\frac{\sqrt{3}}{2} \text{ Wb}$ c) 2 Wb d) $\frac{\sqrt{3}}{2} \text{ Wb}$

421. Which of the following statements is incorrect about hysteresis
- This effect is common to all ferromagnetic substances
 - The hysteresis loop area is proportional to the thermal energy developed per unit volume of the material
 - The hysteresis loop area is independent of the thermal energy developed per unit volume of the material
 - The shape of the hysteresis loop is characteristic of the material
422. Three identical bar magnets each of magnetic moment M are placed in the form of an equilateral triangle as shown. The net magnetic moment of the system is



- Zero
 - $2M$
 - $M\sqrt{3}$
 - $\frac{3M}{2}$
423. A magnet of magnetic moment M and pole strength m is divided in two equal parts, then magnetic moment of each part will be
- M
 - $M/2$
 - $M/4$
 - $2M$

