


Date :
Time :
Marks :

PHYSICS

RAY OPTICS AND OPTICAL INSTRUMENTS

Single Correct Answer Type

1. A dentist has a small mirror of focal length 16 mm. He views the cavity in the tooth of a patient by holding the mirror at a distance of 8 mm from the cavity. The magnification is
a) 1 b) 1.5 c) 2 d) 3
2. In an eye-piece, field lens and eye lens have focal lengths 7.5 cm and 7.3 cm. To eliminate spherical aberration, distance between them would be
a) 0.2 cm b) 0.4 cm c) 0.1 cm d) 0.5 cm
3. When sunlight is scattered by atmospheric atoms and molecules, the amount of scattering of light of wavelength 440 nm is A . The amount of scattering for the light of wavelength 660 nm is approximately
a) $\frac{4}{9}A$ b) $2.25A$ c) $1.5A$ d) $\frac{A}{5}$
4. A double convex lens, lens made of a material of refractive index μ_1 , is placed inside two liquids or refractive indices μ_2 and μ_3 , as shown. $\mu_2 > \mu_1 > \mu_3$. A wide, parallel beam of light is incident on the lens from the left. The lens will give rise to


a) A single convergent beam b) Two different convergent beams
c) Two different divergent beams d) A convergent and a divergent beam
5. A convex lens of focal length 30 cm produces 5 times magnified real image of an object. What is the object distance?
a) 36 cm b) 25 cm c) 30 cm d) 150 cm
6. Light travels in two media A and B with speeds $1.8 \times 10^8 \text{ ms}^{-1}$ and $2.4 \times 10^8 \text{ ms}^{-1}$ respectively. Then the critical angle between them is
a) $\sin^{-1}\left(\frac{2}{3}\right)$ b) $\tan^{-1}\left(\frac{3}{4}\right)$ c) $\tan^{-1}\left(\frac{2}{3}\right)$ d) $\sin^{-1}\left(\frac{3}{4}\right)$
7. We combined a convex lens of focal length f_1 and concave lens of focal lengths f_2 and their combined focal length was F . The combination of these lenses will behave like a concave lens, if
a) $f_1 > f_2$ b) $f_1 < f_2$ c) $f_1 = f_2$ d) $f_1 \leq f_2$
8. A biconvex lens form a real image of an object placed perpendicular to its principal axis. Suppose the radii of curvature of the lens tend to infinity. Then the image would
a) Disappear
b) Remain as real image still
c) Be virtual and of the same size as the object
d) Suffer from aberrations
9. A divergent lens will produce
a) Always a virtual image b) Always real image
c) Sometimes real and sometimes virtual d) None of the above
10. An astronomical telescope has an angular magnification of magnitude 5 for distant objects. The separation between the objective and the eye-piece is 36 cm and the final image is formed at infinity. The focal length f_o of the objective and the focal length f_e of the eye-piece are

- a) $f_o = 45 \text{ cm}$ and $f_e = -9 \text{ cm}$ b) $f_o = -7.2 \text{ cm}$ and $f_e = 5 \text{ cm}$
 c) $f_o = 50 \text{ cm}$ and $f_e = 10 \text{ cm}$ d) $f_o = 30 \text{ cm}$ and $f_e = 6 \text{ cm}$
11. A convex lens is placed between object and a screen. The size of object is 3 cm and an image of height 9 cm is obtained on the screen. When the lens is displaced to a new position, what will be the size of image on the screen?
 a) 2 cm b) 6 cm c) 4 cm d) 1 cm
12. A double convex lens made out of glass (refractive index $\mu = 1.5$) has both radii of curvature of magnitudes 20 cm. Incident light rays parallel to the axis of this lens will converge at a distance d such that
 a) $d = 10 \text{ cm}$ b) $d = \frac{20}{3} \text{ cm}$ c) $d = 40 \text{ cm}$ d) $d = 20 \text{ cm}$
13. The deviation caused in red, yellow and violet colours for crown glass prism are 2.84° , 3.28° and 3.72° respectively. The dispersive power of prism material is
 a) 0.268 b) 0.368 c) 0.468 d) 0.568
14. A beam of monochromatic blue light of wavelength 4200 \AA in air travels in water ($\mu = 4/3$). Its wavelength in water will be
 a) 2800 \AA b) 5600 \AA c) 3150 \AA d) 4000 \AA
15. Refractive index of a medium is μ . The incidence angle is twice that of refracting angle. The angle of incidence is
 a) $\cos^{-1}\left(\frac{\mu}{2}\right)$ b) $\sin^{-1}\left(\frac{\mu}{2}\right)$ c) $2 \cos^{-1}\left(\frac{\mu}{2}\right)$ d) $\sin^{-1} \mu$
16. The far point of a myopia eye is at 40 cm. For removing this defect, the power of lens required will be
 a) 40 D b) -4 D c) -2.5 D d) 0.25 D
17. An object is put at a distance of 5 cm from the first focus of a convex lens of focal length 10 cm. If a real image is formed, its distance from the lens will be
 a) 15 cm b) 20 cm c) 25 cm d) 30 cm
18. A thin rod of length $f/3$ lies along the axis of a concave mirror of focal length f . One end of its magnified image touches an end of the rod. The length of the image is
 a) f b) $\frac{1}{2}f$ c) $2f$ d) $\frac{1}{4}f$
19. The solar spectrum during a complete solar eclipse is
 a) Continuous b) Emission line c) Dark line d) Dark band
20. In an equilateral prism if incident angle is 45° then minimum deviation is
 a) 30° b) 60° c) 45° d) 90°
21. Consider the following two statements A and B and identify the correct choice in the given answers
 A: Line spectra is due to atoms in gaseous state
 B: Band spectra is due to molecules
 a) Both A and B are false b) A is true and B is false
 c) A is false and B is true d) Both A and B are true
22. A person suffering from 'presbyopia' (myopia and hyper metropia both defects) should use
 a) A concave lens
 b) A convex lens
 c) A bifocal lens whose lower portion is convex
 d) A bifocal lens whose upper portion is convex
23. A man having height 6 m, observes image of 2 m height erect, then mirror used is
 a) Concave b) Convex c) Plane d) None of the above
24. An object is viewed through a compound microscope and appears in focus when it is 5 mm away from the objective lens. When a sheet of transparent material 3 mm thick is placed between the objective and the


microscope, the objective lens has to be moved 1 mm to bring the object back into the focus. The refractive index of the transparent material is

- a) 1.5 b) 1.6 c) 1.8 d) 2.0

25. The Cauchy's dispersion formula is

- a) $n = A + B\lambda^{-2} + C\lambda^{-4}$ b) $n = A + B\lambda^2 + C\lambda^{-4}$ c) $n = A + B\lambda^{-2} + C\lambda^4$ d) $n = A + B\lambda^2 + C\lambda^4$

26. A light ray is incident upon a prism in minimum deviation position and suffers a deviation of 34° . If the shaded half of the prism is knocked off, the ray will



- a) Suffer a deviation of 34° b) Suffer a deviation of 68°
c) Suffer a deviation of 17° d) Not come out of the prism

27. The impact of an image on the retina remains for

- a) 0.1 s b) 0.5 s c) 10 s d) 15 s

28. An object is placed at a distance of 10 cm from a convex lens of power 5D. Find the position of the image

- a) -20 cm b) 30 cm c) 20 cm d) -30 cm

29. If eye is kept at a depth h inside water of refractive index and viewed outside, then the diameter of the circle through which the outer objects become visible, will be

- a) $\frac{h}{\sqrt{\mu^2 - 1}}$ b) $\frac{h}{\sqrt{\mu^2 + 1}}$ c) $\frac{2h}{\sqrt{\mu^2 - 1}}$ d) $\frac{h}{\sqrt{\mu^2}}$


30. Magnification at least distance of distinct vision of a simple microscope having its focal length 5 cm is

- a) 2 b) 4 c) 5 d) 6

31. Find the luminous intensity of the sun if it produces the same illuminance on the earth as produced by a bulb of 10000 candela at a distance 0.3 m. The distance between the sun and the earth is $1.5 \times 10^{11} m$

- a) $25 \times 10^{22} cd$ b) $25 \times 10^{18} cd$ c) $25 \times 10^{26} cd$ d) $25 \times 10^{36} cd$

32. Two immiscible liquids of refractive indices 1.5 and $\frac{4}{3}$ are filled in glass jar each of length 6 cm. A light of source S is at the bottom of the jar, the apparent depth of light source will be



- a) 12.5 cm b) 17 cm c) 12 cm d) 8.5 cm

33. A parallel beam of white light falls on a convex lens. Images of blue, yellow and red light are formed on other side of the lens at a distance of 0.20 m, 0.205 m and 0.214 m respectively. The dispersive power of the material of the lens will be

- a) 619/1000 b) 9/200 c) 14/205 d) 5/214

34. A ray of light passes an equilateral prism such that an angle of incidence is equal to the angle of emergence and the latter is equal to $\frac{3}{4}$ th the angle of prism. The angle of deviation is

- a) 45° b) 39° c) 20° d) 30°


35. Two lenses having $f_1 : f_2 = 2 : 3$ has combination to make no dispersion. Find the ratio of dispersive power of glasses used

- a) 2 : 3 b) 3 : 2 c) 4 : 9 d) 9 : 4

36. A ray of light strikes a transparent rectangular slab (of refractive index $\sqrt{2}$) at an angle of incidence of 45° . The angle between the reflected and refracted rays is


- a) 75° b) 90° c) 105° d) 120°

37. What will be the colour of sky as seen from the earth, if there were no atmosphere
 a) Black b) Blue c) Orange d) Red
38. A convex lens of focal length $\frac{1}{3}$ m forms a real, inverted image twice in size of the object. The distance of the object from the lens is
 a) 0.5 m b) 0.166 m c) 0.33 m d) 1 m
39. To get three images of a single object, one should have two plane mirrors at an angle of
 a) 60° b) 90° c) 120° d) 30°
40. Chromatic aberration of lens can be corrected by
 a) Reducing its aperture
 b) Proper polishing of its two surfaces
 c) Suitably combining it with another lens
 d) Providing different suitable curvature to its two surfaces
41. The least angle of deviation for a glass prism is equal to its refracting angle. The refractive index of glass is 1.5. Then the angle of prism is
 a) $2 \cos^{-1}\left(\frac{3}{4}\right)$ b) $\sin^{-1}\left(\frac{3}{4}\right)$ c) $2 \sin^{-1}\left(\frac{3}{2}\right)$ d) $\cot^{-1}\left(\frac{3}{2}\right)$
42. A diverging beam of light from a point source S having divergence angle α , falls symmetrically on a glass slab as shown. The angles of incidence of the two extreme rays are equal. If the thickness of the glass slab is t and the refractive index n , then the divergence angle of the emergent beam is




- a) Zero b) α c) $\sin^{-1}(1/n)$ d) $2 \sin^{-1}(1/n)$
43. A medium shows relation between i and r as shown. If speed of light in the medium is nc then value of n is
-
- a) 1.5 b) 2 c) 2^{-1} d) $3^{-1/2}$
44. What is the angle of incidence for an equilateral prism of refractive index $\sqrt{3}$ so that the ray is parallel to the base inside the prism?
 a) 30° b) 45° c) 60° d) Either 30° or 60°
45. Image formed by a convex mirror is
 a) Virtual b) Real c) Enlarged d) Inverted
46. How much water should be filled in a container 21 cm in height, so that it appears half filled when viewed from the top of the container (given that $\mu_{\text{water}} = 4/3$)
 a) 8.0 cm b) 10.5 cm c) 12.0 cm d) None of the above
47. Fraunhofer spectrum is
 a) Line absorption spectrum b) Band absorption spectrum
 c) Line emission spectrum d) Band emission spectrum
48. A transparent cube of 2.1 m edge contains a small air bubble. Its apparent distance when viewed through one face of the cube is 0.10 m and when viewed from the opposite face is 0.04 m. The actual distance of the bubble from the second face of the cube is
 a) 0.06 m b) 0.17 m c) 0.05 m d) 0.04 m
49. Emission spectrum of CO_2 gas

- a) Is a line spectrum
b) Is a band spectrum
c) Is a continuous spectrum
d) Does not fall in the visible region
50. "Lux" is a unit of
a) Luminous intensity of a source
b) Illuminance on a surface
c) Transmission coefficient of a surface
d) Luminous efficiency of source of light
51. A bi-convex lens is formed with two thin plano-convex lenses as shown in the figure. Refractive index n of the first lens is 1.5 and that of the second lens is 1.2. Both the curved surfaces are of the same radius of curvature $R = 14\text{ cm}$. For this bi-convex lens, for an object distance of 40 cm , the image distance will be




- a) -280.0 cm
b) 40.0 cm
c) 21.5 cm
d) 13.3 cm
52. In a given direction, the intensities of the scattered light by a scattering substance for two beams of light are in the ratio of $256 : 81$. The ratio of the frequency of the first beam to the frequency of the second beam is
a) $64 : 127$
b) $4 : 3$
c) $64 : 27$
d) $2 : 1$
53. Figure shows a cubical room $ABCD$ with the wall CD as a plane mirror. Each side of the room is 3 m . We place a camera at the midpoint of the wall AB . At what distance should the camera be focused to photograph an object placed at A








- a) 1.5 m
b) 3 m
c) 6 m
d) More than 6 m
54. In a compound microscope, the intermediate image is
a) Virtual erect and magnified
b) Real, erect and magnified
c) Real, inverted and magnified
d) Virtual, erect and reduced
55. In the visible region the dispersive powers and the mean angular deviations for crown and flint glass prisms are ω, ω' and d, d' respectively. The condition for getting deviation without dispersion when the two prisms are combined is
a) $\sqrt{\omega d} + \sqrt{\omega' d'} = 0$
b) $\omega' d + \omega d' = 0$
c) $\omega d + \omega' d' = 0$
d) $(\omega d)^2 = (\omega' d')^2 = 0$
56. The separation between two microscopic particles is measured P_A and P_B by two different lights of wavelength 2000 \AA and 3000 \AA respectively, then
a) $P_A > P_B$
b) $P_A < P_B$
c) $P_A < 3/2 P_B$
d) $P_A = P_B$
57. If the critical angle for total internal reflection from a medium to vacuum is 30° , the velocity of light in the medium is
a) $3 \times 10^8\text{ m/s}$
b) $1.5 \times 10^8\text{ m/s}$
c) $6 \times 10^8\text{ m/s}$
d) $\sqrt{3} \times 10^8\text{ m/s}$
58. Sun subtends an angle of 0.5° at the centre of curvature of a concave mirror of radius of curvature 15 m . The diameter of the image of the sun formed by the mirror is
a) 8.55 cm
b) 7.55 cm
c) 6.55 cm
d) 5.55 cm
59. The chromatic aberration in lenses becomes due to
a) Disimilarity of main axis of rays
b) Disimilarity of radii of curvature

- c) Variation of focal length of lenses with wavelength
d) None of these
60. At the time of total solar eclipse, the spectrum of solar radiation would be
a) A large number of dark Fraunhofer lines
b) A less number of dark Fraunhofer lines
c) No lines at all
d) All Fraunhofer lines changed into brilliant colours
61. A man can see the object between 15 cm and 30 cm. He uses the lens to see the far objects. Then due to the lens used, the near point will be at
a) $\frac{10}{3}$ cm b) 30 cm c) 15 cm d) $\frac{100}{3}$ cm
62. A normally incident ray reflected at an angle of 90° . The value of critical angle is
a) 45° b) 90° c) 65° d) 43.2°
63. A ray of light is incident at 50° on the middle of one of the two mirrors arranged at an angle of 60° between them. The ray then touches the second mirror, gets reflected back to the first mirror, making an angle of incidence
a) 50° b) 60° c) 70° d) 80°
64. Glass has refractive index μ with respect to air and the critical angle for a ray of light going from glass to air is θ . If a ray of light is incident from air on the glass with angle of incidence θ , the corresponding angle of refraction is
a) $\sin^{-1}\left(\frac{1}{\sqrt{\mu}}\right)$ b) 90° c) $\sin^{-1}\left(\frac{1}{\mu^2}\right)$ d) $\sin^{-1}\left(\frac{1}{\mu}\right)$
65. A leaf which contains only green pigments, is illuminated by a laser light of wavelength $0.632 \mu\text{m}$. It would appear to be
a) Brown b) Black c) Red d) Green
66. The focal length of the objective and the eye-piece of a microscope are 4 mm and 25 mm respectively. If the final image is formed at infinity and the length of the tube is 16 cm, then the magnifying power of microscope will be
a) -337.5 b) -3.75 c) 3.375 d) 33.75
67. A microscope is focused on a coin lying at the bottom of a beaker. The microscope is now raised up by 1 cm. To what depth should the water be poured into the beaker so that coin is again in focus? (Refractive index of water is $\frac{4}{3}$)
a) 1 cm b) $\frac{4}{3}$ cm c) 3 cm d) 4 cm
68. As shown in figure position of an images I of an object O formed by lens. This is possible if







- a) A convex lens is placed to the left of O b) A concave lens is placed to the left of O
c) A convex lens is placed between O and I d) A concave lens is placed to the right of I
69. The hyper-metropia is a
a) Short-side defect b) Long-side defect
c) Bad vision due to old age d) None of these
70. An object is placed 30 cm to the left of a diverging lens whose focal length is of magnitude 20 cm. Which one of the following correctly states the nature and position of the virtual image formed?
- | Nature of image | Distance from lens |
|---|--|
| a) Inverted, enlarged
60 cm to the right | b) Erect, diminished
12 cm to the left |
| c) Inverted, enlarged
60 cm to the left | d) Erect, diminished
12 cm to the right |

71. A concave mirror of focal length 100cm is used to obtain the image of the sun which subtends an angle of $30'$. The diameter of the image of the sun will be
 a) 1.74cm b) 0.87cm c) 0.435cm d) 100cm
72. In refraction, light waves are bent on passing from one medium to the second medium, because, in the second medium
 a) The frequency is different b) The coefficient of elasticity is different
 c) The speed is different d) The amplitude is smaller
73. When a glass lens with $n = 1.47$ is immersed in a trough of liquid, it looks to be disappeared. The liquid in the trough could be
 a) Water b) Kerosene c) Glycerin d) Alcohol
74. A beaker containing a liquid appears to be half when it is actually two third full. The refractive index of liquid
 a) $7/6$ b) $6/5$ c) $3/2$ d) $4/3$
75. Which of the following diagrams shows correctly the dispersion of white light by a prism
 a)  b)  c)  d) 
76. Two lenses are placed in contact with each other and the focal length of combination is 80 cm . If the focal length of one is 20 cm , then the power of the other will be
 a) 1.66 D b) 4.00 D c) -1.00 D d) -3.75 D
77. A 16 cm long image of an object is formed by a convex lens on a screen. On moving the lens towards the screen, without changing the positions of the object and the screen, a 9 cm long image is formed again on the screen. The size of the object is
 a) 9 cm b) 11 cm c) 12 cm d) 13 cm
78. The dispersion for a medium of wavelength λ is D , then the dispersion for the wavelength 2λ will be
 a) $D/8$ b) $D/4$ c) $D/2$ d) D
79. The resolving power of a telescope depends on
 a) Focal length of eye lens b) Focal length of objective lens
 c) Length of the telescope d) Diameter of the objective lens
80. A point object is placed at distance of 20 cm from a thin planoconvex lens of focal length 15 cm . The plane surface of the lens is now silvered. The image created by the system is at



- a) 60 cm to the left of the system
 b) 60 cm to the right of the system
 c) 12 cm to the left of the system
 d) 12 cm to the right of the system
81. A spectrum is formed by a prism of dispersive power ' ω '. If the angle of deviation is ' δ ', then the angular dispersion is
 a) ω/δ b) δ/ω c) $1/\omega\delta$ d) $\omega\delta$
82. When light travels from one medium to the other of which the refractive index is different, then which of the following will change
 a) Frequency, wavelength and velocity b) Frequency and wavelength
 c) Frequency and velocity d) Wavelength and velocity
83. The astronomical telescope consists of objective and eye-piece. The focal length of the objective is
 a) Equal to that of the eye-piece


- b) Greater than that of the eye-piece
 c) Shorter than that of the eye-piece
 d) Five times shorter than that of the eye-piece
84. If \hat{i} denotes a unit vector along incident light ray, \hat{r} a unit vector along refracted ray into a medium of refractive index μ and \hat{n} unit vector normal to boundary of medium directed towards incident medium, then law of refraction is
 a) $\hat{i} \cdot \hat{n} = \mu(\hat{r} \cdot \hat{n})$ b) $\hat{i} \times \hat{n} = \mu(\hat{r} \times \hat{n})$ c) $\hat{i} \times \hat{n} = \mu(\hat{r} \times \hat{n})$ d) $\mu(\hat{i} \times \hat{n}) = \hat{r} \times \hat{n}$
85. A combination of two thin lenses of the same material with focal length f_1 and f_2 , arranged on a common axis minimizes chromatic aberration. If the distance between them is
 a) $\frac{(f_1 + f_2)}{4}$ b) $\frac{(f_1 + f_2)}{2}$ c) $(f_1 + f_2)$ d) $2(f_1 + f_2)$
86. For a convex lens, if real image is formed the graph between $(u + v)$ and u or v is as follows
 a)  b)  c)  d) 
87. At what distance from a convex lens of focal length 30 cm, an object should be placed, so that the size of the image be $\frac{1}{2}$ th of the object?
 a) 30 cm b) 60 cm c) 15 cm d) 90 cm
88. A beaker containing liquid is placed on a table, underneath a microscope which can be moved along a vertical scale. The microscope is focussed, through the liquid onto a mark on the table when the reading on the scale is a . It is next focused on the upper surface of the liquid and the reading is b . More liquid is added and the observations are repeated, the corresponding readings are c and d . The refractive index of the liquid is
 a) $\frac{d - b}{d - c - b + a}$ b) $\frac{b - d}{d - c - b + a}$ c) $\frac{d - c - b + a}{d - b}$ d) $\frac{d - b}{a + b - c - d}$
89. Which one of the following statements is true
 a) An object situated at the principle focus of a concave lens will have its image formed at infinity
 b) Concave mirror can give diminished virtual image
 c) Given a point source of light, a convex mirror can produce a parallel beam of light
 d) The virtual image formed in a plane mirror can be photographed
90. An object 2.5 cm high is placed at a distance of 10 cm from a concave mirror of radius of curvature 30 cm. The size of the image is
 a) 9.2 cm b) 10.5 cm c) 5.6 cm d) 7.5 cm
91. In compound microscope, magnifying power is 95 and the distance of object from objective lens is $\frac{1}{3.8}$ cm. The focal length of objective lens is $\frac{1}{4}$ cm. What is the magnification of eye piece?
 a) 5 b) 10 c) 100 d) 200
92. Given a point source of light, which of the following can produce a parallel beam of light
 a) Convex mirror b) Concave mirror
 c) Concave lens d) Two plane mirrors inclined at an angle of 90°
93. The refractive index of water is 1.33. The direction in which a man under water should look to see the setting sun is
 a) 49° to the horizontal b) 90° with the vertical c) 49° to the vertical d) Along the horizontal
94. At Kavalur in India, the astronomers using a telescope whose objective had a diameter of one metre started using telescope of diameter 2.54 m. This resulted in
 a) The increase in the resolving power by 2.54 times for the same λ
 b) The increase in the limiting angle by 2.54 times for the same λ

- c) Decrease in the resolving power
d) No effect on the limiting angle
95. A compound microscope has two lenses. The magnifying power of one is 5 and the combined magnifying power is 100. The magnifying power of the other lens is
a) 10 b) 20 c) 50 d) 25
96. Consider an equiconvex lens of radius of curvature R and focal length f . If $f > R$, the refractive index μ of the material of the lens
a) Is greater than zero but less than 1.5 b) Is greater than 1.5 but less than 2.0
c) Is greater than one but less than 1.5 d) None of the above
97. A compound microscope has an eyepiece of focal length 10 cm and an objective of focal length 4 cm. Calculate the magnification, if an object is kept at a distance of 5 cm from the objective, so that final image is formed at the least distance of distinct vision 20 cm.
a) 12 b) 11 c) 10 d) 13
98. For total internal reflection to take place, the angle of incidence i and the refractive index μ of the medium must satisfy the inequality
a) $\frac{1}{\sin i} < \mu$ b) $\frac{1}{\sin i} > \mu$ c) $\sin i < \mu$ d) $\sin i > \mu$
99. Two thin lenses of focal length 20 cm and 25 cm are in contact. The effective power of the combination is
a) 4.5 D b) 18 D c) 45 D d) 9 D
100. A vessel of height $2d$ is half-filled with a liquid of refractive index $\sqrt{2}$ and the other half with a liquid of refractive index n (the given liquids are immiscible). Then the apparent depth of the inner surface of the bottom of the vessel (neglecting the thickness of the bottom of the vessel) will be
a) $\frac{n}{d(n + \sqrt{2})}$ b) $\frac{d(n + \sqrt{2})}{n\sqrt{2}}$ c) $\frac{\sqrt{2}n}{d(n + \sqrt{2})}$ d) $\frac{nd}{(d + \sqrt{2}n)}$
101. A concave lens of focal length 20 cm placed in contact with a plane mirror acts as a convex mirror of focal length
a) 10 cm b) 40 cm c) 60 cm d) 20 cm
102. The focal lengths of the objective and the eye piece of telescope are 100 cm and 10 cm respectively. The magnification of the telescope when final image is formed at infinity is
a) 0.1 b) 10 c) 100 d) ∞
103. A plane mirror reflects a pencil of light to form a real image. Then the pencil of light incident on the mirror is
a) parallel b) convergent c) divergent d) Any of these
104. A converging lens is used to form an image on a screen. When upper half of the lens is covered by an opaque screen
a) Half the image will disappear
b) Complete image will be formed of same intensity
c) Half image will be formed of same intensity
d) Complete image will be formed of decreased intensity
105. Relation between critical angles of water and glass is
a) $C_w > C_g$ b) $C_w < C_g$ c) $C_w = C_g$ d) $C_w = C_g = 0$
106. A thin rod of 5cm length is kept along the axis of a concave mirror of 10cm focal length such that its image is real and magnified and one end touches the rod. Its magnification
a) 1 b) 2 c) 3 d) 4
107. A fish, looking up through the water sees the outside world contained in a circular horizon. If the refractive index of water is $4/3$ and the fish is 12 cm below the surface of water, the radius of the circle in centimetre is
a) $\frac{12 \times 3}{\sqrt{5}}$ b) $12 \times 3 \times \sqrt{5}$ c) $\frac{12 \times 3}{\sqrt{7}}$ d) $12 \times 3 \times \sqrt{7}$

108. One of the refracting surfaces of a prism of angle 30° is silvered. A ray of light incident at an angle of 60° retraces its path. The refractive index of the material of prism is

- a) $\sqrt{3}$ b) $3/2$ c) 2 d) $\sqrt{2}$

109. A source is at 4m height above the centre of a circular table of a circular table of radius 3m. The ratio of illuminance at O and P will be



- a) $\frac{64}{125}$ b) $\frac{125}{64}$ c) 1 d) $\frac{16}{25}$

110. The sun's diameter is $1.4 \times 10^9 m$ and its distance from the earth is $10^{11} m$. The diameter of its image, formed by a convex lens of focal length $2m$ will be

- a) 0.7 cm b) 1.4 cm
c) 2.8 cm d) Zero (i. e. point image)

111. A ray of light incident normally on one face of a right angled isosceles prism. It then grazes the hypotenuse. The refractive index of the material of the prism is

- a) 1.33 b) 1.414 c) 1.5 d) 1.732


112. Each quarter of a vessel of depth H is filled with liquids of the refractive indices n_1, n_2, n_3 and n_4 from the bottom respectively. The apparent depth of the vessel when looked normally is

- a) $\frac{H(n_1 + n_2 + n_3 + n_4)}{4}$ b) $\frac{H\left(\frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} + \frac{1}{n_4}\right)}{4}$ c) $\frac{(n_1 + n_2 + n_3 + n_4)}{4H}$ d) $\frac{H\left(\frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} + \frac{1}{n_4}\right)}{2}$


113. In the formation of primary rainbow, the sunlight rays emerge at minimum deviation from rain-drop after

- a) One internal reflection and one refraction
b) One internal reflection and two refraction
c) Two internal reflection and one refraction
d) Two internal reflection and one refraction

114. Which of the following graphs is the magnification of a real image against the distance from the focus of a concave mirror








115. A triangular prism of glass is shown in the figure. A ray incident normally to one face is totally reflected, if $\theta = 5^\circ$. The index of refraction of glass is



- a) Less than 1.41 b) Equal to 1.41 c) Greater than 1.41 d) None of the above

116. PQR is a right angled prism with other angles as 60° and 30° . Refractive index of prism is 1.5. PQ has a thin layer of liquid. Light falls normally on the face PR . For total internal reflection, maximum refractive index of liquid is



- a) 1.4 b) 1.3 c) 1.2 d) 1.6
117. Which one of the following is not associated with total internal reflection
- a) The mirage formation b) Optical fiber communication
c) The glittering of diamond d) Dispersion of light
118. As the position of an object (u) reflected from a concave mirror is varied, the position of the image (v) also varies. By letting the u changes from 0 to $+\infty$ the graph between v versus u will be
- a)  b)  c)  d) 
119. Dispersive power depends on the following
- a) Material of the prism b) Shape of the prism
c) Size of the prism d) Size, shape and material of the prism
120. The communication using optical fibres is based on the principle of
- a) Total internal reflection b) Brewster angle
c) Polarization d) Resonance
121. A monochromatic beam of light passes from a denser medium into a rarer medium. As a result
- a) Its velocity increases b) Its velocity decreases
c) Its frequency decreases d) Its wavelength decreases
122. In an astronomical telescope in normal adjustment, a straight black line of length L is drawn on the objective lens. The eyepiece forms a real image of this line. The length of this image is l . The magnification of the telescope is
- a) $\frac{L}{l}$ b) $\frac{L}{l} + 1$ c) $\frac{L}{l} - 1$ d) $\frac{L + l}{L - l}$
123. A microscope is focused on an ink mark on the top of a table. If we place a glass slab of 3 cm thick on it, how should the microscope be moved to focus the ink spot again? The refractive index of glass is 1.5.
- a) 2 cm upwards b) 2 cm downwards c) 1 cm upwards d) 1 cm downwards
124. With respect to air critical angle in a medium for light of red colour (λ_1) is θ . Other facts remaining same, critical angle for light of yellow colour [λ_2] will be
- a) θ b) More than θ c) Less than θ d) $\frac{\theta \lambda_1}{\lambda_2}$
125. Total internal reflection of a ray of light is possible when the (i_c = critical angle, i = angle of incidence)
- a) Ray goes from denser medium to rarer medium and $i < i_c$
b) Ray goes from denser medium to rarer medium and $i > i_c$
c) Ray goes from rarer medium to denser medium and $i > i_c$
d) Ray goes from rarer medium to denser medium and $i < i_c$
126. What is the time taken (in seconds) to cross a glass of thickness 4 mm and $\mu = 3$ by light
- a) 4×10^{-11} b) 2×10^{-11} c) 16×10^{-11} d) 8×10^{-10}
127. A compound microscope has an objective and eye-piece as thin lenses of focal lengths 1 cm and 5 cm respectively. The distance between the objective and the eye-piece is 20 cm. The distance at which the

- object must be placed in front of the objective if the final image is located at 25 cm from the eye-piece, it numerically
- a) 95/6 cm b) 5 cm c) 95/89 cm d) 25/6 cm
128. From which source a continuous emission spectrum and a line absorption spectrum are simultaneously obtained
- a) Bunsen burner flame b) The sun
c) Tube light d) Hot filament of an electric bulb
129. In a thin prism of glass (refractive index 1.5), which of the following relations between the angle of minimum deviations δ_m and angle of refraction r will be correct
- a) $\delta_m = r$ b) $\delta_m = 1.5r$ c) $\delta_m = 2r$ d) $\delta_m = \frac{r}{2}$
130. A person cannot see properly beyond 2 m. Power of the lens is
- a) 0.5 D b) 1.5 D c) -2.5 D d) -0.5 D
131. A thin glass (refractive index 1.5) lens has optical power of -5 D in air. Its optical power in a liquid medium with refractive index 1.6 will be
- a) 1 D b) -1 D c) 25 D d) -25 D
132. A bi-convex lens made of glass (refractive index 1.5) is put in a liquid of refractive index 1.7. Its focal length will
- a) Decrease and change sign b) Increase and change sign
c) Decrease and remain of the same sign d) Increase and remain of the same sign
133. If luminous efficiency of a lamp is 2 lumen/watt and its luminous intensity is 42 candela, then power of the lamp is
- a) 62 W b) 76 W c) 1.38 W d) 264 W
134. What cause chromatic aberration?
- a) Non-paraxial rays
b) Paraxial rays
c) Variation of focal length with colour
d) Difference in radii of curvature of the bounding surface of the lens
135. Two plane mirrors inclined to each other at an angle 72° , what is the number of image formed?
- a) 3 b) 5 c) 9 d) 7
136. In an optics experiments, with the position of the object fixed, a student varies the position of a convex lens and for each position, the screen is adjusted to get a clear image of the object. A graph between the object distance u and the image distance v , from the lens, is plotted using the same scale for the two axes. A straight line passing through the origin and making an angle of 45° with the x -axis meets the experimental curve at P . The coordinates of P will be
- a) $(2f, 2f)$ b) $(\frac{f}{2}, \frac{f}{2})$ c) (f, f) d) $(4f, 4f)$
137. The aperture of a telescope is made large, because
- a) To increase the intensity of image b) To decrease the intensity of image
c) To have greater magnification d) To have lesser resolution
138. Four convergent lenses have focal lengths 100 cm, 10 cm, 4 cm and 0.3 cm. For a telescope with maximum possible magnification, we choose the lenses of focal length
- a) 100 cm, 0.3 cm b) 10 cm, 0.3 cm c) 10 cm, 4 cm d) 100 cm, 4 cm
139. The focal length of a convex lens depends upon
- a) Frequency of the light ray b) Wavelength of the light ray
c) Both (a) and (b) d) None of these
140. A virtual image three times the size of the object is obtained with a concave mirror of curvature 36 cm. The distance of the object from the mirror is
- a) 5 cm b) 12 cm c) 10 cm d) 20 cm
141. The spectrum of light emitted by a glowing solid is

- a) $\frac{f}{(m-1)}$ b) $\frac{f}{(m+1)}$ c) $f(m-1)$ d) $f(m+1)$

- a) Myopia b) Presbyopia c) Astigmatism d) Hypermetropia

-

- a) 0.1 m b) 0.4 m c) 0.9 m d) 1 m

-

- a) Total internal reflection
b) Scattering
c) Refraction
d) Dispersion and total internal reflection

- a) 2 cm b) 12 cm c) 4 cm d) 1.2 cm

- a) 120° b) 30° c) 60° d) 90°

- a) $\frac{25}{f}$ b) $\frac{D}{26}$ c) $\frac{f}{25}$ d) $\frac{f}{D+1}$

- a) $+1 D$ b) $-2 D$ c) $+3 D$ d) $+4 D$

- a) Watt b) Horse power c) Diopetre d) Lux


- A telescope of diameter $2m$ uses light of wavelength 5000 \AA for viewing stars. The minimum angular separation between two stars whose image is just resolved by this telescope is

- a) $4 \times 10^{-4} \text{ rad}$ b) $0.25 \times 10^{-6} \text{ rad}$ c) $0.31 \times 10^{-6} \text{ rad}$ d) $5.0 \times 10^{-3} \text{ rad}$

- PHONE NO: 8583042324 Page | 13


- a) 45° b) 60° c) 75° d) 30°

155. A light beam is travelling from Region I to Region IV (refer figure). The refractive index in Region I, II, III and IV are n_0 , $\frac{n_0}{2}$, $\frac{n_0}{6}$ and $\frac{n_0}{8}$, respectively. The angle of incidence θ for which the beam just misses entering Region IV is



- a) $\sin^{-1}\left(\frac{3}{4}\right)$ b) $\sin^{-1}\left(\frac{1}{8}\right)$ c) $\sin^{-1}\left(\frac{1}{4}\right)$ d) $\sin^{-1}\left(\frac{1}{3}\right)$

156. A rectangular glass slab $ABCD$, of refractive index n_1 , is immersed in water of refractive index n_2 ($n_1 > n_2$). A ray of light is incident at the surface AB of the slab as shown. The maximum value of the angle of incidence α_{max} such that the ray comes out only from the other surface CD is given by



- a) $\sin^{-1}\left[\frac{n_1}{n_2} \cos\left(\sin^{-1}\frac{n_2}{n_1}\right)\right]$ b) $\sin^{-1}\left[n_1 \cos\left(\sin^{-1}\frac{1}{n_2}\right)\right]$
c) $\sin^{-1}\left(\frac{n_1}{n_2}\right)$ d) $\sin^{-1}\left(\frac{n_2}{n_1}\right)$

157. A vessel of depth $2d$ cm is half filled with a liquid of refractive index μ_1 and the upper half with a liquid of refractive index μ_2 . The apparent depth of the vessel seen perpendicularly is

- a) $d\left(\frac{\mu_1\mu_2}{\mu_1 + \mu_2}\right)$ b) $d\left(\frac{1}{\mu_1} + \frac{1}{\mu_2}\right)$ c) $2d\left(\frac{1}{\mu_1} + \frac{1}{\mu_2}\right)$ d) $2d\left(\frac{1}{\mu_1\mu_2}\right)$


158. 1% of light of a source with luminous intensity 50 *candela* is incident on a circular surface of radius 10 cm. The average illuminance of surface is

- a) 100 lux b) 200 lux c) 300 lux d) 400 lux

159. If both the object and image are at infinite distance from a refracting telescope its magnifying power will be equal to

- a) The sum of the focal lengths of the objective and the eyepiece
b) The different of the focal lengths of the two lenses
c) The ratio of the focal length of the objective and eyepiece
d) The ratio of the focal length of the eyepiece and objective

160. One side of a glass slab is silvered as shown. A ray of light is incident on the other side at angle of incidence $i = 45^\circ$. Refractive index of glass is given as 1.5, the deviation of the ray of light from its initial path when it comes out of the slab is



- a) 90° b) 180° c) 120° d) 45°


161. The index of refraction of diamond is 2.0. The velocity of light in diamond is approximately


- a) $1.5 \times 10^{10} \text{ cms}^{-1}$ b) $2 \times 10^{10} \text{ cms}^{-1}$ c) $3.0 \times 10^{10} \text{ cms}^{-1}$ d) $6 \times 10^{10} \text{ cms}^{-1}$

162. Image formed by a convex lens is virtual and erect when the object is placed

- a) At F b) Between F and the lens
c) At $2F$ d) Beyond $2F$

163. Dark lines on solar spectrum are due to

- a) Lack of certain elements
b) Black body radiation
c) Absorption of certain wavelengths by outer layers
d) Scattering
164. Where should a person stand straight from the pole of a convex mirror of focal length 2.0 m on its axis so that the image formed become half of his original height?
a) -2.60m b) -4.0m c) -0.5m d) -2.0m
165. Three prisms 1, 2 and 3 have the prism angle $A = 60^\circ$, but their refractive indices are respectively 1.4, 1.5 and 1.6. If $\delta_1, \delta_2, \delta_3$ be their respective angles of deviation then
a) $\delta_3 > \delta_2 > \delta_1$ b) $\delta_1 > \delta_2 > \delta_3$ c) $\delta_1 = \delta_2 = \delta_3$ d) $\delta_2 > \delta_1 > \delta_3$
166. The position of final image formed by the given lens combination from the third lens will be at a distance of [$f_1 = +10$ cm, $f_2 = -10$ cm, $f_3 = +30$ cm]
- 
- a) 15 cm b) Infinity c) 45 cm d) 30 cm
167. The focal length of a convex lens is 10 cm and its refractive index is 1.5. If the radius of curvature of one surface is 7.5 cm, the radius of curvature of the second surface will be
a) 7.5 cm b) 15.0 cm c) 75 cm d) 5.0 cm
168. A convex lens of focal length f is placed some where in between an object and a screen. The distance between object and screen is x . If numerical value of magnification produced by lens is m , focal length of lens is
a) $\frac{mx}{(m+1)^2}$ b) $\frac{mx}{(m-1)^2}$ c) $\frac{(m+1)^2}{m}x$ d) $\frac{(m-1)^2}{m}x$
169. An object of height 1.5 cm is placed on the axis of a convex lens of focal length 25 cm. A real image is formed at a distance of 75 cm from the lens. The size of the image will be
a) 4.5 cm b) 3.0 cm c) 0.75 cm d) 0.5 cm
170. An achromatic combination of lenses is formed by joining
a) 2 convex lenses b) 2 concave lenses
c) 1 convex lens and 1 concave lens d) Convex lens and plane mirror
171. A thin convex lens of focal length 10 cm is placed in contact with a concave lens of same material and of same focal length. The focal length of combination will be
a) Zero b) Infinity c) 10 cm d) 20 cm
172. A thin equiconvex lens of refractive index $3/2$ and radius of curvature 30 m is put in water (refractive index $= \frac{4}{3}$). Its focal length is
a) 0.15 m b) 0.30 m c) 0.45 m d) 1.20 m
173. An astronaut in a spaceship see the outer space as
a) White b) Black c) Blue d) Red
174. Critical angle is that angle of incidence in the denser medium for which the angle of reflection in rarer medium is
a) 0° b) 57° c) 90° d) 180°
175. A ray falls on a prism ABC ($AB = BC$) and travels as shown in figure. The minimum refractive of the prism material should be







a) $\frac{4}{3}$


b) $\sqrt{2}$


c) 1.5

d) $\sqrt{3}$

176. In the position of minimum deviation when a ray of yellow light passes through the prism, then its angle of incidence is
- Less than the emergent angle
 - Greater than the emergent angle
 - Sum of angle incidence and emergent angle is 90°
 - Equal to the emergent angle
177. To increase the magnifying power of telescope (f_o = focal length of the objective and f_e = focal length of the eye lens)
- f_o should be large and f_e should be small
 - f_o should be small and f_e should be large
 - f_o and f_e both should be large
 - f_o and f_e both should be small
178. A lens is placed between a source of light and a wall. It forms images of area A_1 and A_2 on the wall for its two different positions. The area of the source or light is
- $\frac{A_1 + A_2}{2}$
 - $\left[\frac{1}{A_1} + \frac{1}{A_2} \right]^{-1}$
 - $\sqrt{A_1 A_2}$
 - $\left[\frac{\sqrt{A_1} + \sqrt{A_2}}{2} \right]^2$
179. An achromatic combination of lenses produces
- Images in black and white
 - Coloured images
 - Images unaffected by variation of refractive index with wavelength
 - Highly enlarged images are formed
180. A diminished virtual image can be formed only in
- Plane mirror
 - A concave mirror
 - A convex mirror
 - Concave-parabolic mirror
181. Venus looks brighter than other stars because
- It has higher density than other stars
 - It is closer to the earth than other stars
 - It has no atmosphere
 - Atomic fission takes place on its surface
182. A ray of light travelling in water is incident on its surface open to air. The angle of incidence is θ , which is less than the critical angle. Then there will be
- Only a reflected ray and no refracted ray
 - Only a refracted ray and no reflected ray
 - A reflected ray and a refracted ray and the angle between them would be less than $108^\circ - 2\theta$
 - A reflected ray and a refracted ray and the angle between them would be greater than $108^\circ - 2\theta$
183. If the focal length of objective and eye lens are 1.2 cm and 3 cm respectively and the object is put 1.25 cm away from the objective lens and the final image is formed at infinity. The magnifying power of the microscope is
- 150
 - 200
 - 250
 - 400
184. The refractive index of a piece of transparent quartz is the greatest for
- Red light
 - Violet light
 - Green light
 - Yellow light

185. Refractive index for a material for infrared light is
 a) Equal to that of ultraviolet light
 b) Less than for ultraviolet light
 c) Equal to that for red colour of light
 d) Greater than that for ultraviolet light
186. In vacuum the speed of light depends upon
 a) Frequency
 b) Wave length
 c) Velocity of the source of light
 d) None of these
187. A point source of light moves in a straight line parallel to a plane table. Consider a small portion of the table directly below the line of movement of the source. The illuminance at this portion varies with this distance r from the source as
 a) $\propto \frac{1}{r}$
 b) $\propto \frac{1}{r^2}$
 c) $\propto \frac{1}{r^3}$
 d) $\propto \frac{1}{r^4}$
188. When a plane mirror is placed horizontally on a level ground at a distance of $60m$ from the foot of a tower, the top of the tower and its image in the mirror subtend an angle of 90° at the eye. The height of the tower will be
 a) $30m$
 b) $60m$
 c) $90m$
 d) $120m$
189. The distance v of the real image formed by a convex lens is measured for various object distance u . A graph is plotted between v and u . Which one of the following graphs is correct?
 a) 
 b) 
 c) 
 d) 
190. A plano-convex lens of refractive index 1.5 and radius of curvature 30 cm is silvered at the curved surface. Now, this lens has been used to form the image of an object. At what distance from this lens, an object be placed in order to have a real image of the size of the object?
 a) 20 cm
 b) 30 cm
 c) 60 cm
 d) 80 cm
191. A double convex lens of focal length 20 cm is made of glass of refractive index $3/2$. When placed completely in water ($\mu_w = 4/3$), its focal length will be
 a) 80 cm
 b) 15 cm
 c) 17.7 cm
 d) 22.5 cm
192. To a fish under water, viewing obliquely a fisherman standing on the bank of the lake, the man looks
 a) Taller than what he actually is
 b) Shorter than what he actually is
 c) The same height as he actually is
 d) Depends on the obliquity
193. The numerical aperture for a human eye is of the order of
 a) 1
 b) 0.1
 c) 0.01
 d) 0.001
194. Five *lumen/watt* is the luminous efficiency of a lamp and its luminous intensity is 35 candela . The power of the lamp is
 a) 80 W
 b) 176 W
 c) 88 W
 d) 36 W
195. If the speed of light in vacuum is $C\text{ m/sec}$, then the velocity of light in a medium of refractive index 1.5
 a) Is $1.5 \times C$
 b) Is C
 c) Is $\frac{C}{1.5}$
 d) Can have any velocity
196. When sunlight is scattered by minute particles of atmosphere, the intensity of light scattered away is proportional to
 a) $(\text{wavelength of light})^4$
 b) $(\text{frequency of light})^4$
 c) $(\text{wavelength of light})^2$
 d) $(\text{frequency of light})^2$
197. A ray of light is incident at the glass-water interface at an angle i it emerges finally parallel to the surface of water, then the value of μ_g would be



- a) $(4/3) \sin i$ b) $1/\sin i$ c) $4/3$ d) 1
198. The wavelength of light in air and some other medium are respectively λ_a and λ_m . The refractive index of medium is
 a) λ_a/λ_m b) λ_m/λ_a c) $\lambda_a \times \lambda_m$ d) None of these
199. The plane face of a planoconvex lens is silvered. If μ be the refractive index and R , the radius of curvature of curved surface, then the system will behave like a concave mirror of radius of curvature
 a) μR b) $\frac{R}{(\mu - 1)}$ c) $\frac{R^2}{\mu}$ d) $\left[\frac{(\mu + 1)}{(\mu - 1)} \right] R$
200. The slab of a material of refractive index 2 shown in figure has curved surface APB of radius of curvature 10 cm and a plane surface CD . On the left of APB is air and on the right of CD is water with refractive indices as given in figure. An object O is placed at a distance of 15 cm from pole P as shown. The distance of the final image of O from P , as viewed from the left is

 a) 20 cm b) 30 cm c) 40 cm d) 50 cm
201. A lens of refractive index n is put in a liquid of refractive index n' . If focal length of lens in air is f , its focal length in liquid will be
 a) $\frac{fn'(n-1)}{n'-n}$ b) $\frac{f(n'-n)}{n'(n-1)}$ c) $\frac{n'(n-1)}{f(n'-n)}$ d) $\frac{fn'n}{n-n'}$
202. Spherical aberration in a lens
 a) Is minimum when most of the deviation is at the first surface
 b) Is minimum when most of the deviation is at the second surface
 c) Is minimum when the total deviation is equally distributed over the two surfaces
 d) Does not depend on the above considerations
203. The frequency of light in air is 5×10^{14} Hz. What will be the frequency of light, when it enters in the water?
 a) 2.5×10^{14} Hz b) 5×10^{14} Hz c) 10^{15} Hz d) 2.5×10^{12} Hz
204. The refractive index of water and glycerine are 1.33 and 1.47 respectively. What is the critical angle for a light ray going from the latter to the former?
 a) $60^\circ 48'$ b) $64^\circ 48'$ c) $74^\circ 48'$ d) None of these
205. Which of the following is a wrong statement?
 a) $D = 1/f$ where, f is the focal length and D is called the refractive power of a lens
 b) Power is expressed in a diopter when f is in metre
 c) Power is expressed in diopter and does not depend on the system of unit used to measure f
 d) D is positive for convergent lens and negative for divergent lens
206. A telescope has an objective of focal length 50 cm and an eye piece of focal length 5 cm. The least distance of distinct vision is 25 cm. The telescope is focussed for distinct vision on a scale 200 cm away. The separation between the objective and the eye-piece is
 a) 75 cm b) 60 cm c) 71 cm d) 74 cm


207. A point object is placed at a distance of 25 cm from a convex lens of focal length 20 cm. If a glass slab of thickness t and refractive index 1.5 is inserted between the lens and the object, the image is formed at infinity. The thickness t is

- a) 15 cm b) 5 cm c) 10 cm d) 20 cm

208. A concave lens with unequal radii of curvature made of glass ($\mu_g = 1.5$) has a focal length of 40 cm. If it is immersed in a liquid of refractive index $\mu_l = 2$, then

- a) It behave like a convex lens of 80 cm focal length
b) It behave like a concave lens of 20 cm focal length
c) Its focal length becomes 60 cm
d) Nothing can be said

209. A fish is a little away below the surface of a lake. If the critical angle is 49° , then the fish could see things above water surface within an angular range of θ° where



- a) $\theta = 49^\circ$ b) $\theta = 98^\circ$ c) $\theta = 24\frac{1}{4}^\circ$ d) $\theta = 90^\circ$

210. A short sighted person can see distinctly only those objects which lie between 10 cm and 100 cm from him. The power of the spectacle lens required to see a distant object is

- a) +0.5 D b) -1.0 D c) -10 D d) +4.0 D

211. The focal length of a plano convex lens is f and its refractive index is 1.5. It is kept over a plane glass plate with its curved surface touching the glass plate is filled by a liquid. As a result, the effective focal length of the combination becomes $2f$. Then the refractive index of the liquid is

- a) 1.5 b) 2 c) 1.25 d) 1.33

212. The focal length of a concave mirror is f and the distance from the object to the principle focus is x . The ratio of the size of the image to the size of the object is

- a) $\frac{f+x}{f}$ b) $\frac{f}{x}$ c) $\sqrt{\frac{f}{x}}$ d) $\frac{f^2}{x^2}$

213. A circular disc of which $\frac{2}{3}$ part is coated with yellow and $\frac{1}{3}$ part is with blue. It is rotated about its central axis with high velocity, then it will be seen as

- a) Green b) Brown c) White d) Violet


214. The magnifying power of a microscope with an objective of 5 mm focal length is 400. The length of its tube is 20 cm. Then the focal length of the eye-piece is

- a) 200 cm b) 160 cm c) 2.5 cm d) 0.1 cm

215. When the object is self-luminous, the resolving power of a microscope is given by the expression

- a) $\frac{2\mu \sin \theta}{1.22 \lambda}$ b) $\frac{\mu \sin \theta}{\lambda}$ c) $\frac{2\mu \cos \theta}{1.22 \lambda}$ d) $\frac{2\mu}{\lambda}$

216. For an optical arrangement shown in the figure. Find the position and nature of images




- a) 32 cm b) 0.6 cm c) 6 cm d) 0.5 cm


217. A concave mirror of focal length ' f_1 ' is placed at a distance of ' d ' from a convex lens of focal length ' f_2 '. A beam of light coming from infinity and falling on this convex lens-concave mirror combination returns to infinity. The distance ' d ' must equal

- a) $f_1 + f_2$ b) $-f_2 + f_2$ c) $2f_1 + f_2$ d) $-2f_1 + f_2$

218. The graph shows variation of v with change in u for a mirror. Points plotted above the point P on the curve are for values of v




- a) Smaller than f b) Smaller than $2f$ c) Larger than $2f$ d) Larger than f
219. A glass convex lens ($\mu_g = 1.5$) has a focal length of 8 cm when placed in air. What would be the focal length of the lens what it is immersed in water ($\mu_w = 1.33$)
- a) 2 m b) 4 cm c) 16 cm d) 32 cm
220. An electric lamp is fixed at the ceiling of a circular tunnel as shown is figure. What is the ratio the intensities of light at base A and a point B on the wall




- a) $1 : 2$ b) $2 : \sqrt{3}$ c) $\sqrt{3} : 1$ d) $1 : \sqrt{2}$
221. A sitting sun appears to be at an altitude higher than it really is. This is because of
- a) Absorption of light b) Refection of light c) Refraction of light d) Dispersion of light
222. A ray of light falls on the surface of a spherical glass paper weight making an angle α with the normal and is refracted in the medium at an angle β . The angle of deviation of the emergent ray from the direction of the incident ray
- a) $(\alpha - \beta)$ b) $2(\alpha - \beta)$ c) $(\alpha - \beta)/2$ d) $(\beta - \alpha)$
223. If a ray of light in a denser medium enters into a rarer medium at an angle of incidence i , the angle of reflection and refraction are respectively r and r' . If the reflected and refracted rays are at right angles to each other, the critical angle for the given pair of media is
- a) $\sin^{-1}(\tan r')$ b) $\sin^{-1}(\tan r)$ c) $\tan^{-1}(\sin i)$ d) $\cot(\tan i)$
224. Material A has critical angle i_A , and material B has critical angle i_B ($i_B > i_A$). Then which of the following is true?
- (i) Light can be totally internally reflected when it passes from B to A
- (ii) Light can be totally internally reflected when it passes from A to B
- (iii) Critical angle for total internal reflection is $i_B - i_A$
- (iv) Critical angle between A and B is $\sin^{-1}\left(\frac{\sin i_A}{\sin i_B}\right)$
- a) (i) and (iii) b) (i) and (iv) c) (ii) and (iii) d) (ii) and (iv)
225. If angle of incidence is twice the angle of refraction in a medium of refractive index μ , then angle of incidence is
- a) $2 \cos^{-1}\left[\frac{\mu}{2}\right]$ b) $2 \sin^{-1}\left[\frac{\mu}{2}\right]$ c) $2 \cos^{-1}[\mu]$ d) $2 \sin^{-1}[\mu]$
226. An object is at a distance of 0.5 m in front of a plane mirror. Distance between the object and image is
- a) 0.5 m b) 1 m c) 0.25 m d) 1.5 m
227. A hollow double concave lens is made of very thin transparent material. It can be filled with air or either of two liquids L_1 and L_2 having refractive indices n_1 and n_2 respectively ($n_2 > n_1 > 1$). The lens will diverge a parallel beam of light if it is filled with
- a) Air and placed in air b) Air and immersed in L_1
- c) L_1 and immersed in L_2 d) L_2 and immersed in L_1

228. The distance between a convex lens and a plane mirror is 10 cm. The parallel rays incident on the convex lens after reflection from the mirror form image at the optical centre of the lens. Focal length of lens will be








- a) 10 cm
b) 20 cm
c) 30 cm
d) Cannot be determined
229. Which of the prism is used to see infrared spectrum of light
a) Rock salt
b) Nicol
c) Flint
d) Crown
230. A concave mirror is placed on a horizontal table with its axis directed vertically upwards. Let O be the pole of the mirror and C its centre of curvature. A point object is placed at C . It has a real image, also located at C . If the mirror is now filled with water, the image will be
a) Real and will remain at C
b) Real, and located at a point between C and ∞
c) Virtual and located at a point between C and O
d) Real, and located at a point between C and O
231. A point object O is kept at a distance of $OP = u$. The radius of curvature of the spherical surface APB is $CP = R$. The refractive index of the media are n_1 and n_2 which are as shown in diagram. Then,



- (1) If $n_1 > n_2$, image is virtual for all values of u
(2) If $n_2 = 2n_1$, image is virtual when $R > u$
(3) The image is real for all values of u, n_1 and n_2 . Here, the correct statements is/are
a) Only (2)
b) Both (1) and (2)
c) Only (1)
d) (1), (2) and (3)
232. In a compound microscope, if the objective produces an image I_o and the eye piece produces an image I_e , then
a) I_o is virtual but I_e is real
b) I_o is real but I_e is virtual
c) I_o and I_e are both real
d) I_o and I_e are both virtual
233. A Gallilean telescope has objective and eye-piece of focal lengths 200 cm and 2 cm respectively. The magnifying power of the telescope for normal vision is
a) 90
b) 100
c) 108
d) 198
234. Angle of a prism is 30° and its refractive index is $\sqrt{2}$ and one of the surface is silvered. At what angle of incidence, a ray should be incident on one surface so that after reflection from the silvered surface, it retraces its path
a) 30°
b) 60°
c) 45°
d) $\sin^{-1} \sqrt{1.5}$
235. A convex lens made of glass has focal length 0.15 m in air. If the refractive index of glass is $\frac{3}{2}$ and that of water is $\frac{4}{3}$, the focal length of lens when immersed in water is
a) 0.45 m
b) 0.15 m
c) 0.30 m
d) 0.6 m
236. If sound travelling at 340 ms^{-1} enters water where its speed becomes 1480 ms^{-1} , then critical angle for total internal reflection is
a) 13.3°
b) 89.7°
c) 86.7°
d) 10.3°

237. Image of an object approaching a convex mirror of radius of curvature 20 m along its optical axis is observed to move from $\frac{25}{3}$ m to $\frac{50}{7}$ m in 30 s. What is the speed of the object in kmh^{-1} ?
- a) 3 b) 4 c) 5 d) 6
238. The minimum magnifying power of a telescope is M , If the focal length of its eye lens is halved, the magnifying power will become
- a) $M/2$ b) $2M$ c) $3M$ d) $4M$
239. The splitting of white light into several colours on passing through a glass prism is due to
- a) Refraction b) Reflection c) Interference d) Diffraction
240. A ray of light propagates from glass (refractive index = $\frac{3}{2}$) to water (refractive index = $\frac{4}{3}$). The value of the critical angle is
- a) $\sin^{-1}\left(\frac{1}{2}\right)$ b) $\sin^{-1}\left(\sqrt{\frac{9}{8}}\right)$ c) $\sin^{-1}\left(\frac{8}{9}\right)$ d) $\sin^{-1}\left(\frac{5}{9}\right)$
241. A small source of light is to be suspended directly above the centre of a circular table of radius R . What should be the height of the light source above the table so that the intensity of light is maximum at the edges of the table compared to any other height of the source
- a) $\frac{R}{2}$ b) $\frac{R}{\sqrt{2}}$ c) R d) $\sqrt{2}R$
242. A ray of light is incident on a plane mirror at an angle 57° . The resultant polarized light vibrates in a plane which makes an angle with the reflecting surface
- a) 0° b) 90° c) 57° d) 33°
243. In a laboratory four convex lenses L_1, L_2, L_3 and L_4 of focal length 2, 4, 6 and 8 cm respectively are available. Two of these lenses from a telescope of length 10 cm and magnifying power 4. The objective and eye lenses are respectively
- a) L_2, L_3 b) L_1, L_4 c) L_1, L_2 d) L_4, L_1
244. A hypermetropic person having near point at a distance of 0.75 m puts on spectacles of power 2.5 D. The near point now is at
- a) 0.75 m b) 0.83 m c) 0.26 cm d) 0.26 m
245. If the refractive angles of two prisms made of crown glass are 10° and 20° respectively, then the ratio of their colour deviation powers will be
- a) 1 : 1 b) 2 : 1 c) 4 : 1 d) 1 : 2
246. An movie projector forms an image 3.5m long of an object 35mm. Supposing there is negligible absorption of light by aperture then illuminance on slide and screen will be in the ratio of
- a) 100 : 1 b) 10^4 : 1 c) 1 : 100 d) 1 : 10^4
247. Two convex lenses of focal lengths 0.03 m and 0.05 m are used to make a telescope. The distance kept between the two in order to obtain an image at infinity is
- a) 0.35 cm b) 0.25 cm c) 0.175 m d) 0.15 m
248. Angle of deviation (δ) by a prism (refractive index = μ and supposing the angle of prism A to be small) can be given by
- a) $\delta = (\mu - 1)A$ b) $\delta = (\mu + 1)A$ c) $\delta = \frac{\sin \frac{A+\delta}{2}}{\sin \frac{A}{2}}$ d) $\delta = \frac{\mu - 1}{\mu + 1}A$
249. It is desired to make a converging achromatic combination of mean focal length 50 cm by using two lenses of materials A and B . If the dispersive power of A and B are in ratio 1:2, the focal lengths of the convex and the concave lenses are respectively
- a) 25 cm and 50 cm b) 50 cm and 25 cm c) 50 cm and 100 cm d) 100 cm and 50 cm
250. Which of the following is not correct regarding the ratio telescope
- a) It can not work at night
b) It can detect a very faint radio signal

- c) It can be operated even in cloudy weather
d) It is much cheaper than optical telescope
251. The principal section of a glass prism is an isosceles triangle ABC with $AB = AC$. The face AC is silvered. A ray of light is incident normally on the face AB and after two reflections, it emerges from the base BC perpendicular to the base. Angle BAC of the prism is
a) 30° b) 36° c) 60° d) 72°
252. Two thin lenses, one of focal length $+60\text{ cm}$ and the other of focal length -20 cm are put in contact. The combined focal length is
a) $+15\text{ cm}$ b) -15 cm c) $+30\text{ cm}$ d) -30 cm
253. A lamp is hanging at a height of 4 m above a table. The lamp is lowered by 1 m . The percentage increase in illuminance will be
a) 40% b) 64% c) 78% d) 92%
254. The light gathering power of a camera lens depends on
a) Its diameter only b) Ratio of diameter and focal length
c) Product of focal length and diameter d) Wavelength of light used
255. If there had been one eye of the man, then
a) Image of the object would have been inverted
b) Visible region would have decreased
c) Image would have not been seen three dimensional
d) (b) and (c) both
256. Line spectra are due to
a) Hot solids b) Atoms in gaseous state
c) Molecules in gaseous state d) Liquid at low temperature
257. A graph is plotted between angle of deviation (δ) and angle of incidence (i) for a prism. The nearly correct graph is
a)  b)  c)  d) 
258. The ratio of the refractive index of red light to blue light in air is
a) Less than unity
b) Equal to unity
c) Greater than unity
d) Less as well as greater than unity depending upon the experimental arrangement
259. A man has a concave shaving mirror of focal length 0.2 m . How far should the mirror be held from his face in order to give an image of two fold magnification?
a) 0.1 m b) 0.2 m c) 0.3 m d) 0.4 m
260. An isosceles prism of angle 120° has a refractive index of 1.44 . Two parallel monochromatic rays enter the prism parallel to each other in air as shown. The rays emerging from the opposite faces



- a) Are parallel to each other
b) Are diverging
c) Make an angle $2\sin^{-1}(0.72)$ with each other
d) Make an angle $2\{\sin^{-1}(0.72) - 30^\circ\}$ with each other
261. Inverse square law for illuminance is valid for
a) Isotropic point source
b) Cylindrical source
c) Search light
d) All type of sources
262. A person who can see things most clearly at a distance of 10 cm. Requires spectacles to enable to him to see clearly things at a distance of 30 cm. What should be the focal length of the spectacles
a) 15 cm (Concave) b) 15 cm (Convex) c) 10 cm d) 0
263. A thin convex lens of crown glass having refractive index 1.5 has power 1 D. What will be the power of similar convex lens refractive index 1.6?
a) 0.6 D b) 0.8 D c) 1.2 D d) 1.6 D
264. A magnifying glass is to be used at the fixed object distance of 1 inch. If it is to produce an erect image magnified 5 times its focal length should be
a) 0.2 inch b) 0.8 inch c) 1.25 inch d) 5 inch
265. For a telescope to have large resolving power the
a) Focal length of its objective should be large
b) Focal length of its eye piece should be large
c) Focal length of its eye piece should be small
d) Aperture of its objective should be large
266. Angular resolving power of human eye is
a) 3.6×10^3 b) 3.6×10^2 c) 3.6×10^4 d) 3.6×10^6
267. A small lamp is hung at a height of 8 feet above the centre of a round table of diameter 16 feet. The ratio of intensities of illumination at the centre and at points on the circumference of the table will be
a) 1 : 1 b) 2 : 1 c) $2\sqrt{2} : 1$ d) 3 : 2
268. If the focal length of a double convex lens for red light is f_R , its focal length for the violet light is
a) f_R b) Greater than f_R c) Less than f_R d) $2f_R$
269. In the measurement of the angle of a prism using a spectrometer, the readings of first reflected image are Vernier I : $320^\circ 40'$; Vernier II : $140^\circ 30'$ and those of the second reflected image are Vernier I : $80^\circ 38'$; Vernier II : $260^\circ 24'$. Then the angle of the prism is
a) $59^\circ 58'$ b) $59^\circ 56'$ c) $60^\circ 2'$ d) $60^\circ 4'$
270. The focal length of convex lens is 30 cm and the size of image is quarter of the object, then the object distance is
a) 150 cm b) 60 cm c) 30 cm d) 40 cm
271. A car is fitted with a convex side view mirror of focal length 20 cm. A second car 2.8 m behind the first car is overtaking the first car is a relative speed of 15 m/s. The speed of the image of the second car as seen in the mirror of the first one is
a) $\frac{1}{15}$ m/s b) 10 m/s c) 15 m/s d) $\frac{1}{10}$ m/s
272. If the lower half of a concave mirror's reflecting surface is made opaque, which of the following statements describe the image of an object placed in front of the mirror
S1: Intensity of the image will increase
S2: The image will show only half of the object
S3: No change in the image
S4: Intensity of the image will be reduced to half
a) S1 only b) S2 only c) S2 and S3 d) S4 only
273. A concave mirror is used to focus the image of a flower on a nearby wall 120 cm from the flower. If a lateral magnification of 16 is desired, the distance of the flower from the mirror should be
a) 8 cm b) 12 cm c) 80 cm d) 120 cm

- PHONE NO: 8583042324 Page | 25

- a) $(n - 1)f$ b) $(n + 1)f$ c) $\left(\frac{n - 1}{n}\right)f$ d) $\left(\frac{n + 1}{n}\right)f$

288. In a compound microscope, the focal length of the objective and the eye lens are 2.5 cm and 5 cm respectively. An object is placed at 3.75 cm before the objective and image is formed at the least distance of distinct vision, then the distance between two lenses will be (*i.e.* length of the microscopic tube)

- a) 11.67 cm b) 12.67 cm c) 13.00 cm d) 12.00 cm


289. In order to increase the angular magnification of a simple microscope, one should increase

- a) The object size b) The aperture of the lens
c) The focal length of the lens d) The power of the lens

290. In order to obtain a real image of magnification 2 using a converging lens of focal length 20 cm , where should an object be placed

- a) 50 cm b) 30 cm c) -50 cm d) -30 cm

291. For a concave mirror, if virtual image is formed, the graph between m and u is of the form



292. The magnifying power of an astronomical telescope is 10 and the focal length of its eye-piece is 20 cm . The focal length of its object will be

- a) 200 cm b) 2 cm c) 0.5 cm d) $0.5 \times 10^{-2}\text{ cm}$

293. Two point white dots are 1 mm apart on a black paper. They are viewed by eye of pupil diameter 3 mm . Approximately, what is the maximum distance at which these dots can be resolved by the eye? [Take wavelength of light = 500 nm]

- a) 5 m b) 1 m c) 6 m d) 3 m

294. A lamp is hanging at a height of 40 cm from the centre of the table. If its height is increased by 10 cm , the illuminance of the lamp will decreased by

- a) 10% b) 20% c) 27% d) 36%

295. When seen in green light, the saffron and green portions of our National Flag will appear to be

- a) Black b) Black and green respectively
c) Green d) Green and yellow respectively

296. If the wavelength of light in vacuum be λ , the wavelength in a medium of refractive index n will be

- a) $n\lambda$ b) $\frac{\lambda}{n}$ c) $\frac{\lambda}{n^2}$ d) $n^2\lambda$


297. The diameter of the objective of the telescope is 0.1 metre and wavelength of light is 6000 \AA . Its resolving power would be approximately

- a) $7.32 \times 10^{-6}\text{ rad}$ b) $1.36 \times 10^6\text{ rad}$ c) $7.32 \times 10^{-5}\text{ rad}$ d) $1.36 \times 10^5\text{ rad}$

298. A convex lens forms a real image of a point object placed on its principal axis. If the upper half of the lens is painted black, the image will


- a) Be shifted downwards b) Be shifted upwards
c) Not be shifted d) Shift on the principal axis

299. In the figure, an air lens of radii of curvature 10 cm ($R_1 = R_2 = 10\text{ cm}$) is cut in a cylinder of glass ($\mu = 1.5$). The focal length and the nature of the lens is




- a) 15 cm , concave b) 15 cm , convex
c) ∞ , neither concave nor convex d) 0 , concave

300. If a flint lens glass of dispersive power 0.0666 renders achromatic to a convex lens of crown glass of focal length 60 cm and dispersive power 0.033, then its focal length is
 a) -60 cm b) +60 cm c) -120 cm d) +120 cm
301. Two thin equiconvex lenses each of focal length 0.2 m are placed coaxially with their optic centers 0.5 m apart. Then the focal length of the combination is
 a) -0.4 m b) 0.4 m c) -0.1 m d) 0.1 m
302. Resolving power of a microscope depends upon
 a) The focal length and aperture of the eye lens
 b) The focal lengths of the objective and the eye lens
 c) The apertures of the objective and the eye lens
 d) The wavelength of light illuminating the object
303. Two convex lenses placed in contact form the image of a distant object at P . If the lens B is moved to the light, the image will




- a) Move to the left
 b) Move to the right
 c) Remain at P
 d) Move either to the left to right, depending upon focal length of the lenses
304. A ray of light passes from vacuum into a medium of refractive index μ , the angle of incidence is found to be twice the angle of refraction. The angle of incidence is
 a) $\cos^{-1}\left(\frac{\mu}{2}\right)$ b) $2 \cos^{-1}\left(\frac{\mu}{2}\right)$ c) $2 \sin^{-1}(\mu)$ d) $2 \sin^{-1}\left(\frac{\mu}{2}\right)$
305. A light ray travelling in glass medium is incident on glass-air interface at an angle of incidence θ . The reflected (R) and transmitted (T) intensities, both as function of θ , are plotted. The correct sketch is



306. The wavelength of light diminishes μ times ($\mu = 1.33$ for water) in a medium. A diver from inside water looks at an object whose natural colour is green. He sees the object as
 a) Green b) Blue c) Yellow d) Red
307. If the luminous intensity of a 100 W unidirectional bulb is 100 candela, then total luminous flux emitted from the bulb is
 a) 861 lumen b) 986 lumen c) 1256 lumen d) 1561 lumen
308. The refractive index of the material of a prism is $\sqrt{2}$ and the angle of prism is 30° . One of its refracting faces is polished. The incident beam of light will retrace back for angle of incidence
 a) 0° b) 45° c) 60° d) 90°
309. If the angle of minimum deviation is of 60° for an equilateral prism, then the refractive index of the material of the prism is
 a) 1.41 b) 1.5 c) 1.6 d) 1.73
310. An object is placed 12 cm to the left of a converging lens of focal length 8 cm. Another converging of 6 cm focal length is placed at a distance of 30 cm to the right of the first lens. The second lens will produce
 a) No image b) A virtual enlarged image


- c) A real enlarged image
d) A real smaller image
311. Red colour is used for danger signals because
a) It causes fear
b) It undergoes least scattering
c) It undergoes maximum scattering
d) It is in accordance with international convention
312. The wavelength of sodium light in air is 5890\AA . The velocity of light in air is $3 \times 10^8 \text{ ms}^{-1}$. The wavelength of light in a glass of refractive index 1.6 would be close to
a) 5890\AA
b) 3681\AA
c) 9424\AA
d) 15078\AA
313. An object is placed at a distance of 40 cm in front of a concave mirror of focal length 20 cm. The nature of image is
a) Real and inverted and of same size
b) Virtual and erect and of same size
c) Real and erect and of same size
d) Virtual and inverted and of same size
314. A ray of monochromatic light is incident on one refracting face of a prism of angle 75° . It passes through the prism and is incident on the other face at the critical angle. If the refractive index of the material of the prism is $\sqrt{2}$, the angle of incidence on the first face of the prism is
a) 30°
b) 45°
c) 60°
d) 0°
315. Under minimum deviation condition in a prism, if a ray is incident at an angle 30° , the angle between the emergent ray and the second refracting surface of the prism is
a) 0°
b) 30°
c) 45°
d) 60°
316. A ray PQ incident on the refracting face BA is refracted in the prism BAC as shown in the figure and emerges from the other refracting face AC as RS , such that $AQ = AR$. If the angle of prism $A = 60^\circ$ and the refractive index of the material of prism is $\sqrt{3}$, then the angle of deviation of the ray is



- a) 60°
b) 45°
c) 30°
d) None of these
317. The respective angles of the flint and crown glass prisms are A' and A . They are to be used for dispersion without deviation, then the ratio of their angles A'/A will be
a) $-\frac{(\mu_y - 1)}{(\mu_y' - 1)}$
b) $\frac{(\mu_y' - 1)}{(\mu_y - 1)}$
c) $(\mu_y' - 1)$
d) $(\mu_y - 1)$
318. All of the following statements are correct except
a) The total length of an astronomical telescope is the sum of the focal lengths of its two lenses
b) The image formed by the astronomical telescope is always erect because the effect of the combination of the two lenses is divergent
c) The magnification of an astronomical telescope can be increased by decreasing the focal length of the eye-piece
d) The magnifying power of the refracting type of astronomical telescope is the ratio of the focal length of the objective to that of the eye-piece
319. A plane mirror produces a magnification of
a) Zero
b) -1
c) +1
d) Between 0 and +1
320. A plane mirror is approaching you at 10 cms^{-1} . Your image shall approach you with a speed of
a) $+10 \text{ cms}^{-1}$
b) -10 cms^{-1}
c) $+20 \text{ cms}^{-1}$
d) -20 cms^{-1}
321. If a lens is cut into two pieces perpendicular to the principal axis and only one part is used, the intensity of the image

- a) Remains same b) $\frac{1}{2}$ times c) 2 times d) Infinite
322. The distance between a point source of light and a screen which is 60 cm is increased to 180 cm. The intensity on the screen as compared with the original intensity will be
a) (1/9) times b) (1/3) times c) 3 times d) 9 times
323. A plane convex lens is made of refractive index 1.6. The radius of curvature of the curved surface is 60 cm. The focal length of the lens is
a) 50 cm b) 100 cm c) 200 cm d) 400 cm
324. The refractive indices of the crown glass for blue and red light are 1.51 and 1.49 respectively and those of the flint glass are 1.77 and 1.73 respectively. An isosceles prism of angle 6° is made of crown glass. A beam of white light is incident at a small angle on this prism. The other flint glass isosceles prism is combined with the crown glass prism such that there is no deviation of the incident light
(i) Determine the angle of the flint glass prism
(ii) Calculate the net dispersion of the combined system
a) $-4^\circ, 0.04^\circ$, b) $4^\circ, 0.04$ c) $5^\circ, 0.04$ d) $-5, 0.04^\circ$
325. The relative luminosity of wavelength 600 nm is 0.6. Find the radiant flux of 600 nm needed to produce the same brightness sensation as produced by 120 W of radiant flux at 555 nm
a) 50W b) 72W c) $120 \times (0.6)^2 W$ d) 200W
326. Electromagnetic radiation of frequency n , wavelength λ , travelling with velocity v in air, enters a glass slab of refractive index μ . The frequency, wavelength and velocity of light in the glass slab will be respectively
a) $\frac{n}{\mu}, \frac{\lambda}{\mu}, \frac{v}{\mu}$ b) $n, \frac{\lambda}{\mu}, \frac{v}{\mu}$ c) $n, \lambda, \frac{v}{\mu}$ d) $\frac{n}{\mu}, \frac{\lambda}{\mu}, v$
327. White light is passed through a prism whose angle is 5° . If the refractive indices for rays of red and blue colour are respectively 1.64 and 1.66, the angle of deviation between the two colours will be
a) 0.1 degree b) 0.2 degree c) 0.3 degree d) 0.4 degree
328. Two plane mirrors are inclined to each other at an angle θ . A ray of light is reflected first at one mirror and then at the other. The total deviation of the ray is
a) 2θ b) $240^\circ - 2\theta$ c) $360^\circ - 2\theta$ d) $180^\circ - \theta$
329. If in a plano-convex lens, the radius of curvature of the convex surface is 10 cm and the focal length of the lens is 30 cm, then the refractive index of the material of lens will be
a) 1.5 b) 1.66 c) 1.33 d) 3
330. A mark at the bottom of a liquid appears to rise by 0.1 m. The depth of the liquid is 1 m. The refractive index of the liquid is
a) 1.33 b) $\frac{9}{10}$ c) $\frac{10}{9}$ d) 1.5
331. A cut diamond sparkles because of its
a) Hardness b) High refractive index
c) Emission of light by the diamond d) Absorption of light by the diamond
332. A light wave has a frequency of 4×10^{14} Hz and a wavelength of 5×10^{-7} metres in a medium. The refractive index of the medium is
a) 1.5 b) 1.33 c) 1.0 d) 0.66
333. A convex lens
a) Converges light rays b) Diverges light rays
c) Form real images always d) Always forms virtual images
334. A ray of light from a denser medium strikes a rarer medium at angle of incidence i . The reflected and refracted rays make an angle of 90° with each other. The angles of reflection and refraction are r and r' respectively. The critical angle is
a) $\sin^{-1}(\tan r')$ b) $\sin^{-1}(\tan r)$ c) $\tan^{-1}(\tan r')$ d) $\tan^{-1}(\tan i)$
335. A converging lens is to project the image of a lamp 4 times the size of the lamp on a wall at a distance of 10 m from the lamp. The focal length of the lens is

- a) 1.6 m b) 2.67 m c) 4.4 m d) -1.6 m
336. A real object is placed at a distance f from the pole of a convex mirror, in front of the convex mirror. If focal length of the mirror is f , then distance of the image from the pole of the mirror is
- a) $2f$ b) $\frac{f}{2}$ c) $4f$ d) $\frac{f}{4}$
337. The length of the compound microscope is 14 cm. The magnifying power for relaxed eye is 25. If the focal length of eye lens is 5 cm, then the object distance for objective lens will be
- a) 1.8 cm b) 1.5 cm c) 2.1 cm d) 2.4 cm
338. Light takes t_1 second to travel a distance x in vacuum and the same light takes t_2 second to travel $10x$ cm in a medium. Critical angle for corresponding medium will be
- a) $\sin^{-1}\left(\frac{10t_2}{t_1}\right)$ b) $\sin^{-1}\left(\frac{t_2}{10t_1}\right)$ c) $\sin^{-1}\left(\frac{10t_1}{t_2}\right)$ d) $\sin^{-1}\left(\frac{t_1}{10t_2}\right)$
339. Sparking of diamond is due to
- a) Reflection b) Dispersion
c) Total internal reflection d) High refractive index of diamond
340. The field of view is maximum for
- a) Plane mirror b) Concave mirror c) Convex mirror d) Cylindrical mirror
341. A man standing in a swimming pool looks at a stone lying at the bottom. The depth of the swimming pool is h . At what distance from the surface of water is the image of the stone formed (Line of vision is normal; Refractive index of water is n)
- a) h/n b) n/h c) h d) hn
342. A, B and C are the parallel sided transparent media of refractive indices n_1, n_2 and n_3 respectively. They are arranged as shown in the figure. A ray is incident at an angle i on the surface of separation of A and B which is as shown in the figure. After the refraction into the medium B , the ray grazes the surface of separation of the media B and C . Then, $\sin i$ equal to




- a) $\frac{n_3}{n_1}$ b) $\frac{n_1}{n_3}$ c) $\frac{n_2}{n_3}$ d) $\frac{n_1}{n_2}$
343. For a convex mirror, the variation of u versus v is given by
- a) b) c) d)
344. A thick plane mirror shows a number of images of the filament of an electric bulb. Of these, the brightest image is the
- a) First b) Second c) Fourth d) Last
345. The focal length of the objective lens of a compound microscope is
- a) Equal to the focal length of its eye piece b) Less than the focal length of eye piece
c) Greater than the focal length of eye piece d) Any of the above three
346. A man can see the objects upto a distance of one metre from his eyes. For correcting his eye sight so that he can see an object at infinity, he requires a lens whose power is

Or

A man can see upto 100 cm of the distant object. The power of the lens required to see far objects will be

- a) $+0.5 D$ b) $+1.0 D$ c) $+2.0 D$ d) $-1.0 D$

347. Three lenses L_1, L_2, L_3 are placed co-axially as shown in figure. Focal length's of lenses are given $30\text{ cm}, 10\text{ cm}$ and 5 cm respectively. If a parallel beam of light falling on lens L_1 , emerging L_3 as a convergent beam such that it converges at the focus of L_3 . Distance between L_1 and L_2 will be




- a) 40 cm b) 30 cm c) 20 cm d) 10 cm

348. How will the image formed by a convex lens be affected, if the central portion of the lens is wrapped in blank paper, as shown in the figure



- a) No image will be formed
b) Full image will be formed but is less bright
c) Full image will be formed but without the central portion
d) Two images will be formed, one due to each exposed half

349. As shown in figure, the liquid, L_1, L_2 and L_3 have refractive indices $1.55, 1.50$ and 1.20 respectively. Therefore, the arrangement corresponds to




- a) Biconvex lens b) Biconcave lens
c) Concave-convex lens d) Convexo-concave lens

350. For an angle of incidence θ on an equilateral prism of refractive index $\sqrt{3}$, the ray refracted is parallel to the base inside the prism. The value of θ is

- a) 30° b) 45° c) 60° d) 75°

351. A small fish 0.4 m below the surface of a lake, is viewed through a simple converging lens of focal length 3 m . The lens is kept at 0.2 m above the water surface such that fish lies on the optical axis of the lens. The image of the fish seen by observer will be at ($\mu_{\text{water}} = \frac{4}{3}$)




- a) A distance of 0.2 m from the water surface
b) A distance of 0.6 m from the water surface
c) A distance of 0.3 m from the water surface
d) The same location of fish

352. One face of a rectangular glass plate 6 cm thick is silvered. An object held 8 cm in front of the first face, forms an image 12 cm behind the silvered face. The refractive index of the glass is

- a) 0.4 b) 0.8 c) 1.2 d) 1.6

353. A thin lens made of glass of refractive index $\mu = 1.5$ has a focal length equals is 12 cm in air. It is now immersed in water ($\mu = \frac{4}{3}$). Its new focal length is
 a) 48 cm b) 36 cm c) 24 cm d) 12 cm
354. The optical path a monochromatic light is same if it goes through 4.0 cm of glass of 4.5 cm of water. If the refractive index of glass is 1.53, the refractive index of the water is
 a) 1.30 b) 1.36 c) 1.42 d) 1.46
355. A double convex lens of glass of $\mu = 1.5$ has radius of curvature of each of its surface is 0.2 m. The power of the lens is
 a) +10 dioptres b) -10 dioptres c) -5 dioptres d) +5 dioptres
356. A point source of light B is placed at a distance L in front of the centre of a mirror width d hung vertically on a wall. A man walks in front of the mirror along a line parallel to the mirror at a distance $2L$ from it as shown. The greater distance over which he can see the image of the light source in the mirror is



- a) $d/2$ b) d c) $2d$ d) $3d$
357. The speed of light in media M_1 and M_2 is 1.5×10^8 m/s and 2.0×10^8 m/s respectively. A ray of light enters from medium M_1 to M_2 at an incidence angle i . If the ray suffers total internal reflection, the value of i is
 a) Equal to $\sin^{-1}(\frac{2}{3})$ b) Equal to or less than $\sin^{-1}(\frac{3}{5})$
 c) Equal to or greater than $\sin^{-1}(\frac{3}{4})$ d) Less than $\sin^{-1}(\frac{2}{3})$
358. Formula for dispersive power is (where symbols have their usual meanings)


or

If the refractive indices of crown glass for red, yellow and violet colours are respectively μ_r, μ_y and μ_v , then the dispersive power of this glass would be

- a) $\frac{\mu_v - \mu_y}{\mu_r - 1}$ b) $\frac{\mu_v - \mu_r}{\mu_y - 1}$ c) $\frac{\mu_r - \mu_v}{\mu_v - \mu_r}$ d) $\frac{\mu_v - \mu_r}{\mu_v} - 1$
359. If F_o and F_e are the focal length of the objective and eye piece respectively of a telescope, then its magnifying power will be
 a) $F_o + F_e$ b) $F_o \times F_e$ c) F_o/F_e d) $\frac{1}{2} (F_o + F_e)$


360. Ability of the eye to see objects at all distances is called
 a) Binocular vision b) Myopia c) Hypermetropia d) Accommodation
361. The phenomena of total internal reflection is seen when angle of incidence is
 a) 90° b) Greater than critical angle
 c) Equal to critical angle d) 0°

362. A light ray from air is incident (as shown in figure) at one end of a glass fibre (refractive index $\mu = 1.5$) making an incidence angle of 60° on the lateral surface, so that it undergoes a total internal reflection. How much time would it take to traverse the straight fibre of length 1 km



- a) $3.33 \mu s$ b) $6.67 \mu s$ c) $5.77 \mu s$ d) $3.85 \mu s$
363. The refractive index of glass is 1.520 for red light and 1.525 for blue light. Let D_1 and D_2 be angles of minimum deviation for red and blue light respectively in a prism of this glass. Then ,

- a) $D_1 < D_2$
 b) $D_1 = D_2$
 c) D_1 can be less than or greater than D_2 depending upon the angle of prism
 d) $D_1 > D_2$
364. When light is refracted from air into glass
 a) Its wavelength and frequency both increase
 b) Its wavelength increases but frequency remains unchanged
 c) Its wavelength decreases but frequency remains unchanged
 d) Its wavelength and frequency both decrease
365. A Galilean telescope has an objective of focal length 100 cm and magnifying power 50. The distance between the two lenses in normal
 a) 98 cm b) 100 cm c) 150 cm d) 200 cm
366. A concave lens of focal length 20 cm produce an image half in size of the real object. The distance of the real object is
 a) 20 cm b) 30 cm c) 10 cm d) 60 cm
367. A lens of power +2 *dioptries* is placed in contact with a lens of power -1 *dioptry*. The combination will behave like
 a) A divergent lens of focal length 50 cm
 b) A convergent lens of focal length 50 cm
 c) A convergent lens of focal length 100 cm
 d) A divergent lens of focal length 100 cm
368. A convex lens of crown glass ($n = 1.525$) will behave as a divergent lens if immersed in
 a) Water ($n = 1.33$) b) In a medium of $n = 1.525$
 c) Carbon disulphide $n = 1.66$ d) It cannot act as a divergent lens
369. In a plano-convex lens the radius of curvature of the convex lens is 10 cm. If the plane side is polished, then the focal length will be (Refractive index = 1.5)
 a) 10.5 cm b) 10 cm c) 5.5 cm d) 5 cm
370. Radius of curvature of concave mirror is 40 cm and the size of image is twice as that of object, then the object distance is
 a) 60 cm b) 20 cm c) 40 cm d) 30 cm
371. The resolving limit of healthy eye is about
 a) $1'$ or $\left(\frac{1}{60}\right)^\circ$ b) $1''$ c) 1° d) $\frac{1}{60}''$
372. Why sun has elliptical shape on the time when rising and setting? It is due to
 a) Refraction b) Reflection c) Scattering d) Dispersion
373. In an experiment to determine the focal length (f) of a concave mirror by the $u - v$ method, a student places the object pin A on the principle axis at a distance x from the pole P . The student looks at the pin and its inverted image from a distance keeping his/her eye in line with PA . When the student shifts his/her eye towards left, the image appears to the right of the object pin. Then
 a) $x < f$ b) $f < x < 2f$ c) $x = 2f$ d) $x > 2f$
374. A convex lens has a focal length f . If is cut into two parts along the dotted line as shown in the figure. The focal length of each part will be




- a) $\frac{f}{2}$ b) f c) $\frac{3}{2}f$ d) $2f$


375. Continuous emission spectrum is produced by

- a) Incandescent electric lamp b) Mercury vapour lamp
c) Sodium vapour lamp d) The sun
376. The focal lengths of the objective and the eye-piece of a compound microscope are 2.0 cm and 3.0 cm respectively. The distance between the objective and the eye-piece is 15.0 cm . The final image formed by the eye-piece is at infinity. The two lenses are thin. The distances in cm of the object and the image produced by the objective measured from the objective lens are respectively
a) 2.4 and 12.0 b) 2.4 and 15.0 c) 2.3 and 12.0 d) 2.3 and 3.0
377. A ray of light is incident at an angle of incidence i , on one face of a prism of angle A (assumed to be small) and emerges normally from the opposite face. If the refractive index of the prism is μ , the angle of incidence i , is nearly equal to
a) μA b) $\frac{\mu A}{2}$ c) A/μ d) $A/2\mu$
378. All of the following statements are correct except
a) The magnification produced by a convex mirror is always less than one
b) A virtual, erect, same-sized image can be obtained using a plane mirror
c) A virtual, erect, magnified image can be formed using a concave mirror
d) A real, inverted, same-sized image can be formed using a convex mirror
379. A ray of light is incident on a plane mirror along the direction given by vector $A = 2\hat{i} - 3\hat{j} + 4\hat{k}$. Find the unit vector along reflected ray. Take normal to mirror along the direction of vector $B = 3\hat{i} - 6\hat{j} + 2\hat{k}$.
a) $\frac{-94\hat{i} + 237\hat{j} + 68\hat{k}}{49\sqrt{29}}$ b) $\frac{-94\hat{i} + 68\hat{j} - 273\hat{k}}{49\sqrt{29}}$ c) $\frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{7}$ d) None of these
380. When a glass prism of refracting angle 60° is immersed in a liquid its angle of minimum deviation is 30° . The critical angle of glass with respect to the liquid medium is
a) 42° b) 45° c) 50° d) 52°
381. The power of a biconvex lens is 10 dioptre and the radius of curvature of each surface is 10 cm. Then the refractive index of the material of the lens is
a) $3/2$ b) $4/3$ c) $9/8$ d) $5/3$
382. A man runs towards mirror at a speed of 15 m/s . What is the speed of his image
a) 7.5 m/s b) 15 m/s c) 30 m/s d) 45 m/s
383. Amount of light entering into the camera depends upon
a) Focal length of the objective lens
b) Product of focal length and diameter of the objective lens
c) Distance of the object from camera
d) Aperture setting of the camera
384. The image formed by an objective of a compound microscope is
a) Virtual and diminished b) Real and diminished
c) Real and enlarged d) Virtual and enlarged
385. For unit magnification, the distance of an object from a concave mirror of focal length 20 cm will be
a) 20 cm b) 10 cm c) 40 cm d) 60 cm
386. An object is placed in front of a convex mirror of focal length f . Find the maximum and minimum distance of two object from the mirror such that the image is real and magnified.
a) 20 and ∞ b) f and $2f$ c) f and 0 d) None of these
387. An air bubble in sphere having 4 cm diameter appears 1 cm from surface nearest to eye when looked along diameter. If ${}_a\mu_g = 1.5$, the distance of bubble from refracting surface is
a) 1.2 cm b) 3.2 cm c) 2.8 cm d) 1.6 cm
388. A thin equiconvex lens is made of glass of refractive index 1.5 and its focal length is 0.2 m , if it acts as a concave lens of 0.5 m focal length when dipped in a liquid, the refractive index of the liquid is
a) $\frac{17}{8}$ b) $\frac{15}{8}$ c) $\frac{13}{8}$ d) $\frac{9}{8}$

389. The combination of a convex lens ($f = 18 \text{ cm}$) and a thin concave lens ($f = 9 \text{ cm}$) is
- A concave lens ($f = 18 \text{ cm}$)
 - A convex lens ($f = 18 \text{ cm}$)
 - A convex lens ($f = 6 \text{ cm}$)
 - A concave lens ($f = 6 \text{ cm}$)
390. A convex lens is placed with a mirror as shown in figure. If the space between them is filled with water is power will



- Decrease
 - Increase
 - Remain unchanged
 - Increase or decrease depending on the focal length
391. A ray of light on the surface of a glass plate of thickness t . If the angle of incidence θ is small, the emerging ray would be displaced sideways by an amount (Take n = refractive index of glass)
- $t \theta n / (n + 1)$
 - $t \theta (n - 1) / n$
 - $t \theta n / (n - 1)$
 - $t \theta (n + 1) / n$
392. An object is placed at 15 cm from a convex lens of focal length 10 cm. Where should another convex mirror of radius 12 cm be placed such that image will coincide with the object.
- 19.3 cm
 - 18 cm
 - 33 cm
 - 22 cm
393. When a biconvex lens of glass having refractive index 1.47 is dipped in a liquid, it acts as a plane sheet of glass. This implies that the liquid must have refractive index
- Equal to that of glass
 - Less than one
 - Greater than that of glass
 - Less than that of glass
394. A lens behaves as a converging lens in air and a diverging lens in water. The refractive index of the material is
- Equal to unity
 - Equal to 1.33
 - Between unity and 1.33
 - Greater than 1.33
395. A source emits light of wavelength 4700 \AA , 5400 \AA and 6500 \AA . The light passes through red glass before being tested by a spectrometer. Which wavelength is seen in the spectrum
- 6500 \AA
 - 5400 \AA
 - 4700 \AA
 - All the above
396. A plano convex lens is made of glass of refractive index 1.5. The radius of curvature of its convex surface is R . Its focal length is
- $\frac{R}{2}$
 - R
 - $2R$
 - $1.5 R$
397. The reason for shining of air bubble in water is
- Diffraction of light
 - Dispersion of light
 - Scattering of light
 - Total internal reflection of light
398. For a compound microscope, the focal lengths of object lens and eye lens are f_o and f_e respectively, then magnification will be done by microscope when
- $f_o = f_e$
 - $f_o > f_e$
 - $f_o < f_e$
 - None of these
399. The figures represent three cases of a ray passing through a prism of angle A . The case corresponding to minimum deviation is




- 1
- 2
- 3
- None of these

400. A candle placed 25 cm from a lens, forms an image on a screen placed 75 cm on the other end of the lens. The focal length and type of the lens should be

- a) $+18.75\text{ cm}$ and convex lens b) -18.75 cm and concave lens
 c) $+20.25\text{ cm}$ and convex lens d) -20.25 cm and concave lens
401. Finger prints on a piece of paper may be detected by sprinkling fluorescent powder on the paper and then looking it into
 a) Mercury light b) Sunlight c) Infrared light d) Ultraviolet light
402. A neon sign does not produce
 a) Line spectrum b) An emission spectrum
 c) An absorption spectrum d) Photos
403. A convex lens is made of 3 layers of glass of 3 different materials as in the figure. A point object is placed on its axis. The number of images of the object are



- a) 1 b) 2 c) 3 d) 4
404. Which of the following is not the case with the image formed by a concave lens?
 a) It may be erect or inverted
 b) It may be magnified and diminished
 c) It may be real or virtual
 d) Real image may be between the pole and focus or beyond focus
405. Image formed on retina of eye is proportional to
 a) Size of object b) Area of object c) $\frac{\text{Size of object}}{\text{Size of image}}$ d) $\frac{\text{Size of image}}{\text{Size of object}}$
406. A concave and convex lens have the same focal length of 20 cm and are put into contact to form a lens combination. The combination is used to view an object of 5 cm length kept at 20 cm from the lens combination. As compared to the object, the image will be
 a) Magnified and inverted
 b) Reduced and erect
 c) Of the same size as the object and erect
 d) Of the same size as the object but inverted
407. A point object is moving on the principal axis of a concave mirror focal length 24 cm towards the mirror. When it is at a distance of 60 cm from the mirror, its velocity is 9 cm/sec . What is the velocity of the image at that instant
 a) 5 cm/sec towards the mirror b) 4 cm/sec towards the mirror
 c) 4 cm/sec away from the mirror d) 9 cm/sec away from the mirror
408. If an object moves towards a plane mirror with a speed v at an angle θ to the perpendicular to the plane of the mirror, find the relative velocity between the object and the image




- a) v b) $2v$ c) $2v \cos \theta$ d) $2v \sin \theta$
409. A prism ($\mu = 1.5$) has the refracting angle of 30° . The deviation of a monochromatic ray incident normally on its one surface will be ($\sin 48^\circ 36' = 0.75$)
 a) $18^\circ 36'$ b) $20^\circ 30'$ c) 18° d) $22^\circ 1'$

410. Monochromatic light of frequency 5×10^{14} Hz travelling in vacuum enters a medium of refractive index 1.5. Its wavelength in the medium is

- a) 4000Å b) 5000Å c) 6000Å d) 5500Å

411. Which of the following is true for rays coming from infinity?



- a) Two images are formed
b) Continuous image is formed between focal points of upper and lower lens
c) One image is formed
d) None of the above

412. The dispersive powers of glasses of lenses used in an achromatic pair are in the ratio 5 : 3. If the focal length of the concave lens is 15 cm, then the nature and focal length of the other lens would be

- a) Convex, 9 cm b) Concave, 9 cm c) Convex, 25 cm d) Concave, 25 cm

413. Light travels through a glass plate of thickness t and having refractive index n . If c is the velocity of light in vacuum, the time taken by the light to travel this thickness of glass is

- a) $\frac{t}{nc}$ b) tnc c) $\frac{nt}{c}$ d) $\frac{tc}{n}$

414. For a real object, which of the following can produce a real image?

- a) Plane mirror b) Concave lens c) Convex mirror d) Concave mirror

415. The frequency of a light wave in a material is 2×10^{14} Hz and wavelength is 5000Å. The refractive index of material will be

- a) 1.40 b) 1.50 c) 3.00 d) 1.33

416. By placing the prism in minimum deviation position, images of the spectrum

- a) Becomes inverted b) Becomes broader c) Becomes distinct d) Becomes intensive

417. Two thin lenses whose powers are $+2D$ and $-4D$ respectively combine, then the power of combination is

- a) $-2D$ b) $+2D$ c) $-4D$ d) $+4D$

418. A convex lens is in contact with a concave lens. The magnitude of the ratio of their focal lengths is $2/3$. Their equivalent focal length is 30 cm. What are their individual focal lengths?

- a) -75, 50 b) -10, 15 c) 75, 50 d) -15, 10

419. A concave mirror of focal length 15 cm forms an image having twice the linear dimensions of the object. The position of the object when the image is virtual will be

- a) 22.5 cm b) 7.5 cm c) 30 cm d) 45 cm

420. When light enters from air to glass, for which colour is the angle of deviation maximum?

- a) Red b) Yellow c) Blue d) Violet

421. Check the correct statements on scattering of light

S1 : Rayleigh scattering is responsible for the bluish appearance of sky

S2 : Rayleigh scattering is proportional to $1/\lambda^4$ when the size of the scatter is much less than λ

S3 : Clouds having droplets of water (large scattering objects) scatter all wavelengths almost equally and so are generally white

S4 : The sun looks reddish at sunset and sunrise due to Rayleigh scattering


- a) S1 only b) S1 and S2 c) S2 and S3 d) S1, S2, S3 and S4


422. Our eye is most sensitive for which of the following wavelengths

- a) 4500 Å


- b) 5500 Å
c) 6500 Å
d) Equally sensitive for all wave lengths of visible spectrum
423. Which of the following element was discovered by study of Fraunhoffer line
a) Hydrogen b) Oxygen c) Helium d) Ozone
424. When a lens of refractive index n_1 is placed in a liquid of refractive index n_2 , the lens looks to be disappeared only, if
a) $n_1 = n_2/2$ b) $n_1 = 3n_2/4$ c) $n_1 = n_2$ d) $n_1 = 5n_2/4$
425. If the focal length of the objective lens is increased then
a) Magnifying power of microscope will increase but that of telescope will decrease
b) Magnifying power of microscope and telescope both will increase
c) Magnifying power of microscope and telescope both will decrease
d) Magnifying power of microscope will decrease but that of telescope will increase
426. The diameter of the objective of a telescope is a , its magnifying power is m and wavelength of light is λ . The resolving power of the telescope is
a) $(1.22\lambda)/a$ b) $(1.22a)/\lambda$ c) $\lambda m/(1.22a)$ d) $a/(1.22\lambda)$
427. The reason of seeing the Sun a little before the sunrise is
a) Reflection of the light b) Refraction of the light
c) Scattering of the light d) Dispersion of the light
428. A convex lens of focal length 30 cm and a concave lens of 10 cm focal length are placed so as to have the same axis. If a parallel beam of light falling on convex lens leaves concave lens as a parallel beam, then the distance between two lenses will be
a) 40 cm b) 30 cm c) 20 cm d) 10 cm
429. If in compound microscope m_1 and m_2 be the linear magnification of the objective lens and eye lens respectively, then magnifying power of the compound microscope will be
a) $m_1 - m_2$ b) $\sqrt{m_1 + m_2}$ c) $(m_1 + m_2)/2$ d) $m_1 \times m_2$
430. A convex mirror and a concave mirror has radii of curvature 10 cm each are placed 15 cm apart facing each other. An object is placed midway between them. If the reflection first takes place in the concave mirror and then in convex mirror, the position of the final image is
a) on the pole of the convex mirror b) on the pole of the concave mirror
c) at a distance of 10 cm from convex mirror d) at a distance of 5 cm from concave mirror
431. A convergent beam of light is incident on a convex mirror so as to converge to a distance 12 cm from the pole of the mirror. An inverted image of the same size is formed coincident with the virtual object. What is the focal length of the mirror
a) 24 cm b) 12 cm c) 6 cm d) 3 cm
432. The two surfaces of a biconvex lens has same radii of curvatures. This lens is made of glass of refractive index 1.5 and has a focal length 10 cm in air. The lens is cut into two equal halves along a plane perpendicular to its principle axis to yield two plano-convex lenses. The two pieces are glued such that the convex surfaces touch each other. If this combination lens is immersed in water (refractive index = 4/3), its focal length (in cm) is
a) 5 b) 10 c) 20 d) 40
433. If the distance of the far point for a myopia patient is doubled, the focal length of the lens required to cure it will become
a) Half b) Double
c) The same but a convex lens d) The same but a concave lens
434. The head lights of a jeep are 1.2 m apart. If the pupil of the eye of an observer has a diameter of 2mm and light of wavelength 5896 Å is used, what should be the maximum distance of the jeep from the observer if the two head lights are just separated?
a) 33.9 km b) 33.9 m c) 3.34 km d) 3.39 m

435. Which one of the following alternative is FALSE for a prism placed in a position of minimum deviation
 a) $i_1 = i_2$ b) $r_1 = r_2$ c) $i_1 = r_1$ d) All of these
436. A square card of side length 1mm is being seen through a magnifying lens of focal length 10 cm. The card is placed at a distance of 9 cm from the lens. The apparent area of the card through the lens is
 a) 1 cm^2 b) 0.81 cm^2 c) 0.27 cm^2 d) 0.60 cm^2
437. In a photometer, two sources of light when placed at 30 cm and 50 cm respectively produce shadows of equal intensities. Their candle powers are in the ratio of
 a) $\frac{9}{25}$ b) $\frac{16}{25}$ c) $\frac{3}{5}$ d) $\frac{5}{3}$
438. When a white light passes through a hollow prism, then
 a) There is no dispersion and no deviation
 b) Dispersion but no deviation
 c) Deviation but no dispersion
 d) There is dispersion and deviation both
439. A ray incident at 15° on one refracting surface of a prism of angle 60° , suffers a deviation of 55° . What is the angle of emergence
 a) 95° b) 45° c) 30° d) None of these
440. The maximum illumination on a screen at a distance of 2 m from a lamp is 25 lux. The value of total luminous flux emitted by the lamp is
 a) 1256 lumen b) 1600 lumen c) 100 candela d) 400 lumen
441. Two vertical plane mirrors are inclined at an angle of 60° with each other. A ray of light travelling horizontally is reflected first from one mirror and then from the other. The resultant deviation is
 a) 60° b) 120° c) 180° d) 240°
442. A short linear object of a length b lies along the axis of a concave mirror of focal length f at a distance u from the pole of the mirror. The size of the image is equal to
 a) $b \left(\frac{u-f}{f} \right)^{1/2}$ b) $b \left(\frac{f}{u-f} \right)^{1/2}$ c) $b \left(\frac{u-f}{f} \right)$ d) $b \left(\frac{f}{f-u} \right)$
443. Two point light sources are 24 cm apart. Where should a convex lens of focal length 9 cm be put in between them from one source so that the images of both the sources are formed at the same place
 a) 6 cm b) 9 cm c) 12 cm d) 15 cm
444. A wave has velocity u in medium P and velocity $2u$ in medium Q . If the wave is incident in medium P at an angle 30° , then the angle of refraction will be
 a) 30° b) 45° c) 60° d) 90°
445. Line spectrum contains information about
 a) The atoms of the prism b) The atoms of the source
 c) The molecules of the source d) The atoms as well as molecules of the source
446. A convex lens, a glass slab, a glass prism and a solid sphere all are made of the same glass, the dispersive power will be
 a) In the glass slab and prism b) In the lens and solid sphere
 c) Only in prism d) In all the four
447. A beam of light is converging towards a point I on a screen. A plane glass plate whose thickness in the direction of the beam = t , refractive index = μ , is introduced in the path of the beam. The convergence point is shifted by
 a) $t \left(1 - \frac{1}{\mu} \right)$ away b) $t \left(1 + \frac{1}{\mu} \right)$ away c) $t \left(1 - \frac{1}{\mu} \right)$ nearer d) $t \left(1 + \frac{1}{\mu} \right)$ nearer
448. A thin oil layer floats on water. A ray of light making an angle of incidence of 40° shines on oil layer. The angle of refraction of light ray in water is ($\mu_{oil} = 1.45$, $\mu_{water} = 1.33$)
 a) 36.1° b) 44.5° c) 26.8° d) 28.9°
449. A lens when placed on a plane mirror then object needle and its image coincide at 15 cm. The focal length of the lens is



- a) 15 cm b) 30 cm c) 20 cm d) ∞
450. The focal length of the field lens (which is an achromatic combination of two lenses) of telescope is 90 cm. The dispersive powers of the two lenses in the combination are 0.024 and 0.036. The focal lengths of two lenses are
 a) 30 cm and 60 cm b) 30 cm and -45 cm c) 45 cm and 90 cm d) 15 cm and 45 cm
451. A ray is reflected in turn by three plane mirrors mutually at right angles to each other. The angle between the incident and the reflected rays is
 a) 90° b) 60° c) 180° d) none of these
452. A medium is said to be dispersive, if
 a) Light of different wavelengths propagate at different speeds
 b) Light of different wavelengths propagate at same speed but has different frequencies
 c) Light is gradually bent rather than sharply refracted at an interface between the medium and air
 d) Light is never totally internally reflected
453. When a glass slab is placed on a cross made on a sheet, the cross appears raised by 1 cm. The thickness of the glass is 3 cm. The critical angle for glass is
 a) $\sin^{-1}(0.33)$ b) $\sin^{-1}(0.5)$ c) $\sin^{-1}(0.67)$ d) $\sin^{-1}(\sqrt{3}/2)$
454. A luminous object is placed at a distance of 30 cm from the convex lens of focal length 20 cm. On the other side of the lens, at what distance from the lens a convex mirror of radius of curvature 10 cm be placed in order to have an upright image of the object coincident with it
 a) 12 cm b) 30 cm c) 50 cm d) 60 cm
455. A ray of light passes through four transparent medium with refractive indices μ_1, μ_2, μ_3 and μ_4 as shown in the figure. The surfaces of all media are parallel. If the emergent ray CD is parallel to the incident ray AB. We must have

 a) $\mu_1 = \mu_2$ b) $\mu_2 = \mu_3$ c) $\mu_3 = \mu_4$ d) $\mu_3 = \mu_1$
456. The nature of sun's spectrum is
 a) Continuous spectrum with absorption lines b) Line spectrum
 c) The spectrum of the helium atom d) Band spectrum
457. An object is placed in front of a convex mirror at a distance of 50 cm. A plane mirror is introduced covering the lower half of the convex mirror. If the distance between the object and plane mirror is 30 cm, it is found that there is no parallax between the images formed by two mirrors. Radius of curvature of mirror will be
 a) 12.5 cm b) 25 cm c) $\frac{50}{3}$ cm d) 18 cm
458. We wish to see inside an atom. Assuming the atom to have a diameter of 100 pm, this means that one must be able to resolve a width of say 10 pm. If an electron microscope is used, the minimum electron energy required is about
 a) 1.5 keV b) 15 keV c) 150 keV d) 1.5 keV
459. Two thin lenses of focal lengths f_1 and f_2 are placed in contact with each other. The focal length of the combination is
 a) $\frac{f_1 + f_2}{2}$ b) $\sqrt{f_1 f_2}$ c) $\frac{f_1 f_2}{f_1 + f_2}$ d) $\frac{f_1 f_2}{f_1 - f_2}$

460. The dispersive powers of crown and flint glasses are 0.02 and 0.04 respectively. In an achromatic combination of lenses the focal length of flint glass lens is 40 cm. The focal length of crown glass lens will be
 a) -20 cm b) $+20\text{ cm}$ c) -10 cm d) $+10\text{ cm}$
461. The critical angle of the medium with respect to vacuum is 30° . If the velocity of light in vacuum is $3 \times 10^8\text{ ms}^{-1}$, the velocity of light in medium is
 a) $2 \times 10^8\text{ ms}^{-1}$ b) $1.5 \times 10^8\text{ ms}^{-1}$ c) $3 \times 10^8\text{ ms}^{-1}$ d) $\sqrt{2} \times 10^8\text{ ms}^{-1}$
462. A planoconvex lens has a maximum thickness of 6 cm. When placed on a horizontal table with the curved surface in contact with the table surface, the apparent depth of the bottommost point of the lens is found to be 4 cm. If the lens is inverted such that the plane face of the lens is in contact with the surface of the table, the apparent depth of the centre of the plane face is found to be $\left(\frac{17}{4}\right)$ cm. The radius of curvature of the lens is
 a) 34 cm b) 128 cm c) 75 cm d) 68 cm
463. When white light passes through the achromatic combination of prisms, then what is observed
 a) Only deviation b) Only dispersion
 c) Deviation and dispersion d) None of the above
464. The sensation of vision in the retina is carried to the brain by
 a) Ciliary muscles b) Blind spot c) Cylindrical lens d) Optic nerve
465. A compound microscope is used to enlarge an object kept at a distance 0.03m from its objective which consists of several convex lenses in contact and has focal length 0.02m. If a lens of focal length 0.1m is removed from the objective, then by what distance the eye-piece of the microscope must be moved to refocus the image
 a) 2.5 cm b) 6 cm c) 15 cm d) 9 cm
466. Light rays from a source are incident on a glass prism of index of refraction μ and angle of prism a . At near normal incidence, the angle of deviation of the emerging rays is
 a) $(\mu - 2)\alpha$ b) $(\mu - 1)\alpha$ c) $(\mu + 1)\alpha$ d) $(\mu + 2)\alpha$
467. Stars are not visible in the day time because
 a) Stars hide behind the sun
 b) Stars do not reflect sun rays during day
 c) Stars vanish during the day
 d) Atmosphere scatters sunlight into a blanket of extreme brightness through which faint stars cannot be visible
468. A concave mirror is placed at the bottom of an empty tank with face upwards and axis vertical. When sunlight falls normally on the mirror, it is focussed at distance of 32 cm from the mirror. If the tank filled with water $\left(\mu = \frac{4}{3}\right)$ upto a height of 20 cm, then the sunlight will now get focussed at
 a) 16 cm above water level b) 9 cm above water level
 c) 24 cm below water level d) 9 cm below water level
469. A ray of light strikes a material's slab at an angle of incidence 60° . If the reflected and refracted rays are perpendicular to each other, the refractive index of the materials is
 a) $\frac{1}{\sqrt{3}}$ b) $\frac{1}{\sqrt{2}}$ c) $\sqrt{2}$ d) $\sqrt{3}$
470. A ray of light is incident on an equilateral glass prism placed on a horizontal table. For minimum deviation which of the following is true?



- a) PQ is horizontal
 b) QR is horizontal
 c) RS is horizontal
 d) Either PQ or RS is horizontal

471. The angle of minimum deviation for an incident light ray on an equilateral prism is equal to its refracting angle. The refractive index of its material is

- a) $\frac{1}{\sqrt{2}}$
 b) $\sqrt{3}$
 c) $\frac{\sqrt{3}}{2}$
 d) $\frac{3}{2}$

472. A person sees his virtual image by holding a mirror very close to the face. When he moves the mirror away from his face, the image becomes inverted. What type of mirror he is using?

- a) Plane mirror
 b) Convex mirror
 c) Concave mirror
 d) None of these

473. Which of the following statement is true

- a) Velocity of light is constant in all media
 b) Velocity of light in vacuum is maximum
 c) Velocity of light is same in all reference frames
 d) Laws of nature have identical form in all reference frames

474. Transmission of light to large distances through optical fibres is based on

- a) Dispersion
 b) Refraction
 c) Total internal reflection
 d) Interference

475. Pick out the correct statements about optical fibres from the following

S1 : Optical fibres are used for the transmission of optical signals only

S2 : Optical fibres are used for transmitting and receiving electrical signals

S3 : The intensity of light signals sent through optical fibres suffer very small loss

S4 : Optical fibres effectively employ the principle of multiple total internal reflections

S5 : Optical fibres are glass fibres coated with a thin layer of a material with lower refractive index

- a) S1 and S2
 b) S2 and S3
 c) S3 and S4
 d) S2, S3, S4 and S5


476. Given the width of aperture = 3 mm and $\lambda = 500$ nm. For what distance ray optics is good approximation?

- a) 18 m
 b) 18 mm
 c) 18 Å
 d) 18 light years

477. A ray of light makes an angle of 10° with the horizontal above it and strikes a plane mirror which is inclined at an angle θ to the horizontal. The angle θ for which the reflected ray becomes vertical is

- a) 40°
 b) 50°
 c) 80°
 d) 100°

478. Which of the following graphs show appropriate variation of refractive index μ with wavelength λ



479. A plano convex lens of ($f = 20$ cm) is silvered at plane surface. New f will be

- a) 20 cm
 b) 40 cm
 c) 30 cm
 d) 10 cm

480. If h_1 and h_2 are the heights of the images in conjugate position of a convex lens, then the height of the object is


- a) $h_1 + h_2$
 b) $h_1 - h_2$
 c) h_1/h_2
 d) $\sqrt{h_1 h_2}$

481. A thin double convex lens has radii of curvature each of magnitude 40 cm and is made of glass with $\mu = 1.65$. The focal length of the lens is nearly

- a) 30 cm
 b) 31 cm
 c) 40 cm
 d) 41 cm

482. A student can distinctly see the object upto a distance 15 cm. He wants to see the black board at a distance of 3 m. Focal length and power of lens used respectively will be
 a) $-4.8 \text{ cm}, -3.3 \text{ D}$ b) $-5.8 \text{ cm}, -4.3 \text{ D}$ c) $-7.5 \text{ cm}, -6.3 \text{ D}$ d) $-15.8 \text{ cm}, -6.3 \text{ D}$
483. A prism of certain angles deviates the red and blue rays by 8° and 12° respectively. Another prism of the same angle deviates the red and blue rays by 10° respectively. The prisms are small angled and made of different materials. The dispersive powers of the materials of the prism are in the ratio
 a) 5 : 6 b) 9 : 11 c) 6 : 5 d) 11 : 9
484. F_1 and F_2 are focal lengths of objective and eyepiece respectively of the telescope. The angular magnification for the given telescope is equal to
 a) $\frac{F_1}{F_2}$ b) $\frac{F_2}{F_1}$ c) $\frac{F_1 F_2}{F_1 + F_2}$ d) $\frac{F_1 + F_2}{F_1 F_2}$
485. Refractive index of glass with respect to medium is $\frac{4}{3}$. If the differences between velocities of light in medium and glass is $6.25 \times 10^7 \text{ ms}^{-1}$, then velocity of light in medium is
 a) $2.5 \times 10^8 \text{ ms}^{-1}$ b) $0.125 \times 10^8 \text{ ms}^{-1}$ c) $1.5 \times 10^8 \text{ ms}^{-1}$ d) $3 \times 10^8 \text{ ms}^{-1}$
486. A room (cubical) is made of mirrors. An insect is moving along the diagonal on the floor such that the velocity of image of insect on two adjacent wall mirrors is 10 cms^{-1} . The velocity of image of insect in ceiling mirror is
 a) 10 cms^{-1} b) 20 cms^{-1} c) $\frac{10}{\sqrt{2}} \text{ cms}^{-1}$ d) $10\sqrt{2} \text{ cms}^{-1}$
487. A convex mirror forms an image one-fourth the size of the object. If object is at a distance of 0.5 m from the mirror, the focal length of mirror is
 a) 0.17 m b) -1.5 m c) 0.4 m d) -0.4 m
488. A man of length h requires a mirror, to see his own complete image of length at least equal to
 a) $h/4$ b) $h/3$ c) $h/2$ d) h
489. A achromatic combination is made with a lens of focal length f and dispersive power ω with a lens having dispersive power of 2ω . The focal length of second will be
 a) $2f$ b) $f/2$ c) $-f/2$ d) $-2f$
490. The diameter of moon is $3.5 \times 10^3 \text{ km}$ and its distance from the earth is $3.8 \times 10^5 \text{ km}$. The focal length of the objective and eye-piece are 4 m and 10 cm respectively. The diameter of the image of the moon will be approximately
 a) 2° b) 21° c) 40° d) 50°
491. One surface of a lens is convex and the other is concave. If the radii of curvature are r_1 and r_2 respectively, the lens will be convex, if
 a) $r_1 > r_2$ b) $r_1 = r_2$ c) $r_1 < r_2$ d) $r_1 = 1/r_2$
492. A parallel beam of monochromatic light is incident at one surface of an equilateral prism. Angle of incidence is 55° and angle of emergence is 46° . The angle of minimum deviation will be
 a) Less than 41° b) Equal to 41° c) More than 41° d) None of the above
493. A ray of light is incident at an angle of 60° on one face of a prism of angle 30° . The ray emerging out of the prism makes an angle of 30° with the incident ray. The emergent ray is
 a) Normal to the face through which it emerges
 b) Inclined at 30° to the face through which it emerges
 c) Inclined at 60° to the face through which it emerges
 d) None of these

494. The refractive index of the material of the prism and liquid are 1.56 and 1.32 respectively. What will be



the value of θ for the following refraction

- a) $\sin \theta \geq \frac{13}{11}$ b) $\sin \theta \geq \frac{11}{13}$ c) $\sin \theta \geq \frac{\sqrt{3}}{2}$ d) $\sin \theta \geq \frac{1}{\sqrt{2}}$

495. Convergence of concave mirror can be decreased by dipping in

- a) Water b) Oil c) Both d) None of these

496. If a parallel beam of white light is incident on a converging lens, the colour which is brought to focus nearest to the lens is

- a) Violet b) Red
c) The mean colour d) All the colours together

497. The critical angle for diamond (refractive index = 2) is

- a) About 20° b) 60° c) 45° d) 30°


498. Microscope is an optical instrument which

- a) Enlarges the object
b) Increases the visual angle formed by the object at the eye
c) Decreases the visual angle formed by the object at the eye
d) Brings the object nearer

499. Circular part in the centre of retina is called

- a) Blind spot b) Yellow spot c) Red spot d) None of the above

500. When the rectangular metal tank is filled to the top with an unknown liquid, as observer with eyes level with the top of the tank can just see the corner E ; a ray that refracts towards the observer at the top surface of the liquid is shown. The refractive index of the liquid will be



- a) 1.2 b) 1.4 c) 1.6 d) 1.9

501. A double convex lens ($R_1 = R_2 = 100$ cm) having focal length equal to the focal length of a concave mirror. The radius of the concave mirror is

- a) 10 cm b) 20 cm c) 40 cm d) 15 cm


502. A bulb of 100 watt is hanging at a height of one meter above the centre of a circular table of diameter 4 m. If the intensity at a point on its rim is I_0 , then the intensity at the centre of the table will be

- a) I_0 b) $2\sqrt{5}I_0$ c) $2I_0$ d) $5\sqrt{5}I_0$

503. A concave mirror and a converging lens (glass with $\mu = 1.5$) both have a focal length of 3 cm when in air. When they are in water ($\mu = \frac{4}{3}$), their new focal lengths are

- a) $f_{\text{Lens}} = 12$ cm, $f_{\text{Mirror}} = 3$ cm b) $f_{\text{Lens}} = 3$ cm, $f_{\text{Mirror}} = 12$ cm
c) $f_{\text{Lens}} = 3$ cm, $f_{\text{Mirror}} = 3$ cm d) $f_{\text{Lens}} = 12$ cm, $f_{\text{Mirror}} = 12$ cm

504. An observer can see through a pin-hole the top end of a thin rod of height h , placed as shown in the figure. The beaker height is $3h$ and its radius h . When the beaker is filled with a liquid up to a height $2h$, he can see the lower end of the rod. Then the refractive index of the liquid is



- a) $5/2$ b) $\sqrt{(5/2)}$ c) $\sqrt{(3/2)}$ d) $3/2$

505. A lens made of glass whose index of refraction is 1.60 has a focal length of +20 cm in air. Its focal length in water, whose refractive index is 1.33, will be

- a) Three times longer than in air b) Two times longer than in air
c) Same as in air d) None of the above

506. Which of the following is a correct relation

- a) $a\mu_r = a\mu_\omega \times r\mu_\omega$ b) $a\mu_r \times r\mu_\omega = \omega\mu_a$ c) $a\mu_r \times r\mu_a = 0$ d) $a\mu_r / \omega\mu_r = a\mu_\omega$

507. Which of the following colours suffers maximum deviation in a prism

- a) Yellow b) Blue c) Green d) Orange

508. f_v and f_r are the focal lengths of a convex lens for violet and red light respectively and F_v and F_r are the focal lengths of a concave lens for violet and red light respectively, then

- a) $f_v < f_r$ and $F_v > F_r$ b) $f_v < f_r$ and $F_v < F_r$ c) $f_v > f_r$ and $F_v > F_r$ d) $f_v > f_r$ and $F_v < F_r$

509. Fraunhofer lines are obtained in

- a) Solar spectrum
b) The spectrum obtained from neon lamp
c) Spectrum from a discharge tube
d) None of the above

510. A point object is placed mid-way between two plane mirrors distance 'a' apart. The plane mirror forms an infinite number of images due to multiple reflection. The distance between the n th order image formed in the two mirrors is

- a) na b) $2na$ c) $na/2$ d) n^2a


511. The refractive indices of glass and quartz w.r.t. air are $3/2$ and $12/5$ respectively. The refractive index of quartz w.r.t. glass is

- a) $8/5$ b) $5/8$ c) $5/18$ d) $18/5$

512. If an object is placed 10 cm in front of a concave mirror of focal length 20 cm, the image will be

- a) Diminished, upright, virtual b) Enlarged, upright, virtual
c) Diminished, inverted, real d) Enlarged, upright, real

513. A point source of light S is placed at the bottom of a vessel containing a liquid of refractive index $5/3$. A person is viewing the source from above the surface. There is an opaque disc D of radius 1 cm floating on the surface of the liquid. The centre of the disc lies vertically above the source S. The liquid from the vessel is gradually drained out through a tap. The maximum height of the liquid for which the source cannot be seen at all from above is



- a) 1.50 cm b) 1.64 cm c) 1.33 cm d) 1.86 cm

514. A ray of light is incident on a surface of glass slab at an angle 45° . If the lateral shift produced per unit thickness is $\frac{1}{\sqrt{3}}$ m, the angle of refraction produced is

a) $\tan^{-1}\left(\frac{\sqrt{3}}{2}\right)$ b) $\tan^{-1}\left(1 - \sqrt{\frac{2}{3}}\right)$ c) $\sin^{-1}\left(1 - \sqrt{\frac{2}{3}}\right)$ d) $\tan^{-1}\left(\sqrt{\frac{2}{\sqrt{3}-1}}\right)$

515. An achromatic convergent doublet of two lenses in contact has a power of +2D. The convex lens has power +5 D. What is the ratio of the dispersive powers of the convergent and divergent lenses?

- a) 2:5 b) 3:5 c) 5:2 d) 5:3


516. An object is placed at 15 cm in front of a concave mirror whose focal length is 10 cm. The image formed will be

- a) Magnified and inverted b) Magnified and erect
c) Reduced in size and inverted d) Reduced in size and erect

517. Two plane mirrors are at right angles to each other. A man stands between them and combs his hair with his right hand. In how many of the images will he be seen using his right hand

- a) None b) 1 c) 2 d) 3

518. The ray diagram could be correct



- a) If $n_1 = n_2 = n_3$ b) If $n_1 = n_2$ and $n_1 < n_g$
c) If $n_1 = n_2$ and $n_1 > n_g$ d) Under no circumstances

519. Refractive index of the material of a prism is 1.5. If $\delta_m = A$, what will be a value of angle of the given prism?

(where δ_m = minimum deviation; A = angle of prism)

- a) 82.8° b) 41.4° c) 48.6° d) 90°

520. Minimum deviation is observed with a prism having angle of prism A , angle of deviation δ , angle of incidence i and angle of emergence e . We then have generally

- a) $i > e$ b) $i < e$ c) $i = e$ d) $i = e = \delta$

521. An astronomical telescope of ten-fold angular magnification has a length of 44 cm. The focal length of the objective is

- a) 4 cm b) 40 cm c) 44 cm d) 440 cm

522. A piece of plane glass is placed on a word with letters of different colours. The letters which appear minimum raised are

- a) Red b) Green c) Yellow d) Violet

523. A concave lens of glass, refractive index 1.5, has both surfaces of same radius of curvature R . On immersion in a medium of refractive index 1.75, it will behave as a

- a) Convergent lens of focal length $3.5 R$ b) Convergent lens of focal length $3.0 R$
c) Divergent lens of focal length $3.5 R$ d) Divergent lens of focal length $3.0 R$

524. A man's near point is 0.5 m and far point is 3 m. Power spectacle lenses repaired for

(i) reading purposes

(ii) seeing distant objects, respectively

- a) -2 D and +3 D b) +2 D and -3 D
c) +2 D and -0.33 D d) -2 D and +0.33 D


525. A lamp of 250 candela power is hanging at a distance of 6 m from a wall. The illuminance at a point on the wall at a minimum distance from lamp will be

- a) 9.64 lux b) 4.69 lux c) 6.94 lux d) None of these

526. Large aperture of telescope are used for

- a) Large image b) Greater resolution
c) Reducing lens aberration d) Ease of manufacture

527. The angle of minimum deviation measured with a prism is 30° and the angle of prism is 60° . The refractive index of prism material is
 a) $\sqrt{2}$ b) 2 c) $3/2$ d) $4/3$
528. A cube of side 2 m is placed in front of a concave mirror focal length 1 m with its face P at a distance of 3 m and face Q at a distance of 5 m from the mirror. The distance between the images of face P and Q and height of images of P and Q are




- a) $1\text{ m}, 0.5\text{ m}, 0.25\text{ m}$ b) $0.5\text{ m}, 1\text{ m}, 0.25\text{ m}$ c) $0.5\text{ m}, 0.25\text{ m}, 1\text{ m}$ d) $0.25\text{ m}, 1\text{ m}, 0.5\text{ m}$
529. A ray of light passes through an equilateral prism such that the angle of incidence and the angle of emergence are both equal to $3/4$ th of the angle of prism. The angle of minimum deviation is
 a) 15° b) 30° c) 45° d) 60°
530. The refractive index of water, glass and diamond are $1.33, 1.50, 2.40$ respectively. The refractive index of diamond relative to water and of glass relative to diamond, respectively are nearly
 a) $1.80, 0.625$ b) $0.554, 0.625$ c) $1.80, 1.6$ d) $0.554, 1.6$
531. A small object is placed 10 cm in front of a plane mirror. If you stand behind the object, 30 cm from the mirror and look at its image, for what distance must you focus your eyes?
 a) 20 cm b) 60 cm c) 80 cm d) 40 cm
532. The radius of curvature of concave mirror is 24 cm and the image is magnified by 1.5 times. The object distance is
 a) 20 cm b) 8 cm c) 16 cm d) 24 cm
533. If the focal length of the eye piece of a telescope is double, its magnifying power m_1 will be
 a) $2m$ b) $3m$ c) $\frac{m}{2}$ d) $4m$
534. A plano-concave lens is made up of glass of refractive index 1.5 and the radius of the curvature of its curved face is 100 cm . What is the power of the lens?
 a) $+0.5\text{ D}$ b) -0.5 D c) -2 D d) $+2\text{ D}$
535. A rectangular tank of depth 8 meter is full of water ($\mu = 4/3$), the bottom is seen at the depth
 a) 6 m b) $8/3\text{ cm}$ c) 8 cm d) 10 cm
536. As the wavelength is increased from violet to red, the luminosity
 a) Continuously increases b) Continuously decreases
 c) Increases then decreases d) Decreases then increases
537. A light beam is being reflected by using two mirrors, as in a periscope used in submarines. If one of the mirrors rotates by an angle θ , the reflected light will deviate from its original path by the angle
 a) 2θ b) 0° c) θ d) 4θ
538. Two thin lenses of focal length 20 cm and 25 cm are placed in contact. The effective power of the combination is
 a) 9 D b) 2 D c) 3 D d) 7 D
539. The dispersive power is maximum for the material
 a) Flint glass b) Crown glass c) Mixture of both d) None of the above
540. An under water swimmer is at a depth of 12 m below the surface of water. A bird is at a height of 18 m from the surface of water, directly above his eyes. For the swimmer the bird appears to be a distance from the surface of water equal to (Refractive Index of water is $\frac{4}{3}$)
 a) 24 m b) 12 m c) 18 m d) 9 m


541. Parallel beam containing light of $\lambda = 400\text{nm}$ and 500 nm is incident on a prism as shown in figure. The refractive index μ of the prism is given by the relation

$$\mu(\lambda) = 1.20 + \frac{0.8 \times 10^{-14}}{\lambda^2}$$


Which of the following statement is correct?



- a) Light of $\lambda = 400\text{ nm}$ undergoes total internal reflection
 - b) Light of $\lambda = 500\text{ nm}$ undergoes total internal reflection
 - c) Neither of two wavelength undergoes total internal reflection
 - d) Both wavelengths undergoes total internal reflection
542. Figure shows a glowing mercury tube. The illuminances at point A, B and C are related as



- a) $B > C > A$
 - b) $A > C > B$
 - c) $B = C > A$
 - d) $B = C < A$
543. Lenses of power 3 D and -5 D are combined to form a compound lens. An object is placed at a distance of 50 cm from this lens. Its image will be formed at a distance from the lens, will be
- a) 25 cm
 - b) 20 cm
 - c) 30 cm
 - d) 40 cm
544. The dispersive power of the material of lens of focal length 20 cm is 0.08 . The longitudinal chromatic aberration of the lens is
- a) 0.08 cm
 - b) $0.08/20\text{ cm}$
 - c) 1.6 cm
 - d) 0.16 cm
545. When sunlight is incident on a prism, it produces a spectrum due to
- a) Interference of light
 - b) Diffraction of light
 - c) Total internal reflection
 - d) Variation in speeds of different colours of light in the prism
546. Why is refractive index in a transparent medium greater than one?
- a) Because the speed of light in vacuum is medium
 - b) Because the speed of light in vacuum is always greater than speed in a transparent medium
 - c) Frequency of wave changes when it crosses medium
 - d) None of the above
547. A transparent solid cylindrical rod has a refractive index of $\frac{2}{\sqrt{3}}$. It is surrounded by air. A light ray is incident at the mid-point of one end of the rod as shown in the figure.



The incident angle θ for which the light ray grazes along the wall of the rod is

- a) $\sin^{-1}\left(\frac{1}{2}\right)$
 - b) $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right)$
 - c) $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$
 - d) $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$
548. The two lenses of an achromatic doublet should have
- a) Equal powers
 - b) Equal dispersive powers

- c) Equal ratio of their power and dispersive power
 d) Sum of the product of their powers and dispersive power equal to zero

549. The focal lengths of the objective and eyelenses of a microscope are 1.6 cm and 2.5 cm respectively. The distance between the two lenses is 21.7 cm. If the final image is formed at infinity, the distance between the object and the objective lens is


- a) 1.8 cm b) 1.70 cm c) 1.65 cm d) 1.75 cm

550. A beaker contains water up to a height h_1 and kerosene of height h_2 above water so that the total height of (water + kerosene) is $(h_1 + h_2)$. Refractive index of water is u_1 and that of kerosene is u_2 . The apparent shift in the position of the bottom of the beaker when viewed from above is

- a) $\left(1 - \frac{1}{u_1}\right)h_2 + \left(1 - \frac{1}{u_2}\right)h_1$ b) $\left(1 + \frac{1}{u_1}\right)h_1 + \left(1 + \frac{1}{u_2}\right)h_2$
 c) $\left(1 - \frac{1}{u_1}\right)h_2 + \left(1 - \frac{1}{u_2}\right)h_2$ d) $\left(1 + \frac{1}{u_1}\right)h_2 - \left(1 + \frac{1}{u_2}\right)h_1$

551. A ray of light travelling in glass ($\mu = \frac{3}{2}$) is incident on a horizontal glass air surface at the critical angle θ_c .

If thin layer of water ($\mu = \frac{4}{3}$) is now poured on the glass air surface, the angle at which the ray emerges into air the water-air surface is



- a) 60° b) 45° c) 90° d) 180°

552. An object is placed first at infinity and then at 20 cm from the object side focal plane of the convex lens. The two images thus formed are 5 cm apart. The focal length of the lens is

- a) 5 cm b) 10 cm c) 15 cm d) 20 cm

553. Two lenses of power +12 and -2 dioptres are placed in contact. What will the focal length of combination

- a) 10 cm b) 12.5 cm c) 16.6 cm d) 8.33 cm

554. When objects at different distances are seen by the eye, which of the following remains constant

- a) The focal length of the eye lens b) The object distance from the eye lens
 c) The radii of curvature of the eye lens d) The image distance from the eye lens


555. A convex and a concave lens separated by distance d are then put in contact. The focal length of the combination

- a) Decreases b) Increases c) Becomes zero d) Remains the same

556. Four lenses of focal length +15 cm, +20 cm, +150 cm and +250 cm are available for making an astronomical telescope. To produce the largest magnification, the focal length of the eye-piece should be

- a) +15 cm b) +20 cm c) +150 cm d) +250 cm

557. What is the relation between refractive indices μ_1 , μ_2 , and μ_3 if the behavior of light rays is as shown in figure




- a) $\mu_3 < \mu_2, \mu_2 = \mu_1$ b) $\mu_2 < \mu_1, \mu_2 = \mu_3$ c) $\mu_3 < \mu_2 < \mu_1$ d) $\mu_3 > \mu_2 > \mu_1$

558. A person is in a room whose ceiling and two adjacent walls are mirrors. How many images are formed

- a) 5 b) 6 c) 7 d) 8

559. A thin prism P of refracting angle 3° and refractive index 1.5 is combined with another thin prism Q of refractive index 1.6 to produce dispersion without deviation. Then the angle of prism Q is
 a) 3° b) 4° c) 3.5° d) 2.5°
560. The slit of a collimator is illuminated by a source as shown in the adjoining figures. The distance between the slit S and the collimating lens L is equal to the focal length of the lens. The correct direction of the emergent beam will be as shown in figure




- a) 1 b) 3 c) 2 d) None of the figures
561. An endoscope is employed by a physician to view the internal parts of a body organ. It is based on the principle of
 a) Refraction b) Reflection
 c) Total internal reflection d) Dispersion
562. The magnifying power of a telescope is m . If the focal length of the eye-piece is halved, then its magnifying power is
 a) $2m$ b) $\frac{m}{2}$ c) $\frac{1}{2m}$ d) $4m$
563. If the ratio of amounts of scattering of two light waves is 1:4, the ratio of their wavelength is
 a) 1 : 2 b) $\sqrt{2}$: 1 c) 1 : $\sqrt{2}$ d) 1 : 1
564. The black lines in the solar spectrum during solar eclipse can be explained by
 a) Planck's law b) Kirchhoff's law c) Boltzmann's law d) Solar disturbances
565. The length of a telescope is 36 cm. The focal lengths of its lenses can be
 a) 30 cm, 6 cm b) -30 cm, -6 cm c) 30 cm, -6 cm d) -30 cm, 6 cm
566. A man is 180cm tall and his eyes are 10cm below the top of his head. In order to see his entire height right from toe to head, he uses a plane mirror kept at a distance of 1m from him. The minimum length of the plane mirror required is
 a) 180cm b) 90cm c) 85cm d) 170cm
567. A 2.0 cm tall object is placed 15 cm in front of a concave mirror of focal length 10 cm. What is the size and nature of the image
 a) 4 cm, real b) 4 cm, virtual c) 1.0 cm, real d) None of these
568. If the refractive indices of crown glass for red, yellow and violet colours are 1.5140, 1.5170 and 1.5318 respectively and for flint glass these are 1.6434, 1.6499 and 1.6852 respectively, then the dispersive powers for crown and flint glass are respectively
 a) 0.034 and 0.064 b) 0.064 and 0.034 c) 1.00 and 0.064 d) 0.034 and 1.0
569. A telescope has an objective lens of 10 cm diameter and is situated at a distance of one kilometer from two objects. The minimum distance between these two objects, which can be resolved by the telescope, when the mean wavelength of light is 5000 \AA , is of the order of
 a) 0.5 m b) 5 m c) 5 mm d) 5 cm
570. A spherical mirror forms diminished virtual image of magnification $1/3$. Focal length is 18 cm. The distance of the object is
 a) 18 cm b) 36 cm c) 48 cm d) Infinite
571. A light bulb is at a depth of D below the surface of water. An opaque disc of radius R is placed on the surface of water just above the bulb. The bulb is not at all seen through the surface of water, then (n = Refractive index of water)
 a) $R = \frac{D}{\sqrt{n^2 - 1}}$ b) $R > \frac{D}{\sqrt{n^2 - 1}}$ c) $R < \frac{D}{\sqrt{n^2 - 1}}$ d) $R = D\sqrt{n^2 - 1}$

572. Immiscible transparent liquids A, B, C, D and E are placed in a rectangular container of glass with the liquids making layers according to their densities. The refractive index of the liquids are shown in the adjoining diagram. The container is illuminated from the side and small piece of glass having refractive index 1.61 is gently dropped into the liquid layer. The glass piece as it descends downwards will not be visible in

A	1.51
B	1.53
C	1.61
D	1.52
E	1.65

- a) Liquid A and B only
 b) Liquid C only
 c) Liquid D and E only
 d) Liquid A, B, D and E
573. The angular magnification of a simple microscope can be increased by increasing
 a) Focal length of lens b) Size of object c) Aperture of lens d) Power of lens
574. The twinkling effect of star light is due to
 a) Total internal reflection
 b) High dense matter of star
 c) Constant burning of hydrogen in the star
 d) The fluctuating apparent position of the star being slightly different from of the star being different from the actual position of the star
575. The mean distance of sun from the earth is $1.5 \times 10^8 \text{ Km}$ (nearly). The time taken by the light to reach earth from the sun is
 a) 0.12 min b) 8.33 min c) 12.5 min d) 6.25 min
576. A layered lens as shown in figure is made of two types of transparent materials indicated by different shades. A point object is placed on its axis. The object will form




- a) 1 image b) 2 images c) 3 images d) 9 images
577. A double convex thin lens made of glass (refractive index $\mu = 1.5$) has both radii of curvature of magnitude 20 cm. Incident light rays parallel to the axis of the lens, will converge at a distance L such that
 a) $L = \frac{20}{3} \text{ cm}$ b) $L = 40 \text{ cm}$ c) $L = 20 \text{ cm}$ d) $L = 10 \text{ cm}$
578. A boy is trying to start a fire by focusing Sunlight on a piece of paper using an equiconvex lens of focal length 10 cm. The diameter of the Sun is $1.39 \times 10^9 \text{ m}$ and its means distance from the earth is $1.5 \times 10^{11} \text{ m}$. What is the diameter of the Sun's image on the paper
 a) $6.5 \times 10^{-5} \text{ m}$ b) $12.4 \times 10^{-4} \text{ m}$ c) $9.2 \times 10^{-4} \text{ m}$ d) $6.5 \times 10^{-4} \text{ m}$
579. A screen receives 3 watt of radiant flux of wavelength 6000 Å. One lumen is equivalent to $1.5 \times 10^{-3} \text{ watt}$ of monochromatic light of wavelength 5550 Å. If relative luminosity for 6000 Å is 0.685 while that for 5550 Å is 1.00, then the luminous flux of the source is
 a) $4 \times 10^3 \text{ lm}$ b) $3 \times 10^3 \text{ lm}$ c) $2 \times 10^3 \text{ lm}$ d) $1.37 \times 10^3 \text{ lm}$
580. Which has more luminous efficiency
 a) A 40 W bulb b) A 40 W fluorescent tube
 c) Both have same d) Cannot say

581. Sir C.V. Raman was awarded Nobel Prize for his work connected with which of the following phenomenon of radiation

- a) Scattering b) Diffraction c) Interference d) Polarization

582. A ray of light coming. Which of the following figures, shows dispersion of light?



583. Band spectrum is obtained when the source emitted light is in the form of **or**

Band spectrum is characteristic of

- a) Atoms b) Molecules c) Plasma d) None of the above


584. If two mirrors are kept at 60° to each other, then the number of images found by them is

- a) 5 b) 6 c) 7 d) 8

585. A thin plano-convex lens acts like a concave mirror of focal length 0.2 m when silvered from its plane surface. The refractive index of the material of the lens is 1.5. The radius of curvature of the convex surface the lens will be

- a) 0.1 m b) 0.75 m c) 0.4 m d) 0.2 m

586. An object is placed at a point distance x from the focus of a convex lens and its image is formed is I as shown in the figure. The distances x, x' satisfy the relation



- a) $\frac{x + x'}{2} = f$ b) $f = xx'$ c) $x + x' \leq 2f$ d) $x + x' \geq 2f$

587. Angle of prism is A and its one surface is silvered. Light rays falling at an angle of incidence $2A$ on first surface return back through the same path after suffering reflection at second silvered surface. Refractive index of the material of prism is

- a) $2 \sin A$ b) $2 \cos A$ c) $\frac{1}{2} \cos A$ d) $\tan A$

588. The time required for the light to pass through a glass slab (refractive index $= 1.5$) of thickness 4mm is... ($c = 3 \times 10^8 \text{ ms}^{-1}$, speed of light in free space).

- a) $2 \times 10^{-5} \text{ s}$ b) $2 \times 10^{11} \text{ s}$ c) $2 \times 10^{-11} \text{ s}$ d) 10^{-11} s

589. A convex mirror has a focal length f . A real object is placed at a distance f in front of it from the pole produces an image at

- a) Infinity b) f c) $f/2$ d) $2f$

590. A ray of light enters from a rarer to a denser medium. The angle of incidence is i . Then the reflected and refracted rays are mutually perpendicular to each other. The critical angle for the pair of media is

- a) $\sin^{-1}(\tan i)$ b) $\tan^{-1}(\sin i)$ c) $\sin^{-1}(\cot i)$ d) $\cos^{-1}(\tan i)$

591. A point source of light is kept at a depth of h in water of refractive index $4/3$. The radius of the circle at the surface of water through which light emits is

- a) $\frac{3}{\sqrt{7}}h$ b) $\frac{\sqrt{7}}{3}h$ c) $\frac{\sqrt{3}}{7}h$ d) $\frac{7}{\sqrt{3}}h$


592. The power of an achromatic convergent lens of two lenses is $+2 \text{ D}$. The power of convex lens is $+5 \text{ D}$. The ratio of dispersive power of convex and concave lenses will be

- a) 5:3 b) 3:5 c) 2:5 d) 5:2

593. The earth radiates in the infra-red region of the spectrum. The spectrum is correctly given by


- a) Rayleigh Jeans law b) Planck's of law of radiation

- Page | 53




A light beam incident normally on the face AB is totally reflected to reach the face BC if

- a) $\frac{2}{3} < \sin \theta < \frac{8}{9}$ b) $\sin \theta \leq \frac{2}{3}$ c) $\cos \theta \geq \frac{8}{9}$ d) $\sin \theta > \frac{8}{9}$
609. The resolving power of an astronomical telescope is 0.2 seconds. If the central half portion of the objective lens is covered, the resolving power will be
- a) 0.1 sec b) 0.2 sec c) 1.0 sec d) 0.6 sec
610. A boy 1.5 m tall with his eye level at 1.38 m stands before a mirror fixed on a wall. The minimum length of mirror required to view the complete image of boy is
- a) 0.75 m b) 0.06 m c) 0.69 m d) 0.12 m
611. The refractive index of a prism for a monochromatic wave is $\sqrt{2}$ and its refracting angle is 60° . For minimum deviation, the angle of incidence will be
- a) 30° b) 45° c) 60° d) 75°
612. The graph shows how the magnification m produced by a convex thin lens varies with image distance v . What was the focal length of the used lens




- a) b/c b) b/ca c) bc/a d) c/b
613. Least distance of distinct vision is 25 cm. Magnifying power of simple microscope of focal length 5 cm is
- a) $1/5$ b) 5 c) $1/6$ d) 6
614. Retina of eye acts like of camera
- a) Shutter b) Film c) Lens d) None of these
615. The radius of curvature of the convex face of a planoconvex lens is 15 cm and the refractive index of the material is 1.4. Then the power of the lens in diopter is
- a) 1.6 b) 1.66 c) 2.6 d) 2.66
616. The refractive index of a material of a planoconcave lens is $5/3$, the radius of curvature is 0.3 m. The focal length of the lens in air is
- a) -0.45 m b) -0.6 m c) -0.75 m d) -1.0 m
617. What should be the angle between two plane mirrors so that whatever be the angle of incidence, the incident ray and the reflected ray from the two mirrors be parallel to each other
- a) 60° b) 90° c) 120° d) 175°
618. The exposure time of a camera lens at the $\frac{f}{2.8}$ setting is $\frac{1}{200}$ second. The correct time of exposure at $\frac{f}{5.6}$ is
- a) 0.4 s b) 0.02 s c) 0.002 s d) 0.04 s
619. The focal lengths of the lenses of an astronomical telescope are 50 cm and 5 cm. The length of the telescope when the image is formed at the least distance of distinct vision is
- a) 45 cm b) 55 cm c) $\frac{275}{6}$ cm d) $\frac{325}{6}$ cm


620. The focal length of a concave mirror is 20 cm. Where an object must be placed to form an image magnified two times when the image is real?
- a) 30 cm from the mirror b) 10 cm from the mirror
c) 20 cm from the mirror d) 15 cm from the mirror
621. In a compound microscope, the focal lengths of two lenses are 1.5 cm and 6.25 cm. An object is placed at 2 cm from objective and the final image is formed is 25 cm from eye lens. The distance between the two lenses is
- a) 6.00 cm b) 7.75 cm c) 9.25 cm d) 11.00 cm
622. The image of point P when viewed from top of the slabs will be




- a) 2.0 cm above P b) 1.5 cm above P c) 2.0 cm below P d) 1 cm above P
623. In order to increase the magnifying power of a compound microscope
- a) The focal lengths of the objective and the eye piece should be small
b) Objective should have small focal length and the eye piece large
c) Both should have large focal lengths
d) The objective should have large focal length and eye piece should have small
624. A beam of light is travelling from region II to region III (see the figure). The refractive index in region I, II and III are n_0 , $\frac{n_0}{\sqrt{2}}$, and $\frac{n_0}{2}$ respectively. The angle of incidence θ for which the beam just misses entering region III is




- a) 30° b) 45° c) 60° d) $\sin^{-1}(\sqrt{2})$
625. The graph between object distance u and image distance v for lens is given below. The focal length of the lens is







- a) 5 ± 0.1 b) 5 ± 0.05 c) 0.5 ± 0.1 d) 0.5 ± 0.05
626. The length of an astronomical telescope for normal vision (relaxed eye) (f_o = focal length of objective lens and f_e = focal length of eye lens) is
- a) $f_o \times f_e$ b) $\frac{f_o}{f_e}$ c) $f_o + f_e$ d) $f_o - f_e$
627. The magnifying power of a simple microscope is 6. The focal length of its lens in metres will be, if least distance of distinct vision is 25 cm

- a) 0.05 b) 0.06 c) 0.25 d) 0.12
628. A concave mirror gives an image three times as large as the object placed at a distance of 20 cm from it. For the image to be real, the focal length should be
a) 10 cm b) 15 cm c) 20 cm d) 30 cm
629. The spectrum of iodine gas under white light will be
a) Only violet b) Bright lines
c) Only red lines d) Some black bands is continuous spectrum
630. The power of a thin convex lens ($n_g = 1.5$) is + 0.5 D. When it is placed in a liquid of refractive index n_l , then it behaves as a concave lens of focal length 100 cm. The refractive index of the liquid n_l will be
a) 5/3 b) 4/3 c) $\sqrt{3}$ d) 5/4
631. A person 6 feet in length can see his full size erect image in a mirror 2 feet in length. This mirror has to be
a) Plane or convex b) Plane or concave
c) Necessarily convex d) Necessarily concave
632. A prism ABC of angle 30° has its face AC silvered. A ray of light incident at an angle of 45° at the face AB retraces its path after refraction at face AB and reflection at face AC. The refractive index of the material of the prism is
- 
- a) 1.5 b) $3/\sqrt{2}$ c) $\sqrt{2}$ d) 4/3
633. The focal length of the lens of refractive index ($\mu = 1.5$) in air is 10 cm. If air is replaced by water of $\mu = \frac{4}{3}$, its focal length is
a) 20 cm b) 30 cm c) 40 cm d) 25 cm
634. The refractive index of water is 1.33. What will be the speed of light in water
a) 3×10^8 m/s b) 2.25×10^8 m/s c) 4×10^8 m/s d) 1.33×10^8 m/s
635. Lens used to remove long sightedness (hypermetropia) is or
A person suffering from hypermetropia requires which type of spectacle lenses
a) Concave lens b) Plano-concave lens
c) Convexo-concave lens d) Convex lens
636. Focal length of a convex lens of refractive index 1.5 in 2 cm. Focal length of lens when immersed in a liquid of refractive index of 1.25 will be
a) 10 cm b) 2.5 cm c) 5 cm d) 7.5 cm
637. A thin lens made of glass of refractive index 1.5 has a front surface +11 D power and back surface -6 D. If this lens is submerged in a liquid of refractive index 1.6, the resulting power of the lens is
a) -0.5 D b) +0.5 D c) -0.625 D d) +0.625 D
638. A plane mirror makes an angle of 30° with horizontal. If a vertical ray strikes the mirror, find the angle between mirror and reflected ray
a) 30° b) 45° c) 60° d) 90°
639. A ray of light is incident on the hypotenuse of a right-angled prism after travelling parallel to the base inside the prism. If μ is the refractive index of the material of the prism, the maximum value of the base angle for which light is totally reflected from the hypotenuse is
a) $\sin^{-1}\left(\frac{1}{\mu}\right)$ b) $\tan^{-1}\left(\frac{1}{\mu}\right)$ c) $\sin^{-1}\left(\frac{\mu-1}{\mu}\right)$ d) $\cos^{-1}\left(\frac{1}{\mu}\right)$
640. A square wire of side 1 cm is placed perpendicular to the principle axis of a concave mirror of focal length 15 cm at a distance of 20 cm. The area enclosed by the image of the wire is

- a) 4 cm^2 b) 6 cm^2 c) 2 cm^2 d) 9 cm^2
641. The cross-section of a glass prism has the form of an isosceles triangle. One of the refracting faces is silvered. A ray of light falls normally on the other refracting face. After being reflected twice, it emerges through the base of the prism perpendicular to it. The angles of the prism are
- a) $54^\circ, 54^\circ, 72^\circ$ b) $72^\circ, 72^\circ, 36^\circ$ c) $45^\circ, 45^\circ, 90^\circ$ d) $57^\circ, 57^\circ, 76^\circ$
642. The plane faces of two identical plano convex lenses, each with focal length f are pressed against each other using an optical glue to form a usual convex lens. The distance from the optical centre at which an object must be placed to obtain the image same as the size of object is
- a) $\frac{f}{4}$ b) $\frac{f}{2}$ c) f d) $2f$
643. When light of wavelength λ is incident on an equilateral prism kept in its minimum deviation position, it is found that the angle of deviation equals the angle of the prism itself. The refractive index of the material of the prism for the wavelength λ is, then
- a) $\sqrt{3}$ b) $\frac{\sqrt{3}}{2}$ c) 2 d) $\sqrt{2}$
644. The wavelength of a certain colour in air is 600 nm . What is the wavelength and speed of this colour in glass of refractive index 1.5 ?
- a) 500 nm and $2 \times 10^{10} \text{ cms}^{-1}$ b) 400 nm and $2 \times 10^8 \text{ ms}^{-1}$
c) 300 nm and $3 \times 10^9 \text{ cms}^{-1}$ d) 700 nm and $1.5 \times 10^9 \text{ ms}^{-1}$
645. Magnification of a compound microscope is 30 . Focal length of eye-piece is 5 cm and the image is formed at a distance of distinct vision of 25 cm . The magnification of the objective lens is
- a) 6 b) 5 c) 7.5 d) 10
646. The human eye has a lens which has a
- a) Soft portion at its centre b) Hard surface
c) Varying refractive index d) Constant refractive index
647. A source of light emits a continuous stream of light energy which falls on a given area. Luminous intensity is defined as
- a) Luminous energy emitted by the source per second
b) Luminous flux emitted by source per unit solid angle
c) Luminous flux falling per unit area of a given surface
d) Luminous flux coming per unit area of an illuminated surface
648. A prism of a certain angle deviation the red and blue rays by 8° and 12° respectively. Another prism of the same angle deviates the red and blue rays by 10° and 14° respectively. The prisms are small angled and made of different materials. The dispersive powers of the materials of the prisms are in the ratio
- a) $5 : 6$ b) $9 : 11$ c) $6 : 5$ d) $11 : 9$
649. A converging beam of rays is incident on a diverging lens. Having passed through the lens the rays intersect at a point 15 cm from the lens on the opposite side. If the lens is removed the point where the rays meet will move 5 cm closer to the lens. The focal length of the lens is
- a) -30 cm b) 5 cm c) -10 cm d) 20 cm
650. A person is suffering from myopic defect. He is able to see clear objects placed at 15 cm . What type and of what focal length of lens he should use to see clearly the object placed 60 cm away
- a) Concave lens of 20 cm focal length b) Convex lens of 20 cm focal length
c) Concave lens of 12 cm focal length d) Convex lens of 12 cm focal length
651. A point object O is placed in front of a glass rod having spherical end of radius of curvature 30 cm . The image would be formed at




- a) 30 cm left b) Infinity c) 1 cm to the right d) 18 cm to the left
652. For normal vision, what is distance of object from eye?
a) 30 cm b) 25 cm c) Infinite d) 40 cm
653. Three prisms of crown glass, each have angle of prism 9° and two prisms of flint glass are used to make direct vision spectroscope. What will be the angle of flint glass prisms if μ for flint is 1.60 and μ for crown glass is 1.53
a) 11.9° b) 16.0° c) 15.3° d) 9.11°
654. When light enters water from the vacuum, then the wavelength of light
a) Decreases b) Increases c) Remain constant d) Becomes zero
655. A thin prism P_1 with angle 4° made from a glass of refractive index 1.54 is combined with another thin prism P_2 made from glass of refractive index 1.72 to produce dispersion without deviation. The angle of the prism P_2 is
a) 5.33° b) 4° c) 3° d) 2.6°
656. Which one of the followingspherical lenses does not exhibit dispersion? The radii of curvature of the surfaces of the lenses are as given in the diagrams
a)  b)  c)  d) 
 $R_1 \neq R_2$
657. Focal length of a plane mirror is
a) Zero b) Infinite c) Very less d) Indefinite
658. A convex lens of focal length 10 cm and image formed by it, is at least distance of distinct vision then the magnifying power is
a) 3.5 b) 2.5 c) 1.5 d) 1.4
659. A small plane mirror placed at the centre of a spherical screen of radius R . A beam of light is falling on the mirror. If the mirror makes n revolution per second, the speed of light on the screen after reflection from the mirror will be
a) $4\pi nR$ b) $2\pi nR$ c) $\frac{nR}{2\pi}$ d) $\frac{nR}{4\pi}$
660. On heating a liquid, the refractive index generally
a) Decreases
b) Increases or decreases depending on the rate of heating
c) Does not change
d) Increases
661. The spectrum obtained from a sodium vapour lamp is an example of
a) Absorption spectrum b) Emission spectrum c) Continuous spectrum d) Band spectrum
662. Two parallel pillars are 11 km away from an observer. The minimum distance between the pillars so that they can be seen separately will be
a) 3.2 m b) 20.8 m c) 91.5 m d) 183 m
663. The wavelength of red light from He-Ne laser is 633nm in air but 474 nm in the aqueous humor inside the eye ball. Then the speed of red light through the aqueous humor is
a) $3 \times 10^8 \text{ms}^{-1}$ b) $1.34 \times 10^8 \text{ms}^{-1}$ c) $2.25 \times 10^8 \text{ms}^{-1}$ d) $2.5 \times 10^8 \text{ms}^{-1}$
664. The focal length of a simple convex lens used as a magnifier is 10 cm. For the image to be formed at a distance of distinct vision ($D = 25 \text{ cm}$), the object must be placed away from the lens at a distance of
a) 0.5 cm b) 7.14 cm c) 7.20 cm d) 16.16 cm
665. A watch shows time as 3 : 25 when seen through a mirror, time appeared will be
a) 8 : 35 b) 9 : 35 c) 7 : 35 d) 8 : 25
666. If light travels a distance x in $t_1 \text{ sec}$ in air and $10 x$ distance in t_2 in a medium, the critical angle of the medium will be


- a) $\tan^{-1}\left(\frac{t_1}{t_2}\right)$ b) $\sin^{-1}\left(\frac{t_1}{t_2}\right)$ c) $\sin^{-1}\left(\frac{10t_1}{t_2}\right)$ d) $\tan^{-1}\left(\frac{10t_1}{t_2}\right)$
667. The wavelength of light in two liquids 'x' and 'y' is 3500 Å and 7000 Å, then the critical angle of x relative to y will be
 a) 60° b) 45° c) 30° d) 15°
668. Brilliance of diamond is due to
 a) Shape b) Cutting
 c) Reflection d) Total internal reflection
669. A film projector magnifies a 100 cm² film strip on a screen. If the linear magnification is 4, the area of magnified film on the screen is
 a) 1600 cm² b) 400 cm² c) 800 cm² d) 200 cm²
670. A beam of light composed of red and green rays is incident obliquely at a point on the face of a rectangular glass slab. When coming out on the opposite parallel face, the red and green rays emerge from
 a) Two points propagating in two different non-parallel directions
 b) Two point propagating in two different parallel directions
 c) One point propagating in two different directions
 d) One point propagating in the same direction
671. Critical angle of light passing from glass of water is minimum for
 a) Red colour b) Green colour c) Yellow colour d) Violet colour
672. When light emitted by a white hot solid is passed through a sodium flame, the spectrum of the emergent light will show
 a) The D₁ and D₂ bright yellow lines of sodium
 b) Two dark lines in the yellow region
 c) All colours from violet to red
 d) No colours at all
673. An object is approaching a plane mirror at 10 cms⁻¹. A stationary observer sees the image. At what speed will the image approach the stationary observer?
 a) 10 cms⁻¹ b) 5 cms⁻¹ c) 20 cms⁻¹ d) 15 cms⁻¹
674. A lens is made of flint glass (refractive index = 1.5). When the lens is immersed in a liquid of refractive index 1.25, the focal length
 a) Increase by a factor of 1.25 b) Increases by a factor of 2.5
 c) Increases by a factor of 1.2 d) Decreases by a factor of 1.2
675. A convex lens forms an image of an object placed 20 cm away from it at a distance of 20 cm on the other side of the lens. If the object is moved 5 cm towards the lens, the image will move
 a) 5 cm towards the lens b) 5 cm away from the lens
 c) 10 cm towards the lens d) 10 cm away from the lens
676. For a colour of light the wavelength for air is 6000 Å and in water the wavelength is 4500 Å. Then the speed of light in water will be
 a) 5.0×10^{14} m/s b) 2.25×10^8 m/s c) 4.0×10^8 m/s d) Zero
677. The focal lengths of convex lens for red and blue light are 100 cm and 96.8 cm respectively. The dispersive power of material of lens is
 a) 0.325 b) 0.0325 c) 0.98 d) 0.968
678. A thin prism of angle 15° made of glass of refractive index $\mu_1 = 1.5$ is combined with another prism of glass of refractive index $\mu_2 = 1.75$. The combination of the prisms produces dispersion without deviation. The angle of the second prism should be
 a) 12° b) 5° c) 7° d) 10°
679. The velocity of light in a medium is half its velocity in air. If ray of light emerges from such a medium into air, the angle of incidence, at which it will be totally internally reflected, is
 a) 15° b) 30° c) 45° d) 60°

680. The refractive index and the permeability of a medium are respectively 1.5 and $5 \times 10^{-7} \text{ Hm}^{-1}$. The relative permittivity of the medium is nearly
 a) 25 b) 15 c) 81 d) 6
681. A light ray is incident by grazing one of the face of a prism and after refraction ray does not emerge out, what should be the angle of prism while critical angle is C
 a) Equal to $2C$ b) Less than $2C$ c) More than $2C$ d) None of the above
682. A book can be read if it is placed at a distance of 50 cm from a source of 1 cd. At what distance should the book be placed if the source is of 16 cd?
 a) 8 m b) 4 m c) 2 m d) 1 m
683. When diameter of the aperture of the objective of an astronomical telescope is increased, its
 a) Magnifying power is increased and resolving power is decreased
 b) Magnifying power and resolving power both are increased
 c) Magnifying power remains the same but resolving power is increased
 d) Magnifying power and resolving power both are decreased
684. The index of refraction of diamond is 2.0, velocity of light in diamond in cm/s is approximately
 a) 6×10^{10} b) 3.0×10^{10} c) 2×10^{10} d) 1.5×10^{10}
685. Astigmatism (for a human eye) can be removed by using
 a) Concave lens b) Convex lens c) Cylindrical lens d) Prismatic lens
686. A ray of light is incident normally on a plane mirror. The angle of reflection will be
 a) 0° b) 90° c) Will not be reflected d) None of the above
687. An astronaut is looking down on earth's surface from a space shuttle at an altitude of 400 km. Assuming that the astronaut's pupil diameter is 5 mm and the wavelength of visible light is 500 nm. The astronaut will be able to resolve linear object of the size of about
 a) 0.5 m b) 5 m c) 50 m d) 500 m
688. The power of the combination of a convex lens of focal length 50 cm and concave lens of focal length 40 cm is
 a) +1 D b) -1 D c) Zero d) -0.5 D
689. Image is formed for the short sighted person at
 a) Retina b) Before retina
 c) Behind the retina d) Image is not formed at all
690. The magnification produced by the objective lens and the eye lens of a compound microscope are 25 and 6 respectively. The magnification of this microscope is
 a) 25 b) 50 c) 150 d) 200
691. The path of a refracted ray of light in a prism is parallel to the base of the prism only when the
 a) Light is of a particular wavelength b) Ray is incident normally at one face
 c) Ray undergoes minimum deviation d) Prism is made of a minimum deviation
692. Two lamps of luminous intensity of 8 Cd and 32 Cd respectively are lying at a distance of 1.2 m from each other. Where should a screen be placed between two lamps such that its two faces are equally illuminated due to two sources
 a) 10 cm from 8 Cd lamp b) 10 cm from 32 Cd lamp
 c) 40 cm from 8 Cd lamp d) 40 cm from 32 Cd lamp
693. An object is placed at a distance of 10 cm from a concave mirror of radius of curvature 0.6 m. Which of the following statements is incorrect?
 a) The image is formed at a distance for 15 cm from the mirror
 b) The image formed is real
 c) The image is 0.5 times the size of the object
 d) The image is 1.5 times the size of the object

694. Consider the situation shown in figure. Water ($\mu_w = \frac{4}{3}$) is filled in a beaker upto a height of 10 cm. A plane mirror fixed at a height of 5 cm from the surface of water. Distance of image from the mirror after reflection from it of an object O at the bottom of the beaker is




- a) 15 cm b) 12.5 cm c) 7.5 cm d) 10 cm
695. A simple telescope, consisting of an objective of focal length 60 cm and a single eye lens of focal length 5 cm is focussed on a distant object in such a way that parallel rays come out from the eye lens. If the object subtends an angle 2° at the objective, the angular width of the image
- a) 10° b) 24° c) 50° d) $1/6^\circ$
696. If aperture of lens is halved then image will be
- a) No effect on size b) Intensity of image decreases
c) Both (a) and (b) d) None of these
697. Missing lines in a continuous spectrum reveal
- a) Defects of the observing instrument
b) Absence of some elements in the light source
c) Presence in the light source of hot vapours of some elements
d) Presence of cool vapours of some elements around the light source
698. A lamp is hanging along the axis of a circular table of radius r . At what height should the lamp be placed above the table, so that the illuminance at the edge of the table is $\frac{1}{8}$ of that at its center
- a) $\frac{r}{2}$ b) $\frac{r}{\sqrt{2}}$ c) $\frac{r}{3}$ d) $\frac{r}{\sqrt{3}}$
699. A biconvex lens with equal radii of curvature has refractive index 1.6 and focal length 10 cm. Its radius of curvature will be
- a) 20 cm b) 16 cm c) 10 cm d) 12 cm
700. The plane faces of two identical plano-convex lenses each having focal length of 40 cm are pressed against each other to form a usual convex lens. The distance from this lens, at which an object must be placed to obtain a real, inverted image with magnification one is
- a) 80 cm b) 40 cm c) 20 cm d) 162 cm
701. A convex lens made up of a material of refractive index μ_1 is immersed in a medium of refractive index μ_2 as shown in the figure. The relation between μ_1 and μ_2 is




- a) $\mu_1 < \mu_2$ b) $\mu_1 > \mu_2$ c) $\mu_1 = \mu_2$ d) $\mu_1 = \sqrt{\mu_2}$
702. Maximum lateral displacement of a ray of light incident on a slab of thickness t is
- a) $\frac{t}{2}$ b) $\frac{t}{3}$ c) $\frac{t}{4}$ d) t
703. A simple magnifying lens is used in such a way that an image is formed at 25 cm away from the eye. In order to have 10 times magnification, the focal length of the lens should be
- a) 5 cm b) 2 cm c) 25 mm d) 0.1 mm



704. A 4 cm thick layer of water covers a 6 cm thick glass slab. A coin is placed at the bottom of the slab and is being observed from the air side along the normal to the surface. Find the apparent position of the coin from





- a) 7.0 cm b) 8.0 cm c) 10 cm d) 5 cm
705. If μ_0 be the relative permeability and K_0 the dielectric constant of a medium, its refractive index is given by
- a) $\frac{1}{\sqrt{\mu_0 K_0}}$ b) $\frac{1}{\mu_0 K_0}$ c) $\sqrt{\mu_0 K_0}$ d) $\mu_0 K_0$
706. The focal lengths of the objective and of the eye-piece of a compound microscope are f_0 and f_e respectively. If L is the tube length and D , the least distance of distinct vision, then its angular magnification, when the image is formed at infinity, is
- a) $\left(1 - \frac{L}{f_0}\right)\left(\frac{D}{f_e}\right)$ b) $\left(1 + \frac{L}{f_0}\right)\left(\frac{D}{f_e}\right)$ c) $\frac{L}{f_0}\left(1 - \frac{D}{f_e}\right)$ d) $\frac{L}{f_0}\left(\frac{D}{f_e}\right)$
707. A simple microscope consists of a concave lens of power $-10D$ and a convex lens of power $+20D$ in contact. If the image is formed at infinity, then the magnifying power $CD = 25$ cm is
- a) 2.5 b) 3.5 c) 2.0 d) 3.0
708. The wavelength of sodium light in air is 5890 \AA . The velocity of light in air is $3 \times 10^8 \text{ ms}^{-1}$. The wavelength of light in a glass of refractive index 1.6 would be close to
- a) 5890 \AA b) 3681 \AA c) 9424 \AA d) 15078 \AA
709. A plano-convex lens when silvered in the plane side behaves like a concave mirror of focal length 30 cm . However, when silvered on the convex side it behaves like a concave mirror of focal length 10 cm . Then the refractive index of its material will be
- a) 3.0 b) 2.0 c) 2.5 d) 1.5
710. A plane mirror is placed at the bottom of a tank containing a liquid of refractive index μ . P is a small object at a height h above the mirror. An observer O vertically above P outside the liquid sees P and its image in a mirror. The apparent distance between these two will be




- a) $2\mu h$ b) $\frac{2h}{\mu}$ c) $\frac{2h}{\mu - 1}$ d) $h\left(1 + \frac{1}{\mu}\right)$
711. The refractive index of a material of a prism of angles $45^\circ - 45^\circ - 90^\circ$ is 1.5. The path of the ray of light incident normally on the hypotenuse side is shown in





712. How many images are formed by the lens shown, if an object is kept on its axis?



- a) 1 b) 2 c) 3 d) 4


713. The bottom of a container filled with liquid appear slightly raised because of

- a) Refraction b) Interference c) Diffraction d) Reflection

714. The magnifying power of a telescope is 9. When it is adjusted for parallel rays, the distance between the objective and the eye-piece is found to be 20 cm. The focal lengths of the lenses are

- a) 18 cm, 2 cm b) 11 cm, 9 cm c) 10 cm, 10 cm d) 15 cm, 5 cm

715. Two transparent slabs have the same thickness as shown. One is made of material *A* of refractive index 1.5. The other is made of two material *B* and *C* with thickness in the ratio 1 : 2. The refractive index *C* is 1.6. If a monochromatic parallel beam passing through the slabs has the same number of waves inside both, the refractive index of *B* is



- a) 1.1 b) 1.2 c) 1.3 d) 1.4

716. A ray of light passes through an equilateral prism such that an angle of incidence is equal to the angle of emergence and the latter is equal to $\frac{3}{4}$ th the angle of prism. The angle of deviation is

- a) 45° b) 39° c) 20° d) 30°

717. An experiment is performed to find the refractive index of glass using a travelling microscope. In this experiment distance are measured by

- a) A vernier scale provided on the microscope b) A standard laboratory scale
c) A meter scale provided on the microscope d) A screw gauge provided on the microscope

718. Convex lens made up of glass ($\mu_g = 1.5$) and radius of curvature *R* is dipped into water. Its focal length will be (Refractive index of water = $4/3$)

- a) 4*R* b) 2*R* c) *R* d) $\frac{R}{2}$

719. An object is kept at a distance of 16 cm from a thin lens and the image formed is real. If the object is kept at a distance of 6 cm from the same lens, the image formed is virtual. If the sizes of the images formed are equal the focal length of the lens will be

- a) 21 cm b) 11 cm c) 15 cm d) 17 cm


720. Angle of minimum deviation for a prism of refractive index 1.5 is equal to the angle of the prism. The angle of the prism is (given $\cos 41^\circ - 24' - 36'' = 0.75$)

- a) $82^\circ - 49' - 12''$ b) $72^\circ - 48' - 30''$ c) $41^\circ - 24' - 36''$ d) $31^\circ - 49' - 30''$


721. The instrument used by doctors for endoscopy works on the principle of

- a) Total internal reflection b) Reflection
c) Refraction d) None of the above

722. Refractive index of glass is $\frac{3}{2}$ and refractive index of water is $\frac{4}{3}$. If the speed of light in glass is 2.00×10^8 m/s, the speed in water will be
 a) 2.67×10^8 m/s b) 2.25×10^8 m/s c) 1.78×10^8 m/s d) 1.50×10^8 m/s
723. Critical angle for light going from medium (i) to (ii) is θ . The speed of light in medium (i) is v then speed in medium (ii) is
 a) $v(1 - \cos \theta)$ b) $v/\sin \theta$ c) $v/\cos \theta$ d) $v(1 - \sin \theta)$
724. Monochromatic light of wavelength λ_1 travelling in medium of refractive index n_1 enters a denser medium of refractive index n_2 . The wavelength in the second medium is
 a) $\lambda_1 \left(\frac{n_1}{n_2} \right)$ b) $\lambda_1 \left(\frac{n_2}{n_1} \right)$ c) λ_1 d) $\lambda_1 \left(\frac{n_2 - n_1}{n_1} \right)$
725. In a compound microscope cross-wires are fixed at the point
 a) Where the image is formed by the objective
 b) Where the image is formed by the eye-piece
 c) Where the focal point of the objective lies
 d) Where the focal point of the eye-piece lies
726. Consider telecommunication through optical fibres. Which of the following statements is not true
 a) Optical fibres may have homogeneous core with a suitable cladding
 b) Optical fibres can be of graded refractive index
 c) Optical fibres are subject to electromagnetic interference from outside
 d) Optical fibres have extremely low transmission loss
727. In the figure shown, for an angle of incidence 45° , at the top surface, what is the minimum refractive index needed for total internal reflection at vertical face




- a) $\frac{\sqrt{2} + 1}{2}$ b) $\sqrt{\frac{3}{2}}$ c) $\sqrt{\frac{1}{2}}$ d) $\sqrt{2} + 1$
728. Deviation of 5° is observed from a prism whose angle is small and whose refractive index is 1.5. The angle of prism is
 a) 7.5° b) 10° c) 5° d) 3.3°
729. A photograph of the moon was taken with telescope. Later on, it was found that a housefly was sitting on the objective lens of the telescope. In photograph
 a) The image of housefly will be reduced
 b) There is a reduction in the intensity of the image
 c) There is an increase in the intensity of the image
 d) The image of the housefly will be enlarged
730. The graph between sine of angle of refraction ($\sin r$) in medium 2 and sine of angle of incidence ($\sin i$) in medium 1 indicates that ($\tan 36^\circ \approx \frac{3}{4}$)




- a) Total internal reflection can take place b) Total internal reflection cannot take place
 c) Any of (a) and (b) d) Data is incomplete

731. A prism having an apex angle 4° and refractive index 1.5 is located in front of a vertical plane mirror as shown in figure. Through what total angle is the ray deviated after reflection from the mirror




- a) 176° b) 4° c) 178° d) 2°
732. A person can see clearly only upto a distance of 25 cm. He wants to read a book placed at a distance of 50 cm. What kind of lens does he require for his spectacles and what must be its power
- a) Concave, $-1.0 D$ b) Convex, $+1.5 D$ c) Concave, $-2.0 D$ d) Convex, $+2.0 D$
733. When a plane mirror is rotated through an angle θ then the reflected ray turns through the angle 2θ then the size of the image
- a) Is doubled b) Is halved c) Remains the same d) Becomes infinite
734. Two media having speeds of light $2 \times 10^8 \text{ ms}^{-1}$ and $2.4 \times 10^8 \text{ ms}^{-1}$, are separated by a plane surface. What is the angle for a ray going from medium I to medium II?
- a) $\sin^{-1}\left(\frac{5}{6}\right)$ b) $\sin^{-1}\left(\frac{5}{12}\right)$ c) $\sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$ d) $\sin^{-1}\left(\frac{1}{2}\right)$
735. The ratio of angle of minimum deviation of a prism in air and when dipped in water will be (${}_a\mu_g = 3/2$ and ${}_a\mu_w = 4/3$)
- a) $1/8$ b) $1/2$ c) $3/4$ d) $1/4$
736. A student measures the focal length of a convex lens by putting an object pin at a distance u from the lens and measuring the distance v of the image pin. The graph between u and v plotted by the student should look like



737. A light moves from denser to rarer medium. Which of the following is correct?
- a) Energy increases b) Frequency increases
c) Phase changes by 90° d) Velocity increases
738. If the aperture of a telescope is decreased the resolving power will
- a) Increase b) Decrease c) Remain same d) Zero
739. A wire mesh consisting of very small squares is viewed at a distance of 8 cm through a magnifying converging lens of focal length 10 cm, kept close to the eye. The magnification produced by the lens is
- a) 5 b) 8 c) 10 d) 20
740. Light travels with a speed of $2 \times 10^8 \text{ ms}^{-1}$ in crown glass of refractive index 1.5. What is the speed of light in dense flint glass of refractive index 1.8?
- a) $1.33 \times 10^8 \text{ ms}^{-1}$ b) $1.67 \times 10^8 \text{ ms}^{-1}$ c) $2.0 \times 10^8 \text{ ms}^{-1}$ d) $3.0 \times 10^8 \text{ ms}^{-1}$
741. Focal length of a converging lens in air is R . If it is dipped in water of refractive index 1.33, then its focal length will be around (Refractive index of lens material is 1.5)

- a) R b) $2R$ c) $4R$ d) $R/2$

742. A given ray of light suffers minimum deviation in an equilateral prism P . Additional prisms Q and of identical shape and material are now added to P , as shown in the figure. The ray will suffer



- a) Same deviation b) Greater deviation
c) Total internal reflection d) No deviation

743. An electron microscope is superior to an optical microscope in

- a) Having better resolving power b) Being easy to handle
c) Low cost d) Quickness of observation

744. If a thin prism of glass is dipped into water then minimum deviation (with respect to air) of light produced by prism will be left (${}_a\mu_g = \frac{3}{2}$ and ${}_a\mu_w = \frac{4}{3}$)

- a) $1/2$ b) $1/4$ c) 2 d) $1/5$

745. A diminished image of an object is to be obtained on a screen 1.0 m away from it. This can be achieved by approximately placing

- a) A convex mirror of suitable focal length b) A concave mirror of suitable focal length
c) A convex lens of focal length less than 0.25 m d) A concave lens of suitable focal length

746. A symmetric double convex lens is cut in two equal parts by a plane perpendicular to the principle axis. If the power of the original lens is 4D, the power of a cut lens will be

- a) 2D b) 3D c) 4D d) 5D

747. A thin lens has focal length f_1 and its aperture has diameter d . It forms an image of intensity I . Now the central part of the aperture upto diameter $\frac{d}{2}$ is blocked by an opaque paper. The focal length and image intensity will change to

- a) $\frac{f}{2}$ and $\frac{I}{2}$ b) f and $\frac{I}{4}$ c) $\frac{3f}{4}$ and $\frac{I}{2}$ d) f and $\frac{3I}{4}$

748. A combination of two thin lenses with focal lengths f_1 and f_2 respectively forms an image of distant object at distance 60 cm when lenses are in contact. The position of this image shift by 30 cm towards the combination when two lenses are separated by 10 cm. The corresponding values of f_1 and f_2 are

- a) 30 cm, -60 cm b) 20 cm, -30 cm c) 15 cm, -20 cm d) 12 cm, -15 cm

749. The angle of minimum deviation for a prism is 40° and the angle of the prism is 60° . The angle of incidence in this position will be

- a) 30° b) 60° c) 50° d) 100°

750. A plane mirror produces a magnification of

- a) -1 b) +1 c) Zero d) Infinite

751. White light is incident on one of the refracting surfaces of a prism of angle 5° . If the refractive indices for red and blue colours are 1.641 and 1.659 respectively, the angular separation between these two colours when they emerge out of the prism is

- a) 0.9° b) 0.09° c) 1.8° d) 1.2°


752. Diameter of a plano-convex lens is 6 cm and thickness at the centre is 3 mm. If the speed of light in the material of the lens is 2×10^8 m/s, the focal length of the lens is

- a) 15 cm b) 20 cm c) 30 cm d) 10 cm

753. Two plane mirrors are perpendicular to each other. A ray after suffering reflection from the two mirrors will be

- a) Perpendicular to the original ray b) Parallel to the original ray
c) Parallel to the first mirror d) At 45° to the original ray

754. An optical fibre consists of core of μ_1 surrounded by a cladding of $\mu_2 < \mu_1$. A beam of light enters from air at an angle α with axis of fibre. The highest α for which ray can be travelled through fibre is




- a) $\cos^{-1} \sqrt{\mu_2^2 - \mu_1^2}$ b) $\sin^{-1} \sqrt{\mu_1^2 - \mu_2^2}$ c) $\tan^{-1} \sqrt{\mu_1^2 - \mu_2^2}$ d) $\sec^{-1} \sqrt{\mu_1^2 - \mu_2^2}$

755. The angle of prism is 5° and its refractive indices for red and violet colours are 1.5 and 1.6 respectively. The angular dispersion produced by the prism is

- a) 7.75° b) 5° c) 0.5° d) 0.17°

756. A ray of light strikes a plane mirror M at an angle of 45° as shown in the figure. After reflection, the ray passes through a prism of refractive index 1.5 whose apex angle is 4° . The total angle through which the ray is deviated is



- a) 90° b) 91° c) 92° d) 93°

757. Wavelength of light used in an optical instrument are $\lambda_1 = 4000 \text{ \AA}$ and $\lambda_2 = 5000 \text{ \AA}$, then ratio of their respective resolving powers (corresponding to λ_1 and λ_2) is

- a) 16 : 25 b) 9 : 1 c) 4 : 5 d) 5 : 4

758. An equiconvex lens of glass of focal length 0.1 metre is cut along a plane perpendicular to principal axis into two equal parts. The ratio of focal length of new lenses formed is

- a) 1 : 1 b) 1 : 2 c) 2 : 1 d) $2 : \frac{1}{2}$

759. The focal lengths of the objective and eye lenses of a telescope are respectively 200 cm and 5 cm. The minimum magnifying power of the telescope will be

- a) -40 b) -48 c) -60 d) -100

760. The time required for making a print at a distance of 0.25 m from a 60 W lamp is 5 s. If the distance is increased to 40 cm, the time required in second to make a similar print is

- a) 3.1 b) 8 c) 12.8 d) 16

761. An opera glass (Gallilean telescope) measures 9 cm from the objective to the eyepiece. The focal length of the objective is 15 cm. Its magnifying power is

- a) 2.5 b) $\frac{2}{5}$ c) $\frac{5}{3}$ d) 0.4

762. A point source of 3000 lumen is located at the center of a cube of side length 2m. The flux through one side is

- a) 500 lumen b) 600 lumen c) 750 lumen d) 1500 lumen

763. A convex mirror is used to form the image of an object. Then which of the following statements is wrong

- a) The image lies between the pole and the focus
b) The image is diminished in size
c) The image is erect
d) The image is real


764. The light ray is incidence at angle of 60° on a prism of angle 45° . When the light ray falls on the other surface at 90° , the refractive index of the material of prism μ and the angle of deviation δ are given by

- a) $\mu = \sqrt{2}, \delta = 30^\circ$ b) $\mu = 1.5, \delta = 15^\circ$ c) $\mu = \frac{\sqrt{3}}{2}, \delta = 30^\circ$ d) $\mu = \sqrt{\frac{3}{2}}, \delta = 15^\circ$

765. The spectrum of an oil flame is an example for

- a) Line emission spectrum
- b) Continuous emission spectrum
- c) Line absorption spectrum
- d) Band emission spectrum

766. Three right angled prisms of refractive indices n_1, n_2 and n_3 are fixed together using an optical glue as shown in figure. If a ray passes through the prisms without suffering any deviation, then



- a) $n_1 = n_2 = n_3$
- b) $n_1 = n_2 \neq n_3$
- c) $1 + n_1 = n_2 + n_3$
- d) $1 + n_2^2 = n_1^2 = n_3^2$

767. The radius of curvature for a convex lens is 40 cm, for each surface. Its refractive index is 1.5. The focal length will be

- a) 40 cm
- b) 20 cm
- c) 80 cm
- d) 30 cm

768. A white screen illuminated by green and red light appears to be

- a) Green
- b) Red
- c) Yellow
- d) White

769. A terrestrial telescope is made by introducing an erecting lens of focal length f between the objective and eye piece lenses of an astronomical telescope. This causes the length of the telescope tube to increase by an amount equal to

- a) f
- b) $2f$
- c) $3f$
- d) $4f$

770. The critical angle for a medium is 60° . The refractive index of the medium is

- a) $\frac{2}{\sqrt{3}}$
- b) $\frac{\sqrt{2}}{3}$
- c) $\sqrt{3}$
- d) $\frac{\sqrt{3}}{2}$

771. The magnifying power of telescope is high if

- a) Both objective and eye-piece have short focal length
- b) Both objective and eye-piece have long focal length
- c) The objective has a long focal length and the eye piece has a short focal length
- d) The objective has a short focal length and the eye piece has a long focal length

772. Under which of the following conditions will a convex mirror of focal length f produce an image that is erect, diminished and virtual

- a) Only when $2f > u > f$
- b) Only when $u = f$
- c) Only when $u < f$
- d) Always

773. Pick the correct statement from the following

- a) Primary rainbow is a virtual image and secondary rainbow is a real image
- b) Primary rainbow is a real image and secondary rainbow is a virtual image
- c) Both primary and secondary rainbows are virtual images
- d) Both primary and secondary rainbows are real images

774. A beam of light propagating in medium A with index of refraction $n(A)$ passes across an interface into medium B with index of refraction $n(B)$. The angle of incidence is greater than the angle of refraction; $v(A)$ and $v(B)$ denote the speed of light in A and B. Then which of the following is true

- a) $v(A) > v(B)$ and $n(A) > n(B)$
- b) $v(A) > v(B)$ and $n(A) < n(B)$
- c) $v(A) < v(B)$ and $n(A) > n(B)$
- d) $v(A) < v(B)$ and $n(A) < n(B)$

775. Solar spectrum is an example for

- a) Band absorption spectrum
- b) Line absorption spectrum
- c) Line emission spectrum
- d) Continuous emission spectrum

776. When monochromatic red light is used instead of blue light in a convex lens, its focal length will

- a) Does not depend on colour of light
- b) Increase
- c) Decrease
- d) Remain same

777. Speed of light is maximum in

- a) Water
- b) Air
- c) Glass
- d) Diamond






778. In a grease spot photometer, light from a lamp with dirty chimney is exactly balanced by a point source distance 10 cm from the grease spot. On clearing the dirty chimney, the point source is moved 2 cm to obtain a balance again. Then the percentage of light absorbed by the dirty chimney is nearly
 a) 64% b) 36% c) 44% d) 56%
779. The objective lens of a compound microscope produces magnification of 10. In order to get an overall magnification of 100 when image is formed at 25 cm from the eye, the focal length of the eye lens should be
 a) 4 cm b) 10 cm c) $\frac{25}{9}$ cm d) 9 cm
780. When white light passes through a glass prism, one gets spectrum on the other side of the prism. In the emergent beam, the ray which is deviating least is
or
 Deviation by a prism is lowest for
 a) Violet ray b) Green ray c) Red ray d) Yellow ray
781. The ratio of thickness of plates of two transparent mediums *A* and *B* is 6 : 4. If light takes equal time in passing through them, then refractive index of *B* with respect to *A* will be
 a) 1.4 b) 1.5 c) 1.75 d) 1.33
782. The sky would appear red instead of blue if
 a) Atmospheric particles scatter blue light more than red light
 b) Atmospheric particles scatter all colours equally
 c) Atmospheric particles scattered red light more than the blue light
 d) The sun was much hotter
783. If the refractive index of a material of equilateral prism is $\sqrt{3}$, then angle of minimum deviation of the prism is
 a) 30° b) 45° c) 60° d) 75°
784. An astronomical telescope has a magnifying power 10, the focal length of the eye-piece is 20 cm. The focal length of the objective is
 a) $\frac{1}{200}$ cm b) $\frac{1}{2}$ cm c) 200 cm d) 2 cm
785. A boat has green light of wavelength $\lambda = 500$ nm on the mast. What wavelength would be measured and what colour would be observed for this light as seen by a diver submerged in water by the side of the boat?
 Given, $n_w = \frac{4}{3}$.
 a) Green of wavelength 376 nm b) Red of wavelength 665 nm
 c) Green of wavelength 500 nm d) Blue of wavelength 376 nm
786. A substance is behaving as convex lens in air and concave in water, then its refractive index is
 a) Smaller than air b) Greater than both air and water
 c) Greater than air but less than water d) Almost equal to water
787. Two thin lenses when in contact, produce a combination of power +10 D. When they are 0.25 m apart, the power reduces to +6D. The focal lengths of the lenses (in m) are
 a) 0.125 and 0.5 b) 0.125 and 0.125 c) 0.5 and 0.75 d) 0.125 and 0.75
788. The graph between *u* and *v* for a convex mirror is
 a)  b)  c)  d) 
789. Two plane mirrors *A* and *B* are aligned parallel to each other, as shown in the figure. A light ray is incident at an angle of 30° at a point just inside one end of *A*. The plane of incidence coincides with the plane of the

figure. The maximum number of times the ray undergoes reflection (including the first one) before it emerges out is



- a) 28 b) 30 c) 32 d) 34

790. A ball is dropped from a height of 20m above the surface of water in a lake. The refractive index of water is $\frac{4}{3}$. A fish inside the lake, in the line of fall of the ball, is looking at the ball. At an instant, when the ball is 12.8 m above the water surface, the fish sees the speed of ball as

- a) 9 ms^{-1} b) 12 ms^{-1} c) 16 ms^{-1} d) 21.33 ms^{-1}


791. The no. of wavelengths in the visible spectrum

- a) 4000 b) 6000 c) 2000 d) Infinite

792. A 60 watt bulb is hung over the center of a table $4 \text{ m} \times 4 \text{ m}$ at height of 3 m. The ratio of the intensities of illumination at a point on the centre of the edge and on the corner of the table is

- a) $(17/13)^{3/2}$ b) $2/1$ c) $17/13$ d) $5/4$

793. If the space between the lenses in the lens combination shown were filled with water, what would happen to the focal length and power of the lens combination?



Focal Length

Power

- | | | | |
|--------------|--------------|--------------|--------------|
| a) Decreased | b) Decreased | c) Increased | d) Increased |
| Increased | Unchanged | Unchanged | Decreased |

794. A lamp is hanging 1 m above the centre of a circular table of diameter 1m. The ratio of illuminances at the centre and the edge is

- a) $\frac{1}{2}$ b) $\left(\frac{5}{3}\right)^{\frac{3}{2}}$ c) $\frac{4}{3}$ d) $\frac{4}{5}$

795. The spectrum obtained from an electric lamp or red hot heater is

- a) Line spectrum b) Band spectrum
c) Absorption spectrum d) Continuous spectrum

796. The separation between the screen and a plane mirror is $2r$. An isotropic point source of light is placed exactly mid way between the mirror and the screen. Assume that mirror reflects 100% of incident light. Then the ratio of illuminance on the screen with and without the mirror is

- a) 10:1 b) 2:1 c) 10:9 d) 9:1

797. The minimum distance between the object and its real image for concave mirror is

- a) f b) $2f$ c) $4f$ d) Zero

798. In human eye the focussing is done by


- a) To and fro movement of eye lens
b) To and fro movement of the retina
c) Change in the convexity of the lens surface
d) Change the refractive index of the eye fluids

799. When the length of a microscope tube increases, its magnifying power


- a) Decreases b) Increases
c) Does not change d) May decrease or increase

800. An object 15 cm high is placed 10 cm from the optical centre of a thin lens. Its image is formed 25 cm from the optical centre on the same side of the lens as the object. The height of the image is
 a) 2.5 cm b) 0.2 cm c) 16.7 cm d) 37.5 cm
801. For a convex lens the distance of the object is taken on X, axis and the distance of the image is taken on Y-axis, the nature of the graph so obtained is
 a) Straight line b) Circle c) Parabola d) Hyperbola
802. A virtual image larger than the object can be obtained by
 a) Concave mirror b) Convex mirror c) Plane mirror d) Concave lens
803. A glass lens is placed in a medium in which it is found to behave like a glass plate. Refractive index of the medium will be
 a) Greater than the refractive index of glass
 b) Smaller than the refractive index of glass
 c) Equal to refractive index of glass
 d) No case will be possible from above
804. An object has image thrice of its original size when kept at 8 cm and 16 cm from a convex lens. Focal length of the lens is
 a) 8 cm b) 16 cm
 c) Between 8 cm and 16 cm d) Less than 8 cm
805. An object 1cm tall is placed 4 cm in front of a mirror. In order to produce an upright image of 3 cm height one needs a
 a) Convex mirror of radius of curvature 12 cm
 b) Concave mirror of radius of curvature 12 cm
 c) Concave mirror of radius of curvature 4 cm
 d) Plane mirror of height 12 cm
806. When the convergent nature of a convex lens will be less as compared with air
 a) In water b) In oil c) In both (a) and (b) d) None of these
807. A person can see objects clearly only upto a maximum distance of 50 cm. His eye defect, nature of the corrective lens and its focal length are respectively
 a) Myopia, concave, 50 cm b) Myopia, convex, 50 cm
 c) Hypermetropia, concave, 50 cm d) Catract, convex, 50 cm
808. A convex lens makes a real image 4 cm long on a screen. When the lens is shifted to a new position without disturbing the object, we again get a real image on the screen which is 16 cm tall. The length of the object must be
 a) $1/4$ cm b) 8 cm c) 12 cm d) 20 cm
809. Two beams of red and violet colours are made to pass separately through a prism (angle of the prism is 60°). In the position of minimum deviation, the angle of refraction will be
 a) 30° for both the colours b) Greater for the violet colour
 c) Greater for the red colour Equal but not 30° for both the
 d) colours
810. A beam of monochromatic blue light of wavelength 4200 \AA in air travels in water of refractive index $4/3$. Its wavelength in water will be
 a) 4200 \AA b) 5800 \AA c) 4150 \AA d) 3150 \AA
811. What is the ratio of luminous intensity of two sources, which produce shadows of equal intensities at distance 25 cm and 50 cm from the photometer screen?
 a) 1:4 b) 4:1 c) 1:2 d) 2:1
812. In Huygen's eyepiece
 a) The cross wires are outside the eyepiece
 b) Condition for achromatism is satisfied

- c) Condition for minimum spherical aberration is not satisfied
 d) The image formed by the objective is a virtual image
813. Blue colour of sea water is due to
 a) Interference of sunlight reflected from the water surface
 b) Scattering of sunlight by the water molecules
 c) Image of sky in water
 d) Refraction of sunlight
814. The refractive index of the material of a double convex lens is 1.5 and its focal length is 5 cm. If the radii of curvature are equal, the value of the radius of curvature (in cm) is
 a) 5.0 b) 6.5 c) 8.0 d) 9.5
815. Light from sodium lamp is passed through cold sodium vapours, the spectrum of transmitted light consists of
 a) A line at 5890 \AA b) A line at 5896 \AA c) Sodium doublet lines d) No spectral features
816. Given figures show the arrangements of two lenses, The radii of curvature of all the curved surfaces are same. The ratio of the equivalent focal length of combinations P , Q and R is




- a) $1 : 1 : 1$ b) $1 : 1 : -1$ c) $2 : 1 : 1$ d) $2 : 1 : 2$
817. In the given figure, what is the angle of prism?



- a) A b) B c) C d) D
818. An eye specialist prescribes spectacles having a combination of convex lens of focal length 40 cm in contact with a concave lens of focal length 25 cm. The power of this lens combination in dioptres is
 a) +1.5 b) -1.5 c) +6.67 d) -6.67
819. For which of the following colour, the magnifying power of a microscope will be maximum
 a) White colour b) Red colour c) Violet colour d) Yellow colour
820. A point objects is placed at the centre of a glass sphere of radius 6 cm and refractive index 1.5. The distance of the virtual image from the surface of the sphere is
 a) 2 cm b) 4 cm c) 6 cm d) 12 cm
821. A boy stands straight in front of a mirror at a distance of 30 cm away from it. He sees his erect image whose height is $1/5^{\text{th}}$ of his real height. The mirror he is using is
 a) Plane mirror b) Convex mirror c) Concave mirror d) Plano-convex mirror
822. A point object O is placed on the principal axis of a convex lens of focal length 20 cm at a distance of 40 cm to the left of it. The diameter of the lens is 10 cm. If the eye is placed 60 cm to the right of the lens at a distance h below the principal axis, then the maximum value of h to see the image will be
 a) 0 b) 5 cm c) 2.5 cm d) 10 cm
823. A person cannot see distinctly at the distance less than one metre. Calculate the power of the lens that he should use to read a book at a distance of 25 cm

- a) $+3.0 D$ b) $+0.125 D$ c) $-3.0 D$ d) $+4.0 D$
824. Near and far points of human eye are
a) 25 cm and infinite b) 50 cm and 100 cm c) 25 cm and 50 cm d) 0 cm and 25 cm
825. Two mirrors at an angle θ produce 5 images of a point. The number of images produced when θ is decreased to 30° is
a) 9 b) 10 c) 11 d) 12
826. A lens of focal power $0.5 D$ is
a) A convex lens of focal length $0.5 m$ b) A concave lens of focal length $0.5 m$
c) A convex lens of focal length $2 m$ d) A concave lens of focal length $2 m$
827. The power of two convex lenses A and B are 8 dioptres and 4 dioptres respectively. If they are to be used as a simple microscope, the magnification of
a) B will be greater than A b) A will be greater than B
c) The information is incomplete d) None of the above
828. A glass slab ($\mu = 1.5$) of thickness $6 m$ is placed over a paper. What is the shift in the letters?
a) 4 cm b) 2 cm c) 1 cm d) None of the above
829. When light rays from the sun fall on a convex lens along a direction parallel to its axis
a) Focal length for all colours is the same
b) Focal length for violet colour is the shortest
c) Focal length for yellow colour is the longest
d) Focal length red colour is the shortest
830. Beams of red, green and violet light are falling on the refracting face of a prism, all at the same angle of incidence, if their angles of deviation are θ_1 , θ_2 and θ_3 respectively, then
a) $\theta_1 = \theta_2 = \theta_3$ b) $\theta_1 < \theta_2 < \theta_3$ c) $\theta_1 > \theta_2 > \theta_3$ d) $\theta_2 > \theta_1 > \theta_3$
831. Two plane mirrors are inclined to each other such that a ray of light incident on the first mirror and parallel to the second is reflected from the second mirror parallel to the first mirror. The angle between the two mirrors is
a) 30° b) 45° c) 60° d) 75°
832. Two lenses have focal lengths f_1 and f_2 and their dispersive powers are ω_1 and ω_2 respectively. They will together form an achromatic combination if
a) $\omega_1 f_1 = \omega_2 f_2$ b) $\omega_1 f_2 + \omega_2 f_1 = 0$ c) $\omega_1 + f_1 = \omega_2 + f_2$ d) $\omega_1 - f_1 = \omega_2 - f_2$
833. A lens (focal length $50 cm$) forms the image of a distant object which subtends an angle of 1 milliradian at the lens. What is the size of the image
a) $5 mm$ b) $1 mm$ c) $0.5 mm$ d) $0.1 mm$
834. At sun rise or sunset, the sun looks more red than at mid-day because
a) The sun is hottest at these times b) Of the scattering of light
c) Of the effect of refraction d) Of the effect of diffraction
835. Light enters at an angle of incidence in a transparent rod of refractive index n . For what value of the refractive index of the material of the rod the light once entered into it will not leave it through its lateral face whatever be the value of angle of incidence
a) $n > \sqrt{2}$ b) $n = 1$ c) $n = 1.1$ d) $n = 1.3$
836. A ray of light suffers minimum deviation when incident at 60° on a prism of refractive index $\sqrt{2}$. The angle of incidence is
a) $\sin^{-1}(0.8)$ b) 60° c) 45° d) 30°
837. The focal length of objective and eye lens of an astronomical telescope are respectively $2 m$ and $5 cm$. Final image is formed at (1) least distance of distinct vision (2) infinity. Magnifying powers in two cases will be
a) $-48, -40$ b) $-40, 48$ c) $-40, +48$ d) $-48 + 40$
838. Dispersion of light is due to
a) Wavelength b) Intensity of light c) Density of medium d) None of these

839. Light takes 8 min 20 s to reach from sun on the earth. If the whole atmosphere is filled with water, the light will take the time (${}_a\mu_w = 4/3$)
 a) 8 min 20 s b) 8 min c) 6 min 11 s d) 11 min 6 s
840. Velocity of light in glass whose refractive index with respect to air is 1.5 is 2×10^8 m/s and in certain liquid the velocity of light found to be 2.5×10^8 m/s. The refractive index of the liquid with respect to air is
 a) 0.64 b) 0.80 c) 1.20 d) 1.44
841. Magnifying power of a simple microscope is (when final image is formed at $D = 25$ cm from eye)
 a) $\frac{D}{f}$ b) $1 + \frac{D}{f}$ c) $1 + \frac{f}{D}$ d) $1 - \frac{D}{f}$
842. The apparent depth of water in cylindrical water tank of diameter $2R$ cm is reducing at the rate of x cm/minute when water is being drained out at a constant rate. The amount of water drained in c. c. per minute is (n_1 = refractive index of air, n_2 = refractive index of water)
 a) $x \pi R^2 n_1/n_2$ b) $x \pi R^2 n_2/n_1$ c) $2 \pi R n_1/n_2$ d) $\pi R^2 x$
843. A prism of angle 30° is silvered at one side. A ray of light incident at an angle 45° is reflected back from the silvered surface. The refractive index is
 a) $\sqrt{2}$ b) $2\sqrt{2}$ c) $\sqrt{3}$ d) $5\sqrt{3}$
844. The distance between an object and the screen is 100 cm. A lens produces an image on the screen when placed at either of the position 40 cm apart. The power of the lens is
 a) ≈ 3 dioptres b) ≈ 5 dioptres c) ≈ 7 dioptres d) ≈ 9 dioptres
845. A man with defective eyes cannot see distinctly object at the distance more than 60 cm from his eyes. The power of the lens to be used will be
 a) +60 D b) -60 D c) -1.66 D d) $\frac{1}{1.66}$ D
846. A parallel beam of light emerges from the opposite surface of the sphere when a point source of light lies at the surface of the sphere. The refractive index of the sphere is
 a) $\frac{3}{2}$ b) $\frac{5}{3}$ c) 2 d) $\frac{5}{2}$
847. In the adjoining diagram, a wavefront AB, moving in air is incident on a plane glass surface XY. Its position CD after refraction through a glass slab is shown also along with the normal drawn at A and D. The refractive index of glass with respect to air ($\mu = 1$) will be equal to



- a) $\frac{\sin \theta}{\sin \phi'}$ b) $\frac{\sin \theta}{\sin \phi'}$ c) $\frac{\sin \phi'}{\sin \theta}$ d) $\frac{AB}{CD}$
848. A diver at a depth of 12m in water ($\mu = \frac{4}{3}$) sees the sky in a cone of semivertical angle
 a) $\sin^{-1}\left(\frac{4}{3}\right)$ b) $\tan^{-1}\left(\frac{4}{3}\right)$ c) $\sin^{-1}\left(\frac{3}{4}\right)$ d) 90°
849. The plane surface of a plano-convex lens of focal length f is silvered. It will behave as
 a) Plane mirror b) Convex mirror of focal length $2f$
 c) Concave mirror of focal length $\frac{f}{2}$ d) None of the above

850. A large glass slab ($\mu = \frac{5}{3}$) of thickness 8 cm is placed over a point source of light on a plane surface. It is seen that light emerges out of the top surface of the slab from a circular area of radius R cm. What is the value of R ?

- a) 6 cm b) 7 cm c) 8 cm d) 9 cm

851. The focal length of an equi-convex lens is greater than the radius of curvature of any of the surfaces. Then the refractive index of the material of the lens is

- a) Greater than zero but less than 1.5 b) Greater than 1.5 but less than 2.0
c) Greater than 2.0 but less than 2.5 d) Greater than 2.5 but less than 2.0

852. The frequency of a light ray is $6 \times 10^{14} \text{ Hz}$. Its frequency when it propagates in a medium of refractive index 1.5, will be

- a) $1.67 \times 10^{14} \text{ Hz}$ b) $9.10 \times 10^{14} \text{ Hz}$ c) $6 \times 10^{14} \text{ Hz}$ d) $4 \times 10^{14} \text{ Hz}$


853. Which of the following is not due to total internal reflection

- a) Brilliance of diamond
b) Working of optical fibre
c) Difference between apparent and real depth of a pond
d) Mirage on hot summer days

854. The diameter of objective of a telescope is 1 m. its resolving limit for the light of wavelength 4538 \AA , will be


- a) $5.54 \times 10^{-7} \text{ rad}$ b) $2.54 \times 10^{-4} \text{ rad}$ c) $6.54 \times 10^{-7} \text{ rad}$ d) None of the above

855. Shown in the figure here is a convergent lens placed inside a cell filled with a liquid. The lens has focal length $+20 \text{ cm}$ when in air and its material has refractive index 1.50. If the liquid has refractive index 1.60, the focal length of the system is



- a) $+80 \text{ cm}$ b) -80 cm c) -24 cm d) -100 cm

856. While light is incident on the interface of glass and air as shown in the figure. If green light is just totally internally reflected then the emerging ray in air contains



- a) Yellow, orange, red b) Violet, indigo, blue
c) All colours d) All colours except green

857. For compound microscope $f_o = 1 \text{ cm}$, $f_e = 2.5 \text{ cm}$. An object is placed at distance 1.2 cm from object lens. What should be length of microscope for normal adjustment?

- a) 8.5 cm b) 8.3 cm c) 6.5 cm d) 6.3 cm


858. An achromatic prism is made by combining two prisms $P_1 (\mu_v = 1.523, \mu_r = 1.515)$ and $P_2 (\mu_v = 1.666, \mu_r = 1.650)$; where μ represents the refractive index. If the angle of the prism P_1 is 10° , then the angle of the prism P_2 will be

- a) 5° b) 7.8° c) 10.6° d) 20°

859. In a movie hall, the distance between the projector and the screen is increased by 1% illuminates on the screen is

- a) Increased by 1% b) Decreased by 1% c) Increased by 2% d) Decreased by 2%


860. Two points, separated by a distance of 0.1 mm, can just be inspected on a microscope when light of wavelength 6000 \AA is used. If the light of wavelength 4800 \AA is used, the limit of resolution is
 a) 0.8 mm b) 0.08 mm c) 0.1 mm d) 0.04 mm
861. If two +5 D, lenses are mounted at some distance apart, the equivalent power will always be negative, if the distance is
 a) Greater than 40 cm b) Equal to 10 cm c) Equal to 10 cm d) Less than 10 cm
862. Two thin lenses have a combined power of + 9D. When they are separated by a distance of 20 cm, their equivalent power becomes $+\frac{27}{5}$ D. Their individual powers (in dioptre) are
 a) 4, 5 b) 3, 6 c) 2, 7 d) 1, 8
863. Following figure shows the multiple reflections of a light ray along a glass corridor where the walls are either parallel or perpendicular to one another. If the angle of incidence at point P is 30° , what are the angles of reflection of the light ray at points Q, R, S and T respectively




- a) $30^\circ, 30^\circ, 30^\circ, 30^\circ$ b) $30^\circ, 60^\circ, 30^\circ, 60^\circ$ c) $30^\circ, 60^\circ, 60^\circ, 30^\circ$ d) $60^\circ, 60^\circ, 60^\circ, 60^\circ$
864. A ray is incident at an angle of incidence i on one surface of a prism of small angle A and emerges normally from the opposite surface. If the refractive index of the material of the prism is μ , the angle of incidence i is nearly equal to
 a) A/μ b) $A/2\mu$ c) μA d) $\mu A/2$
865. The sun (diameter d) subtends an angle θ radian at the pole of a concave mirror of focal length f . The diameter of the image of sun formed by mirror is
 a) θf b) $\frac{\theta}{2} f$ c) $2\theta f$ d) $\frac{\theta}{\pi} f$
866. A glass hemisphere of radius 0.04 m and R.I. of the material 1.6 is placed centrally over a cross mark on a paper (i) with the flat face; (ii) with the curved face in contact with the paper. In each case the cross mark is viewed directly from above. The position of the images will be
 a) (i) 0.04 m from the flat face; (ii) 0.025 m from the flat face
 b) (i) At the same position of the cross mark; (ii) 0.025 m below the flat face
 c) (i) 0.025 m from the flat face; (ii) 0.04 m from the flat face
 d) For both (i) and (ii) 0.025 m from the highest point of the hemisphere
867. A plano convex lens fits exactly into a plano concave lens. Their plane surfaces are parallel to each other. If the lenses are made of different materials of refractive indices μ_1 and μ_2 and R is the radius of curvature of the curved surface of the lenses, then focal length of the combination is
 a) $\frac{R}{2(\mu_1 + \mu_2)}$ b) $\frac{R}{2(\mu_1 - \mu_2)}$ c) $\frac{R}{(\mu_1 - \mu_2)}$ d) $\frac{2R}{(\mu_2 - \mu_1)}$
868. A point object is placed at a distance of 30 cm from a convex mirror of a focal length 30 cm. The image will form at
 a) Infinity b) Pole
 c) 15 cm behind the mirror d) No image will be formed
869. A person wears glasses of power -2.0 D . The defect of the eye and the far point of the person without the glasses will be
 a) Nearsighted, 50 cm b) Farsighted, 50 cm c) Nearsighted, 250 cm d) Astigmatism, 50 cm
870. On which of the following does the magnifying power of a telescope depends
 a) The focal length of the objective only

- b) The diameter of aperture of the objective only
 c) The focal length of the objective and that of the eye piece
 d) The diameter of aperture of the objective and that of the eye piece
871. The wavelength of emission line spectrum and absorption line spectrum of a substance are related as
 a) Absorption has larger value b) Absorption has smaller value
 c) They are equal d) No relation
872. A ray of light is incident on the surface of separation of a medium at an angle 45° and is refracted in the medium at an angle 30° . What will be the velocity of light in the medium
 a) $1.96 \times 10^8 \text{ m/s}$ b) $2.12 \times 10^8 \text{ m/s}$ c) $3.18 \times 10^8 \text{ m/s}$ d) $3.33 \times 10^8 \text{ m/s}$
873. An achromatic prism is made by crown glass prism ($A_c = 19^\circ$) and flint glass prism ($A_f = 6^\circ$). If ${}^c\mu_v = 1.5$ and ${}^f\mu_v = 1.66$, then resultant deviation for red coloured ray will be
 a) 1.04° b) 5° c) 0.96° d) 13.5°
874. Two plane mirrors are inclined at an angle θ . It is found that a ray incident on one mirror at any angle is rendered parallel to itself after reflection from both the mirrors. The value of θ is
 a) 30° b) 60° c) 90° d) 120°
875. Two plano-concave lenses (1 and 2) of glass of refractive index 1.5 have radii of curvature 25 cm and 20 cm. They are placed in contact with their curved surfaces towards each other and the space between them is filled with liquid of refractive index $\frac{4}{3}$. Then the combination is
-
- a) Convex of focal length 70 cm b) Concave of focal length 70 cm
 c) Concave of focal length 66.6 cm d) Convex of focal length 66.6 cm
876. What will be the height of the image when an object of 2mm is placed at a distance 20 cm in front of the axis of a convex mirror of radius of curvature 40 cm?
 a) 20 mm b) 10 mm c) 6 mm d) 1 mm
877. When a ray of light enters a glass slab from air
 a) Its wavelength decreases
 b) Its wavelength increases
 c) Its frequency increases
 d) Neither its wavelength nor its frequency changes
878. A camera objective has an aperture diameter d . If the aperture is reduced to diameter $d/2$, the exposure time under identical conditions of light should be made
 a) $\sqrt{2}$ fold b) 2 fold c) $2\sqrt{2}$ fold d) 4 fold
879. A satisfactory photographic print is obtained when the exposure time is 10 s at a distance of 2 m from a 60 cd lamp. The time of exposure required for the same quality print at a distance of 4 m from a 120 cd lamp is
 a) 5 s b) 10 s c) 15 s d) 20 s
880. Lux is equal to
 a) 1 lumen/m^2 b) 1 lumen/cm^2 c) 1 candela/m^2 d) 1 candela/cm^2
881. Rising and setting sun appears to be reddish because
 a) Diffraction sends red rays to earth at these times
 b) Scattering due to dust particles and air molecules are responsible
 c) Refraction is responsible
 d) Polarization is responsible


882. Two parallel light rays are incident at one surface of a prism of refractive index 1.5 as shown in figure. The angle between the emergent rays is nearly
 a) 19° b) 37° c) 45° d) 49°
883. When white light enters a prism, it gets split into its constituent colours. This is due to
 a) High density of prism material
 b) Because μ is different for different λ
 c) Diffraction of light
 d) Velocity changes for different frequencies
884. A ray of light is incident normally on one of the face of a prism of angle 30° and refractive index $\sqrt{2}$. The angle of deviation will be
 a) 26° b) 0° c) 23° d) 15°
885. Refractive index of air is 1.0003. The correct thickness of air column which will have one more wavelength of yellow light (6000 \AA) than in the same thickness in vacuum is
 a) 2 mm b) 2 cm c) 2 m d) 2 km
886. A ray of light travelling inside a rectangular glass block of refractive index $\sqrt{2}$ is incident on the glass-air surface at an angle of incidence of 45° . The refractive index of air is 1. Under these conditions the ray
 a) Will emerge into the air without any deviation
 b) Will be reflected back into the glass
 c) Will be absorbed
 d) Will emerge into the air with angle of refraction equal to 90°
887. The band spectra (characteristic of molecular species) is due to emission of radiation
 a) Gaseous state b) Liquid state c) Solid state d) All of three states
888. In a thin spherical fish bowl of radius 10 cm filled with water of refractive index $4/3$ there is a small fish at a distance of 4 cm from the centre C as shown in figure. Where will the image of fish appear, if seen from E



- a) 5.2 cm b) 7.2 cm c) 4.2 cm d) 3.2 cm
889. The intensity of direct sunlight on a surface normal to the rays is I_0 . What is the intensity of direct sunlight on a surface, whose normal makes an angle of 60° with the rays of the sun
 a) I_0 b) $I_0 \left(\frac{\sqrt{3}}{2} \right)$ c) $\frac{I_0}{2}$ d) $2I_0$
890. The focal length of an objective of a telescope is 3 metre and diameter 15 cm . Assuming for a normal eye, the diameter of the pupil is 3 mm for its complete use, the focal length of eye piece must be
 a) 6 cm b) 6.3 cm c) 20 cm d) 60 cm
891. A glass slab of thickness 3 cm and refractive index $3/2$ of placed on ink mark on a piece of paper. For a person looking at the mark at a distance 5.0 cm above it, the distance of the mark will appear to be
 a) 3.0 cm b) 4.0 cm c) 4.5 cm d) 5.0 cm
892. A rod of glass ($\mu = 1.5$) and of square cross section is bent into the shape shown in the figure. A parallel beam of light falls on the plane flat surface A as shown in the figure. If d is the width of a side and R is the radius of circular arc then for what maximum value of $\frac{d}{R}$ light entering the glass slab through surface A emerges from the glass through B




- a) 1.5 b) 0.5 c) 1.3 d) None of these
893. A ray of light falls on a transparent glass slab with refractive index (relative to air) of 1.62. The angle of incidence for which the reflected and refracted rays are mutually perpendicular is
- a) $\tan^{-1}(1.62)$ b) $\sin^{-1}(1.62)$ c) $\cos^{-1}(1.62)$ d) None of these
894. When a ray of light is incident normally on a surface, then
- a) Total internal reflection takes place b) It passes undeviated
- c) It undergoes dispersion d) It gets absorbed by the surface
895. The fine powder of a coloured glass is seen as
- a) Coloured b) White
- c) That of the glass colour d) Black
896. The image of a small electric bulb fixed on the wall of a room is to be obtained on the opposite wall 4 m away by means of a large convex lens. The maximum possible focal length of the lens required for this purpose will be
- a) 0.5 m b) 1.0 m c) 1.5 m d) 2.0 m
897. A convex lens A of focal length 20 cm and a concave lens B of focal length 56 cm are kept along the same axis with the distance d between them. If a parallel beam of light falling on A leaves B as a parallel beam, then distances d in cm will be
- a) 25 b) 36 c) 30 d) 50
898. If the focal length of the lens is 20 cm, what is the distance of the image from the lens in the following figure?




- a) 5.5 cm b) 7.5 cm c) 12.0 cm d) 20.0 cm
899. When a ray of light is incident normally on one refracting surface of an equilateral prism (Refractive index of the material of the prism = 1.5)
- a) Emerging ray is deviated by 30°
- b) Emerging ray is deviated by 45°
- c) Emerging ray just grazes the second refracting surface
- d) The ray undergoes total internal reflection at the second refracting surface
900. A parallel beam of light is incident on a converging lens parallel to its principal axis. As one moves away from the lens on the other side of the principal axis, the intensity of light
- a) First decreases and then increases b) Continuously increases
- c) Continuously decreases d) First increases and then decreases
901. The diameter of the eye-ball of a normal eye is about 2.5 cm. The power of the eye lens varies from

- a) $2D$ to $10D$ b) $40D$ to $32D$ c) $9D$ to $8D$ d) $44D$ to $40D$
902. A transparent plastic bag filled with air forms a concave lens. Now, if this bag is completely immersed in water, then it behaves as
 a) Divergent lens b) Convergent lens c) Equilateral prism d) Rectangular slab
903. The focal length of a convex mirror is 20 cm its radius of curvature will be
 a) 10 cm b) 20 cm c) 30 cm d) 40 cm
904. With a concave mirror, an object is placed at a distance x_1 from the principal focus, on the principal axis. The image is formed at a distance x_2 from the principal focus. The focal length of the mirror is
 a) x_1x_2 b) $\frac{x_1 + x_2}{2}$ c) $\sqrt{\frac{x_1}{x_2}}$ d) $\sqrt{x_1x_2}$
905. A point object is placed on the axis of the concave mirror at a distance of 60 cm from the focal point of the mirror. Its image is formed at the point of object, then focal length of the mirror is
 a) 15 cm b) 30 cm c) 60 cm d) 120 cm
906. In Ramsden eyepiece, the two planoconvex lenses each of focal length f are separated by a distance 12 cm . The equivalent focal length (in cm) of the eyepiece is
 a) 10.5 b) 12.0 c) 13.5 d) 15.5
907. A combination of two thin convex lenses of focal length 0.3 m and 0.1 m will have minimum spherical and chromatic aberrations if the distance between them is
 a) 0.1 m b) 0.2 m c) 0.3 m d) 0.4 m
908. From a spherical mirror, the graph of $1/v$ versus $1/u$ is



909. An electric bulb illuminates a plane surface. The intensity of illumination on the surface at a point 2 m away from the bulb 5×10^{-4} phot (lumen cm^{-2}). The line joining the bulb to the point makes an angle of 60° with the normal to the surface. The intensity of the bulb in candela (candle power) is
 a) 40×10^{-4} b) 40 c) $40\sqrt{3}$ d) 20
910. The radius of the convex surface of plano-convex lens is 20 cm and the refractive index of the material of the lens is 1.5 . The focal length of the lens is
 a) 30 cm b) 50 cm c) 20 cm d) 40 cm
911. A man is suffering from colour blindness for green colour. To remove this defect, he should use goggles of
 a) Green colour glasses b) Red colour glasses c) Smoky colour glasses d) none of the above
912. A concave mirror of focal length f (in air) is immersed in water ($\mu = 4/3$). The focal length of the mirror in water will be
 a) f b) $\frac{4}{3}f$ c) $\frac{3}{4}f$ d) $\frac{7}{3}f$
913. A light ray is incident perpendicular to one face of a 90° prism and is totally internally reflected at the glass-air interface. If the angle of reflection is 45° , we conclude that the refractive index n



- a) $n < \frac{1}{\sqrt{2}}$ b) $n > \sqrt{2}$ c) $n > \frac{1}{\sqrt{2}}$ d) $n < \sqrt{2}$

914. A double convex thin lens made of refractive index 1.6 has radii of curvature 15 cm each. The focal length of this lens when immersed in a fluid of refractive index 1.63, is

- a) 25 cm b) 125 cm c) 250 cm d) -407.5 cm

915. If the angle of prism is 60° and the angle of minimum deviation is 40° , the angle of refraction will be

- a) 30° b) 60° c) 100° d) 120°

916. Dispersion can take place for

- a) Transverse waves only but not for longitudinal waves
b) Longitudinal waves only but not for transverse waves
c) Both transverse and longitudinal waves
d) Neither transverse nor longitudinal

917. A diver inside water ($\mu = 1.33$) should see the sun set at an angle of

- a) 60° b) 90° c) 0° d) 49°


918. A point object is placed at a distance of 10 cm and its real image is formed at a distance of 20 cm from a concave mirror. If the object is moved by 0.1 cm towards the mirror, the image will shift by about

- a) 0.4 cm away from the mirror b) 0.4 cm towards the mirror
c) 0.8 cm away from the mirror d) 0.8 cm towards the mirror

919. Velocity of light in a medium is $1.5 \times 10^8 \text{ m/s}$. Its refractive index will be

- a) 8 b) 6 c) 4 d) 2

920. In an experiment to find the focal length of a concave mirror a graph is drawn between the magnitudes of u and v . The graph looks like




921. Colour of the sky is blue due to

- a) Scattering of light b) Total internal reflection
c) Total emission d) None of the above

922. A bucket contains some transparent liquid and its depth is 40 cm. On looking from above, the bottom appears to be raised up by 8 cm. The refractive index of the liquid is

- a) $5/4$ b) 5 c) $4/5$ d) $8/5$

923. A fish rising vertically up towards the surface of water with speed 3 ms^{-1} observes a bird diving vertically down towards it with speed 9 ms^{-1} . The actual velocity of bird is



- a) 4.5 ms^{-1} b) 5 ms^{-1} c) 3.0 ms^{-1} d) 3.4 ms^{-1}


924. A small piece of wire bent into an L shape with upright and horizontal portions of equal lengths, is placed with the horizontal portion along the axis of the concave mirror whose radius of curvature is 10 cm. If the bend is 20 cm from the pole of the mirror, then the ratio of the lengths of the images of the upright and horizontal portions of the wire is

- a) 1 : 2 b) 3 : 1 c) 1 : 3 d) 2 : 1

925. The distance between an object and a divergent lens is m times the focal length of the lens. The linear magnification produced by the lens is

- a) m b) $1/m$ c) $m + 1$ d) $\frac{1}{m + 1}$
926. A car is fitted with a convex mirror of focal length 20 cm. A second car 2 m broad and 1.6 m height is 6 cm away from the first car. The position of the second car as seen in the mirror or the first car is
a) 19.35 cm b) 17.45 cm c) 21.48 cm d) 15.49 cm
927. How should people wearing spectacles work with a microscope
a) They cannot use the microscope at all
b) They should keep on wearing their spectacles
c) They should take off spectacles
d) (b) and (c) is both way
928. The minimum distance between an object and its real image formed by a convex lens is
a) $1.5 f$ b) $2 f$ c) $2.5 f$ d) $4 f$
929. A prism having refractive index 1.414 and refractive angle 30° has one of the refracting surfaces silvered. A beam of light incident on the other refracting surface will retrace its path, if the angle of incidence is
a) 45° b) 60° c) 30° d) 0°
930. A biconvex lens of focal length f forms a circular image of radius r of sun in focal plane. Then which option is correct?
a) $\pi r^2 \propto f$
b) $\pi r^2 \propto f^2$
c) If lower half part is covered by black sheet, then area of the image is equal to $\pi r^2 / 2$
d) If f is doubled, intensity will increase
931. A ray of light passes through an equilateral prism such that the angle of incidence is equal to the angle of emergence and the latter is equal to $\frac{3}{4}$ the angle of prism. The angle of deviation is
a) 25° b) 30° c) 45° d) 35°
932. The light reflected by a plane mirror may form a real image
a) If the rays incident on the mirror are diverging
b) If the rays incident on the mirror are converging
c) If the object is placed very close to the mirror
d) Under no circumstances
933. A virtual image twice as long as the object is formed by a convex lens when the object is 10 cm away from it. A real image twice as long as the object will be formed when it is placed at a distance.....from the length
a) 40 cm b) 30 cm c) 20 cm d) 15 cm
934. A telescope consists of two thin lenses of focal lengths 0.3 m and 3 cm respectively. It is focused on moon which subtends an angle of 0.5° at the objective. Then, the angle subtended at the eye by the final image will be
a) 5° b) 0.25° c) 0.5° d) 0.35°
935. The refractive index of a material of a planoconcave lens is $5/3$, the radius of curvature is 0.3 m. The focal length of the lens in air is
a) -0.45m b) -0.6m c) -0.75m d) -1.0m
936. The minimum temperature of a body at which it emits light is
a) 1200°C b) 1000°C c) 500°C d) 200°C
937. The refracting angle of a prism is A and the refractive index of the material of the prism is $\cot(A/2)$. The angle of minimum deviation of the prism is
a) $\pi + 2A$ b) $\pi - 2A$ c) $\frac{\pi}{2} + A$ d) $\frac{\pi}{2} - A$
938. If tube length of astronomical telescope is 105 cm and magnifying power is 20 for normal setting, calculate the focal length of objective
a) 100 cm b) 10 cm c) 20 cm d) 25 cm

939. An object is placed at a distance 20 cm from the pole of a convex mirror of focal length 20 cm. The image is produced at
 a) 13.3 cm b) 20 cm c) 25 cm d) 10 cm
940. A thin lens of glass ($\mu = 1.5$) of focal length +10 cm is immersed in water ($\mu = 1.33$). The new focal length is
 a) 20 cm b) 40 cm c) 48 cm d) 12 cm
941. A beam of light consisting of red, green and blue colours is incident on a right-angled prism ABC . The refractive indices of the material of the prism for the above red, green and blue wavelengths are 1.39, 1.44 and 1.47 respectively. The colour/colours transmitted through the face AC of the prism will be




- a) Red only b) Red and green c) All the three d) None of these
942. When the power of eye lens increases, the defect of vision is produced. The defect is known as
 a) Shortsightedness b) Longsightedness c) Colourblindness d) None of the above
943. When an object is kept at a distance of 30 cm from a concave mirror, the image is formed at a distance of 10 cm. If the object is moved with a speed of 9 ms^{-1} , the speed with which images moved is
 a) 0.1 ms^{-1} b) 1 ms^{-1} c) 3 ms^{-1} d) 9 ms^{-1}
944. Wavelength of given light waves in air and in a medium are 6000 \AA and 4000 \AA respectively. The critical angle is
 a) $\sin^{-1}\left(\frac{3}{2}\right)$ b) $\sin^{-1}\left(\frac{2}{3}\right)$ c) $\tan^{-1}\left(\frac{3}{2}\right)$ d) $\tan^{-1}\left(\frac{2}{3}\right)$
945. A candle is placed before a thick plane mirror. When looked obliquely in the mirror, a number of images are seen from the surfaces of the plane mirror. Then
 a) first image is brightest b) second image is brightest
 c) third image is brightest d) all images beyond second are brightest
946. A telescope using light having wavelength 5000 \AA and using lenses of focal 2.5 and 30 cm. If the diameter of the aperture of the objective is 10 cm, then the resolving limit and magnifying power of the telescope is respectively
 a) $6.1 \times 10^{-6} \text{ rad}$ and 12 b) $5.0 \times 10^{-6} \text{ rad}$ and 12
 c) $6.1 \times 10^{-6} \text{ rad}$ and 8.3×10^{-2} d) $5.0 \times 10^{-6} \text{ rad}$ and 8.3×10^{-2}
947. A light ray of 5895 \AA wavelength travelling in vacuum enters a medium of refractive index 1.5. The speed of light in the medium is
 a) $3 \times 10^8 \text{ ms}^{-1}$ b) $2 \times 10^8 \text{ ms}^{-1}$ c) $1.5 \times 10^8 \text{ ms}^{-1}$ d) $6 \times 10^8 \text{ ms}^{-1}$
948. A ray of light is incident at 60° on one face of a prism which has angle 30° . The angle between the emergent ray and incident ray is 30° . What is the angle between the ray and the face from which its emerg?
 a) 0° b) 30° c) 60° d) 90°
949. A convex lens is immersed in a liquid, whose refractive index is equal to the refractive index of the material of the lens. Then its focal length will
 a) Decrease b) Become zero c) Become infinite d) Increase
950. Dispersive power depends upon
 a) The shape of prism b) Material of prism c) Angle of prism d) Height of the prism
951. When a plane electromagnetic wave enters a glass slab, then which of the following will not change?

- a) Wavelength b) Frequency c) Speed d) Amplitude

952. An object moving at a speed of 5 m/s towards a concave mirror of focal length $f = 1$ m is at a distance of 9 m. The average speed of the image is

- a) $\frac{1}{5}$ m/s b) $\frac{1}{10}$ m/s c) $\frac{5}{9}$ m/s d) $\frac{4}{10}$ m/s

953. The graph between the lateral magnification (m) produced by a lens and the distance of the image (v) is given by



954. The size of the image of an object, which is at infinity, as formed by a convex lens of focal length 30 cm is 2 cm. If a concave lens of focal length 20 cm is placed between the convex lens and the image at a distance of 26 cm from the convex lens, calculate the new size of the image

- a) 1.25 cm b) 2.5 cm c) 1.05 cm d) 2 cm


955. The focal length (f) of a spherical (concave or convex) mirror of radius of curvature R is

- a) $\frac{R}{2}$ b) R c) $\left(\frac{3}{2}\right)R$ d) $2R$

956. In a compound microscope the objective of f_o and eyepiece of f_e are placed at distance L such that L equals

- a) $f_o + f_e$ b) $f_o - f_e$
c) Much greater than f_o or f_e d) Much less than f_o or f_e

957. A container is filled with water ($\mu = 1.33$) up to a height of 33.25 cm. A concave mirror is placed 15 cm above the water level and the image of an object placed at the bottom is formed 25 cm below the water level. The focal length of the mirror is



- a) 10 cm b) 15 cm c) 20 cm d) 25 cm

958. A ray of light falls on a transparent glass slab of refractive index 1.62. If the reflected ray and the refracted ray are mutually perpendicular, the angle of incidence is

- a) $\tan^{-1}(1.62)$ b) $\tan^{-1}\left(\frac{1}{1.62}\right)$ c) $\tan^{-1}(1.33)$ d) $\tan^{-1}\left(\frac{1}{1.33}\right)$

959. Monochromatic light of wavelength 589 nm is incident from air on a water surface. The refractive index of water is 1.33. The wavelength of the refracted light is

- a) 589 nm b) 443 nm c) 333 nm d) 221 nm

960. A ray of light travels from an optically denser to rarer medium. The critical angle for the two media is C . The maximum possible deviation of the ray will be

- a) $\left(\frac{\pi}{2} - C\right)$ b) $2C$ c) $\pi - 2C$ d) $\pi - C$

961. The refractive index of a certain glass is 1.5 for light whose wavelength in vacuum is 6000 Å. The wavelength of this light when it passes through glass is

- a) 4000 Å b) 6000 Å c) 9000 Å d) 15000 Å

962. The length of the tube of a microscope is 10 cm. The focal lengths of the objective and eye lenses are 0.5 cm and 1.0 cm. The magnifying power of the microscope is about


a) 5

b) 23


c) 166

d) 500


963. Identify the wrong description of the below figures




1.



2.



3.



4.

a) 1 represents far-sightedness

b) 2 correction for short-sightedness

c) 3 represents far-sightedness


d) 4 correction for far-sightedness

964. Light is incident from a medium X at an angle of incidence i and is refracted into a medium Y at angle of refraction r . The graph $\sin i$ versus $\sin r$ is shown in figure. Which of the following conclusions would fit the situation?

1. Speed of light in medium Y is $\sqrt{3}$ times that in medium X

2. Speed of light in medium Y is $1/\sqrt{3}$ times that in medium X

3. Total internal reflection will occur above a certain i value



a) 2 and 3

b) 1 and 3

c) 2 only

d) 3 only

965. A diver in a swimming pool wants to signal his distress to a person lying on the edge of the pool by flashing his water proof flash light

a) He must direct the beam vertically upwards

b) He has to direct the beam horizontally

c) He has to direct the beam at an angle to the vertical which is slightly less than the critical angle of incidence for total internal reflection

d) He has to direction the beam at an angle to the vertical which is slightly more than the critical angle of incidence for the total internal reflection

966. For the myopic eye, the defect cured by


a) Convex lens

b) Concave lens

c) Cylindrical lens

d) Toric lens

967. Figure given below shows a beam of light converging at point P . When a concave lens of focal length 16 cm is introduced in the path of the beam at a place O shown by dotted line such that OP becomes the axis of the lens, the beam converges at a distance x from the lens. The value x will be value to



- a) 12 cm b) 24 cm c) 36 cm d) 48 cm

968. The aperture of the objective lens of a telescope is made large so as to

- a) Increase the resolving power of the telescope
b) Increase the magnifying power of the telescope
c) To focus on distant objects
d) Make image aberrationless

969. The plano-convex lens of focal length 20 cm and 30 cm are placed together to form a double convex lens. The final focal length will be

- a) 12 cm b) 60 cm c) 20 cm d) 30 cm

970. A fish in water (refractive index n) looks at a bird vertically above in the air. If y is the height of the bird and x is the depth of the fish from the surface, then the distance of the bird as estimated by the fish is

- a) $x + y \left(1 - \frac{1}{n}\right)$ b) $x + ny$ c) $x + y \left(1 + \frac{1}{n}\right)$ d) $y + x \left(1 - \frac{1}{n}\right)$

971. Four lenses are made from the same type of glass. The radius of curvature of each face is given below. What will have the greatest positive power

- a) 10 cm convex and 15 cm concave b) 5 cm convex and 10 cm concave
c) 15 cm convex and plane d) 20 cm convex and 30 cm concave

972. Which source is associated with a line emission spectrum

- a) Electric fire b) Neon street sign c) Red traffic light d) Sun

973. A lens forms a virtual image 4 cm away from it when an object is placed 10 cm away from it. The lens is a... lens of focal length..

- a) Concave, 6.67 cm b) Concave, 2.86 cm
c) Convex, 2.86 cm d) May be concave or convex, 6.67 cm

974. Total flux produced by a source of 1 cd is

- a) $1/4\pi$ b) 8π c) 4π d) $1/8\pi$

975. A person using a lens as a simple microscope sees an

- a) Inverted virtual image b) Inverted real magnified image
c) Upright virtual image d) Upright real magnified image

976. Myopia is due to

- a) Elongation of eye ball b) Irregular change in focal length
c) Shortening of eye ball d) Older age

977. Absolute refractive indices of glass and water are $\frac{3}{2}$ and $\frac{4}{3}$. The ratio of velocity of light in glass and water will be

- a) 4 : 3 b) 8 : 7 c) 8 : 9 d) 3 : 4

978. If ${}_i\mu_j$ represents refractive index when a light ray goes from medium i to medium j , then the product ${}_2\mu_1 \times {}_3\mu_2 \times {}_4\mu_3$ is equal to

- a) ${}_3\mu_1$ b) ${}_3\mu_2$ c) $\frac{1}{{}_1\mu_4}$ d) ${}_4\mu_2$

979. Which of the following is not a correct statement

- a) The wavelength of red light is greater than the wavelength of green light
b) The wavelength of blue light is smaller than the wavelength of orange light
c) The frequency of green light is greater than the frequency of blue light
d) The frequency of violet light is greater than the frequency of blue light

980. A ray of light travelling from glass to air (refractive index of glass=1.5). The angle of incidence is 50° . The deviation of the ray is


- PHONE NO: 8583042324 Page | 87

- c) Polarization d) Total internal reflection
993. The focal length of the objective and eye-piece of a telescope are respectively 100 cm and 2 cm. The moon subtends an angle of 0.5° at the eye. If it is looked through the telescope, the angle subtended by the moon's image will be
 a) 100° b) 50° c) 25° d) 10°
994. Focal length of objective and eyepiece of telescope are 200 cm and 4 cm respectively. What is length of telescope for normal adjustment?
 a) 196 cm b) 204 cm c) 250 cm d) 225 cm
995. 'Mirage' is a phenomenon due to
 a) Reflection of light b) Refraction of light
 c) Total internal reflection of light d) Diffraction of light
996. P is a point on the axis of a convex mirror. The image of P formed by the mirror, coincides with P . A rectangular glass slab of thickness t and refractive index μ is now introduced between P and the mirror. For the image of P to coincide with P again, the mirror must be moves
 a) Towards P by $(\mu - 1)t$ b) Away from P by $(\mu - 1)t$
 c) Towards P by $t\left(1 - \frac{1}{\mu}\right)$ d) Away from P by $t\left(1 - \frac{1}{\mu}\right)$
997. An object placed 10 cm in front of a lens has an image 20 cm behind the lens. What is the power of the lens (in dioptres)
 a) 1.5 b) 3.0 c) -15.0 d) $+15.0$
998. The light takes in travelling a distance of 500 m in water. Given that μ for water is $\frac{4}{3}$ and the velocity of light in vacuum is $3 \times 10^{10} \text{ cms}^{-1}$. Calculate equivalent optical path
 a) 566.64 m b) 666.64 m c) 586.45 m d) 576.64 m
999. Two similar plano-convex lenses are combined together in three different ways as shown in the adjoining figure. The ratio of the focal lengths in three cases will be



- a) 2 : 2 : 1 b) 1 : 1 : 1 c) 1 : 2 : 2 d) 2 : 1 : 1


- 100 An equiconvex lens is cut into two halves along (i) XOX' and (ii) YOY' as shown in the figure. Let f, f', f'' be the focal lengths of the complete lens, of each half in case (i), and of each half in case (ii), respectively.




Choose the correct statement from the following

- a) $f' = 2f, f'' = f$ b) $f' = f, f'' = f$ c) $f' = 2f, f'' = 2f$ d) $f' = f, f'' = 2f$
- 100 A biconvex lens of focal length 15 cm is in front of a plane mirror. The distance between the lens and the mirror is 10 cm. A small object is kept at a distance of 30 cm from the lens. The final image is
 a) Virtual and at a distance of 16 cm from the mirror
 b) Real and at a distance of 16 cm from the mirror
 c) Virtual and at a distance of 20 cm from the mirror
 d) None of the above
- 100 A lamp rated at 100 cd hangs over the middle of a round table with diameter 3 m at a height of 2 m. It is replaced by a lamp of 25 cd and the distance to the table is changed so that the illumination at the centre of the table remains as before. The illumination at edge of the table becomes X times the original. Then X is

- a) $1/3$ b) $16/27$ c) $1/4$ d) $1/9$
- 100 A plano-convex lens has a thickness of 4 cm. When placed on a horizontal table, with the curved surface in contact with it, the apparent depth of the bottom most point of the lens is found to be 3 cm. If the lens is inverted such that the plane face is in contact with the table, the apparent depth of the centre of the plane face is found to be $25/8$ cm. Find the focal length of the lens. Assume thickness to be negligible
- a) 85 cm b) 59 cm c) 75 cm d) 7.5 cm
- 100 The focal lengths for violet, green and red light rays are f_V, f_G and f_R respectively. Which of the following is the true relationship
- a) $f_R < f_G < f_V$ b) $f_V < f_G < f_R$ c) $f_G < f_R < f_V$ d) $f_G < f_V < f_R$
- 100 If I_1 and I_2 be the size of the images respectively for the two positions of lens in the displacement method, then the size of the object is given by
- a) I_1/I_2 b) $I_1 \times I_2$ c) $\sqrt{I_1 \times I_2}$ d) $\sqrt{I_1/I_2}$
- 100 Resolving power of a microscope depends upon
- a) Wavelength of light used, directly b) Wavelength of light used, inversely
- c) Frequency of light used d) Focal length of objective
- 100 Three glass prisms A, B and C of same refractive index are placed in contact with each other as shown in figure, with no air gap between the prisms. Monochromatic ray of light OP passes through the prism assembly and emerges as QR . The conditions of minimum deviation is satisfied in the prisms



- a) A and C b) B and C
- c) A and B d) In all prisms A, B and C
- 100 Figure shows a mixture of blue, green and red coloured rays incident normally on a right angled prism. The critical angles of the material of the prism for red, green and blue are $46^\circ, 44^\circ$ and 43° respectively. The arrangement will separate




- a) Red colour from blue and green b) Blue colour from red and green
- c) Green colour from red and blue d) All the three colours
- 100 In Gallilean telescope, if the powers of an objective and eye lens are respectively $+1.25 D$ and $-20 D$, then for relaxed vision, the length and magnification will be
- a) 21.25 cm and 16 b) 75 cm and 20 c) 75 cm and 16 d) 8.5 cm and 21.25
- 101 To remove myopia (short sightedness) a lens of power $0.66 D$ is required. The distance point of the eye is approximately
- a) 100 cm b) 150 cm c) 50 cm d) 25 cm
- 101 An astronomical telescope has a large aperture to
- a) Reduce spherical aberration b) Have high resolution
- c) Increase span of observation d) Have low dispersion

101 An object is immersed in a fluid. In order that the object becomes invisible, it should

- Behave as a perfect reflector
- Absorb all light falling on it
- Have refractive index one
- Have refractive index exactly matching with that of the surrounding fluid


101 A ray of light travels from a medium of refractive index μ to air. Its angle of incidence in the medium is i , measured from the normal to the boundary, and its angle of deviation is δ . δ is plotted against i which of the following best represents the resulting curve



101 A hypermetropic person has to use a lens of power +5 D to normalize his vision. The near point of the hypermetropic eye is

- 1 m
- 1.5 m
- 0.5 m
- 0.66 m

101 A light source is located at P_1 as shown in the figure. All sides of the polygon are equal. The intensity of illumination at P_2 is I_0 . What will be the intensity of illumination at P_3



- $\frac{3\sqrt{3}}{8} I_0$
- $\frac{I_0}{8}$
- $\frac{3}{8} I_0$
- $\frac{\sqrt{3}}{8} I_0$


101 Correct exposure for a photographic print is 10 seconds at a distance of one metre from a point source of 20 candela. For an equal fogging of the print placed at a distance of 2 m from a 16 candela source, the necessary time for exposure is

- 100 s
- 25 s
- 50 s
- 75 s

101 A spherical mirror forms an image of magnification 3. The object distance, if focal length of mirror is 24 cm, may be

- 32 cm, 24 cm
- 32 cm, 16 cm
- 32 cm only
- 16 cm only

101 Which of the following ray diagram show physically possible refraction



- (i)
- (ii)
- (iii)
- None of these

101 When light travels from glass to air, the incident angle is θ_1 and the refracted angle is θ_2 . The true relation is


- $\theta_1 = \theta_2$
- $\theta_1 < \theta_2$
- $\theta_1 > \theta_2$
- Not predictable

102 To increase both the resolving power and magnifying power of a telescope

- Both the focal length and aperture of the objective has to be increased
- To focal length of the objective has to be increased

- c) The aperture of the objective has to be increased
d) The wavelength of light has to be decreased
- 102 A focal length of a thin biconvex lens is 20 cm. When an object is moved from a distance of 25cm in front of it to 50cm, the magnification of its image changes from m_{25} to m_{50} . The ratio $\frac{m_{25}}{m_{50}}$ is
1. a) 6 b) 7 c) 8 d) 9
- 102 A beam of parallel rays is brought to focus by a plano-convex lens. A then concave lens of the same focal length is joined to the first lens. The effect of this is
2. a) The focus shifts to infinity
b) The focal point shifts towards the lens by a small distance
c) The focal point shifts away from the lens by a small distance
d) The focus remains undisturbed
- 102 The critical angle of a medium with respect to air is 45° . The refractive index of that medium will be
3. a) 1.72 b) 1.414 c) 2.12 d) 1.5
- 102 For a given lens, the magnification was found to be twice as large as when the object was 0.15 m distant from it as when the distance was 0.2 m. The focal length of the lens is
4. a) 1.5 m b) 0.20 m c) 0.10 m d) 0.05 m
- 102 Optical fibres are related with
5. a) Communication b) Light c) Computer d) None of these
- 102 When a convergent beam of light is incident on a plane mirror, the image formed is
6. a) upright and real b) upright and virtual
c) inverted and virtual d) inverted and real
- 102 An astronomical telescope has objective and eye-piece lenses of powers 0.5 D and 20 D respectively. What will be its magnifying power?
7. a) 30 b) 10 c) 40 d) 20
- 102 A person's near point is 50 cm and his far point is 3 m. Power of the lenses he requires for
8. (i) reading and
(ii) for seeing distant stars are
- a) $-2 D$ and $0.33 D$ b) $2 D$ and $-0.33 D$ c) $-2 D$ and $3 D$ d) $2 D$ and $-3 D$
- 102 A convex mirror of radius of curvature 1.6 m has an object placed at a distance of 1 m from it. The image is
9. formed at a distance of
- a) $8/13$ m in front of the mirror b) $8/13$ m behind the mirror
c) $4/9$ m in front of the mirror d) $4/9$ m behind the mirror
- 103 A spherical surface of radius of curvature R separates air (refractive index 1.0) from glass (refractive index 1.5). The centre of curvature is in the glass. A point object P placed in air is found to have a real image Q in the glass. The line PQ cuts the surface at a point O , and $PO = OQ$. The distance PO is equal to
0. a) $5 R$ b) $3 R$ c) $2 R$ d) $1.5 R$
- 103 A point source of light is placed 4 m below the surface of water of refractive index $5/3$. The minimum diameter of a disc, which should be placed over the source, on the surface of water to cut-off all light coming out of water
1. a) Infinite b) 6 m c) 4 m d) 3m
- 103 Relative difference of focal lengths of objective and eye lens in the microscope and telescope is given as
2. a) It is equal in both b) It is more in telescope
c) It is more in microscope d) It may be more in any one
- 103 A person is suffering from the defect astigmatism. Its main reason is
- 3.

- a) Distance of the eye lens from retina is increased
 b) Distance of the eye lens from retina is decreased
 c) The cornea is not spherical
 d) Power of accommodation of the eye is decreased
- 103 An infinitely long rod lies along the axis of concave mirror of focal length f . The near end of the rod is at a distance $x > f$ from the mirror. Then the length of the image of the rod is
- a) $\frac{f^2}{x+f}$ b) $\frac{f^2}{x}$ c) $\frac{xf}{x-f}$ d) $\frac{f^2}{x-f}$
- 103 The magnification of the image when an object is placed at a distance x from the principle focus of a mirror of focal length f is
- a) $\frac{x}{f}$ b) $1 + \frac{f}{x}$ c) $\frac{f}{x}$ d) $1 - \frac{f}{x}$
- 103 An object 2.4 m in front of a lens forms a sharp image on a film 12 cm behind the lens. A glass plate 1 cm thick, of refractive index 1.50 is interposed between lens and film with its plane faces parallel to film. At what distance (from lens) should object shifted to be in sharp focus on film
- a) 7.2 m b) 2.4 m c) 3.2 m d) 5.6 m
- 103 For a prism of refractive index 1.732 , the angle of minimum deviation is equal to the angle of prism. Then the angle of the prism is
- a) 50° b) 60° c) 70° d) None of these
- 103 The plane faces of two identical plano-convex lenses each having a focal length of 50 cm are placed against each other to form a usual biconvex lens. The distance from this lens combination at which an object must be placed to obtain a real, inverted image which has the same size as the object is
- a) 50 cm b) 25 cm c) 100 cm d) 40 cm
- 103 In absorption spectrum of Na the missing wavelength (λ) are
- a) 589 nm b) 589.6 nm c) Both d) None of these
- 104 In fog, photographs of the objects taken with infrared radiations are more clear than those obtained during visible light because
- a) $I - R$ radiation has lesser wavelength than visible radiation
 b) Scattering of $I - R$ light is more than visible light
 c) The intensity of $I - R$ light from the object is less
 d) Scattering of $I - R$ light is less than visible light
- 104 If the red light is replaced by blue light illuminating object in a microscope the resolving power of the microscope
- a) Decreases b) Increases c) Gets halved d) Remains unchanged
- 104 A fish at a depth of 12 cm in water is viewed by an observer on the bank of a lake. To what height the image of the fish is raised?
 (Refractive index of water = $4/3$)
- a) 9 cm b) 12 cm c) 3.8 cm d) 3 cm
- 104 Focal length of a convex lens will be maximum for
- a) Blue light b) Yellow light c) Green light d) Red light
- 104 The relation between n_1 and n_2 if the behavior of light ray is as shown in the figure




- a) $n_2 > n_1$ b) $n_1 \gg n_2$ c) $n_1 > n_2$ d) $n_1 = n_2$

- 104 If ϵ_0 and μ_0 are respectively, the electric permittivity and the magnetic permeability of free space, ϵ and μ
5. the corresponding quantities in a medium, the refractive index of the medium is

a) $\sqrt{\frac{\mu\epsilon}{\mu_0\epsilon_0}}$ b) $\frac{\mu\epsilon}{\mu_0\epsilon_0}$ c) $\sqrt{\frac{\mu_0\epsilon_0}{\mu\epsilon}}$ d) $\sqrt{\frac{\mu\mu_0}{\epsilon\epsilon_0}}$

- 104 A ray of light incident normally on an isosceles right angled prism travels as shown in the figure. The least
6. value of the refractive index of the prism must be



- a) $\sqrt{2}$ b) $\sqrt{3}$ c) 1.5 d) 2.0

- 104 Two beams of red and violet colours are made to pass separately through a prism of $A = 60^\circ$. In the
7. minimum deviation position, the angle of refraction inside the prism will be

- a) Greater for red colour b) Equal but not 30° for both the colours
c) Greater for violet colour d) 30° for both the colours

- 104 A defective eye cannot see close objects clearly because their image is formed
8.

- a) On the eye lens b) Between eye lens and retina
c) On the retina d) Beyond retina

- 104 The critical angle between an equilateral prism and air is 45° . If the incident ray is perpendicular to the
9. refractive surface, then

- a) After deviation it will emerge from the second refracting surface
b) It is totally reflected on the second surface and emerges out perpendicularly from third surface in air
c) It is totally reflected from the second and third refracting surfaces and finally emerges out from the first surface
d) It is totally reflected from all the three sides of prism and never emerges out

- 105 An object is placed asymmetrically between two plane mirrors inclined at an angle of 72° . The number of
0. images formed is

- a) 5 b) 4 c) 2 d) Infinite

- 105 A monochromatic light is passed through a prism.....colours shows minimum deviation
1.

- a) Red b) Violet c) Yellow d) Green