

GPLUS EDUCATION

Date :
Time :
Marks :

CHEMISTRY


EQUILIBRIUM

Single Correct Answer Type

- Which may be added to one litre of water to act as a buffer?
 - One mole of $\text{HC}_2\text{H}_3\text{O}_2$ and one mole of HCl
 - One mole of NH_4OH and one mole of NaOH
 - One mole of NH_4Cl and one mole of HCl
 - One mole of $\text{HC}_2\text{H}_3\text{O}_2$ and 0.5 mole of NaOH
- An aqueous solution of 1 M NaCl and 1 M HCl is
 - not a buffer but $\text{pH} < 7$
 - not a buffer but $\text{pH} > 7$
 - a buffer with $\text{pH} < 7$
 - a buffer with $\text{pH} > 7$
- In the following reversible reaction,
 $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3 + Q \text{ cal}$
Most suitable condition for the higher production of SO_3 is
 - Low temperature and high pressure
 - Low temperature and low pressure
 - High temperature and high pressure
 - High temperature and low pressure
- Select the $\text{p}K_a$ value of the strongest acid from the following
 - 1.0
 - 3.0
 - 2.0
 - 4.5
- The pH of a 0.1 M solution of NH_4OH (having $K_b = 1.0 \times 10^{-5}$) is equal to
 - 10
 - 6
 - 11
 - 12
- In the reaction, $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons 2\text{HCl}(\text{g})$
 - $K_p \neq K_c$
 - $K_p = K_c$
 - $K_p > K_c$
 - $K_p < K_c$
- The total number of different kind of buffers obtained during the titration of H_3PO_4 with NaOH are:
 - 3
 - 1
 - 2
 - Zero
- Which will not affect the degree of ionisation?
 - Temperature
 - Concentration
 - Type of solvent
 - Current
- Which of the following has highest pH ?
 - $\frac{\text{M}}{4} \text{KOH}$
 - $\frac{\text{M}}{4} \text{NaOH}$
 - $\frac{\text{M}}{4} \text{NH}_4\text{OH}$
 - $\frac{\text{M}}{4} \text{Ca}(\text{OH})_2$
- Solubility product constant [K_{sp}] of salts of types MX , MX_2 and M_3X at temperature ' T ' are 4.0×10^{-8} , 3.2×10^{-14} and 2.7×10^{-15} respectively. Solubilities (mol, dm^{-3}) of the salts at temperature ' T ' are in the order
 - $\text{MX} > \text{MX}_2 > \text{M}_3\text{X}$
 - $\text{M}_3\text{X} > \text{MX}_2 > \text{MX}$
 - $\text{MX}_2 > \text{M}_3\text{X} > \text{MX}$
 - $\text{MX} > \text{M}_3\text{X} > \text{MX}_2$
- Which of the following base is weakest?
 - NH_4OH ; $K_b = 1.6 \times 10^{-6}$
 - $\text{C}_6\text{H}_5\text{NH}_2$; $K_b = 3.8 \times 10^{-10}$
 - $\text{C}_2\text{H}_5\text{NH}_2$; $K_b = 5.6 \times 10^{-4}$
 - $\text{C}_9\text{H}_7\text{N}$; $K_b = 6.3 \times 10^{-10}$
- One litre of water contains 10^{-7} mole H^+ ions. Degree of ionisation of water is:
 - $1.8 \times 10^{-7}\%$
 - $1.8 \times 10^{-9}\%$
 - $3.6 \times 10^{-7}\%$
 - $3.6 \times 10^{-9}\%$
- A precipitate is formed when
 - The ionic product is nearly equal to the solubility product
 - A solution becomes saturated
 - The ionic product exceeds the solubility product
 - The ionic product is less than solubility product

14. The precipitation is noticed when an aqueous solution of HCl is added to an aqueous solution of:
- a) NaNO_2 b) $\text{Ba}(\text{NO}_3)_2$ c) ZnSO_4 d) HgNO_3
15. Which of the following is not a Lewis base?
- a) NH_3 b) H_2O c) AlCl_3 d) None of these
16. Solubility of BaF_2 in a solution of $\text{Ba}(\text{NO}_3)_2$ will be represented by the concentration term
- a) $[\text{Ba}^{2+}]$ b) $[\text{F}^-]$ c) $\frac{1}{2}[\text{F}^-]$ d) $2[\text{NO}_3^-]$
17. Which of the following is a buffer?
- a) $\text{NaOH} + \text{CH}_3\text{COOH}$ b) $\text{NaOH} + \text{Na}_2\text{SO}_4$ c) $\text{K}_2\text{SO}_4 + \text{H}_2\text{SO}_4$ d) $\text{NH}_4\text{OH} + \text{NaOH}$
18. For the following three reactions I, II and III, equilibrium constants are given
- I. $\text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}_2(\text{g}) + \text{H}_2(\text{g}); K_1$
 II. $\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + 3\text{H}_2(\text{g}); K_2$
 III. $\text{CH}_4(\text{g}) + 2\text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}_2(\text{g}) + 4\text{H}_2(\text{g}); K_3$
- Which of the following relations is correct?
- a) $K_1\sqrt{K_2} = K_3$ b) $K_2K_3 = K_1$ c) $K_3 = K_1K_2$ d) $K_3K_2^3 = K_1^2$
19. 0.1 mole of $\text{N}_2\text{O}_4(\text{g})$ was sealed in a tube under one atmospheric conditions at 25°C . Calculate the number of moles of $\text{NO}_2(\text{g})$ present, if the equilibrium $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g}) (K_p = 0.14)$ is reached after some time
- a) 0.036 b) 36.00 c) 360.0 d) 3.600
20. A buffer solution is prepared by mixing 0.1 M ammonia and 1.0 M ammonium chloride. At 298 K, the $\text{p}K_b$ of NH_4OH is 5.0. The pH of the buffer is
- a) 10.0 b) 9.0 c) 6.0 d) 8.0
21. Which of the following molecules acts as a Lewis acid?
- a) $(\text{CH}_3)_3\text{N}$ b) $(\text{CH}_3)_3\text{B}$ c) $(\text{CH}_3)_2\text{O}$ d) $(\text{CH}_3)_3\text{P}$
22. Which among the following is an electron deficient compound?
- a) NF_3 b) PF_3 c) BF_3 d) AsF_3
23. Identify the correct order of acidic strength of $\text{CO}_2, \text{CuO}, \text{CaO}, \text{H}_2\text{O}$:
- a) $\text{CaO} < \text{CuO} < \text{H}_2\text{O} < \text{CO}_2$
 b) $\text{H}_2\text{O} < \text{CuO} < \text{CaO} < \text{H}_2\text{O}$
 c) $\text{CaO} < \text{H}_2\text{O} < \text{CuO} < \text{CO}_2$
 d) $\text{H}_2\text{O} < \text{CO}_2 < \text{CaO} < \text{CuO}$
24. Which of the following is a strong acid?
- a) HClO_4 b) HBrO_4 c) HIO_4 d) HNO_3
25. According to Arrhenius concept the, strength of an acid depends on:
- a) Hydrolysis
 b) Concentration of acid
 c) H^+ ions furnished by acid
 d) Number of mole of base used for neutralization
26. $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$
 In the above equilibrium system, if the concentration of the reactants at 25°C is increased, the value of K_c will
- a) Increase b) Decrease
 c) Remains the same d) Depends on the nature of the reactants
27. 0.04 g of pure NaOH is dissolved in 10 litre of distilled water. The pH of the solution is:
- a) 9 b) 10 c) 11 d) 12
28. What is the equilibrium expression for the reaction, $\text{P}_4(\text{s}) + 5\text{O}_2(\text{g}) \rightleftharpoons \text{P}_4\text{O}_{10}(\text{s})$?
- a) $K_c = \frac{1}{[\text{O}_2]^5}$ b) $K_c = [\text{O}_2]^5$ c) $K_c = \frac{[\text{P}_4\text{O}_{10}]}{5[\text{P}_4][\text{O}_2]}$ d) $K_c = \frac{[\text{P}_4\text{O}_{10}]}{[\text{P}_4][\text{O}_2]^5}$
29. When 10^{-8} mole of HCl is dissolved in one litre of water, the pH of the solution will be:
- a) 8 b) 7 c) Above 8 d) Below 7

30. A physician wishes to prepare a buffer solution at $\text{pH} = 3.58$ that efficiently resists a change in pH yet contains only small conc. of the buffering agents. Which one of the following weak acid together with its sodium salt would be best to use?
- m -chloro benzoic acid ($\text{p}K_a = 3.98$)
 - p -chlorocinnamic acid ($\text{p}K_a = 4.41$)
 - 2,5-dihydroxy benzoic acid ($\text{p}K_a = 2.97$)
 - Acetoacetic acid ($\text{p}K_a = 3.58$)
31. The pH of 10^{-8}M HCl solution is
- 8
 - More than 8
 - Between 6 and 7
 - Slightly more than 7
32. A certain buffer solution contains equal concentration of X^- and HX . The K_a for HX is 10. The pH of the buffer is:
- 7
 - 8
 - 11
 - 14
33. 100 mL of 0.01 M solution of NaOH is diluted to 1 dm^3 . What is the pH of the diluted solution?
- 12
 - 11
 - 2
 - 3
34. Which of the following salt does not get hydrolysed in water?
- KClO_4
 - NH_4Cl
 - CH_3COONa
 - None of these
35. A higher value for equilibrium constant, K shows that:
- The reaction has gone to near completion towards right
 - The reaction has not yet started
 - The reaction has gone to near completion towards left
 - None of the above
36. Which one is least basic?
- CH_3NH_2
 - NH_3
 - $\text{C}_2\text{H}_5\text{NH}_2$
 - $\text{C}_6\text{H}_5\text{NH}_2$
37. The aqueous solution of disodium hydrogen phosphate is:
- Acidic
 - Neutral
 - Basic
 - None of these
38. 3.2 moles of hydrogen iodide were heated in a sealed bulb at 444°C till the equilibrium state was reached. Its degree of dissociation at this temperature was found to be 22%. The number of moles of hydrogen iodide present at equilibrium are
- 1.876
 - 2.496
 - 3.235
 - 4.126
39. In the reactions, $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$, the amounts of PCl_5 , PCl_3 and Cl_2 at equilibrium are 2 mole each and the total pressure is 3 atm. The equilibrium constant K_p is :
- 1.0 atm
 - 2.0 atm
 - 3.0 atm
 - 6.0 atm
40. Which of the following is correct for the reaction?
- $$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$$
- $K_p = K_c$
 - $K_p < K_c$
 - $K_p > K_c$
 - Pressure is required to predict the correlation
41. The graph relates $\ln K_{eq}$ vs $\frac{1}{T}$ for a reaction. The reaction must be :



- Exothermic

54. Which one is hard base?
 a) Ag^+ b) Cr^{3+} c) I_2 d) F^-
55. Which species acts as an acid and also a conjugate base of another acid?
 a) HSO_4^- b) CO_3^{2-} c) SO_4^{2-} d) H_3O^+
56. Predict the conditions for forward reaction on the basis of Le-Chatelier's principle for : $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$; $\Delta H = -198 \text{ kJ}$.
 a) Lowering the temperature and increasing pressure
 b) Any value of temperature and pressure
 c) Lowering of temperature as well as pressure
 d) Increasing temperature as well as pressure
57. The solubility of AgCl in water at 10°C is $6.2 \times 10^{-6} \text{ mol/litre}$. The K_{sp} of AgCl is:
 a) $[6.2 \times 10^{-6}]^{1/2}$ b) $6.2 \times (10^{-6})^2$ c) $(6.2)^2 \times 10^{-6}$ d) $[6.2 \times 10^{-6}]^2$
58. When pressure is applied to the equilibrium system ice r water. Which of the following phenomenon will happen?
 a) More ice will be formed b) Water will evaporate
 c) More water will be formed d) Equilibrium will not be formed
59. At constant temperature in one litre vessel, when the reaction, $2\text{SO}_3(\text{g}) \rightleftharpoons 2\text{SO}_2(\text{g}) + \text{O}_2(\text{g})$ is at equilibrium, the SO_2 concentration is 0.6 M , initial concentration of SO_3 is 1 M . The equilibrium constant is :
 a) 2.7 b) 1.36 c) 0.34 d) 0.675
60. When 20g of CaCO_3 were put into 10 litre flask and heated to 800°C , 35% of CaCO_3 remained unreacted at equilibrium. K_p for decomposition of CaCO_3 is :
 a) 1.145 atm b) 0.145 atm c) 2.145 atm d) 3.145 atm
61. For the reaction equilibrium,
 $2\text{NOBr}(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}) + \text{Br}_2(\text{g})$, if $P_{\text{Br}_2} = \frac{P}{9}$ at equilibrium and P is total pressure. The ratio K_p/P is equal to:
 a) $1/9$ b) $1/81$ c) $1/27$ d) $1/3$
62. $K_{sp} = 1.2 \times 10^{-5}$ of M_2SO_4 (M^+ is monovalent metal ion) at 298 K. The maximum concentration of M^+ ions that could be attained in a saturated solution of this solid at 298 K is:
 a) $3.46 \times 10^{-3} \text{ M}$ b) $7.0 \times 10^{-3} \text{ M}$ c) $2.88 \times 10^{-2} \text{ M}$ d) $14.4 \times 10^{-3} \text{ M}$
63. Which of the following describes correct sequence for decreasing Lewis acid nature?
 a) $\text{BCl}_3 > \text{BF}_3 > \text{BBr}_3$ b) $\text{BBr}_3 > \text{BCl}_3 > \text{BF}_3$ c) $\text{BBr}_3 > \text{BF}_3 > \text{BCl}_3$ d) $\text{BF}_3 > \text{BCl}_3 > \text{BBr}_3$
64. What should be the pH of solution to dissolve the $\text{Cr}(\text{OH})_3$ precipitate?
 [Given, $[\text{Cr}^{3+}] = 1.0 \text{ mol/L}$, $K_{sp} = 6 \times 10^{-31}$)
 a) 2.0 b) 3.0 c) 5.0 d) 4.0
65. Which one of the following salts on being dissolved in water gives $\text{pH} > 7$ at 25°C ?
 a) KCN b) KNO_3 c) NH_4Cl d) NH_4CN
66. Aqueous solution of which salt has the lowest pH?
 a) NaOH b) NH_4Cl c) Na_2CO_3 d) NaCl
67. In a gaseous reversible reaction,
 $\text{N}_2 + \text{O}_2 \rightleftharpoons 2\text{NO} + \text{heat}$
 If pressure is increased then the equilibrium constant would be
 a) Unchanged
 b) Increased
 c) Decreased
 d) Sometimes increased, sometimes decreased
68. Glycine is:
 a) Arrhenius acid b) Lewis base c) Simplest amino acid d) All of these
69. On a given condition, the equilibrium concentration of HI , H_2 and I_2 are 0.80, 0.10 and 0.10 mol/L. The

82. Which among the following is the strongest acid?
 a) $\text{H}(\text{ClO})\text{O}_2$ b) $\text{H}(\text{ClO})\text{O}_3$ c) $\text{H}(\text{ClO})\text{O}$ d) $\text{H}(\text{ClO})$
83. Which one of the following is not an amphoteric substance?
 a) HNO_3 b) HCO_3^- c) H_2O d) NH_3
84. For the chemical reaction $3\text{X}(\text{g}) + \text{Y}(\text{g}) \rightleftharpoons \text{X}_3\text{Y}(\text{g})$, that amount of X_3Y at equilibrium is affected by
 a) Temperature and pressure b) Temperature only
 c) Pressure only d) Temperature, pressure and catalyst
85. K_p/K_c for the reaction,
 $\text{CO}(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightleftharpoons \text{CO}_2(\text{g})$ is:
 a) RT b) $1/\sqrt{RT}$ c) \sqrt{RT} d) 1
86. Densities of diamond and graphite are 3.5 and 2.3 g/mL respectively. Increase of pressure on the equilibrium $\text{C}_{\text{diamond}} \rightleftharpoons \text{C}_{\text{graphite}}$:
 a) Favours backward reaction
 b) Favours forward reaction
 c) Have no effect
 d) Increases the reaction rate
87. The solubility product of BaCl_2 is 4×10^{-9} . Its solubility in mol/L is
 a) 4×10^{-3} b) 4×10^{-9} c) 1×10^{-3} d) 1×10^{-9}
88. Addition of sodium acetate to 0.1 M acetic acid will cause
 a) Increase in pH b) Decrease in pH
 c) No change in pH d) Change in pH that cannot be predicted
89. The solubility in water of a sparingly soluble salt A_2B is $1.0 \times 10^{-3} \text{ mol L}^{-1}$. Its solubility product will be
 a) 4×10^{-9} b) 4×10^9 c) 1×10^9 d) 1×10^{-9}
90. NaHCO_3 and NaOH can not co-exist in a solution because of:
 a) Common ion effect
 b) Acid-base neutralisation
 c) Le – Chatelier’s principle
 d) Redox change
91. Formation of SO_3 from SO_2 and O_2 is favoured by
 a) Increase in pressure b) Decrease in pressure
 c) Increase in temperature d) Decrease in temperature
92. A definite amount of solid NH_4HS is placed in a flask already containing NH_3 gas at certain temperature and 0.50 atm pressure. NH_4HS decomposes to give NH_3 and H_2S and total equilibrium pressure in flask is 0.84 atm. The equilibrium constant for the reaction is :
 a) 0.30 b) 0.18 c) 0.17 d) 0.11
93. Hydroxyl ion concentration of 10^{-2} M HCl is
 a) $1 \times 10^1 \text{ mol dm}^{-3}$ b) $1 \times 10^{-12} \text{ mol dm}^{-3}$ c) $1 \times 10^{-1} \text{ mol dm}^{-3}$ d) $1 \times 10^{-14} \text{ mol dm}^{-3}$
94. For a reaction in equilibrium :
 a) There is no volume change
 b) The reaction has stopped completely
 c) The rate of forward reaction is equal to the rate of backward reaction
 d) The forward reaction is faster than reverse reaction
95. A solution of CuSO_4 in water will:
 a) Turn red litmus blue
 b) Turn blue litmus red
 c) Show no effect on litmus

110. Ammonia under a pressure of 15 atm at 27°C is heated to 347°C in a closed vessel in the presence of catalyst. Under the conditions, NH_3 is partially decomposed according to the equation, $2\text{NH}_3 \rightleftharpoons \text{N}_2 + 3\text{H}_2$. The vessel is such that the volume remains effectively constant whereas pressure increases to 50 atm. Calculate the percentage of NH_3 actually decomposed
- a) 61.3% b) 63.5% c) 65.3% d) 66.6%
111. For the system; $3A + 2B \rightleftharpoons C$, the expression for equilibrium constant is
- a) $\frac{[A]^3[B]^2}{[C]}$ b) $\frac{[C]}{[A]^3[B]^2}$ c) $\frac{[3A][2B]}{[C]}$ d) $\frac{[C]}{[3A][2B]}$
112. A monoprotic acid in a 0.1 M solution ionises to 0.001%. Its ionisation constant is
- a) 1×10^{-11} b) 1×10^{-3} c) 1×10^{-6} d) 1×10^{-8}
113. For the reaction, $\text{C}(s) + \text{CO}_2(g) \rightleftharpoons 2\text{CO}(g)$, the principle pressure of CO_2 and CO are 2.0 and 4.0 atm respectively at equilibrium. The K_p for the reaction is
- a) 2.0 b) 4.0 c) 8.0 d) 1.6
114. The vapour density of completely dissociated NH_4Cl would be:
- a) Slightly less than half of that of ammonium chloride
 b) Half of that of ammonium chloride
 c) Double that of ammonium chloride
 d) Determined by the amount of solid ammonium chloride used in the experiment
115. Mg^{2+} is ... than Al^{3+} .
- a) Strong Lewis acid b) Strong Lewis base c) Weak Lewis acid d) Weak Lewis base
116. The equilibrium constant for the reaction, $\text{N}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}(g)$ is 4×10^{-4} at 2000 K. In presence of a catalyst the equilibrium is attained ten times faster. Therefore, the equilibrium constant, in present of the catalyst, at 2000 K is:
- a) 40×10^{-4}
 b) 4×10^{-4}
 c) 4×10^{-3}
 d) Difficult to compute without more data
117. The activation energies of forward and backward reaction: $A_2 + B_2 \rightleftharpoons 2AB$ are 180 kJ mol^{-1} and 200 kJ mol^{-1} respectively. The presence of a catalyst lowers the activation energy of both (forward and backward) reactions by 100 kJ mol^{-1} . The enthalpy change of the reaction in the presence of catalyst will be (in kJ mol^{-1}):
- a) -20
 b) -300
 c) +120
 d) -280
118. How will increase of pressure affect the equation?
 $\text{C}(s) + \text{H}_2\text{O}(g) \rightleftharpoons \text{CO}(g) + \text{H}_2(g)$
- a) Shift in the forward direction b) Shift in the reverse direction
 c) Increase in the yield of hydrogen d) No effect
119. If the pressure of N_2/H_2 mixture in a closed apparatus is 100 atm and 20% of the mixture then reacts, the pressure at the same temperature would be :
- a) 100 b) 90 c) 85 d) 80
120. $\text{C}_2\text{H}_5\text{ONa}$ acts as ... in $\text{C}_2\text{H}_5\text{OH}$.
- a) Strong acid b) Weak acid c) Strong base d) Weak base
121. A solution of sodium borate has a pH of approximately:
- a) > 7 b) < 7 c) = 7 d) Between 4 and 5
122. A certain buffer solution contains equal concentration of X^- and HX . The K_a for HX is 10^{-8} . The pH of the buffer is
- a) 3 b) 8 c) 11 d) 14

123. Study the following table.

Buffer solution	Volume (in mL) of 1 M weak acid	Volume (in mL) of 0.1 M sodium salt of weak acid
I	4.0	4.0
II	4.0	40.0
III	40.0	4.0
IV	0.1	10.0

Which of the two sets of buffer solutions have least pH?

- a) I and II b) I and III c) II and III d) II and IV
124. Which indicator works in the pH range 8-9.8?
 a) Phenolphthalein b) Methyl orange c) Methyl red d) Litmus
125. 100 mL of 0.015 M HCl solution is mixed with 100 mL of 0.005 M HCl. What is the pH of the resultant solution?
 a) 2.5 b) 1.5 c) 2 d) 1
126. The solubility of A_2X_3 is y mol dm^{-3} . Its solubility product is:
 a) $6y^4$ b) $64y^4$ c) $36y^5$ d) $108y^5$
127. The volume of water needed to dissolve 1 g of $BaSO_4$ ($K_{sp} = 1.1 \times 10^{-10}$) at $25^\circ C$ is:
 a) 820 litre
 b) 410 litre
 c) 205 litre
 d) None of these
128. In a vessel containing SO_3 , SO_2 , and O_2 at equilibrium, some helium gas is introduced so that, the total pressure increase, while temperature and volume remain constant. According to Le-Chatelier's principle the dissociation of SO_3 :
 a) Increases
 b) Decreases
 c) Remains unaltered
 d) Changes unpredictably
129. Given the equilibrium system
 $NH_4Cl(s) \rightleftharpoons NH_4^+(aq) + Cl^-(aq)$
 $(\Delta H = +3.5 \text{ kcal/mol})$.
 What change will shift the equilibrium to the right?
 a) Decreasing the temperature
 b) Increasing the temperature
 c) Dissolving NaCl crystals in the equilibrium mixture
 d) Dissolving NH_4NO_3 crystals in the equilibrium mixture
130. The solubility product of $BaSO_4$ is 1.5×10^{-9} . The precipitation in a 0.01 M Ba^{2+} solution will start, on adding H_2SO_4 of concentration
 a) 10^{-9} M b) 10^{-8} M c) 10^{-7} M d) 10^{-6} M
131. The solubility of $Pb(OH)_2$ in water is 6.7×10^{-6} M. Its solubility in a buffer solution of pH=8 would be
 a) 1.2×10^{-2} b) 1.6×10^{-3} c) 1.6×10^{-2} d) 1.2×10^{-3}
132. In which of the following reactions is $K_p < K_c$?
 a) $I_2(g) \rightleftharpoons 2I(g)$ b) $2BrCl(g) \rightleftharpoons Cl_2(g) + Br_2(g)$


- c) $\text{CO(g)} + 3\text{H}_2\text{(g)} \rightleftharpoons \text{CH}_4\text{(g)} + \text{H}_2\text{O(g)}$ d) All of the above
133. Any precipitate is formed when
 a) Solution becomes saturated
 b) The value of ionic product is less than the value of solubility product
 c) The value of ionic product is equal to the value of solubility product
 d) The value of ionic product is greater than the value of solubility product
134. At 25°C , K_b for a base BOH is 1.0×10^{-12} . The $[\text{OH}^-]$ in 0.01M aqueous solution of base is:
 a) $1.0 \times 10^{-6}\text{ M}$ b) $1.0 \times 10^{-7}\text{ M}$ c) $1.0 \times 10^{-5}\text{ M}$ d) $2.0 \times 10^{-6}\text{ M}$
135. The pH of a 10^{-9} M solution of HCl in water is
 a) 8 b) -8 c) Between 7 and 8 d) Between 6 and 7
136. If pH of a saturated solution of Ba(OH)_2 is 12, the value of its K_{sp} is:
 a) $4.0 \times 10^{-6}\text{ M}^3$ b) $4.0 \times 10^{-7}\text{ M}^3$ c) $5.0 \times 10^{-6}\text{ M}^3$ d) $5.0 \times 10^{-7}\text{ M}^3$
137. Liquid ammonia ionises to a slight extent. At -50°C , its self ionisation constant, $K_{\text{NH}_3} = [\text{NH}_4^+][\text{NH}_2^-] = 10^{-30}$. How many amide ions, are present per cm^3 of pure liquid ammonia? (Assume $N = 6.0 \times 10^{23}$)
 a) 6×10^6 ions b) 6×10^5 ions c) 6×10^{-5} ions d) 6×10^{-6} ions
138. The first and second dissociation constants of an acid H_2A are 1.0×10^{-5} and 5.0×10^{-10} respectively. The overall dissociation constant of the acid will be
 a) 5.0×10^{-5} b) 5.0×10^{15} c) 5.0×10^{-15} d) 0.2×10^5
139. Which is the strongest acid?
 a) CH_3COOH b) CH_2ClCOOH c) CHCl_2COOH d) CCl_3COOH
140. A 0.01 M ammonia solution is 5% ionized. The concentration of $[\text{OH}^-]$ ion is:
 a) 0.005 M b) 0.0001 M c) 0.0005 M d) 0.05 M
141. Nucleophiles are:
 a) Lewis acids b) Lewis bases c) Bronsted acids d) Bronsted bases
142. Theory of ionisation was given by
 a) Rutherford b) Graham c) Faraday d) Arrhenius
143. 0.01 mole of lime (CaO) was dissolved in 100 cm^3 of water. Assuming the base is completely ionised in the solution, the pH of the solution will be
 a) 13.3 b) 8.5 c) 6 d) 8
144. Consider the following solutions of equal concentrations
 $A = \text{NH}_4\text{Cl}$ $B = \text{CH}_3\text{COONa}$
 $C = \text{NH}_4\text{OH}$ $D = \text{CH}_3\text{COOH}$
 A buffer solution can be obtained by mixing equal volumes of
 a) C and D b) A and B c) A and C d) C and D
145. At 600°C , K_p for the following reaction is 1 atm.

$$\text{X(g)} \rightleftharpoons \text{Y(g)} + \text{Z(g)}$$
 At equilibrium, 50% of X(g) is dissociated. The total pressure of the equilibrium system is p atm. What is the partial pressure (in atm) of X(g) at equilibrium?
 a) 1 b) 4 c) 2 d) 0.5
146. Equilibrium constants K_1 and K_2 for the following equilibria are related as :

$$\text{NO(g)} + \frac{1}{2}\text{O}_2\text{(g)} \rightleftharpoons \text{NO}_2\text{(g)}; K_1$$

$$2\text{NO}_2\text{(g)} \rightleftharpoons 2\text{NO(g)} + \text{O}_2\text{(g)}; K_2$$
 a) $K_2 = \frac{1}{K_1^2}$ b) $K_2 = \frac{1}{K_1}$ c) $K_2 = K_1^2$ d) $K_2 = \frac{K_1}{2}$
147. If K_1 and K_2 are equilibrium constants for reactions (I) and (II) respectively for,

$$\text{N}_2 + \text{O}_2 \rightleftharpoons 2\text{NO} \quad \dots \text{(i)}$$



Then:

a) $K_2 = K_1$

b) $K_2 = \sqrt{K_1}$

c) $K_1 = 2K_2$


d) $K_1 = \frac{1}{2}K_2$

148. All reactions which have chemical disintegration are

- a) Exothermic
- b) Reversible
- c) Reversible and exothermic
- d) Reversible or irreversible and endothermic or exothermic

149. For which of the following sparingly soluble salt, the solubility (s) and solubility product (K_{sp})


are related by the expression $s = (K_{sp}/4)^{1/3}$?



150. For $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$, initial concentration of each reactant and product is 1 M. If $K_{eq} = 0.41$ then

- a) More PCl_3 will form
- b) More Cl_2 will form
- c) More PCl_5 will form
- d) No change

151. The exothermic formation of ClF_3 is represented by the equation



$$\Delta H = -329 \text{ kJ}$$

Which of the following will increase the quantity of ClF_3 in an equilibrium mixture of Cl_2 , F_2 and ClF_3 ?





a) Adding F_2

b) Increasing the volume of the container

c) Removing Cl_2

d) Increasing the temperature

152. For which of the following reactions, does the equilibrium constant depend on the units of concentration?



153. If the solubility product of lead iodide (PbI_2) is 3.2×10^{-8} , its solubility will be:


a) $2 \times 10^{-3} \text{ M}$

b) $4 \times 10^{-4} \text{ M}$

c) $1.6 \times 10^{-5} \text{ M}$

d) $1.8 \times 10^{-5} \text{ M}$

154. At 30°C the solubility of Ag_2CO_3 ($K_{sp} = 8 \times 10^{-12}$) would be greatest in 1 L of



155. The interfering radicals interfere in the test of usual inorganic analysis after II group analysis due to:

- a) Their solubility in acid medium
- b) Their solubility in alkaline medium
- c) Their insoluble nature in alkaline medium
- d) None of the above

156. The $\text{p}K_b$ value of NH_3 is 5. Calculate the pH of the buffer solution, 1 L of which contains 0.01 M NH_4Cl and 0.10 M NH_4OH

a) 4

b) 6

c) 8

d) 10

157. The equilibrium constant K for the reaction $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$ at room temperature 300 K is 2.85 and at 698 K 1.84×10^{-2} . Hence the reason that HI exists as a stable compound at room temperature is because:

- a) It decomposes so slowly that equilibrium is not readily achieved
- b) The HI bond has a large covalent contribution
- c) The heat of reaction at room temperature is -5.31 kcal
- d) It is uncatalytic reaction

158. A mixture of 0.3 mole of H_2 and 0.3 mole of I_2 is allowed to react in a 10 L evacuated flask at 500°C . The

reaction is $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$, the K is found to be 64. The amount of unreacted I_2 at equilibrium is

- a) 0.03 mol b) 0.06 mol c) 0.09 mol d) 3.6 mol

159. In a solution of a weak electrolyte at infinite dilution we have:

- a) Only cations and electrolyte in 10% dissociated
 b) Only anions and electrolyte is 10% dissociated
 c) Both cations and anions and electrolyte is 100% dissociated
 d) Cations, anions and unionised electrolyte


160. In the reaction, $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$, the equilibrium concentration of PCl_5 and PCl_3 are 0.4 and 0.2 mol/L respectively. If the value of K_c is 0.5 what is the concentration of Cl_2 in mol/L?

- a) 0.5 b) 1.0 c) 1.5 d) 2.0

161. The reaction that proceeds in the forward direction is :

- a) $\text{SnCl}_4 + \text{Hg}_2\text{Cl}_2 \rightarrow \text{SnCl}_2 + 2\text{HgCl}_2$
 b) $\text{NH}_4\text{Cl} + \text{NaOH} \rightarrow \text{H}_2\text{O} + \text{NH}_3 + \text{NaCl}$
 c) $\text{Mn}^{2+} + 2\text{H}_2\text{O} + \text{Cl}_2 \rightarrow \text{MnO}_2 + 4\text{H}^+ + 2\text{Cl}^-$
 d) $\text{S}_4\text{O}_6^{2-} + 2\text{I}^- \rightarrow 2\text{S}_2\text{O}_3^{2-} + \text{I}_2$

162. Which is a Lewis base



- a) I_2 b) I_3^- c) I^- d) None of these

163. A solution contains 10 mL of 0.1 N NaOH and 10 mL of 0.05 N H_2SO_4 , pH of this solution is

- a) Less than 7 b) 7 c) Zero d) Greater than 7

164. The solubility of PbCl_2 in water is 0.01 M at 25°C. Its maximum concentration in 0.1 M NaCl will be:

- a) $2 \times 10^{-3} \text{ M}$ b) $1 \times 10^{-4} \text{ M}$ c) $1.6 \times 10^{-2} \text{ M}$ d) $4 \times 10^{-4} \text{ M}$

165. HX is a weak acid ($K_a = 10^{-5}$). It forms a salt NaX (0.1 M on reacting with caustic soda. The degree of hydrolysis of NaX is

- a) 0.01% b) 0.0001 % c) 0.1 % d) 0.5 %

166. Which species acts as stronger acid than formic acid in aqueous solution?

- a) CH_3COOH b) H_2SO_4 c) NH_4^+ d) HPO_4^{2-}

167. In a reaction at equilibrium 'X' mole of the reactant A decompose to give 1 mole each of C and D. If the fraction of A decomposed at equilibrium is independent of initial concentration of A, then the value of 'X' is :

- a) 1 b) 3 c) 2 d) 4

168. Starting with 1 mole of N_2O_4 , if α is the degree of dissociation of N_2O_4 for the reaction, $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$ then at equilibrium the total number of moles of N_2O_4 and NO_2 present is

- a) 2 b) $(1 - \alpha)$ c) $(1 - \alpha)^2$ d) $(1 + \alpha)$

169. A saturated solution of $\text{Mg}(\text{OH})_2$ in water at 25°C contains 0.11 g $\text{Mg}(\text{OH})_2$ per litre of solution. The solubility product of $\text{Mg}(\text{OH})_2$ is:

- a) $(0.11)^2$ b) $(0.11)^3$ c) $4 \times (0.11)^3$ d) $4 \times (0.11)^3 / (58)^3$

170. For the reaction,



$$(K_c = 1.8 \times 10^{-6} \text{ at } 184^\circ\text{C})$$

$$(R = 0.00831 \text{ kJ}/(\text{mol K}))$$

When K_p and K_c are compared at 184°C, it is found that

- a) Whether K_p is greater than less than or equal to K_c depends upon the total gas pressure b) $K_p = K_c$
 c) K_p is less than K_c d) K_p is greater than K_c

171. Which is the best choice for weak base-strong acid titration?

- a) Methyl red b) Litmus c) Phenol red d) Phenolphthalein

172. The value of the ionic product of water depends

- a) On volume of water
c) Changes by adding acid or alkali
- b) On temperature
d) Always remain constant
173. The formation of SO_3 takes place according to the following reaction, $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3, \Delta H = -45.2 \text{ kcal}$.
The formation of SO_3 is favoured by
a) Increase of volume
b) Increase in pressure
c) Increase in temperature
d) Removal of oxygen
174. Which one is strongest electrolyte in the following?
a) NaCl
b) CH_3COOH
c) NH_4OH
d) $\text{C}_6\text{H}_{12}\text{O}_6$
175. For which of the following reactions, $K_p = K_c$?
a) $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$
b) $\text{N}_2 + \text{O}_2 \rightleftharpoons 2\text{NO}$
c) $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$
d) $2\text{SO}_3 \rightleftharpoons 2\text{SO}_2 + \text{O}_2$
176. The solubility of AgI in NaI solution is less than that in pure water because
a) AgI forms complex with NaI
b) Of common ion effect
c) Solubility product of AgI is less
d) The temperature of the solution decreases
177. The partial pressure of $\text{CH}_3\text{OH}(\text{g})$, $\text{CO}(\text{g})$ and $\text{H}_2(\text{g})$ in equilibrium mixture for the reaction, $\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH}(\text{g})$ are 2.0, 1.0 and 0.1 atm respectively at 427°C . The value of K_p for the decomposition of CH_3OH to CO and H_2 is :
a) 10^2 atm
b) $2 \times 10^2 \text{ atm}^{-1}$
c) 50 atm^2
d) $5 \times 10^{-3} \text{ atm}^2$
178. What happens to pH of a solution when NH_4Cl crystal is added to a dilute solution of NH_4OH ?
a) Decreases
b) Increases
c) Remains unaffected
d) All of these
179. What mole of $\text{Ca}(\text{OH})_2$ is dissolved in 250 mL aqueous solution to give a solution of pH 10.65, assuming full dissociation?
a) 0.47×10^{-4}
b) 0.48×10^{-4}
c) 0.56×10^{-4}
d) 0.58×10^{-4}
180. The volume of the reaction vessel containing an equilibrium mixture in the reaction, $\text{SO}_2\text{Cl}_2(\text{g}) \rightleftharpoons \text{SO}_2(\text{g}) + \text{Cl}_2(\text{g})$ is increased. When equilibrium is reestablished:
a) The amount of $\text{SO}_2(\text{g})$ will decrease
b) The amount of $\text{SO}_2\text{Cl}_2(\text{g})$ will increase
c) The amount of $\text{Cl}_2(\text{g})$ will increase
d) The amount of $\text{Cl}_2(\text{g})$ will remain unchanged
181. The acidic nature of zinc oxide is shown from the formation of salt:
a) NaZnO_2
b) Na_2ZnO_2
c) $\text{Na}_2\text{Zn}_2\text{O}_2$
d) None of these
182. Consider the following reaction equilibrium
$$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$$

Initially, 1 mole of N_2 and 3 moles of H_2 are taken in a 2 L flask. At equilibrium state if, the number of moles of N_2 is 0.6, what is the total number of moles of all gases present in the flask?
a) 0.8
b) 1.6
c) 3.2
d) 6.4
183. If 0.1 mole of I_2 is introduced into 1.0 litre flask at 1000 K, at equilibrium ($K_c = 10^{-6}$), which one is correct?
a) $[\text{I}_2(\text{g})] > [\text{I}(\text{g})]$
b) $[\text{I}_2(\text{g})] < [\text{I}(\text{g})]$
c) $[\text{I}_2(\text{g})] = [\text{I}(\text{g})]$
d) $[\text{I}_2(\text{g})] = \frac{1}{2}[\text{I}(\text{g})]$
184. The equilibrium constant K_c for $A(\text{g}) \rightleftharpoons B(\text{g})$ is 1.1, Gas B will have molar concentration greater than 1 if :
a) $(A) = 0.91$
b) $(A) > 0.91$
c) $(A) > 1$
d) At all these
185. The equilibrium which remains unaffected by change in pressure of the reactants is
a) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$
b) $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$
c) $2\text{O}_3(\text{g}) \rightleftharpoons 3\text{O}_2(\text{g})$
d) $2\text{NO}_2 \rightleftharpoons \text{N}_2\text{O}_4(\text{g})$
186. The pH value of 0.001 M aqueous solution of NaCl is
a) 7
b) 4
c) 11
d) unpredictable
187. The solution of AgCl is unsaturated if:
a) $[\text{Ag}^+][\text{Cl}^-] < K_{sp}$
b) $[\text{Ag}^+][\text{Cl}^-] > K_{sp}$
c) $[\text{Ag}^+][\text{Cl}^-] = K_{sp}$
d) None of these
188. A decimolar solution of ammonium hydroxide is ionised to the extent of 1.3%. If $\log 1.3 = 0.11$,

what is the pH of the solution?

- a) 11.11 b) 9.11 c) 8.11 d) Unpredictable
189. The equivalent conductance of $\frac{M}{32}$ solution of a weak monobasic acid is 8.0 mhos cm^2 and at infinite dilution is 400 mhos cm^2 . The dissociation constant of this acid is:
 a) 1.25×10^{-4} b) 1.25×10^{-5} c) 1.25×10^{-6} d) 6.25×10^{-4}
190. Hydrolysis of oxide ion in water produces:
 a) H^+ b) OH^- c) O_2 d) H_2O
191. A weak acid HX has the dissociation constant 1×10^{-5} M. It forms a salt NaX on reaction with alkali. The degree of hydrolysis of 0.1 M solution of NaX is
 a) 0.0001% b) 0.01% c) 0.1% d) 0.15%
192. The species among the following which can act as an acid and a base is:
 a) HSO_4^- b) SO_4^{2-} c) H_3O^+ d) Cl^-
193. For the reactions,
 $A \rightleftharpoons B; K_c = 2$
 $B \rightleftharpoons C; K_c = 4$
 $C \rightleftharpoons D; K_c = 6$
 K_c for the reaction, $A \rightleftharpoons D$ is:
 a) $(2 + 4 + 6)$ b) $(2 \times 4)/6$ c) $(4 \times 6)/2$ d) $2 \times 4 \times 6$
194. 0.365 g of HCl gas was passed through 100 cm^3 of 0.2 M NaOH solution. The pH of the resulting solution would be
 a) 1 b) 5 c) 8 d) 13
195. The pH of a 0.0001 N solution of KOH will be
 a) 4 b) 6 c) 10 d) 12
196. The equilibrium constant for a reaction is 1×10^{20} at 300 K. The standard Gibbs energy change for this reaction is :
 a) - 115 kJ b) + 115 kJ c) + 166 kJ d) - 116 kJ
197. The equilibrium constant for the reaction ; $\text{P}_4(\text{s}) + 5\text{O}_2(\text{g}) \rightleftharpoons \text{P}_4\text{O}_{10}(\text{s})$ is :
 a) $K_c = \frac{1}{[\text{O}_2]^5}$ b) $K_c = [\text{O}_2]^5$ c) $K_c = \frac{[\text{P}_4\text{O}_{10}]}{5[\text{P}_4][\text{O}_2]}$ d) $K_c = \frac{[\text{P}_4\text{O}_{10}]}{[\text{P}_4][\text{O}_2]^5}$
198. The correct relation for hydrolysis constant of NH_4CN is:
 a) $\sqrt{\frac{K_w}{K_a}}$ b) $\frac{K_w}{K_a \times K_b}$ c) $\frac{\sqrt{K_H}}{c}$ d) $\frac{K_a}{K_b}$
199. The gaseous reaction,
 $A + B \rightleftharpoons 2C + D + Q$ is most favoured at
 a) Low temperature and high pressure b) High temperature and low high pressure
 c) High temperature and low pressure d) Low temperature and low pressure
200. An aqueous solution of 0.1 M NH_4Cl will have a pH closer to:
 a) 9.1 b) 8.1 c) 7.1 d) 5.1
201. If the concentration of OH^- ions is the reaction $\text{Fe}(\text{OH})_3(\text{s}) \rightleftharpoons \text{Fe}^{3+}(\text{aq}) + 3\text{OH}^-(\text{aq})$, is decreased by $\frac{1}{4}$ times, then equilibrium concentration of Fe^{3+} will increase by :
 a) 16 times b) 64 times c) 4 times d) 8 times
202. $A(\text{g}) + 3B(\text{g}) \rightleftharpoons 4C(\text{g})$.
 Initially concentration of A is equal to that of B. The equilibrium concentrations of A and C are equal. K_c is
 a) 0.08 b) 0.08 c) 8 d) 80
203. 18 mL of mixture of acetic acid and sodium acetate required 6 mL of 0.1 M NaOH for neutralization of the acid and 12 mL of 0.1 M HCl for reaction with salt, separately. If $\text{p}K_a$ of the acid is 4.75, what is the pH of the mixture?

- a) 5.05 b) 4.75 c) 4.5 d) 4.6
204. 50 mL of 0.1 M HCl and 50 mL of 0.2 M NaOH are mixed. The pH of the resulting solution is
a) 1.30 b) 4.2 c) 12.70 d) 11.70
205. K_c for the reaction : $[\text{Ag}(\text{CN})_2]^- \rightleftharpoons \text{Ag}^+ + 2\text{CN}^-$, the equilibrium constant at 25°C is 4.0×10^{-19} , then the silver ion concentration in a solution which was originally 0.1 molar in KCN and 0.03 molar in AgNO_3 is :
a) 7.5×10^{18} b) 7.5×10^{-18} c) 7.5×10^{19} d) 7.5×10^{-19}
206. The $\text{p}K_a$ for acid *A* is greater than $\text{p}K_a$ for acid *B*. The strong acid is:
a) Acid *A* b) Acid *B* c) Are equally strong d) None of these
207. When 100 mL of 1 M NaOH solution is mixed with 10 mL of 10 M H_2SO_4 , the resulting mixture will be
a) Acidic b) Alkaline c) HClO_3 d) H_3PO_3
208. The $[\text{H}_3\text{O}^+]$ in the rain water of $\text{pH} = 4.35$ is:
a) $4.5 \times 10^{-5} \text{ M}$ b) $6.5 \times 10^{-5} \text{ M}$ c) $9.5 \times 10^{-5} \text{ M}$ d) $12.5 \times 10^{-5} \text{ M}$
209. For which salt the pH of its solution does not change with dilution?
a) NH_4Cl b) $\text{CH}_3\text{COONH}_4$ c) CH_3COONa d) None of these
210. When hydrogen molecules decomposed into it's atoms which conditions gives maximum yield of H atom?
a) High temperature and low pressure b) Low temperature and high pressure
c) High temperature and high pressure d) Low temperature and low pressure
211. Which is not and acid salt?
a) NaH_2PO_2 b) NaH_2PO_3 c) NaH_2PO_4 d) NaHSO_3
212. Which is a Lewis base?
a) B_2H_6 b) LiAlH_4 c) AlH_3 d) NH_3
213. Final pressure is higher than initial pressure of a container filled with an ideal gas at constant temperature. What will be the value of equilibrium constant?
a) $K = 1.0$ b) $K = 10.0$ c) $K > 1.0$ d) $K < 1.0$
214. In which of the following cases, does not reaction go farthest to completion?
a) $K = 10^3$ b) $K = 10^{-2}$ c) $K = 10$ d) $K = 1$
215. For the reaction, $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightarrow 2\text{HI}(\text{g})$, the equilibrium constant K_p changes with
a) Total pressure b) Catalyst
c) The amount H_2 and I_2 d) Temperature
216. The equilibrium constant for the reaction,

$$\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$$
At temperature T is 4×10^{-4} . The value of K_c for the reaction

$$\text{NO}(\text{g}) \rightleftharpoons \frac{1}{2}\text{N}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g})$$
at the same temperature is
a) 2.5×10^2 b) 50 c) 4×10^{-4} d) 0.02
217. The reaction, $2\text{A}(\text{g}) + \text{B}(\text{g}) \rightleftharpoons 3\text{C}(\text{g}) + \text{D}(\text{g})$ is begun with the concentration of *A* and *B* both at an initial value of 1.00M. When equilibrium is reached, the concentration of *D* is measured and found to be 0.25 M. The value for the equilibrium constant for this reaction is given by the expression :
a) $[(0.75)^3 (0.25)] \div [(1.00)^2 (1.00)]$
b) $[(0.75)^3 (0.25)] \div [(0.50)^2 (0.75)]$
c) $[(0.75)^3 (0.25)] \div [(0.50)^2 (0.25)]$
d) $[(0.75)^3 (0.25)] \div [(0.75)^2 (0.25)]$
218. In HS^- , I^- , $\text{R} - \text{NH}_2$, NH_3 order of proton accepting tendency will be:
a) $\text{I}^- > \text{NH}_3 > \text{RNH}_2 > \text{HS}^-$
b) $\text{NH}_3 > \text{RNH}_2 > \text{HS}^- > \text{I}^-$
c) $\text{RNH}_2 > \text{NH}_3 > \text{HS}^- > \text{I}^-$

- d) $\text{HS}^- > \text{RNH}_2 > \text{NH}_3 > \text{I}^-$
219. Strong electrolytes are those which:
 a) Dissolve readily in non-polar solvent
 b) Conduct electricity in aqueous solution
 c) Dissociate into ions at high concentration
 d) None of the above
220. The pH of 0.1 N HCl solution is:
 a) 1.0 b) 7.0 c) 14.0 d) 4.0
221. A solution of FeCl_3 in water acts as acidic due to:
 a) Acidic impurities b) Ionisation c) Hydrolysis of Fe^{3+} d) Dissociation
222. The concept that an acid is a proton donor and a base is a proton acceptor was introduced by:
 a) Arrhenius b) Bronsted-Lowry c) Lewis d) Faraday
223. Which is decreasing order of strength of bases?
 $\bar{\text{O}}\text{H}, \bar{\text{N}}\text{H}_2, \text{HC} \equiv \text{C}^-$ and CH_3CH_2^-
 a) $\text{H}_3\text{CCH}_2^- > \text{NH}_2^- > \text{HC} \equiv \text{C}^- > \text{OH}^-$ b) $\text{HC} \equiv \text{C}^- > \text{CH}_3\text{CH}_2^- > \text{NH}_2^- > \text{OH}^-$
 c) $\text{OH}^- > \text{NH}_2^- > \text{CH} \equiv \text{C}^- > \text{H}_3\text{CCH}_2^-$ d) $\text{NH}_2^- > \text{HC} \equiv \text{C}^- > \text{OH}^- > \text{H}_3\text{CCH}_2^-$
224. The strength of an acid depends on its tendency to
 a) Accept protons b) Donate protons c) Accept electrons d) Donate electrons
225. The following reactions are known to occur in the body,
 $\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3 \rightleftharpoons \text{H}^+ + \text{HCO}_3^-$
 If CO_2 escapes from the system, then:
 a) pH will decrease
 b) Hydrogen ion concentration will diminish
 c) H_2CO_3 concentration will be unaltered
 d) The forward reaction will be promoted
226. The common ion effect is shown by which of the following sets of solutions?
 a) $\text{BaCl}_2 + \text{BaNO}_3$ b) $\text{NaCl} + \text{HCl}$ c) $\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$ d) None of these
227. In the reaction, $\text{C}(s) + \text{CO}_2(g) \rightleftharpoons 2\text{CO}(g)$, the equilibrium pressure is 12 atm. If 50% of CO_2 reacts, K_p for the change is :
 a) 12 atm b) 16 atm c) 20 atm d) 6 atm
228. For a given solution pH = 6.9 at 60°C , where $K_w = 10^{-12}$. The solution is:
 a) Acidic b) Basic c) Neutral d) Unpredictable
229. A quantity of PCl_5 was heated in a 10 litre vessel at 250°C to show $\text{PCl}_5(g) \rightleftharpoons \text{PCl}_3(g) + \text{Cl}_2(g)$. At equilibrium the vessel contains 0.1 mole of PCl_5 , 0.20 mole of PCl_3 and 0.20 mole of Cl_2 . The equilibrium constant of the reaction is :
 a) 0.02 b) 0.05 c) 0.04 d) 0.025
230. One mole of ethyl alcohol was treated with one mole of acetic acid at 25°C . $2/3$ of the acid changes into ester at equilibrium. The equilibrium constant for the reaction will be:
 a) 1 b) 2 c) 3 d) 4
231. 9.2 g of $\text{N}_2\text{O}_4(g)$ is taken in a closed 1 L vessel and heated till the following equilibrium is reached
 $\text{N}_2\text{O}_4(g) \rightleftharpoons 2\text{NO}_2(g)$
 At equilibrium, 50% $\text{N}_2\text{O}_4(g)$ is dissociated. What is the equilibrium constant (in molL^{-1})? (Molecular weight of $\text{N}_2\text{O}_4 = 92$)
 a) 0.1 b) 0.2 c) 0.3 d) 0.4
232. Assuming complete dissociation which of the following aqueous solutions will have the same pH value?
 (i) 100 mL of 0.01 M HCl
 (ii) 100 mL of 0.01 M H_2SO_4
 (iii) 50 mL of 0.01 M HCl
 (iv) Mixture of 50 mL of 0.02 M H_2SO_4 and 50 mL of 0.02 M NaOH

- a) (i), (ii) b) (i), (iii) c) (ii), (iv) d) (i), (iv)
233. At 3000 K, the equilibrium pressure of CO_2 , CO and O_2 are 0.6, 0.4 and 0.2 atm respectively. K_p for the reaction $2\text{CO}_2 \rightleftharpoons 2\text{CO} + \text{O}_2$, is
 a) 0.089 b) 0.098 c) 0.189 d) 0.198
234. The $\text{p}K_a$ of weak acid H_A is 4.5. The pOH of an aqueous buffer solution of HA in which 50% of the acid is ionised:
 a) 7.0 b) 4.5 c) 2.5 d) 9.5
235. An amphoteric buffer solution in which conc. of H^+ and HX is same. The value of K_a of HX is 10^{-8} , then pH of buffer solution is
 a) 3 b) 8 c) 10 d) 14
236. In the reaction, $3A + 2B \rightarrow 2C$, the equilibrium constant K_c is given by
 a) $\frac{[3A] \times [2B]}{[C]}$ b) $\frac{[A]^3 \times [B]}{[C]}$ c) $\frac{[C]^2}{[A]^3 \times [B]^2}$ d) $\frac{[C]}{[3A][2B]}$
237. Which reaction is not affected by change in pressure?
 a) $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$ b) $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$
 c) $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$ d) $2\text{C} + \text{O}_2 \rightleftharpoons 2\text{CO}$
238. Three reactions involving H_2PO_4^- are given below
 (i) $\text{H}_3\text{PO}_4 + \text{H}_2\text{O} \rightarrow \text{H}_3\text{O}^+ + \text{H}_2\text{PO}_4^-$
 (ii) $\text{H}_2\text{PO}_4^- + \text{H}_2\text{O} \rightarrow \text{HPO}_4^{2-} + \text{H}_3\text{O}^+$
 (iii) $\text{H}_2\text{PO}_4^- + \text{OH}^- \rightarrow \text{H}_3\text{PO}_4 + \text{O}^{2-}$
 In which of the above does H_2PO_4^- act as an acid?
 a) (ii) only b) (i) and (ii) c) (iii) only d) (i) only
239. pH for the solution of salt undergoing anionic hydrolysis (say CH_3COONa) is given by:
 a) $\text{pH} = \frac{1}{2}[\text{p}K_w + \text{p}K_a + \log c]$
 b) $\text{pH} = \frac{1}{2}[\text{p}K_w + \text{p}K_a - \log c]$
 c) $\text{pH} = \frac{1}{2}[\text{p}K_w + \text{p}K_b - \log c]$
 d) None of the above
240. For the reactions, $A + B + Q \rightleftharpoons C + D$, if the temperature is increased then concentration of the products will
 a) Increase b) Decrease c) Remains the same d) Become zero
241. Under what conditions of temperature and pressure, the formation of atomic hydrogen from molecular hydrogen will be favoured most?
 a) High temperature and high pressure b) High temperature and low pressure
 c) Low temperature and low pressure d) Low temperature and high pressure
242. Mohr's salt is a:
 a) Normal salt b) Acid salt c) Basic salt d) Double salt
243. pH of 0.05 M $\text{Mg}(\text{OH})_2$ is:
 a) 13 b) 10 c) 1 d) Zero
244. In which of the following reactions, the concentration of product is higher than the concentration of reactant at equilibrium? (K = equilibrium constant)
 a) $A \rightleftharpoons B; K = 0.001$ b) $M \rightleftharpoons N; K = 10$ c) $X \rightleftharpoons Y; K = 0.005$ d) $R \rightleftharpoons P; K = 0.01$
245. The values of dissociation constant of bases are given below. Which is the weakest base?
 a) 1.8×10^{-5} b) 4.8×10^{-10} c) 7.2×10^{-11} d) 7.07×10^{-7}
246. The dissociation equilibrium of a gas AB_2 can be represented as :

$$2\text{AB}_2(\text{g}) \rightleftharpoons 2\text{AB}(\text{g}) + \text{B}_2(\text{g})$$
 The degree of dissociation is ' x ' and is small compared to 1. The expression relating the degree of

dissociation (x) with equilibrium constant K_p and total pressure p is :

- a) $(2K_p/P)^{1/3}$ b) $(2K_p/P)^{1/2}$ c) (K_p/P) d) $(2K_p/P)$
247. In which one of the following gaseous equilibria, K_p is less than K_c ?
- a) $N_2O_4 \rightleftharpoons 2NO_2$ b) $2SO_2 + O_2 \rightleftharpoons 2SO_3$ c) $2HI \rightleftharpoons H_2 + I_2$ d) $N_2 + O_2 \rightleftharpoons 2NO$
248. K_{sp} for $Cr(OH)_3$ is 2.7×10^{-31} . What is its solubility in mol/L?
- a) 1×10^{-8} b) 8×10^{-8} c) 1.1×10^{-8} d) 0.18×10^{-8}
249. N_2O_4 is dissociated to 33% and 40% at total pressure P_1 and P_2 atm respectively. Then the ratio P_1/P_2 is:
- a) 7/4 b) 7/3 c) 8/3 d) 8/5
250. In the reactions, $A + 2B \rightleftharpoons 2C$, if 2 moles of A , 3.0 moles of B and 2.0 moles of C are placed in a 2 L flask and the equilibrium concentration of C is 0.5 mol/L, the equilibrium constant (K_c) for the reactions is
- a) 0.21 b) 0.50 c) 0.75 d) 0.025
251. The pH value of 1/1000 N KOH solution is
- a) 3 b) 10^{-11} c) 2 d) 11
252. The pH of tears coming out of a person's eye is:
- a) 7.4 b) 6.4 c) 7.0 d) 2.36
253. The solubility of CaF_2 is 2×10^{-4} mol/L. Its solubility product (K_{sp}) is
- a) 2.0×10^{-4} b) 4.0×10^{-3} c) 8.0×10^{-12} d) 3.2×10^{-11}
254. The solubility product of a salt having general formula MX_2 in water is 4×10^{-12} . The concentration of M^{2+} ions in the aqueous solution of the salt is
- a) $2.0 \times 10^{-6}M$ b) $1.0 \times 10^{-4}M$ c) $1.6 \times 10^{-4}M$ d) $4.0 \times 10^{-10}M$
255. The solubility product of barium sulphate is 1.5×10^{-9} at $18^\circ C$. Its solubility in water at $18^\circ C$ is
- a) 1.5×10^{-9} b) 1.5×10^{-5} c) 3.9×10^{-9} d) 3.9×10^{-5}
256. The strongest Bronsted base is
- a) ClO_3^- b) ClO_2^- c) ClO_4^- d) ClO^-
257. The reaction quotient (Q) at equilibrium is:
- a) = 1 b) = K c) $> K$ d) $< K$
258. The concentration of oxalic acid is ' x ' mol L^{-1} . 40 mL of this solution reacts with 16 mL of 0.05 M acidified $KMnO_4$. What is the pH of ' x ' M oxalic acid solution?
(Assume that oxalic acid dissociates completely)
- a) 1.3 b) 1.699 c) 1 d) 2
259. Metal ions like Ag^+ , Cu^{2+} etc. act as
- a) Bronsted acids b) Bronsted bases c) Lewis acids d) Lewis bases
260. The pK_a of acetylsalicylic acid (aspirin) is 3.5. The pH of gastric juice in human stomach is about 2 – 3 and the pH in the small intestine is about 8. Aspirin will be
- a) Unionised in the small intestine and in the stomach
b) Completely ionised in the small intestine and in the stomach
c) Ionised in the stomach and almost unionised in the small intestine
d) Ionised in the small intestine and almost unionised in the stomach
261. A solution is called saturated if:
- a) Ionic concentration product $<$ solubility product
b) Ionic concentration product $>$ solubility product
c) Ionic concentration product \geq solubility product
d) None of the above
262. The auto protonation constant of H_2O is:
- a) 1×10^{-14} b) 3.23×10^{-18} c) 1.8×10^{-18} d) 3.23×10^{-20}
263. K_c for $m_1A + m_2B = n_1C + n_2D$ is given by:
- a) $K_c = \frac{[A]^{m_1}[B]^{m_2}}{[C] \times [D]}$ b) $K_c = \frac{[A]^{n_1}[B]^{n_2}}{[C]^{m_1}[D]^{m_2}}$ c) $K_c = \frac{[C]^{n_1}[D]^{n_2}}{[A]^{m_1}[B]^{m_2}}$ d) $K_c = \frac{[C]^{m_1} \times [D]^{m_2}}{[A]^{n_1} \times [B]^{n_2}}$

264. The pH of millimolar HCl is
 a) 1 b) 3 c) 2 d) 4
265. Partial pressure of A, B, C and D on the basis of gaseous system, $A + 2B \rightleftharpoons C + 3D$, are $A = 0.20, B = 0.10, C = 0.30$ and $D = 0.50$ atm. The numerical value of equilibrium constant is
 a) 3.75 b) 18.75 c) 17.85 d) 15.87
266. Which equilibrium can be described as Lewis acid-base reaction but not Bronsted acid-base reaction?
 a) $H_2O + CH_3COOH \rightleftharpoons H_3O^+ + CH_3COO^-$
 b) $2NH_3 + H_2SO_4 \rightleftharpoons 2NH_4^+ + SO_4^{2-}$
 c) $NH_3 + CH_3COOH \rightleftharpoons NH_4^+ + CH_3COO^-$
 d) $[Cu(H_2O)_4]^{2+} + 4NH_3 \rightleftharpoons [Cu(NH_3)_4]^{2+} + 4H_2O$
267. $SnCl_2$ and $HgCl_2$ cannot co-exist in a solution because of:
 a) Common ion effect
 b) Le – Chatelier’s principle
 c) Conc. of Cl^- increases to precipitate both
 d) Redox change
268. The species which acts as a Lewis but not a Bronsted acid is
 a) NH_2^- b) O^{2-} c) BF_3 d) OH^-
269. What is the best description of the change that occurs when $Na_2O(s)$ is dissolved in water?
 a) Oxidation number of sodium decreases
 b) Oxide ion accepts sharing in a pair of electrons
 c) Oxide ion donates a pair of electrons
 d) Oxidation number of oxygen increases
270. pH of 0.005 M calcium acetate is
 (pK_a of $CH_3COOH = 4.74$)
 a) 7.04 b) 9.37 c) 9.26 d) 8.2195
271. Relation between hydrolysis constant and dissociation constant are given. Which is the correct formula for $MgCl_2$?
 a) $K_h = \frac{K_w}{K_a}$ b) $K_h = \frac{K_w}{K_b}$ c) $K_h = \frac{K_w}{K_a \times K_b}$ d) $K_w = \frac{K_h}{K_b}$
272. Theory’s ‘active mass’ indicates that the rate of chemical reaction is directly proportional to the
 a) Equilibrium constant b) Volume of apparatus
 c) Properties of reactants d) Concentration of reactants
273. In which of the following reactions, the value of K_p will be equal to K_c ?
 a) $PCl_5 \rightleftharpoons PCl_3 + Cl_2$ b) $2NH_3 \rightleftharpoons N_2 + 3H_2$ c) $H_2 + I_2 \rightleftharpoons 2HI$ d) $2SO_2 + O_2 \rightleftharpoons 2SO_3$
274. In the hydrolysis of a salt of weak acid and weak base, the hydrolysis constant K_h is equal to
 a) $\frac{K_w}{K_b}$ b) $\frac{K_w}{K_a}$ c) $\frac{K_w}{K_a \cdot K_b}$ d) $K_a \cdot K_b$
275. In which reaction ammonia acts as an acid?
 a) $NH_3 + HCl \rightarrow NH_4Cl$
 b) $NH_3 + H^+ \rightarrow NH_4^+$
 c) $NH_3 + Na \rightarrow NaNH_2 + \frac{1}{2}H_2$
 d) NH_3 cannot act as an acid
276. The compounds A and B are mixed in equimolar proportion to form the products, $A + B \rightleftharpoons C + D$. At equilibrium, one third of A and B are consumed. The equilibrium constant for the reaction is
 a) 0.5 b) 4.0 c) 2.5 d) 0.25
277. 40% of a mixture of 0.2 mole of N_2 and 0.6 mole of H_2 react to give NH_3 according to the equation, $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ at constant temperature and pressure. Then the ratio of the final volume to the initial

volume of gases is :

- a) 4 : 5 b) 5 : 4 c) 7 : 10 d) 8 : 5

278. An aqueous solution contains a substance which yields 4×10^{-3} mol litre⁻¹ ion of H_3O^+ . If $\log 2 = 0.3010$, the pH of the solution is:

- a) 1.5 b) 2.398 c) 3.0 d) 3.4

279. For preparing a buffer solution of pH 6 by mixing sodium acetate and acetic acid, the ration of concentration of salt and acid ($K_a = 10^{-5}$) should be:

- a) 1 : 10 b) 10 : 1 c) 100 : 1 d) 1 : 100

280. The concentration of hydrogen ion $[\text{H}^+]$ and pH in 10 M HCl is:

- a) 10^1 , zero b) 10^1 , -1 c) 10^2 , 1 d) 10^1 , 1

281. Solubility product of $\text{Mg}(\text{OH})_2$ at ordinary temperature is 1.96×10^{-11} . pH of a saturated solution of $\text{Mg}(\text{OH})_2$ will be

- a) 10.53 b) 8.47 c) 6.94 d) 3.47

282. For the reaction $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$:

- a) $K_c = 2K_p$ b) $K_c > K_p$ c) $K_c = K_p$ d) $K_c < K_p$

283. When CaCO_3 is heated at a constant temperature in a closed container, the pressure due to CO_2 produced will:

- a) Change with the amount of CaCO_3 taken
 b) Change with the size of the container
 c) Remain constant so long as temperature is constant
 d) Remain constant even if temperature is changed

284. Four species are listed below

IV. HCO_3^-

V. H_3O^+

VI. HSO_4^-

VII. HSO_3F

Which one of the following is the correct sequence of their acid strength?

- a) (iv) < (ii) < (iii) < (i) b) (ii) < (iii) < (i) < (iv)
 c) (i) < (iii) < (ii) < (iv) d) (iii) < (i) < (iv) < (ii)

285. 1 dm³ solution containing 10^5 moles each of Cl^- ions and CrO_4^{2-} ions is treated with 10^4 moles of silver nitrate. Which one of the following observation is made?

$$[K_{sp}\text{Ag}_2\text{CrO}_4 \quad 4 \quad 10^{12}]$$

$$[K_{sp}\text{AgCl} \quad 1 \quad 10^{10}]$$

- a) Precipitation does not occur
 b) Silver chromate gets precipitated first
 c) Silver chloride gets precipitated first
 d) Both silver chromate and silver chloride start precipitating simultaneously

286. Which is a basic salt?

- a) PbS b) PbCO_3 c) PbSO_4 d) $2\text{PbCO}_3\text{Pb}(\text{OH})_2$

287. A reversible reaction, $\text{H}_2 + \text{Cl}_2 \rightleftharpoons 2\text{HCl}$ is carried out in one litre flask. If the same reaction is carried out in two litre flask, the equilibrium constant will be:

- a) Doubled b) Decreased c) Halved d) Same

288. In the system, $\text{CaF}_2(s) \rightleftharpoons \text{Ca}^{2+}(aq) + 2\text{F}^-(aq)$, increasing the concentration of Ca^{2+} ions 4 times will cause the equilibrium concentration of F^- ions to change to :

- a) $\frac{1}{4}$ of the initial value
 b) $\frac{1}{2}$ of the initial value
 c) 2 times of the initial value
 d) None of the above

289. Hydrogen ion concentration in mol/L in a solution of pH = 5.4 will be
 a) 3.98×10^8 b) 3.88×10^6 c) 3.68×10^{-6} d) 3.98×10^{-6}
290. The strongest conjugate base is
 a) NO_3^- b) Cl^- c) SO_4^{2-} d) CH_3COO^-
291. In the reaction $\text{I}_2 + \text{I}^- = \text{I}_3^-$, the Lewis base is:
 a) I_2 b) I^- c) I_3^- d) None of these
292. HI was heated in a sealed tube at 440°C till the equilibrium was reached, HI was found to be 22% decomposed. The equilibrium constant for dissociation is :
 a) 0.282 b) 0.0796 c) 0.0199 d) 1.99
293. Which one is amphoteric oxide?
 a) SO_2 b) B_2O_3 c) ZnO d) Na_2O
294. For which reaction K_p is less than K_c ?
 a) $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$ b) $2\text{HI} \rightleftharpoons \text{H}_2 + \text{I}_2$ c) $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3$ d) $\text{N}_2 + \text{O}_2 \rightleftharpoons 2\text{NO}$
295. For the reactions, $\text{H}_2(\text{g}) + \text{CO}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g})$ if the initial concentration of $[\text{H}_2] = [\text{CO}_2]$ and x mol/L of hydrogen is consumed at equilibrium, the correct expression of K_p is
 a) $\frac{x^2}{(1-x)^2}$ b) $\frac{x^2}{(2+x)^2}$ c) $\frac{x^2}{1-x^3}$ d) $\frac{(1+x)^2}{(1-x)^2}$
296. In the given reaction,

$$2\text{X}(\text{g}) + \text{Y}(\text{g}) \rightleftharpoons 2\text{Z}(\text{g}) + 80 \text{ kcal,}$$
 Which combination of pressure and temperature will give the highest yield of Z at equilibrium?
 a) 1000 atm and 200°C b) 500 atm and 500°C
 c) 1000 atm and 100°C d) 500 atm and 100°C
297. Equimolar solutions of the following were prepared in water separately. Which one of the solutions will record the highest pH?
 a) BaCl_2 b) MgCl_2 c) CaCl_2 d) SrCl_2
298. Which is not correct for Lewis acids?
 a) They contain at least one vacant orbital
 b) They have a tendency to accept electrons
 c) The smaller ion has greater acidic strength
 d) In case of ions, the strength of acid is inversely proportional to its charge
299. The vapour density of N_2O_4 at a certain temperature is 30. What is the percentage dissociation of N_2O_4 at this temperature?
 a) 46.5% b) 36.2% c) 53.3% d) 64.2%
300. For which reaction $K_p \neq K_c$?
 a) $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2(\text{g}) + \text{O}_2(\text{g})$
 b) $\text{SO}_2(\text{g}) + \text{NO}_2(\text{g}) \rightleftharpoons \text{SO}_3(\text{g}) + \text{NO}(\text{g})$
 c) $\text{I}_2(\text{g}) + \text{H}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$
 d) $2\text{C}(\text{s}) + \text{O}_2(\text{g}) \rightarrow 2\text{CO}(\text{g})$
301. A buffer mixture of acetic acid and potassium acetate has pH = 5.24. The ratio of $[\text{CH}_3\text{COO}^-]/[\text{CH}_3\text{COOH}]$ in this buffer is, ($\text{p}K_a = 4.74$):
 a) 3 : 1 b) 1 : 3 c) 1 : 1 d) 1 : 2
302. $\text{p}K_a$ of acetic acid is 4.74. The concentration of CH_3COONa is 0.01 M. The pH of CH_3COONa is
 a) 3.37 b) 4.37 c) 4.74 d) 0.474
303. If 1 M CH_3COONa is added to 1 M CH_3COOH :
 a) pH of the solution increases
 b) pH decreases
 c) pH does not change
 d) None of the above
304. 2.5 mL of $\frac{2}{5}$ M weak monoacidic base ($K_b = 1 \times 10^{-12}$ at 25°C) is titrated with $\frac{2}{15}$ M HCl in water

at 25°C. The concentration of H^+ at equivalence point is

($K_w = 1 \times 10^{-14}$ at 25°C)

- a) 3.7×10^{-13} M b) 3.2×10^{-7} M c) 3.2×10^{-2} M d) 2.7×10^{-2} M

305. Solubility product of a salt AB is $1 \times 10^{-8} M^2$ in a solution in which the concentration of A^+ ions is 10^{-3} M. The salt will precipitate when the concentration of B^- ions is kept

- a) Between 10^{-8} to 10^{-7} M b) Between $10^{-7} M$ to 10^{-8} M
c) $> 10^{-5}$ M d) $< 10^{-8}$ M

306. For the gaseous reaction, $C_2H_4 + H_2 \rightleftharpoons C_2H_6$, $\Delta H = -130 \text{ kJ mol}^{-1}$ carried in a closed vessel, the equilibrium concentration of the C_2H_6 can definitely be increased by

- a) Increasing temperature and decreasing pressure b) Decreasing temperature and increasing pressure
c) Increasing temperature and pressure both d) Decreasing temperature and pressure only

307. Chemical equilibrium is dynamic in nature because:

- a) The equilibrium is maintained rapidly
b) The concentration of reactants and products become same at equilibrium
c) The concentration of reactants and products decrease with time
d) Both forward and backward reactions occur at all times with same speed

308. What happens to the yield on application of high pressure in the Haber's synthesis of ammonia?

- a) Increases b) Decreases c) Unaffected d) Reaction stops

309. The buffering action of an acidic buffer is maximum when its pH is equal to

- a) 5 b) 7 c) 1 d) pK_a



310. HA is a weak acid. The pH of 0.1 M HA solution is 2. What is the degree of dissociation (α) of HA ?



- a) 0.5 b) 0.2 c) 0.1 d) 0.301

311. Which of the following is a wrong statement about equilibrium state?

- a) Rate of forward reaction = Rate of backward reaction
b) Equilibrium is dynamic
c) Catalysts increase value of equilibrium constant
d) Free energy change is zero

312. In an experiment to determine the enthalpy of neutralization of sodium hydroxide with sulphuric acid, 50 cm^3 of 0.4 M sodium hydroxide were titrated thermometrically with 0.25 M sulphuric acid. Which of the following plots gives the correct representation?





313. H⁺ ion produces common ion effect in the wet analysis of:
 a) Group I metals b) Group II metals c) Group III metals d) Group IV metals
314. 15 moles of H₂ and 5.2 moles of I₂ are mixed and allowed to attain equilibrium at 500°C. At equilibrium, the concentration of HI is found to be 10 moles. The equilibrium constant for the formation of HI is
 a) 50 b) 15 c) 100 d) 25
315. 10⁻⁶ M HCl is diluted to 100 times. Its pH is:
 a) 6.0 b) 8.0 c) 6.95 d) 9.5
316. For the reaction, PCl₃(g) + Cl₂(g) ⇌ PCl₅(g), the position of equilibrium can be shifted to the right by
 a) Doubling the volume
 b) Increasing the temperature
 c) Addition of Cl₂ at constant volume
 d) Addition of equimolar quantities of PCl₃ and PCl₅
317. The pH of an aqueous solution containing [H⁺] concentration = 3.0 × 10⁻³ M. The pH of the solution is
 a) 2.523 b) 3.0 c) 2.471 d) None of these
318. The addition of which salt will decrease the H⁺ concentration of HCN solution?
 a) NH₄Cl b) Al₂(SO₄)₃ c) AgNO₃ d) NaCN
319. The pH of the solution obtained by mixing 10 mL of 10⁻¹ N HCl and 10 mL of 10⁻¹ N NaOH is:
 a) 8 b) 2 c) 7 d) None of these
320. The solubility product of PbCl₂ is 2.3 × 10⁻³². Its solubility will be
 a) 1.78 × 10⁻¹¹ g/L b) 2.95 × 10⁻⁹ g/L c) 3.42 × 10⁻⁹ g/L d) 4.95 × 10⁻⁹ g/L
321. A white salt is readily soluble in water and gives a colourless solution with a pH of about 9. The salt would be:
 a) NH₄NO₃ b) CH₃COONa c) CH₃COONH₄ d) CaCO₃
322. The dissociation constant of NH₄OH is 1.8 × 10⁻⁵. The hydrolysis constant of NH₄Cl would be:
 a) 1.8 × 10⁻¹⁹ b) 1.8 × 10⁻⁵ c) 5.55 × 10⁻⁵ d) 5.55 × 10⁻¹⁰
323. 50 mL of H₂O is added to 50 mL of 1 × 10⁻³ M barium hydroxide solution. What is the pH of the resulting solution?
 a) 3.0 b) 3.3 c) 11.0 d) 11.7
324. The indicator used in titration of oxalic acid with caustic soda solution is
 a) Methyl orange b) Methyl red c) Fluorescein d) Phenolphthalein
325. For H₂ + I₂ ⇌ 2HI, at equilibrium some I₂ is added. What happens to the equilibrium?
 a) It is shifted to the right b) It gets shifted to the left c) It remains unchanged d) None of the above
326. Which of the following is a characteristic of a reversible reaction?

- a) It can never proceed to completion
 b) It can be influenced by a catalyst
 c) Number of moles of reactants and products are equal
 d) None of the above
327. An aqueous solution of hydrogen sulphide shows the equilibrium,

$$\text{H}_2\text{S} \rightleftharpoons \text{H}^+ + \text{HS}^-$$
 If dilute hydrochloric acid is added to an aqueous solution of hydrogen sulphide without any change in temperature, then:
 a) The equilibrium constant will change
 b) The concentration of HS^- will increase
 c) The concentration of undissociated hydrogen sulphide will decrease
 d) The concentration of HS^- will decrease
328. Le-Chatelier's principle is not applicable to:
 a) Homogeneous reactions
 b) Heterogeneous reactions
 c) Homogeneous or heterogeneous systems in equilibrium
 d) Systems not in equilibrium
329. If $\text{p}K_a$ values of four acids are given below at 25°C , the strongest acid is
 a) 2.0 b) 2.5 c) 3.0 d) 4.0
330. Weakest base among the following is:
 a) NaOH b) $\text{Ca}(\text{OH})_2$ c) $\text{Zn}(\text{OH})_2$ d) KOH
331. A solution of pH 9.0 is one thousand times as basic as a solution of pH:
 a) 6 b) 7 c) 4 d) 10
332. Aprotic solvent is:
 a) CCl_4 b) C_6H_6 c) SO_2 d) All of these
333. The hydroxide with highest solubility product is:
 a) $\text{Al}(\text{OH})_3$ b) $\text{Co}(\text{OH})_2$ c) $\text{Cr}(\text{OH})_3$ d) $\text{Fe}(\text{OH})_3$
334. In the absence of formation of complex ions by the addition of a common ion, the solubility of a given salt is:
 a) Increased
 b) Decreased
 c) Unaffected
 d) First increased and then decreased
335. The pH of 0.1 M NaHS is, K_{a_1} and K_{a_2} for H_2S are 1.3×10^{-7} and 7.1×10^{-15} respectively:
 a) 10.52 b) 9.52 c) 12.52 d) 13.52
336. $A + B \rightleftharpoons C + D$
 Initially moles of A and B are equal. At equilibrium, moles of C are three times that of A. the equilibrium constant of the reaction will be
 a) 1 b) 3 c) 4 d) 9
337. The strongest acid among the following is:
 a) $\text{ClO}_3(\text{OH})$
 b) $\text{ClO}_2(\text{OH})$
 c) $\text{SO}(\text{OH}_2)$
 d) $\text{SO}_2(\text{OH})_2$
338. The equilibrium constant in a reversible reaction at a given temperature
 a) Does not depend on the initial concentrations
 b) Depends on the initial concentrations of the reactants
 c) Depends on the concentration of the products at equilibrium
 d) It is not characteristic of the reaction

339. For the reaction, $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ at 720 K, the value of equilibrium constant is 50, when equilibrium concentration of both H_2 and I_2 is 0.5 M. K_p under the same conditions will be :
- a) 0.02 b) 0.2 c) 50 d) 50 RT
340. If 340 g of a mixture of N_2 and H_2 in the correct ratio gave a 20% yield of NH_3 . The mass produced would be :
- a) 16 g b) 17 g c) 20 g d) 68 g
341. The conjugate acid of CO_3^{2-} is:
- a) H_2O b) H_2CO_3 c) OH^- d) HCO_3^-
342. Calculate the partial pressure of carbon monoxide from the following datas
- $$\text{CaCO}_3 \xrightarrow{\Delta} \text{CaO}(\text{g}) + \text{CO}_2 \uparrow \quad K_p = 8 \times 10^{-2}$$
- $$\text{O}_2(\text{g}) + \text{C}(\text{s}) \rightarrow 2\text{CO}(\text{g}), \quad K_p = 2$$
- a) 0.2 b) 0.4 c) 1.6 d) 4
343. In aqueous solution, the ionisation constants for carbonic acid are,
- $$K_1 = 4.2 \times 10^{-7} \quad \text{and} \quad K_2 = 4.8 \times 10^{-11}$$
- Select the correct statement for a saturated 0.034 M solution of the carbonic acid.
- a) The concentration of CO_3^{2-} is 0.034 M b) The concentration of CO_3^{2-} is greater than that of HCO_3^-
- c) The concentration of H^+ and HCO_3^- are approximately equal d) The concentration of H^+ is double that of CO_3^{2-}
344. The rapid change of pH near the stoichiometric point of an acid base titration is the basis of indicator detection. pH of the solution is related to the ratio of the concentration of the conjugate acid (HIn) and base (In^-) forms of the indicator given by the expression
- a) $\log \frac{[\text{In}^-]}{[\text{HIn}]} = \text{p}K_{\text{In}} - \text{pH}$ b) $\log \frac{[\text{HIn}]}{[\text{In}^-]} = \text{p}K_{\text{In}} - \text{pH}$
- c) $\log \frac{[\text{HIn}]}{[\text{In}^-]} = \text{pH} - \text{p}K_{\text{In}}$ d) $\log \frac{[\text{In}^-]}{[\text{HIn}]} = \text{pH} - \text{p}K_{\text{In}}$
345. The number of mole of hydroxide [OH^-] ion in 0.3 litre of 0.005 M solution of $\text{Ba}(\text{OH})_2$ is:
- a) 0.0075 b) 0.0015 c) 0.0030 d) 0.0050
346. 4.5 moles each of hydrogen and iodine heated in a sealed 10 L vessel. At equilibrium 3 moles of HI were found. The equilibrium constant for $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ is
- a) 1 b) 5 c) 10 d) 0.5
347. The degree of hydrolysis in hydrolytic equilibrium $\text{A}^- + \text{H}_2\text{O} \rightleftharpoons \text{HA} + \text{OH}^-$ at salt concentration of 0.001 M is ($K_a = 1 \times 10^{-5}$)
- a) 1×10^{-3} b) 1×10^{-4} c) 5×10^{-4} d) 1×10^{-6}
348. For a hypothetical equilibrium:
- $$4\text{A} + 5\text{B} \rightleftharpoons 4\text{x} + 6\text{y}; \text{ the equilibrium constant } K_c \text{ has the unit:}$$
- a) $\text{mol}^2\text{litre}^{-2}$ b) litre mol^{-1} c) $\text{litre}^2\text{mol}^{-2}$ d) mol litre^{-1}
349. Salting out action of soap is based on:
- a) Complex ion formation
- b) Common ion effect
- c) Solubility product
- d) Acid-base neutralization
350. The equilibrium constant for the reaction,
- $$\text{SO}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightleftharpoons \text{SO}_3(\text{g}) \text{ is } 5 \times 10^{-2} \text{ atm. The equilibrium constant of the reaction}$$
- $$2\text{SO}_3(\text{g}) \rightleftharpoons 2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \text{ would be}$$
- a) 100 atm b) 200 atm c) 4×10^2 atm d) 6.25×10^4 atm
351. Which can be explained as applications of Le-Chatelier's principle?

- a) Transport of oxygen by haemoglobin in blood
 b) Removal of CO₂ from tissues by blood
 c) Tooth decay due to use of sweet substances
 d) All of the above
352. Which equilibrium in gaseous phase would be unaffected by an increase in pressure?
 a) N₂O₄ ⇌ 2NO₂ b) N₂ + O₂ ⇌ 2NO c) N₂ + 3H₂ ⇌ 2NH₃ d) CO + $\frac{1}{2}$ O₂ ⇌ CO₂
353. The aqueous solution of AlCl₃ is acidic due to the hydrolysis of
 a) Aluminum ion b) Chloride ion
 c) Both aluminium and chloride ion d) None of the above
354. The percentage error in [H⁺] made by neglecting the ionisation of water in 1.0 × 10⁻⁶ M NaOH is:
 a) 1% b) 2% c) 3% d) 4%
355. The colour of CuCr₂O₇ solution in water is green because:
 a) Cu²⁺ ions is green
 b) Cr₂O₇²⁻ ion is green
 c) Both the ions are green
 d) Cu²⁺ ion is blue and Cr₂O₇²⁻ ion is yellow
356. Ammonium carbonate decomposes as
 NH₂COONH₄(s) ⇌ 2NH₃(g) + CO₂(g)
 For the reaction, K_p = 2.9 × 10⁻⁵ atm⁻³. If we start with 1 mole of the compound, the total pressure at equilibrium would be
 a) 0.0766 atm b) 0.0582 atm c) 0.388 atm d) 0.0194 atm
357. Ionic product of water increases if
 a) Pressure is reduced b) H⁺ is added
 c) OH⁻ is added d) Temperature increase
358. In which of the following reactions, increases in the volume at constant temperature do not affect the number of moles at equilibrium?
 a) 2NH₃ ⇌ N₂ + 3H₂ b) C(s) + $\frac{1}{2}$ O₂(g) → CO(g)
 c) H₂(g) + O₂(g) → H₂O₂(g) d) None of the above
359. Which one of the following is least likely to act as a Lewis base?
 a) I⁺ b) I c) SCl₂ d) PCl₃
360. An aqueous solution of ammonium acetate is:
 a) Faintly acidic b) Fair acidic c) Faintly alkaline d) Almost neutral
361. The strongest Lewis base in the following is
 a) CH₃⁻ b) F⁻ c) NH₂⁻ d) OH⁻
362. For anionic hydrolysis, pH is given by
 a) $\text{pH} = \frac{1}{2}\text{p}K_w - \frac{1}{2}\text{p}K_b - \frac{1}{2}\log C$ b) $\text{pH} = \frac{1}{2}\text{p}K_w + \frac{1}{2}\text{p}K_a - \frac{1}{2}\text{p}K_b$
 c) $\text{pH} = \frac{1}{2}\text{p}K_w + \frac{1}{2}\text{p}K_a + \frac{1}{2}\log C$ d) $\text{pH} = -\frac{1}{2}(\text{p}K_w - \text{p}K_a - \text{p}K_b)$
363. Which of the following is a conjugated acid-base pair?
 a) HCl, NaOH b) NH₄Cl, NH₄OH c) H₂SO₄, HSO₄⁻ d) KCN, HCN
364. In the hydrolytic equilibrium,
 A⁻ + H₂O ⇌ HA + OH⁻
 K_a = 1.0 × 10⁻⁵. The degree of hydrolysis of 0.001 M solution of the salt is:
 a) 10⁻³ b) 10⁻⁴ c) 10⁻⁵ d) 10⁻⁶
365. The equilibrium constant (K_c) for the reaction, N₂(g) + O₂(g) ⇌ 2NO(g) at room temperature T is 4 × 10⁻⁴. The value of K_c for NO(g) ⇌ $\frac{1}{2}$ N₂(g) + $\frac{1}{2}$ O₂(g) at the same T is :
 a) 0.02 b) 50 c) 4 × 10⁻⁴ d) 2.5 × 10⁻²

366. For the reaction,
 $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}(s) \rightleftharpoons \text{CuSO}_4 \cdot 3\text{H}_2\text{O}(s) + 2\text{H}_2\text{O}(l)$ which one is correct representation?
 a) $K_p = (P_{\text{H}_2\text{O}})^2$ b) $K_c = [\text{H}_2\text{O}]^2$ c) $K_p = K_c(RT)^2$ d) All of these
367. The correct order of increasing $[\text{H}_3\text{O}^+]$ in the following aqueous solutions is:
 a) $0.01\text{ M H}_2\text{S} < 0.01\text{ M H}_2\text{SO}_4 < 0.01\text{ M NaCl} < 0.01\text{ M NaNO}_2$
 b) $0.01\text{ M NaCl} < 0.01\text{ M NaNO}_2 < 0.01\text{ M H}_2\text{S} < 0.01\text{ M H}_2\text{SO}_4$
 c) $0.01\text{ M NaNO}_2 < 0.01\text{ M NaCl} < 0.01\text{ M H}_2\text{S} < 0.01\text{ M H}_2\text{SO}_4$
 d) $0.01\text{ M H}_2\text{S} < 0.01\text{ M NaNO}_2 < 0.01\text{ M NaCl} < 0.01\text{ M H}_2\text{SO}_4$
368. K_c for $A + B \rightleftharpoons C + D$ is 10 at 25°C . If a container contains 1, 2, 3 and 4 mole per litre of A, B, C and D respectively at 25°C , the reaction shall:
 a) Proceed from left to right
 b) Proceed from right to left
 c) Be at equilibrium
 d) None of the above
369. The compound whose 0.1 M solution is basic is
 a) Ammonium acetate b) Ammonium chloride
 c) Ammonium sulphate d) Sodium acetate
370. Isoelectric point is defined as the pH at which:
 a) An amino acid becomes acidic
 b) An amino acid becomes basic
 c) Zwitter ion has positive charge
 d) Zwitter ion has zero charge
371. The equilibrium constant of a reaction is 300. If the volume of reaction flask is tripled, the equilibrium constant is
 a) 300 b) 600 c) 900 d) 100
372. The oxoacid of SO_2 is:
 a) H_2SO_3 b) H_2SO_4 c) $\text{H}_2\text{S}_2\text{O}_8$ d) None of these
373. The solubility of CaF_2 in pure water is $2.3 \times 10^{-6}\text{ mol dm}^{-3}$. Its solubility product will be
 a) 4.8×10^{-18} b) 48.66×10^{-18} c) 4.9×10^{-11} d) 48.66×10^{-15}
374. pH value of which one of the following is not equal to one?
 a) 0.1 M HNO_3 b) 0.05 M H_2SO_4
 c) 0.1 M CH_3COOH d) 50 cm^3 of 0.4 M HCl + 50 cm^3 of 0.2 M NaOH
375. 50 mL of 2 N acetic mixed with 10 mL of 1N sodium acetate solution will have an approximate pH of ($K_a = 10^{-5}$):
 a) 4 b) 5 c) 6 d) 7
376. The solubility of AgCl in 0.2 M NaCl is (K_{sp} of $\text{AgCl} = 1.8 \times 10^{-10}$)
 a) $1.8 \times 10^{-11}\text{ M}$ b) $9 \times 10^{-10}\text{ M}$ c) $6.5 \times 10^{-12}\text{ M}$ d) $5.6 \times 10^{-11}\text{ M}$
377. The dissociation of water at 25°C is 1.9×10^{-7} percent and the density of water is 1.0 g/cm^3 . The ionisation constant of water is:
 a) 3.42×10^{-6} b) 3.42×10^{-8} c) 1.00×10^{-14} d) 2.00×10^{-16}
378. $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3 + \text{heat}$. What is the effect of the increase of temperature on the equilibrium of the reaction?
 a) Equilibrium is shifted to the left b) Equilibrium is shifted to the right
 c) Equilibrium is unaltered d) Reaction rate does not change
379. The expression for the solubility product of Ag_2CO_3 will be
 a) $K_{sp} s^2$ b) $K_{sp} 4s^3$ c) $K_{sp} 27s^4$ d) $K_{sp} s$

380. The solubility of AgCl in 0.2 M NaCl solution is
 $[K_{sp} \text{ of AgCl } 1.2 \times 10^{10}]$
 a) $6.0 \times 10^{10} \text{ M}$ b) 0.2 M c) $1.2 \times 10^{10} \text{ M}$ d) $0.2 \times 10^{10} \text{ M}$
381. Which of the following will not function as a buffer solution?
 (i) NaCl and NaOH
 (ii) NaOH and NH_4OH
 (iii) $\text{CH}_3\text{COONH}_4$ and HCl
 (iv) Borax and boric acid
 a) (i), (ii), (iii) b) (ii), (iii), (iv) c) (i), (iii), (iv) d) (i), (ii), (iii), (iv)
382. K_{sp} of salts AB , AB_2 and A_3B are 4.0×10^{-8} , 3.2×10^{-14} and 2.7×10^{-15} respectively at temperature T . The solubility order of these salts in water at temperature T (in mol litre $^{-1}$) is:
 a) $AB > AB_2 > A_3B$ b) $A_3B > AB_2 > AB$ c) $AB_2 > A_3B > AB$ d) $AB > A_3B > AB_2$
383. Which does not act as Bronsted acid?
 a) NH_4^+ b) CH_3COO^- c) HCO_3^- d) HSO_3^-
384. Which of the following solutions will have pH=9 at 298 K?
 a) $1 \times 10^{-9} \text{ M HCl solution}$ b) $1 \times 10^{-5} \text{ M NaOH solution}$
 c) $1 \times 10^{-9} \text{ M KOH solution}$ d) Both (a) and (b)
385. Acidosis is diagnosed when blood pH:
 a) Falls below 7.35 b) Rises above 7.45 c) Both (a) and (b) d) None of these
386. Which statement is false? (Assume complete dissociation in each case)
 a) If 2.0 L of a solution of H_2SO_4 contains 0.1 mole, then pH of the solution is 2
 b) The concentration of OH^- in 0.005 M HNO_3 is $2.0 \times 10^{-12} \text{ mol/L}$
 c) The pH of 0.01 M KOH is 12
 d) In a 0.001 M solution of NaOH the concentration of H^+ is 10^{-3} mol/L
387. 10 mL of a solution contains 0.1 M NH_4Cl + 0.01 M NH_4OH . Which addition would not change the pH of the solution?
 a) Adding 1 mL water
 b) Adding 5 mL of 0.1 M NH_4Cl
 c) Adding 5 mL of 0.1 M NH_4OH
 d) Adding 10 mL of 0.1 M NH_4Cl
388. Arrhenius theory of acid-base is not valid for:
 a) Aqueous solution
 b) In presence of water
 c) Non-aqueous solution
 d) None of these
389. The solubility in water of a sparingly soluble salt AB_2 is $1 \times 10^{-5} \text{ mol L}^{-1}$. Its solubility product number will be
 a) 4×10^{-15} b) 4×10^{-10} c) 1×10^{-15} d) 1×10^{-10}
390. The equilibrium constant (K_p) for the reaction, $\text{PCl}_5(\text{g}) \rightarrow \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$ is 16. If the volume of the container is reduced to one half its original volume, the value of K_p for the reaction at the same temperature will be
 a) 8 b) 16 c) 32 d) 64
391. The indicators used in the titration of iodine against sodium thiosulphate is
 a) Starch b) $\text{K}_3\text{Fe}(\text{CN})_6$ c) K_2CrO_4 d) Potassium
392. For the reaction,
 $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g}) - Q \text{ kJ}$, the equilibrium constant depends upon
 a) Temperature b) Pressure c) Catalyst d) Volume
393. In the dissociation of $2\text{HI} \rightleftharpoons \text{H}_2 + \text{I}_2$, the degree of dissociation will be influenced by the:

- d) Dissociation constant of weak base
407. If in the reaction $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$, x is that part of N_2O_4 which dissociates, then the number of molecules at equilibrium will be
 a) 1 b) 3 c) $1 + x$ d) $(1 + x)^2$
408. The $\text{p}K_a$ of a weak acid (HA) is 4.5. The pOH of an aqueous buffered solution of HA in which 50% of the acid ionised is
 a) 4.5 b) 2.5 c) 9.5 d) 7.0
409. Number of H^+ ions present in 10 mL of a solution of $\text{pH} = 3$ is:
 a) 10^{13} b) 6.02×10^{18} c) 6.02×10^{13} d) 6.02×10^{10}
410. The pH of gastric juice is normally:
 a) Greater than 1.5 and less than 1.2
 b) Less than 1.5
 c) Greater than 1 and less than 3
 d) Less than 1 and greater than zero
411. With reference to protonic acids, the correct statement is:
 a) PH_3 is more basic than NH_3
 b) PH_3 is less basic than NH_3
 c) PH_3 is amphoteric while NH_3 is basic
 d) None of the above
412. In a 500 mL flask, the degree of dissociation of PCl_5 at equilibrium is 40% and the initial amount is 5 moles. The value of equilibrium constant in mol L^{-1} for the decomposition of PCl_5 is
 a) 2.33 b) 2.66 c) 5.32 d) 4.66
413. The unit of ionic product of water (K_w) is:
 a) $\text{mol}^{-1}\text{litre}^{-1}$ b) $\text{mol}^{-1}\text{litre}^{-2}$ c) $\text{mol}^{-2}\text{litre}^{-1}$ d) $\text{mol}^2\text{litre}^{-2}$
414. In a reversible reaction two substance are in equilibrium. If the concentration each one is doubled, the equilibrium constant will be
 a) Reduced to one fourth of its original value b) Reduced to half of its original value
 c) Constant d) Doubled
415. Le-Chatelier principle is applicable only to a
 a) System in equilibrium b) System not in equilibrium
 c) Homogeneous reaction d) Heterogeneous reaction
416. The dissociation constant of a substituted benzoic acid at 25°C is 1.0×10^{-4} . The pH of 0.01 M solution of its sodium salt is
 a) 3 b) 7 c) 8 d) 6
417. A mixture of sodium oxide and calcium oxide are dissolved in water and saturated with excess carbon dioxide gas. The resulting solution is It contains
 a) Basic; NaOH and $\text{Ca}(\text{OH})_2$ b) Neutral; Na_2CO_3 and CaCO_3
 c) Basic; Na_2CO_3 and CaCO_3 d) Acidic; NaOH and CaCO_3
418. Conjugate acid of HF_2^- is
 a) H^+ b) HF c) F_2^- d) H_2F_2
419. Consider the reaction, $\text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH} \rightleftharpoons \text{CH}_3\text{COOC}_2\text{H}_5 + \text{H}_2\text{O}$. One mole each of CH_3COOH and $\text{C}_2\text{H}_5\text{OH}$ are heated in the presence of little concentrated H_2SO_4 . On equilibrium:
 a) 1 mole of ethyl acetate is formed
 b) 2 mole of ethyl acetate are formed
 c) $1/2$ mole of ethyl acetate is formed
 d) $2/3$ mole of ethyl acetate is formed
420. K_b for the hydrolysis reaction,
 $\text{B}^+ + \text{H}_2\text{O} \rightleftharpoons \text{BOH} + \text{H}^+$ is 1.0×10^{-6} , the hydrolysis constant of the salt is:

- a) 10^{-6} b) 10^{-7} c) 10^{-8} d) 10^{-9}
421. The pH of a 0.001 M NaOH will be
a) 3 b) 2 c) 11 d) 12
422. In lime kiln, the reversible reaction, $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$ proceeds to completion because :
a) Of high temperature
b) CO_2 escapes out
c) CaO is removed
d) Of low temperature
423. Solid $\text{Ba}(\text{NO}_3)_2$ is gradually dissolved in a 1.0×10^{-4} M Na_2CO_3 solution. At what concentration of Ba^{2+} will a precipitate begin to form?
(K_{sp} for $\text{BaCO}_3 = 5.1 \times 10^{-9}$)
a) 4.1×10^{-5} M b) 5.1×10^{-5} M c) 8.1×10^{-8} M d) 8.1×10^{-7} M
424. The solubility of AgCl is 0.0015 g/litre. The solubility product of AgCl will be:
a) 2×10^{-10} b) 1.1×10^{-10} c) 3.1×10^{-10} d) 4.1×10^{-10}
425. Although CO is neutral but it shows acidic nature on reaction with high P and T :
a) $\text{Ca}(\text{OH})_2$ b) NaOH c) $\text{Mg}(\text{OH})_2$ d) LiOH
426. The acidic character order for given oxy-acids of halogens is:
a) $\text{HOCl} > \text{HOBr} > \text{HOI}$ b) $\text{HOI} > \text{HOBr} > \text{HOCl}$ c) $\text{HOBr} > \text{HOCl} > \text{HOI}$ d) $\text{HOI} > \text{HOCl} > \text{HOBr}$
427. The stronger Bronsted base is:
a) ClO^- b) ClO_2^- c) ClO_3^- d) ClO_4^-
428. The indicator used in the titration of sodium carbonate with sulphuric acid is
a) Potassium ferrocyanide
b) Potassium ferricyanide
c) Methyl orange
d) Phenolphthalein
429. A cylinder fitted with a movable piston contains liquid water in equilibrium with water vapour at 25°C . Which operation result in a decrease in the equilibrium vapour pressure?
a) Moving the piston downward a short distance
b) Removing a small amount of vapour
c) Removing a small amount of the liquid water
d) Dissolving salt in the water
430. If acetic acid is mixed with sodium acetate then H^+ ion concentration will
a) Increase b) Decrease c) Remain unchanged d) pH decrease
431. The reverse process of neutralization is:
a) Hydrolysis b) Decomposition c) Dehydration d) Synthesis
432. The hydrogen ion concentration of a 10^{-8} M HCl aqueous solution at 298 K ($K_w = 10^{-14}$) is:
a) 9.525×10^{-8} M
b) 1.0×10^{-8} M
c) 1.0×10^{-6} M
d) 1.0525×10^{-7} M
433. The number of ions formed when cuprammonium sulphate dissolves in water is:
a) One b) Two c) Four d) Zero
434. A reversible chemical reaction have two reactants in equilibrium. If the concentrations of the reactants are doubled then the equilibrium constant will
a) Be halved b) Also be doubled c) Remains the same d) None of these
435. According to Arrhenius concept, base is a substance that:
a) Gives a pair of protons
b) Donates a proton
c) Accepts an electron pair

453. For the reaction
 $\text{CO(g)} + 0.5\text{O}_2\text{(g)} \rightarrow \text{CO}_2\text{(g)}$ K_p/K_c is equal to
 a) \sqrt{RT} b) $\frac{1}{\sqrt{RT}}$ c) 1 d) RT^2
454. Central metal ion in a complex or a cation acts as:
 a) Lewis base b) Lewis acid c) Bronsted acid d) Arrhenius acid
455. The degree of dissociation of CH_3COOH is influenced by:
 a) HCl b) CH_3COONa c) NH_4OH d) Either of these
456. Solubility of Ca(OH)_2 is $s \text{ mol L}^{-1}$. The solubility product (K_{sp}) under the same condition is
 a) $4s^3$ b) $3s^4$ c) $4s^2$ d) s^3
457. Which one of the following substances has the highest proton affinity?
 a) H_2O b) H_2S c) NH_3 d) PH_3
458. Which of the following is not a conjugate acid base pair?
 a) $\text{HPO}_3^{2-}, \text{PO}_3^{3-}$ b) $\text{H}_2\text{PO}_4^-, \text{HPO}_4^{2-}$ c) $\text{H}_3\text{PO}_4, \text{H}_2\text{PO}_4^-$ d) $\text{H}_2\text{PO}_4^-, \text{PO}_3^{3-}$
459. In an equilibrium reaction, if $\Delta G^\circ = 0$ the equilibrium constant, K should be equal to:
 a) 0 b) 1 c) 2 d) 10
460. The gastric juice in our stomach contains enough hydrochloride acid to make the hydrogen ion concentration about 0.01 mol/litre. The pH of the gastric juice is:
 a) 0.01 b) 1 c) 2 d) 14
461. At 550 K, the K_c for the following reaction is $10^4 \text{ mol}^{-1}\text{L}$
 $\text{X(g)} + \text{Y(g)} \rightleftharpoons \text{Z(g)}$
 At equilibrium, it was observed that
 $[\text{X}] = \frac{1}{2}[\text{Y}] = \frac{1}{2}[\text{Z}]$
 What is the value of $[\text{Z}]$ (in mol L^{-1}) at equilibrium?
 a) 2×10^{-4} b) 10^{-4} c) 2×10^4 d) 10^4
462. A vessel at 1000 K contains CO_2 with a pressure of 0.5 atm. Some of the CO_2 is converted into CO on the addition of graphite. If the total pressure at equilibrium is 0.8 atm, the value of K_p is
 a) 1.8 atm b) 3 atm c) 0.3 atm d) 0.18 atm
463. Ammonium carbonate decomposes as
 $\text{NH}_2\text{COONH}_4\text{(s)} \rightleftharpoons 2\text{NH}_3\text{(g)} + \text{CO}_2\text{(g)}$
 For the reaction, $K_p = 2.9 \times 10^{-5} \text{ atm}^3$. If we start with 1 mole of the compound, the total pressure at equilibrium would be
 a) 0.766 atm b) 0.0582 atm c) 0.0388 atm d) 0.0194 atm
464. The $\text{p}K_a$ of a weak acid, HA, is 4.80. The $\text{p}K_b$ of a weak base, BOH is 4.78. The pH of an aqueous solution of the corresponding salt, BA, will be
 a) 9.58 b) 4.79 c) 7.01 d) 9.22
465. The species among the following, which can act as an acid and a base is
 a) HSO_4^- b) SO_4^{2-} c) H_3O^+ d) Cl^-
466. A monoprotic weak acid (HA) is ionised 5% in 0.1 M aqueous solution. What is the equilibrium constant for its ionisation?
 $\text{HA(aq)} + \text{H}_2\text{O(l)} \rightleftharpoons \text{H}_3\text{O}^+\text{(aq)} + \text{A}^-\text{(aq)}$
 a) 2.63×10^4 b) 2.63×10^3 c) 2.63×10^{-4} d) 2.63×10^{-3}
467. Strongest acid is
 a) C_2H_6 b) CH_3OH c) $\text{HC} \equiv \text{HC}$ d) C_6H_6
468. Which of the following will have maximum pH?
 a) $\frac{\text{M}}{10} \text{ HCl}$ b) $\frac{\text{M}}{100} \text{ HCl}$ c) $\frac{\text{M}}{10} \text{ NaOH}$ d) $\frac{\text{M}}{100} \text{ NaOH}$

469. The pH of 1.0 M aqueous solution of a weak acid HA is 6.0. Its dissociation constant is:
 a) 10^{-6} b) 10^{-12} c) 1.0 d) 6.0
470. In which of the following, the reaction proceeds towards completion?
 a) $K = 1$ b) $K = 10$ c) $K = 10^2$ d) $K = 10^3$
471. The mixed salt among the following is:
 a) $\begin{matrix} \text{CHOHCOOK} \\ | \\ \text{CHOHCOONa} \end{matrix}$ b) NaKSO_4 c) CaOCl_2 d) All of these
472. What volume of M/10 NaOH added in 50 mL, M/10 acetic acid solution to get a buffer solution having highest buffer capacity?
 a) 50 mL b) 25 mL c) 10 mL d) 40 mL
473. The pH value of an acid is 5 and its concentration is 1 M. What is the value of K_a for the acid?
 a) 10^{-7} b) 10^{-5} c) 10^{-10} d) 10^{-8}
474. The pH of a neutral water sample is 6.5. Then the temperature of water
 a) is 25°C b) is more than 25°C
 c) is less than 25°C d) can be more or less than 25°C
475. The formation of phosgene is represented as, $\text{CO} + \text{Cl}_2 \rightleftharpoons \text{COCl}_2$. The reaction is carried out in 500 mL flask. At equilibrium 0.3 mole of phosgene, 0.1 mole of CO and 0.1 mole of Cl_2 are present. The equilibrium constant of the reaction is:
 a) 30 b) 15 c) 5 d) 3
476. In qualitative analysis, in order to detect second group basic radical, H_2S gas is passed in the presence of dilute HCl to
 a) Increase the dissociation of H_2S b) Decrease the dissociation of salt solution
 c) Decrease the dissociation of H_2S d) Increase the dissociation of salt solution
477. For two acids A and B $pK_a = 1.2$, and 2.8 respectively in value then which is true?
 a) A and B both are equally acidic b) A is stronger than B
 c) B is stronger than A d) Neither A nor B is strong
478. The weakest base among the following is:
 a) H^- b) CH_3^- c) CH_3O^- d) Cl^-
479. The hydrogen ion concentration in mol/litre of a solution of pH = 0 is:
 a) Zero b) 10^{-7} c) 1 M d) None of these
480. For which reaction is $K_p = K_c$?
 a) $2\text{NOCl}(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}) + \text{Cl}_2(\text{g})$
 b) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
 c) $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons 2\text{HCl}(\text{g})$
 d) $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$
481. For the reaction,
 $\text{H}_2(\text{g}) + \text{CO}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g})$, if the initial concentration of
 $[\text{H}_2] = [\text{CO}_2]$ and x mol/L of hydrogen is consumed at equilibrium, the correct expression of K_p is
 a) $\frac{x^2}{(1-x)^2}$ b) $\frac{(1+x)^2}{(1-x)^2}$ c) $\frac{x^2}{(2+x)^2}$ d) $\frac{x^2}{1-x^2}$
482. $K_{sp} = [A]^3[B]^2$ for the salt where A and B are the cation and anion as the case may be stand true for:
 a) $\text{Ca}_3(\text{PO}_4)_2$ b) As_2S_3 c) Bi_2S_3 d) All are correct
483. The dissociation constant for acetic acid and HCN at 25°C are 1.5×10^{-3} and 4.5×10^{-10} respectively, the equilibrium constant for the equilibrium,
 $\text{CN}^- + \text{CH}_3\text{COOH} \rightleftharpoons \text{HCN} + \text{CH}_3\text{COO}^-$ would be:
 a) 3.0×10^4 b) 3.0×10^5 c) 3.0×10^{-5} d) 3.0×10^{-4}
484. The pH of the solution

- b) Favours the backward rate only
 c) Favours both the forward and backward rates as the case may be
 d) Favours neither the forward nor backward rates
499. Zn salt is mixed with $(\text{NH}_4)_2\text{S}$ of molarity 0.021 M . The amount of Zn^{2+} remains unprecipitated in 12 mL of this solution (K_{sp} of $\text{ZnS} = 4.51 \times 10^{-24}$) which is:
 a) $1.677 \times 10^{-22}\text{g}$ b) $1.767 \times 10^{-22}\text{g}$ c) $2.01 \times 10^{-23}\text{g}$ d) None of these
500. Pure ammonia is placed in a vessel at temperature where its dissociation constant (α) is appreciable. At equilibrium,
 a) K_p does not change significantly with pressure
 b) α does not change with pressure
 c) Concentration of NH_3 does not change with pressure
 d) Concentration of H_2 is less than that of N_2
501. A chemical reaction $A \rightleftharpoons B$ is said to be at equilibrium when:
 a) Complete conversion of A to B has taken place
 b) Conversion of A to B is only 50% complete
 c) Only 10% conversion of A to B has taken place
 d) The rate of transformation of A and B is just equal to the rate of transformation of B to A in the system
502. As the temperature increases, the pH of a KOH solution
 a) Will decrease
 b) Will increase
 c) Remains constant
 d) Depends upon the concentration of KOH solution
503. Tribasic acid furnishes...type of anions.
 a) 2 b) 1 c) 3 d) 4
504. Which of the following pairs constitutes buffer?
 a) HNO_3 and NH_4NO_3 b) HCl and KCl c) HNO_2 and NaNO_2 d) NaOH and NaCl
505. What will be the pH and % α respectively for the salt BA of 0.1 M concentration? Given, K_a for $HA = 10^{-6}$ and K_b for $BOH = 10^{-6}$
 a) 7, 10% b) 5, 10% c) 5, 0.1% d) 7, 1%
506. The indicator used in titrating oxalic acid with caustic soda solution is:
 a) Methyl orange b) Methyl red c) Fluorescein d) Phenolphthalein
507. The pH of a 0.02 M solution of hydrochloric acid is:
 a) 2.0 b) 1.7 c) 0.3 d) 2.2
508. For the reaction, $\text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}_2(\text{g}) + \text{H}_2(\text{g})$, at a given temperature, the equilibrium amount of $\text{CO}_2(\text{g})$ can be increased by
 a) Increasing the amount of $\text{CO}(\text{g})$ b) Decreasing the volume of the container
 c) Adding a suitable catalyst d) Adding an inert gas
509. $\text{Ag}^+ + \text{NH}_3 \rightleftharpoons [\text{Ag}(\text{NH}_3)]^+$; $K_1 = 3.5 \times 10^{-3}$
 $[\text{Ag}(\text{NH}_3)]^+ + \text{NH}_3 \rightleftharpoons [\text{Ag}(\text{NH}_3)_2]^+$;
 $K_2 = 1.7 \times 10^{-3}$
 Then the formation constant of $[\text{Ag}(\text{NH}_3)_2]^+$ is
 a) 6.08×10^{-6} b) 6.08×10^6 c) 6.08×10^{-9} d) None of these
510. 1 mL of 0.1 N HCl is added to 999 mL solution of NaCl . The pH of the resulting solution will be:
 a) 7 b) 4 c) 2 d) 1
511. The role of catalyst in reversible reaction is :
 a) To increase the rate of forward reaction
 b) Decrease the rate after equilibrium
 c) Allow equilibrium to be achieved quickly
 d) None of the above

512. Which one is correct representation for,
 $2\text{SO}_3 \rightleftharpoons 2\text{SO}_2 + \text{O}_2$?
- a) $K_p = \frac{(p_{\text{SO}_2})^2 (p_{\text{O}_2})}{(p_{\text{SO}_3})^2}$
 b) $K_c = \frac{[\text{SO}_2]^2 [\text{O}_2]}{[\text{SO}_3]^2}$
 c) $K_p = \frac{(\text{mole of SO}_2)^2 \times (\text{mole of O}_2)}{(\text{mole of SO}_3)^2} \times \left[\frac{P}{\text{total mole at equilibrium}} \right]$
 d) All of the above
513. The solubility of $\text{Ca}_3(\text{PO}_4)_2$ in water is y moles/litre. Its solubility product is
 a) $6y^4$ b) $36y^4$ c) $64y^5$ d) $108y^5$
514. For the reaction $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$, the equilibrium constant is K_1 , the equilibrium constant is K_2 , for the reaction $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$. What is K for the reaction $\text{NO}_2(\text{g}) \rightleftharpoons \frac{1}{2}\text{N}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g})$?
 a) $\frac{1}{(K_1 K_2)}$ b) $\frac{1}{(2K_1 K_2)}$ c) $\frac{1}{(4K_1 K_2)}$ d) $\left(\frac{1}{(K_1 K_2)} \right)^{1/2}$
515. The equilibrium constant for the reaction $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$ at temperature T is 4×10^{-4} . The value of K_c for the reaction $\text{NO}(\text{g}) \rightleftharpoons \frac{1}{2}\text{N}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g})$ at the same temperature is
 a) 25 b) 50 c) 75 d) 100
516. If the value of K_c for an equilibrium reaction is 10^{-4} , then the reaction is in
 a) Backward direction b) Forward direction
 c) Equilibrium d) Reaction is not possible
517. 5 moles of X are mixed with 3 moles of Y . At equilibrium for the reaction, $X + Y \rightleftharpoons Z$, 2 moles of Z are formed. The equilibrium constant for the reaction will be
 a) $\frac{2}{3}$ b) $\frac{1}{2}$ c) $\frac{3}{2}$ d) $\frac{1}{4}$
518. If in the reaction : $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$, α is degree of dissociation of N_2O_4 , then the number of molecules at equilibrium will be:
 a) 3 b) 1 c) $(1 - \alpha)^2$ d) $(1 + \alpha)$
519. The hydroxide having the lowest value of K_{sp} at 25°C is:
 a) $\text{Mg}(\text{OH})_2$ b) $\text{Ca}(\text{OH})_2$ c) $\text{Ba}(\text{OH})_2$ d) $\text{Be}(\text{OH})_2$
520. The effect of increasing the pressure on the equilibrium $2A + 3B \rightleftharpoons 3A + 2B$ is
 a) Forward reaction is favoured b) Backward reaction is favoured
 c) No effect d) None of the above
521. Conjugate base of hydrazoic acid is:
 a) HN_3^- b) N_3^- c) N^{3-} d) N_2^-
522. A reaction is, $A + B \rightarrow C + D$. Initially we start with equal concentrations of A and B . At equilibrium, we find the moles of C are two times of A . What is the equilibrium constant of the reaction?
 a) 2 b) 4 c) $\frac{1}{2}$ d) $\frac{1}{4}$
523. Which of the following is the strongest conjugate base?
 a) Cl^- b) CH_3COO^- c) SO_4^{2-} d) NO_2^-
524. Consider the following equilibrium in a closed container
 $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$
 At a fixed temperature, the volume of the reaction container is halved. For this change which of the following statement holds true regarding the equilibrium constant (K_p) and degree of dissociation (α) ?
 a) Neither K_p nor α changes b) Both K_p and α changes
 c) K_p changes but α does not d) K_p does not change but α changes

525. If K_a for a weak acid is 10^{-5} , pK_b value of its conjugate base is:
 a) 5 b) 6 c) 7 d) 9
526. According to law of mass action, for the reaction
 $2A + B \rightarrow \text{Products}$
 a) Rate = $k[A][B]$ b) Rate = $k[A]^2[B]$ c) Rate = $k[A][B]^2$ d) Rate = $k[A]^{1/2}[B]$
527. The equilibrium constant for the reaction, $2X(g) + Y(g) \rightleftharpoons 2Z(g)$ is $2.25 \text{ litre mol}^{-1}$. What would be the concentration of Y at equilibrium with 2.0 mole of X and 3.0 mole of Z in one litre vessel?
 a) 1.0 M b) 2.25 M c) 2.0 M d) 4.0 M
528. $A(g) + B(g) \rightleftharpoons AB(g)$ is a reversible reaction. At equilibrium 0.4 mole of AB is formed when each A and B are taken one mole. How much of A changes into AB ?
 a) 20% b) 40% c) 60% d) 4%
529. A solute undergoes complex formation with ions of sparingly soluble salt, the solubility of salt:
 a) Increases b) Decreases c) Is unaffected d) Either of these
530. The pH of the solution containing 0.1 N NaOH and 0.1N NH_4OH is:
 a) 1 b) 7 c) 2 d) 13
531. Which buffer solution comprising of the following has its pH value greater than 7?
 a) $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$ b) $\text{HCOOH} + \text{HCOOK}$
 c) $\text{CH}_3\text{COONH}_4$ d) $\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$
532. Which of the following behaves as both Lewis and Bronsted base?
 a) BF_3 b) Cl^- c) CO d) None of these
533. If solubility of calcium hydroxide is $\sqrt{3}$, then its solubility product will be
 a) 27 b) 3 c) 9 d) $12\sqrt{3}$
534. Pure water is kept in a vessel and it remains exposed to atmospheric CO_2 which is absorbed. Then the pH will be:
 a) Greater than 7
 b) Less than 7
 c) 7
 d) Depends on ionic product of water
535. The decomposition of N_2O_4 to NO_2 is carried out at 280 K in chloroform. When equilibrium has been established, 0.2 mole of N_2O_4 and 2×10^{-3} mole of NO_2 are present in 2 L solution. The equilibrium constant for reaction $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$ is
 a) 1×10^{-2} b) 1×10^{-3} c) 1×10^{-4} d) 1×10^{-5}
536. The most favourable condition for the manufacture of NH_3 is
 a) high temperature and high pressure b) low temperature and low pressure
 c) high temperature and low pressure d) low temperature and high pressure
537. The value of K_p for the following reaction $2\text{H}_2\text{S}(g) \rightleftharpoons 2\text{H}_2(g) + \text{S}_2(g)$, is 1.2×10^{-2} at 106.5°C . The value of K_c for this reaction is
 a) $= 1.2 \times 10^{-2}$ b) $< 1.2 \times 10^{-2}$ c) $> 1.2 \times 10^{-2}$ d) None of these
538. In hydrolysis of a salt of weak acid and strong base $A^- + \text{H}_2\text{O} \rightleftharpoons \text{HA} + \text{OH}^-$, the hydrolysis constant (K_h) is equal to...
 a) $\frac{K_w}{K_a}$ b) $\frac{K_w}{K_b}$ c) $\sqrt{\frac{K_a}{C}}$ d) $\frac{K_w}{K_a \times K_b}$
539. The pH of $7 \times 10^{-8} \text{ M}$ CH_3COOH is:
 a) 8.1 b) 7.9 c) 7.1 d) 6.85
540. Water acts as an acid in presence of:
 a) NH_3 b) H_2SO_4 c) C_6H_6 d) HCl
541. Which statement is/are correct?
 a) All Bronsted bases are also Lewis bases

- b) All Bronsted acids are not Lewis acids
 c) All cations are acids and all anions are bases
 d) All of the above
542. Three moles of PCl_5 , three moles of PCl_3 and two moles of Cl_2 are taken in a closed vessel. If at equilibrium the vessel has 1.5 moles of PCl_5 , the number of moles of PCl_3 present in it is
 a) 5 b) 3 c) 6 d) 4.5
543. Ice and water are in equilibrium at 273 K, which of the following statements is correct?
 a) $G_{(\text{ice})} > G_{(\text{H}_2\text{O})}$ b) $G_{(\text{ice})} < G_{(\text{H}_2\text{O})}$ c) $G_{(\text{ice})} = G_{(\text{H}_2\text{O})} = 0$ d) $G_{(\text{ice})} = G_{(\text{H}_2\text{O})} \neq 0$
544. For an equilibrium reaction involving gases, the forward reaction is 1st order while the reverse reaction is 2nd order. The units of K_p for the forward equilibrium is:
 a) atm b) atm^2 c) atm^{-1} d) atm^{-2}
545. A buffer solution has equal volumes of 0.2 M NH_4OH and 0.02 M NH_4Cl . The pK_b of the base is 5. The pH is
 a) 10 b) 9 c) 4 d) 7
546. Which of the following will suppress the ionisation of acetic acid in aqueous solution?
 a) NaCl b) HCl c) KCl d) Unpredictable
547. The Henderson's equation used to calculate the pOH of basic buffer is:
 a) $\text{pOH} = \text{p}K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$
 b) $\text{pOH} = \text{p}K_b - \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$
 c) $\text{pOH} = \text{p}K_b + \log \frac{[\text{Base}]}{[\text{Conjugate acid}]}$
 d) $\text{pOH} = \text{p}K_b - \log \frac{[\text{Base}]}{[\text{Conjugate acid}]}$
548. Which of the following is the weakest acid?
 a) HCl b) HF c) H_2SO_4 d) HNO_3
549. How many gram of NaOH must be present in one litre of the solution to give it a pH = 12?
 a) 0.20 g litre⁻¹ b) 0.4 g litre⁻¹ c) 4.0 g litre⁻¹ d) 0.10 g litre⁻¹
550. Approximate pH of 0.10 M aqueous H_2S solution having K_1 and K_2 for H_2S at 25°C 10^{-7} and 10^{-13} respectively, is:
 a) 4 b) 5 c) 9 d) 8
551. An aqueous solution whose pH is zero will be called as
 a) Acidic b) Basic c) Neutral d) Amphoteric
552. The concentration of KI and KCl in a certain solution containing both is 0.001 M each. If 20 mL of this solution is added to 20 mL of a saturated solution of AgI in water. What will happen? ($K_{\text{sp}} \text{AgCl} = 10^{-10}$, $K_{\text{sp}} \text{AgI} = 10^{-16}$)
 a) AgI will be precipitated b) AgCl will be precipitated
 c) There will be no precipitate d) Both AgCl and AgI will be precipitated
553. The equilibrium reaction that is not influenced by volume change at constant temperature is
 a) $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ b) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
 c) $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ d) $2\text{NO}(\text{g}) + \text{O}_2 \rightleftharpoons 2\text{NO}_2(\text{g})$
554. Group IA metals react violently with water to produce ... nature in solution.
 a) Acidic b) Basic c) Amphoteric d) Neutral
555. $A + B = C + D$. If initially the concentration of A and B are both equal but at equilibrium, concentration of D will be twice of that of A, then what will be the equilibrium constant of reaction?

- a) $\frac{4}{9}$ b) $\frac{9}{4}$ c) $\frac{1}{9}$ d) 4

556. The phenomenon of interaction of anions and cations furnished by an electrolyte with the H^+ and OH^- ions of water to produce acidic nature or alkalinity is known as hydrolysis. In hydrolysis:

- a) The pH may either increase or decrease
 b) All the salts (except those made up with strong acid and base) undergo hydrolysis
 c) The variation of pH depends upon the nature of salts as well as on the temperature
 d) All of the above

557. $A_2(g) + B_2(g) \rightleftharpoons 2AB(g)$, $\Delta H = -ve$

The reaction

- a) Remains unaffected by pressure b) Occurs at 1000 atm pressure
 c) Occurs at high temperature d) Occurs at high pressure and high temperature

558. 20 mL of 0.5 N HCl and 35 mL of 0.1 N NaOH are mixed. The resulting solution will

- a) Be neutral b) Be basic
 c) Turn phenolphthalein solution pink d) Turn methyl orange red

559. A solution which is $10^{-3}M$ each in Mn^{2+} , Fe^{2+} , Zn^{2+} and Hg^{2+} is treated with $10^{-16}M$ sulphide ion. If K_{sp} of MnS, FeS, ZnS and HgS are 10^{-15} ,

10^{-23} , 10^{-20} and 10^{-54} respectively, which one will precipitate first?

- a) FeS b) MgS c) HgS d) ZnS

560. Ostwald's dilution law is applicable for

- a) Weak electrolyte b) Strong electrolyte
 c) Both weak and strong electrolyte d) Non-electrolyte

561. The equilibrium $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$ shows that K_p is 24.63 times the value of K_c at a particular temperature T . Then T (in K) is:

- a) 200 b) 100 c) 300 d) 400


562. At $90^\circ C$, pure water has $[H^+] = 10^{-6}M$, if 100 mL of 0.2 M HCl is added to 200 mL of 0.1 M KOH at $90^\circ C$ then pH of the resulting solution will be

- a) 7 b) 8 c) 4 d) 6

563. In a reaction, the rate of reactions is proportional to its active mass. This statement is known as

- a) Law of mass-action b) Le-Chatelier principle
 c) Faraday law of electrolysis d) Law of constant proportion

564. The values of K_{p1} and K_{p2} for the reactions



are in the ratio 9 : 1. If degree of dissociation of X and A be equal, then total pressure at equilibrium (1) and (2) are in the ratio :

- a) 1 : 9 b) 36 : 1 c) 1 : 1 d) 3 : 1

565. Which of the following is not a characteristic property of chemical equilibrium?

- a) Rate of forward reaction is equal to rate of backward reaction at equilibrium
 b) After reaching the chemical equilibrium, the concentrations of reactants and products remain unchanged with time
 c) For $A(g) \rightleftharpoons B(g)$, K_c is 10^{-2} . If this reaction is carried out in the presence of catalyst, the value of K_c decreases
 d) After reaching the equilibrium, both forward and backward reactions continue to take place

566. Which molecule is an electron donor?

- a) NH_3 b) BF_3 c) PF_5 d) AsF_5

567. The correct expression for the solubility product of $Ca_3(PO_4)_2$ is

- a) $108 s^5$ b) $27 s^5$ c) $16 s^4$ d) $81 s^4$
568. pH of solution can be expressed as
 a) $-\log_e[H^+]$ b) $-\log_{10}[H^+]$ c) $\log_e[H^+]$ d) $\log_{10}[H^+]$
569. Which aqueous solution will have pH less than 7?
 a) KNO_3 b) $NaOH$ c) $NaCN$ d) $FeCl_3$
570. If the salts M_2X , QY_2 and PZ_3 have the same solubilities, K_{sp} values are related as:
 a) $K_{sp}(M_2X) = K_{sp}(QY_2) < K_{sp}(PZ_3)$
 b) $K_{sp}(M_2X) > K_{sp}(QY_2) = K_{sp}(PZ_3)$
 c) $K_{sp}(M_2X) = K_{sp}(QY_2) = K_{sp}(PZ_3)$
 d) $K_{sp}(M_2X) > K_{sp}(QY_2) > K_{sp}(PZ_3)$
571. The pH values of 0.1 M solution of HCl , CH_3COOH , NH_4Cl and CH_3COONa will have the order
 a) $HCl < CH_3COOH < NH_4Cl < CH_3COONa$ b) $CH_3COONa < NH_4Cl < CH_3COOH < HCl$
 c) $NH_4Cl < CH_3COONa < CH_3COOH < HCl$ d) All will have same of pH value
572. When rain is accompanied by a thunderstorm the collected rain water will have a pH value:
 a) Uninfluenced by occurrence of thunderstorm
 b) Depending on the amount of dust in air
 c) Slightly lower than that of rain water without thunderstorm
 d) Slightly higher than that when the thunderstorm is not there
573. Ammonia gas dissolves in water to form NH_4OH . In this reaction water acts as
 a) A conjugate base b) A non-polar solvent c) An acid d) A base
574. pH scale was introduced by
 a) Arrhenius b) Sorensen c) Lewis d) Lowry
575. Given : $[Ag(NH_3)_2]^+ \rightleftharpoons Ag(NH_3)^+ + NH_3$; $K_1^0 = 1.4 \times 10^{-4}$
 $Ag(NH_3)^+ \rightleftharpoons Ag^+ + NH_3$; $K_2^0 = 4.3 \times 10^{-4}$
 The instability constant of the complex $Ag(NH_3)_2^+$ is equal to :
 a) 7.14×10^3 b) 2.33×10^3 c) 6.02×10^{-8} d) 1.66×10^7
576. An aqueous solution contains Ni^{2+} , Co^{2+} and Pb^{2+} ions at equal concentrations. The solubility product of NiS , PbS and CoS in water at $25^\circ C$ are 1.4×10^{-24} , 3.4×10^{-28} and 3×10^{-26} , respectively. Indicate which of these ions will be precipitated first and last when sulphide concentration is progressively increased from zero?
 a) NiS and PbS b) NiS and CoS c) CoS and NiS d) PbS and NiS
577. 10^{-6} M $NaOH$ is diluted 100 times. The pH of the diluted base is
 a) Between 7 and 8 b) Between 5 and 6 c) Between 6 and 7 d) Between 10 and 11
578. An acid with molecular formula $C_7H_6O_3$ forms three types of sodium salts, i. e., $C_7H_5O_3Na$, $C_7H_4O_3Na_2$ and $C_7H_3O_3Na_3$. The basicity of the acid is:
 a) One b) Two c) Three d) Four
579. If the concentration of CrO_4^{2-} ion in a saturated solution of silver chromate be 2×10^{-4} M, solubility product of silver chromate will be:
 a) 4×10^{-8} b) 8×10^{-12} c) 32×10^{-12} d) 6×10^{-12}
580. What would be the effect on addition of $CaCO_3$ on the equilibrium of $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$ in a closed container:
 a) conc. of CO_2 increases
 b) conc. of CO_2 decreases
 c) Remains unaffected
 d) Cannot be predicted
581. In a chemical equilibrium, the rate constant of the backward reaction is 7.5×10^{-4} and the equilibrium constant is 1.5. So, the rate constant of the forward reaction is
 a) 1.125×10^{-3} b) 2.225×10^{-3} c) 3.335×10^{-5} d) 1.125×10^{-1}

582. 0.1 M acetic acid solution is titrated against 0.1 M NaOH solution. What would be the difference in pH between 1/4 and 3/4 stages of neutralization of acid?
 a) $2 \log 3/4$ b) $2 \log 1/4$ c) $\log 1/3$ d) $2 \log 3$
583. For a concentrated solution of a weak electrolyte A_xB_y of concentration 'C', the degree of dissociation ' α ' is given as
 a) $\alpha = \sqrt{K_{eq}/C(x+y)}$ b) $\alpha = \sqrt{K_{eq}C/(xy)}$
 c) $\alpha = (K_{eq}/C^{x+y-1} x^x y^y)^{1/(x+y)}$ d) $\alpha = (K_{eq}/Cxy)$
584. The values of dissociation constant of some acids (at 25°C) are given below. The strongest acid in water is:
 a) 1.4×10^{-2} b) 1.6×10^{-4} c) 4.4×10^{-10} d) 4.3×10^{-7}
585. In a mixture of acetic acid and sodium acetate the ratio of concentration of the salt to the acid is increased ten times. Then, the pH of the solution
 a) Increases by one b) Decreases by one c) Decreases ten fold d) Increased ten fold
586. Which of the following salts is most soluble?
 a) $Bi_2S_3(K_{sp} = 1 \times 10^{-17})$ b) $MnS(K_{sp} = 7 \times 10^{-16})$
 c) $CuS(K_{sp} = 8 \times 10^{-37})$ d) $Ag_2S(K_{sp} = 6 \times 10^{-51})$
587. van't Hoff's equation giving the effect of temperature on chemical equilibrium is represented as:
 a) $\frac{d \ln F}{dT} = \frac{\Delta H}{RT^2}$ b) $\frac{d \ln K_p}{dT} = \frac{\Delta HT^2}{R}$ c) $\frac{d \ln K_p}{dT} = \frac{\Delta H}{RT^2}$ d) $\frac{d \ln K_p}{dT} = \frac{RT^2}{\Delta H}$
588. Hydrogen and oxygen were heated together in a closed vessel. The equilibrium constant is found to decrease after 2000°C. Which is responsible for this?
 a) Backward reaction predominates
 b) Forward reaction predominates
 c) Both forward and backward reactions have same rate
 d) It is a property of the system, hence no reason for lower value
589. The dissociation constant of HCN is 1.3×10^{-9} . The value of hydrolysis constant of KCN will be:
 a) 1.3×10^{-9} b) 10^{-14} c) 7.7×10^{-5} d) 0.77×10^{-5}
590. Solubility product of silver bromide is 5.0×10^{-13} . The quantity of potassium bromide (molar mass taken as 120 g mol^{-1}) to be added to 1 L of 0.05 M solution of silver nitrate to start the precipitation of AgBr is
 a) $1.2 \times 10^{10} \text{ g}$ b) $1.2 \times 10^9 \text{ g}$ c) $6.2 \times 10^5 \text{ g}$ d) $5.0 \times 10^8 \text{ g}$
591. In the thermal decomposition of potassium chlorate given as $2KClO_3 \rightarrow 2KCl + 3O_2$, law of mass action
 a) Can be applied
 b) Cannot be applied
 c) Can be applied at low temperature
 d) Can be applied at high temperature and pressure
592. The solubility product of a sparingly soluble salt AB at room temperature is 1.21×10^{-6} . Its molar solubility is
 a) 1.21×10^{-6} b) 1.21×10^{-3} c) 1.1×10^{-4} d) 1.1×10^{-3}
593. What is the correct representation for the solubility product of SnS_2 ?
 a) $[Sn^{2+}][S^{2-}]^2$ b) $[Sn^{4+}][S^{2-}]^2$ c) $[Sn^{2+}][2S^{2-}]$ d) $[Sn^{4+}][2S^{2-}]^2$
594. Which of the following pK_a values, represent the strongest acid?
 a) 10^{-4} b) 10^{-8} c) 10^{-5} d) 10^{-2}
595. An electrolyte
 a) Gives complex ions in solution b) Dissolves in water to give ions
 c) Is ionised in the solid state d) Generates ions on passing electric current
596. The strongest base among the following is:
 a) CH_3^- b) F^- c) NH_2^- d) OH^-

608. Two system $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$ and $\text{COCl}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{Cl}_2(\text{g})$ are simultaneously in equilibrium in a vessel at constant volume. If some CO is introduced into the vessel then at the new equilibrium the concentration of:
- PCl_5 is greater
 - PCl_3 remains unchanged
 - PCl_5 is less
 - Cl_2 is greater
609. The solubility product of Ag_2CrO_4 in water at 298 K is 3.2×10^{-11} . What will be the concentration of CrO_4^{2-} ions in the saturated solution of Ag_2CrO_4 ?
- $2 \times 10^{-4} \text{ M}$
 - $5.7 \times 10^{-5} \text{ M}$
 - $5.7 \times 10^{-6} \text{ M}$
 - $3.2 \times 10^{-11} \text{ M}$
610. Sulphide ions in alkaline solution react with solid sulphur to form polyvalent sulphide ions. The equilibrium constant for the formation of S_2^{2-} and S_3^{2-} from S and S^{2-} ions are 1.7 and 5.3 respectively. Equilibrium constant for the formation of S_3^{2-} from S_2^{2-} and S is:
- 1.33
 - 3.11
 - 4.21
 - 1.63
611. The pH of water is 7 at 25°C. If water is heated to 50°C, which of the following should be true?
- pH will decrease
 - pH will increase
 - pH will remain seven
 - $[\text{H}^+]$ will increase but $[\text{OH}^-]$ will decrease
612. The conjugate acid of NH_2^- is
- N_2H_4
 - NH_4^+
 - NH_2OH
 - NH_3
613. pH of a solution of the mixture of 0.1 N HCl and 0.1 N CH_3COOH is:
- 1
 - 2
 - 1.7
 - None of these
614. A buffer solution is prepared by mixing 10 mL of 1.0 M acetic and 20 mL of 0.5 M sodium acetate and then diluted to 100 mL with distilled water. If the $\text{p}K_a$ of CH_3COOH is 4.76, what is the pH of the buffer solution prepared?
- 5.21
 - 4.76
 - 4.34
 - 5.21
615. $\text{CH}_3\text{COOH}(\text{l}) + \text{C}_2\text{H}_5\text{OH}(\text{l}) \rightleftharpoons \text{CH}_3\text{COOC}_2\text{H}_5(\text{l}) + \text{H}_2\text{O}(\text{l})$
In the above reaction, one mole of each of acetic acid and alcohol are heated in the presence of little conc. H_2SO_4 . On equilibrium being attained
- 1 mole of ethyl acetate is formed
 - 2 mole of ethyl acetate is formed
 - $\frac{2}{3}$ mole of ethyl acetate is formed
 - $\frac{1}{2}$ mole of ethyl acetate is formed
616. When the pH of a solution is 2, the hydrogen ion concentration is:
- $1 \times 10^{-14} \text{ M}$
 - $1 \times 10^{-2} \text{ M}$
 - $1 \times 10^{-7} \text{ M}$
 - $1 \times 10^{-12} \text{ M}$
617. On adding A to the reaction at equilibrium, $\text{AB}(\text{s}) \rightleftharpoons \text{A}(\text{g}) + \text{B}(\text{g})$, the new equilibrium concentration of A becomes double, the equilibrium concentration of B would become :
- 1/2 of its original value
 - 1/4 of its original value
 - 1/8 of its original value
 - Twice of its original value
618. The change in pressure will not affect the equilibrium constant for
- $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$
 - $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$
 - $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$
 - All of these
619. Equilibrium concentration of HI, I_2 and H_2 is 0.7, 0.1 and 0.1 M respectively. The equilibrium constant for the reaction, $\text{I}_2 + \text{H}_2 \rightleftharpoons 2\text{HI}$ is :
- 0.36
 - 36
 - 49
 - 0.49
620. The dissociation constant of acetic acid K_a is 1.74×10^{-5} at 298 K. The pH of a solution of 0.1 M acetic acid is
- 2.88
 - 3.6
 - 4.0
 - 1.0
621. What is the effect of increasing pressure on the dissociation of PCl_5 according to the equation

- $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g}) - x \text{ cal?}$
- a) Dissociation decreases
b) Dissociation increases
c) Dissociation does not change
d) None of the above
622. The ionic product of water at 60°C is 9.61×10^{-14} . The pH of water at 60°C is:
a) 6.51
b) 6.70
c) 9.61
d) 7.0
623. For the reaction equilibrium, $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$, the concentration of N_2O_4 and NO_2 at equilibrium are 4.8×10^{-2} and 1.2×10^{-2} mol/L respectively. The value of K_c for the reaction is
a) 3×10^{-3} mol/L
b) 3.3×10^{-3} mol/L
c) 3×10^{-1} mol/L
d) 3.3×10^{-1} mol/L
624. If first dissociation of $\text{X}(\text{OH})_3$ is 100% where as second dissociation is 50% and third dissociation is negligible then the pH of 4×10^{-3} M $\text{X}(\text{OH})_3$ is
a) 7.5
b) 9.54
c) 11.78
d) 13.25
625. The equilibrium constant for the reaction, $3\text{C}_2\text{H}_2 \rightleftharpoons \text{C}_6\text{H}_6$ is 4.0 at T K. If the equilibrium concentration of C_2H_2 is 0.5 mol/litre, the concentration of C_6H_6 is :
a) 0.5 M
b) 1.5 M
c) 5×10^{-2} M
d) 0.25 M
626. For the reaction, $\text{CH}_4(\text{g}) + 2\text{O}_2(\text{g}) \rightleftharpoons \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$,
 $\Delta_r H = -170.8 \text{ kJ mol}^{-1}$
- Which of the following statements is not true?
a) Addition of $\text{CH}_4(\text{g})$ or $\text{O}_2(\text{g})$ at equilibrium will cause a shift to the right
b) The reaction is exothermic
c) At equilibrium, the concentrations of $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\text{l})$ are not equal
d) The equilibrium constant for the reaction is given by $K_p = \frac{[\text{CO}_2]}{[\text{CH}_4][\text{O}_2]}$
627. The law of mass action was enunciated by
a) Graham
b) Bodestein
c) Rutherford
d) Guldberg and Waage
628. The correct statement about buffer solution is:
a) It contains a weak acid and its conjugate base
b) It contains a weak base and its conjugate acid
c) It shows little change in pH on adding small amount of an acid or base
d) All of the above
629. Which reaction has $\Delta n = 2$?
a) $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
b) $3\text{Fe}(\text{s}) + 4\text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{Fe}_3\text{O}_4(\text{s}) + 4\text{H}_2(\text{g})$
c) $\text{NH}_4\text{Cl}(\text{g}) \rightleftharpoons \text{NH}_3(\text{g}) + \text{HCl}(\text{g})$
d) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}(\text{s}) \rightleftharpoons \text{CuSO}_4 \cdot 3\text{H}_2\text{O}(\text{s}) + 2\text{H}_2\text{O}(\text{g})$
630. In the reaction, $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$, α is that part of N_2O_4 which dissociates, then the number of moles at equilibrium will be
a) 1
b) 3
c) $(1 + \alpha)$
d) $(1 - \alpha)^2$
631. Decreasing acid strengths of HI, HBr, HCl and HF is:
a) $\text{HF} > \text{HCl} > \text{HBr} > \text{HI}$
b) $\text{HI} > \text{HBr} > \text{HCl} > \text{HF}$
c) $\text{HI} > \text{HCl} > \text{HBr} > \text{HF}$
d) $\text{HI} > \text{HF} > \text{HCl} > \text{HBr}$
632. A monoprotic acid in 0.1 M solution has $K_a = 1.0 \times 10^{-5}$. The degree of dissociation acid is:
a) 1.0%
b) 99.9%
c) 0.1%
d) 99%
633. In what manner will increase of pressure affect the following equation?
 $\text{C}(\text{s}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{H}_2(\text{g})$
a) Shift in the reverse direction
b) Shift in the forward direction
c) Increase in the yield of hydrogen
d) No effect
634. Which one of the following pair shows buffer's solution?
a) $\text{NaCl} + \text{NaOH}$
b) $\text{CH}_3\text{COONa} + \text{CH}_3\text{COOH}$

- c) These give unionised acid or base on reaction with added acid or alkali
 d) Acids and alkalies in these solutions are shielded from attack by other ions
647. A precipitate of AgCl is formed when equal volumes of the following are mixed [K_{sp} for AgCl = 10^{-10}]
 a) 10^{-4} M AgNO₃ and 10^{-7} M HCl
 b) 10^{-5} M AgNO₃ and 10^{-6} M HCl
 c) 10^{-5} M AgNO₃ and 10^{-4} M HCl
 d) 10^{-6} M AgNO₃ and 10^{-6} M HCl
648. When different types of salts have nearly same solubility product constant K_{sp} but less than one the most soluble salt is that:
 a) Which produces maximum number of ions
 b) Which produces minimum number of ions
 c) Which produces more charge on ion
 d) None of the above
649. In a flask, colourless N₂O₄ is in equilibrium with brown coloured NO₂. At equilibrium, when the flask is heated at 100°C the brown colour deepens and on cooling it becomes less coloured. The change in enthalpy ΔH , for the system is :
 a) Negative
 b) Positive
 c) Zero
 d) Undefined
650. 56 g of nitrogen and 8 g of hydrogen gas heated in a closed vessel. At equilibrium, 34 g of ammonia are present. The equilibrium number of moles of nitrogen, hydrogen and ammonia are, respectively
 a) 1, 1, 2
 b) 1, 2, 2
 c) 2, 1, 1
 d) 2, 2, 1
651. Calculate pOH of 0.001 M NH₄OH, when it is 1 % dissociated in the solution
 a) 5
 b) 2.96
 c) 9.04
 d) 11.4
652. On addition of an inert gas at constant volume to the reaction, N₂ + 3H₂ \rightleftharpoons 2NH₃ at equilibrium:
 a) The reaction halts
 b) Forward reaction is favoured
 c) The reaction remains unaffected
 d) Backward reaction is favoured
653. The reaction quotient (Q) for the reaction is given by:

$$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g}); \quad Q = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$

 The reaction will proceed from right to left if:
 a) $Q = 0$
 b) $Q = K_c$
 c) $Q < K_c$
 d) $Q > K_c$
654. Sulphuric acid is a dibasic acid. Hence, it forms:
 a) Acidic salt
 b) Basic and acidic salt
 c) Acidic and normal salt
 d) Double salt
655. Solubility of AgCl at 20°C is 1.435×10^{-3} g/L. The solubility product of AgCl is
 a) 1×10^{-5}
 b) 1×10^{-10}
 c) 1.435×10^{-5}
 d) 108×10^{-3}
656. Le-Chatelier's principle is not applicable to:
 a) Fe(s) + S(s) \rightleftharpoons FeS(s)
 b) H₂(g) + I₂(g) \rightleftharpoons 2HI(g)
 c) N₂(g) + 3H₂(g) \rightleftharpoons 2NH₃(g)
 d) N₂(g) + O₂(g) \rightleftharpoons 2NO(g)
657. 4 moles each of SO₂ and O₂ gases are allowed to react to form SO₃ in a closed vessel. At equilibrium 25% of O₂ is used up. The total number of moles of all the gases at equilibrium is
 a) 6.5
 b) 7.0
 c) 8.0
 d) 2.0
658. The pH of a 0.005 M aqueous solution of sulphuric acid is approximately:
 a) 0.005
 b) 2
 c) 1
 d) 0.01
659. When ammonium chloride is added to ammonia solution, the pH of the resulting solution will be
 a) Increased
 b) Seven
 c) Decreased
 d) Unchanged

phenomenon is called:

- a) Colloidal b) Buffer action c) Acidity d) Salt balance


675. Weak electrolytes are only partly dissociated in solutions. The extent to which they are dissociated in a given solution is known as the 'Degree of dissociation'. Which of the following statements for this degree of dissociation in aqueous solution is true?

- a) It is inversely proportional to the square root of initial concentration of the electrolyte
 b) It is directly proportional to the initial concentration of the electrolyte
 c) It is independent of the initial concentration of the electrolyte
 d) It depends on the equilibrium concentration of the electrolyte

676. At a certain temperature, the dissociation constants of formic acid and acetic acid are 1.8×10^{-4} and 1.8×10^{-5} respectively. The concentration of acetic acid solution in which the hydrogen ion has the same concentration as in 0.001 M formic acid solution is equal to

- a) 0.01 M b) 0.001 M c) 0.1 M d) 0.0001 M

677. Partial pressure of O_2 in the reaction



- a) K_p b) $\sqrt{K_p}$ c) $3\sqrt{K_p}$ d) $2K_p$

678. The solubility product of a salt having general formula MX_2 , in water is 4×10^{-12} . The concentration of M^{2+} ions in the aqueous solution of the salt is

- a) 4.0×10^{-10} M b) 1.6×10^{-4} M c) 1.0×10^{-4} M d) 2.0×10^{-6} M


679. In a lime kiln, to get higher yield of CO_2 , the measure that can be taken is

- a) To remove CaO b) To add more $CaCO_3$
 c) To maintain high temperature d) To pump out CO_2

680. Which of the following is a Bronsted acid as well as Bronsted base:

- a) Na_2CO_3 b) H_2O c) NH_3 d) BF_3

681. The rate constant for forward reaction and backward reaction of hydrolysis of ester are 1.1×10^{-2} and 1.5×10^{-3} per minute respectively. Equilibrium constant for the reaction is



- a) 33.7 b) 7.33 c) 5.33 d) 33.3

682. Which acid is involved in digestion process?

- a) HF b) HCl c) HBr d) HI

683. For $NH_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$, the observed pressure for reaction mixture in equilibrium is 1.12 atm at $106^\circ C$. The value of K_p for the reaction is :

- a) 3.136 atm^2 b) 0.3136 atm^2 c) 31.36 atm^2 d) 6.98 atm^2


684. A solution of NaCl in contact with atmosphere has a pH of about:

- a) 3.2 b) 5.4 c) 7.0 d) 14

685. Amines behave as

- a) Lewis acid b) Lewis base c) Aprotic acid d) Neutral compound

686. The two Bronsted bases in the reaction are



- a) $HC_2O_4^-$ and PO_4^{3-} b) HPO_4^{2-} and $C_2O_4^{2-}$ c) PO_4^{3-} and $C_2O_4^{2-}$ d) $HC_2O_4^-$ and HPO_4^{2-}

687. 1 M NaCl and 1 M HCl are present in an aqueous solution. The solution is

- a) Not a buffer solution with $pH < 7$ b) Not a buffer solution with $pH > 7$
 c) A buffer solution with $pH < 7$ d) A buffer solution with $pH > 7$

688. The use of methyl orange as an indicator in the volumetric determination of the equivalent weight of a weak acid would lead to:

- a) A low value for the equivalent weight
 b) A high value for the equivalent weight
 c) No error in the value

- d) Improved accuracy
689. Arrange NH_4^+ , H_2O , H_3O^+ , HF and OH^- in increasing order of acidic nature
- a) $\text{H}_3\text{O}^+ < \text{NH}_4^+ < \text{HF} < \text{OH}^- < \text{H}_2\text{O}$ b) $\text{NH}_4^+ < \text{HF} < \text{H}_3\text{O}^+ < \text{H}_2\text{O} < \text{OH}^-$
 c) $\text{OH}^- < \text{H}_2\text{O} < \text{NH}_4^+ < \text{HF} < \text{H}_3\text{O}^+$ d) $\text{H}_3\text{O}^+ > \text{HF} > \text{H}_2\text{O} > \text{NH}_4^+ > \text{OH}^-$
690. Which of the following is a Lewis base?
- a) CH_4 b) $\text{C}_2\text{H}_5\text{OH}$ c) Acetone d) Secondary
691. An aqueous solution of sodium carbonate is alkaline because sodium carbonate is a salt of
- a) Weak acid and weak base b) Strong acid and weak base
 c) Weak acid and strong base d) Strong acid and strong base
692. Which of the following is least likely to behave as Lewis acid?
- a) OH^- b) H_2O c) NH_3 d) BF_3
693. Which statement is false in case of equilibrium state?
- a) There is no apparent change in properties with time
 b) It is dynamic in nature
 c) It can be attained from either side of the reaction
 d) It can be attained from the side of the reactants only
694. The active mass of 45 g of KCl in a 3 L flask would be
- a) 0.20 b) 2.0 c) 3 d) 4
695. A litre of solution is saturated with AgCl. To this solution if 1.0×10^{-4} mole of solid NaCl is added, what will be the $[\text{Ag}^+]$ assuming no volume change?
- a) More b) Less c) Equal d) Zero
696. 9.2g N_2O_4 is heated in a 1L vessel till equilibrium state is established
- $$\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$$
- In equilibrium state 50% N_2O_4 was dissociated, Equilibrium constant will be (mol. wt. of $\text{N}_2\text{O}_4 = 92$)
- a) 0.1 b) 0.4 c) 0.3 d) 0.2
697. At 3000 K, the equilibrium partial pressure of CO_2 , CO and O_2 are 0.6, 0.4 and 0.2 atm respectively. K_p for the reaction, $2\text{CO}_2 \rightleftharpoons 2\text{CO} + \text{O}_2$ is
- a) 0.088 b) 0.0533 c) 0.133 d) 0.177
698. Aqueous solution of sodium cyanide is
- a) Acidic b) Amphoteric c) Basic d) Neutral
699. A mixture of N_2 and H_2 in the mole ratio 1:3 is allowed to attain equilibrium when 50% of mixture has reacted. If P is the equilibrium pressure, then partial pressure of NH_3 formed is:
- a) $P/6$ b) $P/2$ c) $P/3$ d) $P/4$
700. CH_3COOH is weaker acid than H_2SO_4 . It is due to
- a) More ionisation b) Less ionisation c) Covalent bond d) Electrovalent bond
701. Equal volume of three acid solutions of pH 3, 4 and 5 are mixed in a vessel. What will be the H^+ ion concentration in the mixture?
- a) $3.7 \times 10^{-4} \text{ M}$ b) $3.7 \times 10^{-3} \text{ M}$ c) $1.11 \times 10^{-3} \text{ M}$ d) $1.11 \times 10^{-4} \text{ M}$
702. The ionisation of strong electrolytes in acetic acid, compared to in water, is
- a) Weak, low b) Strong, more
 c) Medium, the same d) No ionisation, 100%
703. A saturated solution of CaF_2 is $2 \times 10^{-4} \text{ mol/L}$. Its solubility product constant is
- a) 2.6×10^{-9} b) 4×10^{-8} c) 4×10^{-12} d) 3.2×10^{-11}
704. The vapour density of PCl_5 is 104.16, but when heated at 230°C . Its vapour density is reduced to 62. The percentage of dissociation of PCl_5 at this temperature will be
- a) 6.8% b) 68% c) 46% d) 64%
705. Which is not amphoteric?

- a) HSO_4^- b) HCO_3^- c) H_2PO_4^- d) HCOO^-
706. An amount of solid NH_4HS is placed in a flask already containing ammonia gas at a certain temperature and 0.50 atm pressure. Ammonium hydrogen sulphide decomposes to yield NH_3 and H_2S gases in the flask. When the decomposition reaction reaches equilibrium the total pressure in the flask rises to 0.84 atm? The equilibrium constant for NH_4HS decomposition at this temperature is
- a) 0.11 b) 0.17 c) 0.18 d) 0.30
707. The correct order of acetic strength is:
- a) $\text{Cl}_2\text{O}_7 > \text{SO}_2 > \text{P}_4\text{O}_{10}$
 b) $\text{CO}_2 > \text{N}_2\text{O}_5 > \text{SO}_3$
 c) $\text{Na}_2\text{O} > \text{MgO} > \text{Al}_2\text{O}_3$
 d) $\text{K}_2\text{O} > \text{CaO} > \text{MgO}$
708. The equilibrium constant K_p for the reaction,
 $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ is
- a) More than one b) Less than one c) Equal to K_c d) Zero
709. Which of the following is least soluble?
- a) $\text{MnS}(K_{sp} = 7 \times 10^{-16})$
 b) $\text{FeS}(K_{sp} = 7 \times 10^{-19})$
 c) $\text{PtS}(K_{sp} = 8 \times 10^{-73})$
 d) $\text{NiS}(K_{sp} = 3 \times 10^{-12})$
710. The best explanation for the solubility of MnS in dil. HCl is that:
- a) Solubility product of MnCl_2 is less than that of MnS
 b) Concentration of Mn^{2+} is lowered by the formation of complex ions with chloride ions
 c) Concentration of sulphide ions is lowered by oxidation to free sulphur
 d) Concentration of sulphide ions is lowered by formation of the weak acid H_2S
711. If the pOH of a solution is 6.0, then pH is:
- a) 6 b) 8 c) 10 d) 14
712. If the pH of a solution is 2, its normality will be:
- a) 2 N b) 0.5 N c) 0.01 N d) None of these
713. The pH of solution, whose hydrogen ion concentration is one mole per litre, will be:
- a) 1.0 b) 0.1 c) Zero d) 1.5
714. Consider the reaction equilibrium
 $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g}), \Delta H^\circ = -198 \text{ kJ}$
 On the basis of Le-Chatelier's principle, the condition favourable for the forward reaction is
- a) Lowering of temperature as well as pressure
 b) Lowering the temperature and increasing the pressure
 c) Increasing the temperature and pressure
 d) Any value of temperature and pressure
715. NH_4Cl is acidic because
- a) On hydrolysis NH_4Cl gives weak base NH_4OH and strong acid HCl
 b) Nitrogen donates a pair of electrons
 c) It is a salt of weak acid and strong base
 d) On hydrolysis NH_4Cl gives strong base and weak acid
716. In a buffer solution consisting of a weak acid and its conjugate base, the ratio of concentration of conjugate base to acid is increased tenfold; then the pH of the solution will:
- a) Increase by one b) Increase tenfold c) Decrease by one d) Decrease tenfold
717. The solubility in water of a sparingly soluble salt AB_2 is 1.0×10^{-5} mol/litre. Its solubility product is:
- a) 1×10^{-15} b) 1×10^{-10} c) 4×10^{-15} d) 4×10^{-10}

718. If ClO_3^- is chlorate ion, then HClO_3 is:
 a) Hydrochloric acid b) Chlorous acid c) Chloric acid d) Chlorate acid
719. Which of the following is not a Lewis base?
 a) H_2O b) Ag^+ c) NH_3 d) OH^-
720. For a system in equilibrium, $\Delta G = 0$, under conditions of constant
 a) Temperature and pressure b) Energy and volume
 c) Temperature and volume d) Pressure and volume
721. In 100 mL of an aqueous HCl of pH 1.0, 900 mL of distilled water is added, the pH of the resultant solution becomes:
 a) 1.0 b) 2.0 c) 4.0 d) 7.0
722. 20 mL of a 0.1 N HCl is mixed with 20 mL of a 0.1 N KOH solution. The pH of the solution would be:
 a) Zero b) 7 c) 2 d) 9
723. In a system : $A(s) \rightleftharpoons 2B(g) + 3C(g)$. If the concentration of C at equilibrium is increased by a factor 2, it will cause the equilibrium concentration of B to change to:
 a) Two times of its original value
 b) One half of its original value
 c) $2\sqrt{2}$ times of its original value
 d) $\frac{1}{2\sqrt{2}}$ times of its original value
724. The Bronsted acid which gives the weakest conjugate base is:
 a) HF b) H_2S c) H_2O d) HCl
725. How much sodium acetate should be added to 0.1 M solution of CH_3COOH to give a solution of pH = 5.5 ($\text{p}K_a$ of $\text{CH}_3\text{COOH} = 4.5$)?
 a) 0.1 M b) 0.01 M c) 1.0 M d) 10.0 M
726. Which solution has pH equal to 10?
 a) 10^{-4} M KOH b) 10^{-10} M KOH c) 10^{-10} M HCl d) 10^{-4} M HCl
727. A saturated solution prepared by dissolved $\text{CaF}_2(s)$ in water, has $[\text{Ca}^{2+}] = 3.3 \times 10^{-4}$ M. What is the K_{sp} of CaF_2 ?
 a) 1.44×10^{-10} b) 2.24×10^{-8} c) 1.58×10^{-8} d) 1.67×10^{-8}
728. When 1.0 mL of dil. HCl acid is added to 100 mL of a buffer solution of pH 4.0, the pH of the solution:
 a) Becomes 7 b) Does not change c) Becomes 2 d) Becomes 10
729. The pH of blood is:
 a) Less than 6
 b) Greater than 7 and less than 8
 c) Greater than 8 and less than 9
 d) Greater than 10
730. The unit of equilibrium constant, K for the reaction, $A + B \rightleftharpoons C$, would be
 a) mol L^{-1} b) mol L c) L mol^{-1} d) Dimensionless
731. Which statement is correct?
 a) An acid and its conjugate base react to form salt and water
 b) The acid H_2O is its own conjugate base
 c) The conjugate base of a weak acid is a strong base
 d) The conjugate base of a strong acid is strong base
732. From separate solutions of four sodium salts NaW, NaX, NaY and NaZ had pH 7.0, 9.0, 10.0 and 11.0 respectively. When each solution was 0.1 M, the strongest acid is:
 a) HW
 b) HX
 c) HY
 d) HZ
733. Which information can be obtained from Le-Chatelier's principle?

- a) Shift in equilibrium position on changing P , T and concentration
 b) Dissociation constant of a weak acid
 c) Energy change in a reaction
 d) Equilibrium constant of a chemical reaction
734. The solubility product of Hg_2I_2 is equal to:
 a) $[\text{Hg}_2^{2+}][\text{I}^-]$ b) $[\text{Hg}^{2+}][\text{I}^-]$ c) $[\text{Hg}_2^{2+}][\text{I}^-]^2$ d) $[\text{Hg}^{2+}]^2[\text{I}^-]^2$
735. The pH of a solution formed by mixing 40 mL of 0.10 M HCl and 10 mL of 0.45 M NaOH is:
 a) 5 b) 8 c) 12 d) 10
736. The correct relationship between K_c and K_p in gaseous equilibrium is :
 a) $K_c = K_p (RT)^{\Delta n}$
 b) $K_p = K_c (RT)^{\Delta n}$
 c) $\frac{K_c}{RT} = (K_p)^{\Delta n}$
 d) $\frac{K_p}{RT} = (K_c)^{\Delta n}$
737. In a mixture of CH_3COOH and CH_3COONa , the ratio of salt to acid concentration is increased by ten folds. The pH of the solution will increase by:
 a) Zero b) 1 c) 2 d) 3
738. $\text{NaOH}(aq)$, $\text{HCl}(aq)$ and $\text{NaCl}(aq)$ concentration of each is 10^{-3} M. Their pH will be respectively
 a) 10, 6, 2 b) 11, 3, 7 c) 10, 2, 6 d) 3, 4, 7
739. At 25°C , the equilibrium K_1 , K_2 and K_3 of three reactions are :
 $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3; K_1$
 $\text{N}_2 + \text{O}_2 \rightleftharpoons 2\text{NO}; K_2$
 $\text{H}_2 + \frac{1}{2}\text{O}_2 \rightleftharpoons \text{H}_2\text{O}; K_3$
 The equilibrium constants for the oxidation of NH_3 by oxygen to give NO is :
 a) K_3^2/K_1 b) $K_2^2 K_3/K_1$ c) $K_1 K_2/K_3$ d) $K_2 K_3^3/K_1$
740. Which of the following is the strongest base?
 a) C_2H_5^- b) $\text{C}_2\text{H}_5\text{COO}^-$ c) $\text{C}_2\text{H}_5\text{O}^-$ d) H_2PO_4
741. The equilibrium constant, K for the reaction
 $2\text{HI}(g) \rightleftharpoons \text{H}_2(g) + \text{I}_2(g)$
 At room temperature is 2.85 and that at 698 K, it is 1.4×10^{-2} . This implies that
 a) HI is resonance stabilised b) HI is exothermic compound
 c) HI is stable at room temperature d) HI is less stable than H_2 and I_2
742. If the solubility product of BaSO_4 is 1.5×10^{-10} in water. Its solubility, in moles per litre, is
 a) 1.5×10^{-9} b) 3.9×10^{-5} c) 7.5×10^{-5} d) 1.5×10^{-5}
743. Soda water has a pH value:
 a) Less than 7 b) More than 7 c) 7 d) Greater than 14
744. According to Le-Chatelier's principle, if heat is given to solid-liquid system, then
 a) Quantity of solid will reduce b) Quantity of liquid will reduce
 c) Temperature will increase d) Temperature will decrease
745. The salt that does not hydrolyse, is:
 a) SnCl_2 b) FeCl_3 c) SnCl_4 d) CaCl_2
746. An acid-base indicator has $K_a = 3.0 \times 10^{-5}$. The acid form of the indicator is red and the basic form is blue. The $[\text{H}^+]$ required to change the indicator from 75% red to 75% blue is:
 a) $8 \times 10^{-5} \text{ M}$ b) $9 \times 10^{-5} \text{ M}$ c) $1 \times 10^{-5} \text{ M}$ d) $3 \times 10^{-4} \text{ M}$
747. When ammonium chloride is added to a solution of ammonium hydroxide,
 a) Dissociation of NH_4OH increases b) Concentration of OH^- increases
 c) Concentration of NH_4^+ and OH^- increases d) Concentration of NH_4^+ decreases
748. The pH of an aqueous solution having hydroxide ion concentration as 1×10^{-5} is
 a) 5 b) 9 c) 4.5 d) 11

749. In the manufacture of NH_3 by Haber's process, the condition which would give maximum yield is
 $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3 + Q \text{ kcal}$
 a) Low temperature and high pressure
 b) Low temperature, low pressure and low concentration of H_2
 c) High temperature, low pressure and low concentration
 d) High temperature, high pressure and high concentration
750. In water, the acid HClO_4 , HCl , H_2SO_4 and HNO_3 exhibit the same strength as they are completely ionised in water (a base). This is called ... of the solvent water.
 a) Strength b) Capacity c) Buffer effect d) Levelling effect
751. Which of the following solutions will have pH close to 1.0?
 a) 100 mL of $\frac{M}{10}$ HCl + 100 mL of $\frac{M}{10}$ NaOH b) 55 mL of $\frac{M}{10}$ HCl + 45 mL of $\frac{M}{10}$ NaOH
 c) 10 mL of $\frac{M}{10}$ HCl + 90 mL of $\frac{M}{10}$ NaOH d) 75 mL of $\frac{M}{5}$ HCl + 25 mL of $\frac{M}{5}$ NaOH
752. For the reaction $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g}); \Delta H = -93.6 \text{ kJ mol}^{-1}$, the concentration of NH_3 at equilibrium can be increased by
 (I) lowering the temperature
 (II) low pressure
 (III) excess of N_2
 (IV) excess of H_2
 a) (II) and (IV) are correct b) (II) only is correct
 c) (I), (II) and (III) are correct d) (III) and (IV) are correct
753. Some salts although containing two different metallic elements give test for only one of them in solution. Such salts are:
 a) Double salts b) Normal salts c) Complex salts d) None of these
754. Which statement is correct?
 a) NH_4OH is a strong base
 b) CH_3COONa given acidic solution in water
 c) CH_3COOH is a weak acid
 d) NH_4Cl gives alkaline solution in water
755. Which one of the following species acts as both Bronsted acid and base?
 a) H_2PO_2^- b) HPO_3^{2-} c) HPO_4^{2-} d) All of these
756. Which one is the strongest base?
 a) OH^- b) RO^- c) NH_2^- d) R^-
757. To a mixture of acetic acid and sodium acetate a further amount of sodium acetate is added. The pH of the mixture:
 a) Increases b) Decreases c) Remains unchanged d) Not predictable
758. Ionisation constant of CH_3COOH is 1.7×10^{-5} and $[\text{H}^+]$ ions is 3.4×10^{-4} . Then, initial concentration of CH_3COOH molecules is
 a) 6.8×10^{-3} b) 2.5×10^{-4} c) 3.5×10^{-3} d) 4.5×10^{-3}
759. 0.023 g of sodium metal is reacted with 100 cm^3 of water. The pH of the resulting solution is
 a) 10 b) 11 c) 9 d) 12
760. For the reaction, $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ in a vessel, after the addition of equal number of mole of N_2 and H_2 , equilibrium state is formed. Which of the following is correct?
 a) $[\text{H}_2] = [\text{N}_2]$ b) $[\text{H}_2] < [\text{N}_2]$ c) $[\text{H}_2] > [\text{N}_2]$ d) $[\text{H}_2] > [\text{NH}_3]$
761. 1.1 mole of A are mixed with 2.2 mole of B and the mixture is then kept in one litre flask till the equilibrium is attained $A + 2B \rightleftharpoons 2C + D$. At the equilibrium 0.2 mole of C are formed. The equilibrium constant of the reaction is :
 a) 0.001 b) 0.002 c) 0.003 d) 0.004

762. 50% neutralization of a solution of formic acid ($K_a = 2 \times 10^{-4}$) with NaOH would result in a solution having a hydrogen ion concentration of:
 a) 2×10^{-4} b) 3.7 c) 2.7 d) 1.85
763. pH of K_2S solution is:
 a) > 7 b) < 7 c) 7 d) Zero
764. If pressure is applied to the following equilibrium, Liquid \rightleftharpoons Vapour. The boiling point of the liquid:
 a) Will decrease b) Will increase c) May not change d) Will not change
765. The extent of ionisation increases
 a) With the increase in concentration b) On addition of excess water to the solution
 c) On decreasing the temperature of solution d) On stirring the solution vigorously
766. Which one of the following salts will produce an alkaline solution while dissolving in water?
 a) NH_4Cl b) Na_2CO_3 c) $NaNO_3$ d) Na_2SO_4
767. Which addition would not change the pH of 10 mL of dilute hydrochloric acid?
 a) 20 mL of the same dilute hydrochloric acid
 b) 5 mL of pure water
 c) 20 mL of pure water
 d) 10 mL of concentrated hydrochloric acid
768. Which does not contribute significantly to acid rains?
 a) SO_3 b) NO_2 c) CO_2 d) CO
769. Given that the equilibrium constant for the reaction

$$2SO_{2(g)} + O_{2(g)} \rightleftharpoons 2SO_{3(g)}$$
 has a value of 278 at a particular temperature. What is the value of the equilibrium constant for the following reaction at the same temperature?

$$SO_{3(g)} \rightleftharpoons SO_{2(g)} + \frac{1}{2}O_{2(g)}$$
 a) 1.8×10^{-3} b) 3.6×10^{-3} c) 6.0×10^{-2} d) 1.3×10^{-5}
770. The equilibrium constant for the reaction, $HONO + CN^- \rightarrow HCN + ONO^-$ is 1×10^{-14} . The magnitude of the equilibrium constant suggests that:
 a) CN^- is stronger base than ONO^-
 b) HCN is stronger acid than HONO
 c) ONO^- is the conjugate base of HONO
 d) The conjugate acid of CN^- is HCN
771. A buffer solution contains 0.1 mole of sodium acetate dissolved in 1000 cm^3 of 0.1 M acetic acid. To the above buffer solution, 0.1 mole of sodium acetate is further added and dissolved. The pH of the resulting buffer is
 a) pK_a b) $pK_a + 2$ c) $pK_a - \log 2$ d) $pK_a + \log 2$
772. Some chemists at ISRO wished to prepare a saturated solution of a silver compound and they wanted it to have the highest concentration of silver ion possible. Which of the following compounds, would they use?
 $K_{sp}(AgCl) = 1.8 \times 10^{-10}$
 $K_{sp}(AgBr) = 5.0 \times 10^{-13}$
 $K_{sp}(Ag_2CrO_4) = 2.4 \times 10^{-12}$
 a) AgCl b) AgBr c) Ag_2CrO_4 d) None of these
773. Ostwald's dilution law is applicable in the case of the solution of:
 a) CH_3COOH b) NaCl c) NaOH d) H_2SO_4
774. H_2S is passed into one dm^3 of a solution containing 0.1 mole of Zn^{2+} and 0.01 mole of Cu^{2+} till the sulphide ion concentration reaches 8.1×10^{-10} moles. Which one of the following statements is true?
 $[K_{sp}$ of Zn and CuS are 3×10^{-22} and 8×10^{-36}

respectively]

- a) Only ZnS precipitates b) Both CuS and ZnS precipitate c) Only CuS precipitates d) No precipitation occurs

775. The degree of dissociation of 0.1 M HCN solution is 0.01% its ionisation constant would be

- a) 10^{-3} b) 10^{-5} c) 10^{-7} d) 10^{-9}

776. Solubility of a gas in liquid increases on:

- a) Addition of a catalyst
b) Increasing the pressure
c) Decreasing the pressure
d) Increasing the temperature

777. In chemical reaction, $A \rightleftharpoons B$, the system will be known in equilibrium when

- a) 50% of A changes to B
b) A completely changes to B
c) Only 10% of A changes to B
d) The rate of change of A to B and B to A on both the sides are same

778. For a polyprotic acid say H_3PO_4 , its three dissociation constant K_1, K_2 and K_3 are in the order:

- a) $K_1 < K_2 < K_3$ b) $K_1 > K_2 > K_3$ c) $K_1 = K_2 = K_3$ d) $K_1 = K_2 > K_3$

779. The reaction, $A + 2B \rightleftharpoons 2C + D$ was studied using an initial concentration of B which was 1.5 times that of A. But the equilibrium concentration of A and C were found to be equal. Then the K_c for the equilibrium is :

- a) 4 b) 8 c) 6 d) 0.32

780. The expression for the solubility product of $Al_2(SO_4)_3$ is

- a) $K_{sp} = [Al^{3+}][SO_4^{2-}]$ b) $K_{sp} = [Al^{3+}]^2[SO_4^{2-}]^3$ c) $K_{sp} = [Al^{3+}]^3[SO_4^{2-}]^2$ d) $K_{sp} = [Al^{3+}]^2[SO_4^{2-}]^2$

781. If the dissociation constant of an acid HA is 1×10^{-5} , the pH of a 0.1 M solution of the acid HA will be approximately:

- a) 3 b) 5 c) 1 d) 6

782. Calculate the pOH of a solution at 25°C that contains 1×10^{-10} M of hydronium ions *i.e.*, H_3O^+ :

- a) 4 b) 9 c) 1 d) 7




783. When a strong acid-strong base or their salt are dissolved in water, they are completely ionised. If a strong acid is added to a strong base, H^+ ions from the former combines with OH^- ions of the latter forming water. The formation of each water molecule liberates a certain quantity of energy and the reaction is exothermic. The heat liberated when one mole of water is formed by combining hydrochloric acid and sodium hydroxide is 13.7 kcal. The heat liberated when one mole of water is formed by combining sulphuric acid and sodium hydroxide is:

- a) 34 kcal b) 13.7 kcal c) 8.5 kcal d) 25.5 kcal

784. K_a for HCN is 5×10^{-10} at 25°C. For maintaining a constant pH = 9, the volume of 5 M KCN solution required to be added to 10 mL of 2 M HCN solution is

- a) 2 mL b) 4 mL c) 8.2 mL d) 6.4 mL

785. The equilibrium constants for the reactions are:



The equilibrium constant for $H_3PO_4 \rightleftharpoons 3H^+ + PO_4^{3-}$ will be:

- a) K_1/K_2K_3 b) $K_1 \times K_2 \times K_3$ c) K_2/K_1K_3 d) $K_1 + K_2 + K_3$

786. Four moles of PCl_5 are heated in a closed 4 dm³ container to reach equilibrium at 400 K. at equilibrium 50% of PCl_5 is dissociated. What is the value of K_c for the dissociation of PCl_5 into

PCl₃ and Cl₂ at 400 K?

- a) 0.50 b) 1.00 c) 1.15 d) 0.05

787. Favourable conditions for manufacture of ammonia by the reaction.

$N_2 + 3H_2 \rightleftharpoons 2NH_3$; $\Delta H = -21.9$ kcal are:

- a) Low temperature, low pressure and catalyst
 b) Low temperature, high pressure and catalyst
 c) High temperature, low pressure and catalyst
 d) High temperature, high pressure and catalyst

788. If K_{sp} of Ag₂S is 10^{-17} , the solubility of Ag₂S in 0.1 M solution of Na₂S will be

- a) 10^{-8} b) 5×10^{-9} c) 10^{-15} d) 10^{-16}

789. 5 moles of SO₂ and 5 moles of O₂ are allowed to react to form SO₃ in a closed vessel. At the equilibrium stage 60% of SO₂ is used up. The total number of moles of SO₂, O₂ and SO₃ in the vessel now is

- a) 8.5 b) 9.5 c) 10 d) 10.5


790. How many gram of CaC₂O₄ will dissolve in one litre of saturated solution? K_{sp} of CaC₂O₄ is $2.5 \times 10^{-9} M^{-2}$ and its molecular weight is 128:

- a) 0.0064 g b) 0.0128 g c) 0.0032 g d) 0.0640 g

791. In the iodometric estimation in laboratory which process is involved?

- a) $Cr_2O_7^{2-} + H^+ + I^- \rightarrow 2Cr^{5+} + I_2$ b) $MnO_4^- + H^+ + I^- \rightarrow Mn^{2+} + I_2$
 $I_2 + S_2O_3^{2-} \rightarrow S_4O_6^{2-} + I^-$ $I_2 + S_2O_3^{2-} \rightarrow S_4O_6^{2-} + I_2$
 c) $MnO_4^- + OH^- + I^- \rightarrow MnO_2 + I_2$ d) $Cr_2O_7^{2-} + OH^- + I^- \rightarrow 2Cr^{3+} + I_2$
 $I_2 + S_2O_3^{2-} \rightarrow S_4O_6^{2-} + I^-$ $I_2 + S_2O_3^{2-} \rightarrow S_4O_6^{2-} + I^-$

792. Equilibrium constant K_1 and K_2 for the following equilibria



and, $2NO_2(g) \rightleftharpoons 2NO(g) + O_2(g)$ are related as

- a) $K_1 = \frac{1}{K_2}$ b) $K_2 = \frac{1}{K_1}$ c) $K_2 = \frac{1}{K_1^2}$ d) $K_1 = \frac{1}{K_2^2}$

793. A sample of Na₂CO₃ · H₂O weighing 0.62 g is added to 100 mL of 0.1 N (NH₄)₂SO₄ solution. What will be the resulting solution?

- a) Acidic b) Neutral c) Basic d) None of these


794. For which one of the following reactions $K_p = K_c$?

- a) $PCl_5 \rightleftharpoons PCl_3 + Cl_2$ b) $N_2 + O_2 \rightleftharpoons 2NO$ c) $N_2 + 3H_2 \rightleftharpoons 2NH_3$ d) $2SO_3 \rightleftharpoons 2SO_2 + O_2$

795. K_{sp} of AgCl at 18°C is 1.8×10^{-10} . If Ag⁺ of sodium is 4×10^{-3} mol/litre the Cl⁻ that must exceed before AgCl is precipitated would be:

- a) 4.5×10^{-8} mol/litre
 b) 7.2×10^{-13} mol/litre
 c) 4.0×10^{-3} mol/litre
 d) 4.5×10^{-7} mol/litre

796. The equilibrium constant (K_p) for the decomposition of gaseous H₂O



is related to degree of dissociation (α) at a total pressure p is given by

- a) $K_p = \frac{\alpha^3 p^{1/2}}{(1+\alpha)(2+\alpha)^{1/2}}$ b) $K_p = \frac{\alpha^3 p^{3/2}}{(1-\alpha)(2+\alpha)^{1/2}}$ c) $K_p = \frac{\alpha^3 p^2}{(1-\alpha)(2+\alpha)^{1/2}}$ d) $K_p = \frac{\alpha^{3/2} p^{1/2}}{(1-\alpha)(2+\alpha)^{1/2}}$

797. A monoprotic acid in 1.00 M solution is 0.01 % ionised. The dissociation constant of this acid is


- a) 1×10^{-8} b) 1×10^{-4} c) 1×10^{-6} d) 1×10^{-5}

798. pH of 1 M NH₃ aqueous solution is

($K_b = 1.8 \times 10^{-5}$)

- a) 11.13 b) 12.5 c) 13.42 d) 11.55
799. K_a for formic acid and acetic acid are 2.1×10^{-4} and 1.1×10^{-5} respectively. The relative strength of acids is:
- a) 2 : 1 b) 2.3 : 1 c) 1 : 2.1 d) 4.36 : 1
800. Would gaseous HCl be considered as an Arrhenius acid?
- a) Yes b) No
c) Not known d) Gaseous HCl does not exist
801. According to Le-Chatelier's principle, the addition of temperature to the following reaction
- $$\text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g}) \rightarrow \text{CH}_4(\text{g}) + 2\text{O}_2(\text{g})$$
- will cause it to the right. This reaction is, therefore
- a) Exothermic b) Unimolecular c) Endothermic d) Spontaneous
802. The degree of dissociation of $\text{PCl}_5(\alpha)$ obeying the equilibrium,
- $$\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g}),$$
- is approximately related to the pressure at equilibrium by:
- a) $\alpha \propto P$ b) $\alpha \propto \frac{1}{\sqrt{P}}$ c) $\alpha \propto \frac{1}{P^2}$ d) $\alpha \propto \frac{1}{P^4}$
803. Solubility product of MX_2 at 298 K is 4×10^{-12} . At this temperature concentration of M^{2+} ion in mol per litre is:
- a) $2 \times 10^{-6} \text{M}$ b) $1 \times 10^{-4} \text{M}$ c) $1.6 \times 10^{-4} \text{M}$ d) $4 \times 10^{-6} \text{M}$
804. A reaction attains equilibrium when the Gibbs energy change accompanying the reaction is:
- a) Positive and large b) Zero c) Negative and large d) Negative and small
805. $\text{NH}_4\text{HS}(\text{s}) \rightleftharpoons \text{NH}_3(\text{g}) + \text{H}_2\text{S}(\text{g})$
- In the above reaction, if the pressure at equilibrium and at 300 K is 100 atm then what will be the equilibrium constant K_p ?
- a) 2500 atm² b) 50 atm² c) 100 atm² d) 200 atm²
806. The solubility product of a sparingly salt AX_2 is 3.2×10^{-11} . Its solubility (in mol/L) is
- a) 5.6×10^{-6} b) 3.1×10^{-4} c) 2×10^{-4} d) 4×10^{-4}
807. If the solubility product of AgBrO_3 and Ag_2SO_4 are 2×10^{-5} and 5.5×10^{-5} respectively, the relationship between the solubilities of these can be correctly represented as:
- a) $s_{\text{AgBrO}_3} > s_{\text{Ag}_2\text{SO}_4}$ b) $s_{\text{AgBrO}_3} < s_{\text{Ag}_2\text{SO}_4}$ c) $s_{\text{AgBrO}_3} = s_{\text{Ag}_2\text{SO}_4}$ d) $s_{\text{AgBrO}_3} \equiv s_{\text{Ag}_2\text{SO}_4}$
808. The conjugate acid of HPO_4^{2-} is:
- a) H_2PO_4^- b) PO_4^{3-} c) H_3PO_4 d) H_3PO_3
809. The colour of an electrolyte solution depends on:
- a) The nature of the anion
b) The nature of the cation
c) The nature of both the ions
d) The nature of the solvent
810. The resultant pH of a solution on mixing 200 mL of an aqueous solution of HCl (pH = 2.0) is mixed with 300 mL of an aqueous solution of NaOH (pH = 12.0) is:
- a) 11.0310 b) 11.3010 c) 10.000 d) None of these
811. If ΔG^0 for the reaction given below is 1.7 kJ; the equilibrium constant of the reaction, $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$ at 25°C is :
- a) 24.0 b) 3.9 c) 2.0 d) 0.5
812. In a reaction, $\text{A} + \text{B} \rightleftharpoons \text{C} + \text{D}$, the concentrations of A, B, C and D (in mol/L) are 0.5, 0.8, 0.4 and 1.0 respectively. The equilibrium constant is
- a) 0.1 b) 1.0 c) 0.5 d) 5.0
813. The solvent which is neither proton donor nor proton acceptor is called:
- a) Amphoteric b) Neutral c) Aprotic d) Protonic
814. The equilibrium constant of a reaction is 20.0. At equilibrium, the rate constant of forward reaction is 10.0. The rate constant for backward reaction is :

- a) 0.5 b) 2.0 c) 10.0 d) 200.0
815. For the reversible reaction,
 $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ at $500^\circ C$, the value of K_p is 1.44×10^{-5} . When partial pressure is measured in atmospheres. The corresponding value of K_c with concentration in mol L^{-1} is
 a) $1.44 \times 10^{-5} / (0.082 \times 773)^{-3}$ b) $1.44 \times 10^{-5} / (0.082 \times 500)^{-2}$
 c) $1.44 \times 10^{-5} / (8.314 \times 773)^2$ d) $1.44 \times 10^{-5} / (0.082 \times 773)^{-2}$
816. The pH of the solution produced when an aqueous solution of strong acid pH 5 is mixed with equal volume of an aqueous solution of strong acid of pH 3 is:
 a) 3.3 b) 3.5 c) 4.5 d) 4.0
817. Given, $HF + H_2O \xrightarrow{K_a} H_3O^+ + F^-$,
 $F^- + H_2O \xrightarrow{K_b} HF + OH^-$
 Which relation is correct?
 a) $K_b = K_w$ b) $K_b = \frac{1}{K_w}$ c) $K_a \times K_b = K_w$ d) $\frac{K_a}{K_b} = K_w$
818. The pH of a solution is 4. The hydrogen ion concentration of the solution in mol/L is
 a) 9.5 b) 10^{-4} c) 10^4 d) 10^{-2}
819. Which one of the following statements is not true?
 a) The conjugate base of $H_2PO_4^-$ is HPO_4^{2-}
 b) $pH + pOH = 14$ for all aqueous solutions
 c) The pH of 1×10^{-8} M HCl is 8
 d) 96,500 C of electricity when passed through a $CuSO_4$ solution deposits 1 g equivalent of copper at the cathode
820. The correct representation for the solubility product constant of Ag_2CrO_4 is:
 a) $[Ag^+]^2[CrO_4^{2-}]$ b) $[Ag^+][CrO_4^{2-}]$ c) $[2Ag^+][CrO_4^{2-}]$ d) $[2Ag^+]^2[CrO_4^{2-}]$
821. For a reversible reaction, the rate constant for the forward reaction is 2.38×10^{-4} and for the backward reaction is 8.15×10^{-5} . The K_c for the reaction is :
 a) 0.342 b) 2.92 c) 0.292 d) 3.42
822. The pH of a soft drink is 3.82. Its H^+ ion concentration will be:
 a) 1.96×10^{-2} mol/litre b) 1.96×10^{-3} mol/litre c) 1.5×10^{-4} mol/litre d) 1.96×10^{-1} mol/litre
823. The decreasing order of strength of the bases,
 OH^- , NH_2^- , $H-C \equiv C^-$ and $CH_3-CH_2^-$, is:
 a) $CH_3-CH_2^- > NH_2^- > H-C \equiv C^- > OH^-$
 b) $H-C \equiv C^- > CH_3-CH_2^- > NH_2^- > OH^-$
 c) $OH^- > NH_2^- > H-C \equiv C^- > CH_3-CH_2^-$
 d) $NH_2^- > H-C \equiv C^- > OH^- > CH_3-CH_2^-$
824. The metallic sulphide not precipitated if H_2S gas is passed through an aqueous solution containing cuprous chloride, bismuth chloride, mercuric chloride and sodium chloride is:
 a) CuS b) Bi_2SO_3 c) HgS d) Na_2S
825. pH of 1×10^{-8} M nitric acid solution will be
 a) 6 b) 6.96 c) 7.96 d) 8
826. Indicate the correct answer out of the following for the reaction,
 $NH_4Cl + H_2O \rightleftharpoons NH_4OH + HCl$
 a) The reaction is retarded by the addition of KOH
 b) The reaction is favoured by the addition of NH_4OH
 c) The reaction is retarded by the addition of hydrogen ion
 d) None of the above
827. The salt of strong acid and weak base ($FeCl_2$) is
 a) Acidic b) Basic c) Neutral d) None of these
828. For the following reaction in gaseous phase



K_c/K_p is

- a) $(RT)^{1/2}$
 b) $(RT)^{-1/2}$
 c) (RT)
 d) $(RT)^{-1}$

829. For a reaction if $K_p > K_c$ the forward reaction is favoured by
 a) Low pressure b) High pressure c) High temperature d) Low temperature
830. What will be the pH value of 0.05M $\text{Ba}(\text{OH})_2$ solution?
 a) 12 b) 13 c) 1 d) 12.96
831. pH of a saturated solution of $\text{Ba}(\text{OH})_2$ is 12. The value of solubility product (K_{sp}) of $\text{Ba}(\text{OH})_2$ is:
 a) 4.0×10^{-6} b) 5.0×10^{-6} c) 3.3×10^{-7} d) 5.0×10^{-7}
832. The conjugate base of H_2SO_4 is
 a) SO_4^{2-} b) HSO_4^- c) HSO_4^+ d) H_3SO_4
833. For the reaction, $\text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons \text{PCl}_5(\text{g})$, the value of K_c at 250°C is 26. The value of K_p at this temperature will be
 a) 0.41 b) 0.51 c) 0.61 d) 0.71
834. The pH of a 10^{-10} M NaOH solution is nearest to
 a) 10 b) 7 c) 4 d) -10
835. Which can act as buffer?
 a) $\text{NH}_4\text{Cl} + \text{NH}_4\text{OH}$
 b) $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$
 c) 40 mL of 0.1 M NaCN + 20 mL of 0.1 M HCl
 d) All of the above
836. The pH indicators are
 a) Salts of strong acids and strong bases b) Salts of weak acid and weak bases
 c) Either weak acids or weak bases d) Either strong acid or strong bases
837. Phenolphthalein is a:
 a) Weak acid b) Weak base c) Strong acid d) Strong base
838. The solubility product of Hg_2I_2 is equal to
 a) $[\text{Hg}_2^{2+}][\text{I}^-]$ b) $[\text{Hg}_2^{2+}][\text{I}^-]^2$ c) $[\text{Hg}_2^{2+}][\text{I}^-]^2$ d) $[\text{Hg}_2^{2+}][\text{I}^-]^2$
839. The rate at which a substance reacts depends on its
 a) Atomic weight b) Atomic number c) Molecular weight d) Active mass
840. The compound HCl behaves as ... in the reaction,
 $\text{HCl} + \text{HF} \rightarrow \text{H}_2\text{F}^+\text{Cl} + \text{F}^-$
 a) Strong acid b) Strong base c) Weak acid d) Weak base
841. At temperature T , a compound AB_2 (g) dissociated according to the reaction $2\text{AB}_2(\text{g}) \rightleftharpoons 2\text{AB}(\text{g}) + \text{B}_2(\text{g})$ with a degree of dissociation x , which is small compared with unity. The expression for K_p , in terms of x and the total pressure P , is:
 a) $\frac{Px^3}{2}$ b) $\frac{Px^2}{3}$ c) $\frac{Px^3}{3}$ d) $\frac{Px^2}{2}$
842. Which of the following is not a Lewis acid?
 a) BF_3 b) AlCl_3 c) SO_2 d) H_2O
843. The pH of a 0.1 molar solution of the acid HQ is 3. The value of the ionization constant, K_a of this acid is:
 a) 1×10^{-3} b) 1×10^{-5} c) 1×10^{-7} d) 3×10^{-1}
844. For the precipitation of IIInd group cations only the solution is made acidic so that:
 a) The sulphide ion concentration may increase

- b) The sulphide ion concentration may decrease
 c) The H^+ ion concentration may increase
 d) The cations concentration may increase
845. Let the solubility of an aqueous solution of $Mg(OH)_2$ be X then its K_{sp} is
 a) $4x^3$ b) $108x^5$ c) $27x^4$ d) $9x$
846. In the equilibrium, $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$, the partial pressure of SO_2 , O_2 and SO_3 are 0.662, 0.101 and 0.331 atm respectively. What should be the partial pressure of oxygen so that the equilibrium concentration of SO_2 and SO_3 are equal?
 a) 0.4 atm b) 1.0 atm c) 0.8 atm d) 0.25 atm
847. In the following reaction,
 $AgCl + KI \rightleftharpoons KCl + AgI$
 as KI is added, the equilibrium is shifted towards right giving more AgI precipitate. because
 a) Both AgCl and AgI are sparingly soluble
 b) The K_{sp} of AgI is lower than K_{sp} of AgCl
 c) The K_{sp} of AgI is higher than K_{sp} of AgCl
 d) Both AgCl and AgI have same solubility product
848. Which of the following is a Lewis base?
 a) NaOH b) NH_3 c) BCl_3 d) All of these
849. The ionisation constant of NH_4^+ in water is 5.6×10^{-10} at $25^\circ C$. The rate constant for the reaction of NH_4^+ and OH^- to form NH_3 and H_2O at $25^\circ C$ is $3.4 \times 10^{10} L mol^{-1} s^{-1}$. The rate constant for proton transfer from water to NH_3 is:
 a) $6.07 \times 10^5 s^{-1}$ b) $6.07 \times 10^{-10} s^{-1}$ c) $6.07 \times 10^{-5} s^{-1}$ d) $6.07 \times 10^{10} s^{-1}$
850. K_{sp} for AgCl in water at $25^\circ C$ is 1.8×10^{-10} . If 10^{-5} mol of Ag^+ ions are added to this solution. K_{sp} will be:
 a) 1.8×10^{-16} b) 1.8×10^{-10} c) 1.8×10^{-5} d) None of these
851. The reaction which proceeds in the forward direction is :
 a) $Fe_2O_3 + 6HCl = 2FeCl_3 + 3H_2O$
 b) $SnCl_4 + Hg_2Cl_2 = SnCl_2 + 2HgCl_2$
 c) $NH_3 + H_2O + NaCl = NH_4Cl + NaOH$
 d) $2CuI + I_2 + 4K^+ = 2Cu^{2+} + 3KI$
852. The chemical equilibrium of a reversible reaction is not influenced by
 a) Pressure b) Catalyst
 c) Concentration of the reactants d) Temperature
853. A 0.01 M ammonia solution is 5 % ionised, its pH will be
 a) 11.80 b) 10.69 c) 7.22 d) 12.24
854. For the decomposition reaction
 $NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$
 The $K_p = 2.9 \times 10^{-5} atm^3$. The total pressure of gases at equilibrium when 1 mole of $NH_2COONH_4(g)$ was taken to start with would be
 a) 0.0766 atm b) 0.0582 atm c) 0.0388 atm d) 0.0194 atm
855. 3 moles of A and 4 moles of B are mixed together and allowed to come into equilibrium according to the following reaction.
 $3A(g) + 4B(g) \rightleftharpoons 2C(g) + 3D(g)$
 When equilibrium is reached, there is 1 mole of C.
 The equilibrium extent of the reaction is
 a) $\frac{1}{4}$ b) $\frac{1}{3}$ c) $\frac{1}{2}$ d) 1
856. For the reaction, $C_2H_4 + H_2 \rightleftharpoons C_2H_6$
 The correct relation is

- a) $K_p = K_c$ b) $K_p = K_c[RT]$ c) $K_p = K_c[RT]^{-2}$ d) $K_p = K_c[RT]^{-1}$
857. Which solution will have pH closer to 1.0?
 a) 100 mL of (M/10) HCl + 100 mL of (M/10) NaOH
 b) 55 mL of (M/10) HCl + 45 mL of (M/10) NaOH
 c) 10 mL of (M/10) HCl + 90 mL of (M/10) NaOH
 d) 75 mL of (M/5) HCl + 25 mL of (M/5) NaOH
858. Which of the following is a Lewis acid?
 a) Cl^- b) H_3O^+ c) PF_3 d) $\text{C}_2\text{H}_5\text{OH}$
859. A buffer solution can be prepared from a mixture of
 (i) sodium acetate and acetic acid in water
 (ii) sodium acetate and hydrochloric acid in water
 (iii) ammonia and ammonium chloride in water
 (iv) ammonia and sodium hydroxide in water
 a) (i), (ii) b) (ii), (iii) c) (iii), (iv) d) (i), (iii)
860. An equilibrium mixture of the reaction, $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ contains 0.120 mole of NO_2 , 0.080 mole of NO and 0.640 mole of O_2 in a 4 litre flask at constant temperature. The value of K_c for the reaction at this temperature is:
 a) 14 b) 24 c) 7 d) 28
861. For the reaction,
 $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$, the value of K_c at 800°C is 0.1. When the equilibrium concentration of both the reactants is 0.5 mole, what is the value of K_p at the same temperature?
 a) 0.5 b) 0.1 c) 0.01 d) 0.025
862. The equilibrium,
 $\text{P}_4(\text{s}) + 6\text{Cl}_2(\text{g}) \rightleftharpoons 4\text{PCl}_3(\text{g})$ attained by mixing equal moles of P_4 and Cl_2 in a evacuated vessel. Then, at equilibrium,
 a) $[\text{Cl}_2] > [\text{PCl}_3]$ b) $[\text{Cl}_2] > [\text{P}_4]$ c) $[\text{P}_4] > [\text{Cl}_2]$ d) $[\text{PCl}_3] > [\text{P}_4]$
863. The yield of product in the reaction $\text{A}_2(\text{g}) + 2\text{B}(\text{g}) \rightleftharpoons \text{C}(\text{g}) + \text{Q}$ kJ would be high at
 a) High temperature and high pressure b) High temperature and low pressure
 c) Low temperature and high pressure d) Low temperature and low pressure
864. According to law of mass action, rate of a chemical reaction is proportional to
 a) Molar concentration of reactants b) Concentration of reactants
 c) Concentration of products d) Molar concentration of products
865. If NaOH is added to a solution of acetic acid:
 a) H^+ ions increases b) pH decreases c) $[\text{C}_2\text{H}_3\text{O}_2]^-$ increases d) $[\text{HC}_2\text{H}_3\text{O}_2]$ increases
866. What is the effect of having the pressure by doubling the volume on the following system at 500°C ?
 $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$
 a) Shift to reactant side b) Shift to product formation
 c) Liquefaction of HI d) No effect
867. The solubility product (K_{sp}) of the following compounds are given at 25°C

Compound	K_{sp}
AgCl	1.1×10^{10}
AgI	1.0×10^{16}
PbCrO_4	4.0×10^{14}
Ag_2CO_3	8.0×10^{12}

The most soluble and least soluble compounds are

- a) AgCl and PbCrO_4 b) AgI and Ag_2CO_3 c) AgCl and Ag_2CO_3 d) Ag_2CO_3 and AgI

868. The pH of 10^{-10} M NaOH solution is nearest to
 a) -4 b) -10 c) 4 d) 7
869. The conjugate base of H_2PO_4^- is
 a) H_3PO_4 b) P_2O_5 c) PO_4^{3-} d) HPO_4^{2-}
870. The ionization constant of ammonium hydroxide is 1.77×10^{-5} at 298 K. Hydrolysis constant of ammonium chloride is:
 a) 5.65×10^{-12} b) 5.65×10^{-10} c) 6.50×10^{-12} d) 5.65×10^{-13}
871. 1 M solution of an acid has a pH of 5. Which of the following is the most reasonable explanation for this acid?
 a) The acid is too dilute
 b) It is a strong acid
 c) It reacts with water to produce a high concentration of hydronium ions
 d) It is a weak acid
872. The solubility product K_{sp} , of a sparingly soluble salt AgIO_3 is 1.0×10^{-8} at a given temperature. What is the mass of AgIO_3 (mol. Mass = 283) contained in 100 mL solution at this temperature is:
 a) 1.0×10^{-4} g b) 28.3×10^{-2} g c) 2.83×10^{-3} g d) 1.0×10^{-7} g
873. The pH of a buffer solution containing equal molal concentration of a weak base and its chloride (K_b for weak base = 2×10^{-5}) is
 a) 5 b) 9 c) 4.7 d) 9.3
874. In qualitative analysis, in III group NH_4Cl is added before NH_4OH because
 a) To increase the concentration of NH_4^+ ions b) To increase the concentration of Cl^- ions
 c) To reduce the concentration of OH^- ions d) To increase the concentration of OH^- ions
875. The K_{sp} for $\text{Cr}(\text{OH})_3$ is 1.6×10^{-30} . The molar solubility of this compound in water is
 a) $\sqrt[2]{1.6 \times 10^{30}}$ b) $\sqrt[4]{1.6 \times 10^{30}}$ c) $\sqrt[4]{1.6 \times 10^{30}/27}$ d) $1.6 \times 10^{30}/27$
876. ΔG° for the reaction $X + Y \rightleftharpoons Z$ is - 4.606 kcal. The equilibrium constant for the reaction at 227°C is :
 a) 100 b) 10 c) 2 d) 0.01
877. For the reaction, $2A(g) \rightleftharpoons 3C(g) + D(s)$, the value of K_c will be equal to
 a) $K_p(RT)$ b) K_p/RT c) $= K_p$ d) None of these
878. Reaction between barium chloride and sodium sulphate goes to completion because:
 a) Barium sulphate is almost insoluble
 b) The reaction is reversible
 c) The solubility of barium chloride decreases
 d) None of the above
879. Which of the following cannot act as a Lewis or Bronsted acid?
 a) BF_3 b) AlCl_3 c) SnCl_4 d) CCl_4
880. Consider the reaction,

$$\text{NO}_2 \rightleftharpoons \frac{1}{2}\text{N}_2 + \text{O}_2, K_1; \text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2, K_2$$
 Give the equilibrium constant for the formation of N_2O_4 from N_2 and O_2 .
 a) $\frac{1}{K_1^2} + \frac{1}{K_2}$ b) $\frac{1}{K_1 K_2}$ c) $\sqrt{\frac{1}{K_1 K_2}}$ d) $\frac{K_2}{K_1}$
881. A weak monobasic acid is 1% ionised in 0.1 M solution at 25°C . The percentage of ionisation in its 0.025 M solution is
 a) 1 b) 2 c) 3 d) 4
882. In $K_p = K_c[RT]^{\Delta n}$, Δn may have:
 a) +ve values
 b) -ve values
 c) Integer or fractional values

- d) Either of the above
883. The conjugate acid of H^- ion is:
 a) H_3O^+ b) H_2 c) OH^- d) H_2O
884. The addition of HCl does not suppresses the ionisation of:
 a) Acetic acid b) Benzoic acid c) H_2S d) H_2SO_4
885. A colourless solution liberates CO_2 gas when added to a metal bicarbonate. The solution is:
 a) Basic b) Acidic c) Amphoteric d) Neutral
886. Which one of the following statement is not true?
 a) The conjugate base of H_2PO_4^- is HPO_4^{2-}
 b) $\text{pH} + \text{pOH} = 14$ for all aqueous solutions
 c) The pH of 1×10^{-8} M HCl is 8
 d) 96500 C of electricity when passed through a CuSO_4 solution deposit 1 g equivalent of copper at the cathode
887. Two moles of PCl_5 is heated in a closed vessel of 2 L capacity. When the equilibrium is attained 40% of it has been found to be dissociated. What is the K_c in mol/dm^3 ?
 a) 0.532 b) 0.266 c) 0.133 d) 0.174
888. A liquid is in equilibrium with its vapour at it's boiling point. On the average, the molecules in two phases have equal
 a) Inter-molecular forces b) Potential energy
 c) Kinetic energy d) Total energy
889. At equilibrium, the amount of HI in a 3 litre vessel was 12.8 g. Its equilibrium concentration is :
 a) 4.267 M b) 0.033 M c) 0.1 M d) 0.2 M
890. Which one of the following salts give an acidic solution in water?
 a) CH_3COONa b) NH_4Cl c) NaCl d) $\text{CH}_3\text{COONH}_4$
891. For which reaction does the equilibrium constant depend on the units of concentration?
 a) $\text{NO}(\text{g}) \rightleftharpoons \frac{1}{2}\text{N}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g})$
 b) $\text{Zn}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \rightleftharpoons \text{Cu}(\text{s}) + \text{Zn}^{2+}(\text{aq})$
 c) $\text{C}_2\text{H}_5\text{OH}(\text{l}) + \text{CH}_3\text{COOH}(\text{l}) \rightleftharpoons \text{CH}_3\text{COOC}_2\text{H}_5(\text{l}) + \text{H}_2\text{O}(\text{l})$
 d) $\text{COCl}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{Cl}_2(\text{g})$
892. The solubility product of As_2S_3 is 2.8×10^{-72} . What is the solubility of As_2S_3 ?
 a) 1.92×10^{-15} mol/L b) 1.72×10^{-15} mol/L
 c) 2.3×10^{-16} mol/L d) 1.65×10^{-36} mol/L
893. When CO_2 dissolves in water, the following equilibrium is established, $\text{CO}_2 + 2\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HCO}_3^-$, for which the equilibrium constant is 3.8×10^{-7} and $\text{pH} = 6.0$. The ratio of $[\text{HCO}_3^-]/[\text{CO}_2]$ is :
 a) 3.8×10^{-18} b) 3.8 c) 0.38 d) 13.8
894. The blood buffers are most often involved in stabilizing the pH in presence of metabolically produced:
 a) Acids b) Bases c) Salts d) None of these
895. If concentration of N_2 , H_2 and NH_3 are 1, 2 and 3 respectively, their concentrations at equilibrium will be:
 $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$
 a) $(1-x)$ $(2-3x)$ b) $(1-x/3)$ $(2-x)$ $2x$ $(1-x)$ $(2-x)$ c) $(1-x)$ $(2-3x)$ d) $(1-x)$ $(2-3x)$ $(2-3x)$
896. For the reaction, $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ at 500°C , the value of K_p is 1.44×10^{-5} . What will be the value of K_p at low pressure where the gases are behaving almost ideally?
 a) 1.44×10^{-5} b) $(0.082 \times 773)^2 \times 1.44 \times 10^{-5}$ c) $1.44 \times 10^{-5} \times (0.082 \times 500)^2$ d) $1.44 \times 10^{-5} \times (0.082 \times 773)^3$
897. The range of pH in which methyl orange works as indicator:
 a) 3-4 b) 10-12 c) 8-10 d) 6-8

898. For the reaction : $\text{CO(g)} + \frac{1}{2}\text{O}_2\text{(g)} \rightleftharpoons \text{CO}_2\text{(g)}$, K_p/K_c is:
 a) RT b) $(RT)^{-1}$ c) $(RT)^{-1/2}$ d) $(RT)^{1/2}$
899. K_{sp} of CuS , Ag_2S and HgS are 10^{-31} , 10^{-44} and 10^{-54} respectively. Select the correct order for their solubility in water:
 a) $\text{Ag}_2\text{S} > \text{HgS} > \text{CuS}$ b) $\text{HgS} > \text{CuS} > \text{Ag}_2\text{S}$ c) $\text{HgS} > \text{Ag}_2\text{S} > \text{CuS}$ d) $\text{Ag}_2\text{S} > \text{CuS} > \text{HgS}$
900. pH of a 0.0001 M HCl solution is
 a) 4.0 b) 2.0 c) 6.0 d) 7.0
901. The pH of 0.1 M HCl is approximately 1. The approximate pH of 0.05 M H_2SO_4 is:
 a) 0.05 b) 0.5 c) 1 d) 2
902. Phenolphthalein shows ... in acid medium.
 a) Red colour b) Yellow colour c) Pink colour d) No colour
903. The $[\text{OH}^-]$ in 100 mL of 0.015 M HCl (aq.) is:
 a) $5 \times 10^{-12} \text{ M}$ b) $3 \times 10^{-10} \text{ M}$ c) $6.7 \times 10^{-13} \text{ M}$ d) $2.0 \times 10^{-9} \text{ M}$
904. For an equilibrium reaction if the value of $K_c \gg 1$, then the reaction favoured more towards
 a) Backward b) Forward
 c) Equilibrium will be obtained d) Reaction will stop
905. K_c for $A + B \rightleftharpoons 3C$ is 20 at 25°C. If a 2 litre vessel contains 1,2 and 4 mole of A, B and C respectively, the reaction at 25°C shall :
 a) Proceed from left to right
 b) Proceed from right to left
 c) Be at equilibrium
 d) Not occur
906. Solution prepared by dissolving equal number of mole of HOCl ($K_a = 3.2 \times 10^{-8}$) and NaOCl is a buffer of pH:
 a) 8.0 b) 3.2 c) 7.5 d) 4.8
907. An increase in the temperature of an equilibrium system:
 a) Favours the exothermic reaction
 b) Favours the endothermic reaction
 c) Favours both the exothermic and endothermic reactions
 d) Favours neither the exothermic nor endothermic reactions
908. Which of these is a Lewis acid?
 a) AlCl_3 b) NCl_3 c) HCl d) ROR
909. The pH of a solution is 4. The hydrogen ion concentration of the solution if pH is to be increased to 5 is:
 a) Halved
 b) Doubled
 c) Decreased by 10 times
 d) Decreased to half of its original value of concentration
910. The oxo-acid of anhydride P_2O_5 is:
 a) H_3PO_4 b) $\text{H}_4\text{P}_2\text{O}_7$ c) HPO_3 d) All of these
911. In a saturated solution of the sparingly soluble strong electrolyte AgIO_3 (molecular mass=283) the equilibrium which sets in is

$$\text{AgIO}_3\text{(s)} \rightleftharpoons \text{Ag}^+\text{(aq)} + \text{IO}_3^-\text{(aq)}$$
 If the solubility product constant K_{sp} of AgIO_3 at a given temperature is 1.0×10^{-8} , what is the mass of AgIO_3 contained in 100 mL of its saturated solution?
 a) $28.3 \times 10^{-2} \text{ g}$ b) $2.83 \times 10^{-3} \text{ g}$ c) $1.0 \times 10^{-7} \text{ g}$ d) $1.0 \times 10^{-4} \text{ g}$
912. H_3BO_3 is:
 a) Monobasic and weak Lewis acid b) Monobasic and weak Bronsted acid c) Monobasic and strong Lewis acid d) Tribasic and weak Bronsted acid

913. All Lewis acids are not necessarily:
 a) Proton donor b) Bronsted acids c) Arrhenius acids d) All of these
914. In the reaction,
 $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$. Which of the following is correct?
 a) $K_p = K_c$ b) $K_p > K_c$ c) $K_p < K_c$ d) $K_p \geq K_c$
915. Which of the following is most soluble in water?
 a) Bi_2S_3 ($K_{sp} = 10^{-70}$) b) MnS ($K_{sp} = 7 \times 10^{-16}$) c) CuS ($K_{sp} = 8 \times 10^{-37}$) d) Ag_2S ($K_{sp} = 6 \times 10^{-51}$).
916. The equilibrium constant for the reaction
 $SO_3(g) \rightleftharpoons SO_2(g) + \frac{1}{2}O_2(g)$
 is $K_c = 4.9 \times 10^{-2}$. The value of K_c for the reaction
 $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ will be
 a) 416 b) 2.40×10^{-3} c) 9.8×10^{-2} d) 4.9×10^{-2}
917. pK_a or a weak acid is defined as
 a) $\log K_a$ b) $\frac{1}{\log K_a}$ c) $\log \frac{1}{K_a}$ d) $-\log \frac{1}{K_a}$
918. In the equilibrium, $AB \rightleftharpoons A + B$, if the equilibrium concentration of A is doubled, the equilibrium concentration of B would become
 a) Half b) Twice c) $\frac{1}{4}$ th d) $\frac{1}{8}$ th
919. Two moles of PCl_5 were heated in a closed vessel of 2 L. At equilibrium 40% of PCl_5 is dissociated into PCl_3 and Cl_2 . The value of equilibrium constant is
 a) 0.53 b) 0.267 c) 2.63 d) 5.3
920. When NaOH is dissolved in water, heat is evolved. If the temperature is raised, the solubility of NaOH:
 a) Increases
 b) Decreases
 c) Remains the same
 d) Cannot be predicted
921. In which of the following equilibrium, change in volume of the system does not alter the number of moles?
 a) $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$
 b) $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$
 c) $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$
 d) $SOCl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$
922. Conjugate base of HSO_4^- is
 a) SO_4^{2-} b) H_2SO_4 c) $H_3SO_4^+$ d) None of these
923. Which favours the backward reaction in a chemical equilibrium?
 a) Increasing the concentration of one of the reactants
 b) Removal of at least one of the products at regular intervals
 c) Increasing the concentration of one or more of the products
 d) None of the above
924. A weak acid HX ($K_a = 1 \times 10^{-5}$) on reaction with NaOH gives NaX. For 0.1 M aqueous solution of NaX, the % hydrolysis is:
 a) 0.001% b) 0.01% c) 0.15% d) 1%
925. For the reaction: $2NO_2(g) \rightleftharpoons 2NO(g) + O_2(g)$; $K_c = 1.8 \times 10^{-6}$ at $184^\circ C$ and $R = 0.083$ litre atm $K^{-1}mol^{-1}$. When K_p and K_c are compared at $184^\circ C$, it is found that :
 a) $K_p > K_c$ b) $K_p < K_c$ c) $K_p = K_c$ d) $K_p \geq K_c$
926. $Mg(OH)Cl$ is an example of:
 a) Acidic salt b) Basic salt c) Neutral salt d) Amphoteric salt
927. The degree of dissociation of a weak acid is 1.34% at 0.1 M concentration. Its dissociation constant is:
 a) 8×10^{-6} b) 1.79×10^{-5} c) 0.182×10^{-5} d) 8×10^{-5}

928. In the reaction $\text{HCl} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{Cl}^-$
- a) H_2O is the conjugate base of HCl acid
 b) Cl^- is the conjugate base of HCl acid
 c) Cl^- is the conjugate acid of H_2O base
 d) H_3O^+ is the conjugate base of HCl
929. Slope and intercepts of the plots $\log_{10} K$ vs $\frac{1}{T}$ are given respectively by :
- a) $-\frac{\Delta H^\circ}{R}, \frac{\Delta S^\circ}{R}$
 b) $\frac{-\Delta H^\circ}{2.303R}, \frac{\Delta S^\circ}{2.303}$
 c) $\frac{-\Delta H^\circ}{2.303R}, \frac{\Delta S^\circ}{2.303R}$
 d) $\frac{\Delta H^\circ}{2.303}, \frac{-\Delta S^\circ}{2.303R}$
930. On adding 0.1 M solution each of $[\text{Ag}^+]$, $[\text{Ba}^{2+}]$, $[\text{Ca}^{2+}]$ in a Na_2SO_4 solution, species first precipitated is
- $[K_{\text{sp}} \text{BaSO}_4 = 10^{-11}, K_{\text{sp}} \text{CaSO}_4 = 10^{-6}, K_{\text{sp}} \text{Ag}_2\text{SO}_4 = 10^{-5}]$
- a) Ag_2SO_4
 b) BaSO_4
 c) CaSO_4
 d) All of these
931. K for the synthesis of HI is 50. K for the dissociation of HI is
- a) 0.2
 b) 0.02
 c) 0.4
 d) 0.04
932. Which of the following factors will favour the reverse reaction in a chemical equilibrium?
- a) Increase in the concentration of one of the reactants
 b) Increase in the concentration of one or more products
 c) Removal of at least one of the product at regular time intervals
 d) None of the above
933. The anhydride of HNO_3 is:
- a) P_2O_5
 b) N_2O_3
 c) NO
 d) N_2O_5
934. According to Le-Chatelier principle, adding heat a solid and liquid in equilibrium will cause the
- a) Amount of solid to decrease
 b) Amount of liquid to decrease
 c) Temperature to rise
 d) Temperature to fall
935. What is the conjugate base of OH^- ?
- a) O^{2-}
 b) O^-
 c) H_2O
 d) O_2
936. The solubility of $\text{Al}(\text{OH})_3$ is 's' mol per litre, the solubility product of $\text{Al}(\text{OH})_3$ is:
- a) s^3
 b) $27s^4$
 c) s^2
 d) $4s^2$
937. The pH of 0.1 M solution of the following salts increases in the order
- a) $\text{NaCl} < \text{NH}_4\text{Cl} < \text{NaCN} < \text{HCl}$
 b) $\text{HCl} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{NaCN}$
 c) $\text{NaCN} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{HCl}$
 d) $\text{HCl} < \text{NaCl} < \text{NaCN} < \text{NH}_4\text{Cl}$
938. For the Haber's process for the formation of NH_3 at 298 K is :
- $$\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3; \Delta H = -460 \text{ kJ}$$
- Which of the following is correct?
- a) The condition for equilibrium is $G_{\text{N}_2} + 3G_{\text{H}_2} = 2G_{\text{NH}_3}$, where G is Gibbs energy per mole of gaseous species measured at that partial pressure
 b) On addition N_2 , the equilibrium will shift to forward direction because according to II law of thermodynamics the entropy must decrease in the direction of spontaneous reaction.
 c) The catalyst will increase the rate of forward reaction by 2 times and that of backward reaction by 1.5 times
 d) Name of the above
939. Consider the reversible reaction,
- $$\text{HCN}(aq) \rightleftharpoons \text{H}^+(aq) + \text{CN}^-(aq)$$
- At equilibrium, the addition of $\text{CN}^-(aq)$ would:
- a) Reduce $\text{HCN}(aq)$ concentration
 b) Decrease the $\text{H}^+(aq)$ ion concentration
 c) Increase the equilibrium constant
 d) Decrease the equilibrium constant
940. The solubility product of iron (III) hydroxide is 1.6×10^{-39} . If X is the solubility of iron (III) hydroxide, which one of the following expression can be used to calculate X ?

- a) $K_{sp} = X^4$ b) $K_{sp} = 9X^4$ c) $K_{sp} = 27X^3$ d) $K_{sp} = 27X^4$
941. Baking soda is a/an:
 a) Basic salt b) Double salt c) Complex salt d) Acidic salt
942. An aqueous solution of sodium carbonate has a pH greater than 7 because
 a) It contains more carbonate ions than H_2O molecules
 b) Contains more hydroxide ions than carbonate ions
 c) Na^+ ions react with water
 d) Carbonate ions react with H_2O
943. The pH of a solution obtained by mixing 10 mL of 0.1 M HCl and 40 mL of 0.2 M H_2SO_4 is:
 a) 1.4865 b) 0.4865 c) 0.4685 d) 3
944. Just before a reversible reaction attains equilibrium it is found that:
 a) The velocity of both forward reaction and backward reaction is also increasing
 b) The velocity of the forward reaction is decreasing and that of backward reaction is increasing
 c) The velocity of both forward and backward reactions is decreasing
 d) All of the above
945. How many mole of HCl are required to prepare one litre of buffer solution (containing NaCN + HCl) of pH 8.5 using 0.01 g formula weight of NaCN ($K_{HCN} = 4.1 \times 10^{-10}$)?
 a) 8.85×10^{-3} b) 8.75×10^{-2} c) 8.85×10^{-4} d) 8.85×10^{-2}
946. For the reaction $A + B \rightleftharpoons 3C$ at $25^\circ C$, a 3 litre vessel contains 1, 2, 4 mole of A, B and C respectively. If K_c for the reaction is 10, the reaction will proceed in :
 a) Forward direction b) Backward direction c) In either direction d) In equilibrium
947. What is the pH of a 1M CH_3COONa solution? K_a of acetic acid = 1.8×10^{-5} , $K_w = 10^{-14} \text{ mol}^2 \text{ litre}^{-2}$.
 a) 2.4 b) 3.6 c) 4.8 d) 9.4
948. Formaldehyde polymerizes to form glucose according to the reaction,

$$6 \text{ HCHO} \rightleftharpoons \text{C}_6\text{H}_{12}\text{O}_6$$
 The theoretically computed equilibrium constant for this reaction is found to be 6×10^{22} . If 1 M solution of glucose dissociates according to the above equilibrium, the concentration of formaldehyde in the solution will be
 a) $1.6 \times 10^{-2} \text{ M}$ b) $1.6 \times 10^{-4} \text{ M}$ c) $1.6 \times 10^{-6} \text{ M}$ d) $1.6 \times 10^{-8} \text{ M}$
949. The polyprotic acid is:
 a) HCl
 b) $HClO_4$
 c) H_3PO_4
 d) HNO_3
950. The solubility of Sb_2S_3 in water is $1.0 \times 10^{-5} \text{ mol/L}$ at 298 K. What will be its solubility product?
 a) 108×10^{-25} b) 1.0×10^{-25} c) 144×10^{-25} d) 126×10^{-24}
951. The pH of 1/1000 N KOH solution is:
 a) 10^{-11} b) 3 c) 2 d) 11
952. Which acts both as Lowry Bronsted acid and base?
 a) OH^- b) Na_2CO_3 c) NH_3 d) HSO_4^-
953. By applying law of mass action, the equilibrium constant, K for the reaction
 $HA + H_2O \rightleftharpoons H_3O^+ + A^-$, is given as
 a) $K = \frac{[HA][H_2O]}{[H_3O^+][A^-]}$ b) $K = \frac{[H_3O^+][A^-]}{[HA][H_2O]}$ c) $K = \frac{[H_3O^+][H_2O]}{[A^-][HA]}$ d) $K = \frac{[HA][A^-]}{[H_2O][H_3O^+]}$
954. The Haber's process for the manufacture of ammonia is usually carried out at about $500^\circ C$. If a temperature of about $250^\circ C$ was used instead of $500^\circ C$:
 a) Ammonia would not be formed at all
 b) The percentage of ammonia in the equilibrium mixture would be too low

- c) A catalyst would be of no use at all at this temperature
 d) The rate of formation of ammonia would be too slow
955. The equilibrium constant for, $\text{H}_2(\text{g}) + \text{CO}_2(\text{g}) \rightleftharpoons \text{H}_2\text{O}(\text{g}) + \text{CO}(\text{g})$ is 1.80 at 1000°C . If 1.0 mole of H_2 and 1.0 mole of CO_2 are placed in one litre flask, the final equilibrium concentration of CO at 1000°C will be:
 a) 0.573 M b) 0.385 M c) 5.73 M d) 0.295 M
956. For a reaction $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$ at 721 K, the value of equilibrium constant is 50. If 0.5 moles each of H_2 and I_2 is added to the system the value of equilibrium constant will be
 a) 40 b) 60 c) 50 d) 30
957. Among BMe_3 , BF_3 , BCl_3 and B_2H_6 which one will be the best Lewis acid?
 a) BCl_3 b) BMe_3 c) B_2H_6 d) BF_3
958. Potassium ferrocyanide is a
 a) Mixed salt b) Normal salt c) Complex salt d) Double salt
959. The pH of pure water at 25°C and 35°C are 7 and 6 respectively. The heat of formation of water from H^+ and OH^- is:
 a) $84.55 \text{ kcal mol}^{-1}$
 b) $-84.55 \text{ kcal mol}^{-1}$
 c) $74.55 \text{ kcal mol}^{-1}$
 d) $-74.55 \text{ kcal mol}^{-1}$
960. The pH of solution A, B, C, D are 9.5, 2.5, 3.5 and 5.5 respectively. The most acidic solution is:
 a) A b) B c) C d) D
961. According to Debye-Hückel theory of strong electrolytes, increase in conductivity on dilution is due to:
 a) Increase in number of ions
 b) Increase in the mobility of ions
 c) Decrease in the number of ions
 d) Decrease in the mobility of ions
962. If K_1 and K_2 are the respective equilibrium constants for the two reactions,
 $\text{XeF}_6(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{XeOF}_4(\text{g}) + 2\text{HF}(\text{g})$
 $\text{XeO}_4(\text{g}) + \text{XeF}_6(\text{g}) \rightleftharpoons \text{XeOF}_4(\text{g}) + \text{XeO}_3\text{F}_2(\text{g})$
 The equilibrium constant for the reaction,
 $\text{XeO}_4(\text{g}) + 2\text{HF}(\text{g}) \rightleftharpoons \text{XeO}_3\text{F}_2(\text{g}) + \text{H}_2\text{O}(\text{g})$ is:
 a) K_1K_2 b) K_1/K_2^2 c) K_2/K_1 d) K_1/K_2
963. The solubility of a saturated solution of calcium fluoride is 2×10^{-4} moles per litre. Its solubility product is:
 a) 32×10^{-10} b) 32×10^{-8} c) 32×10^{-14} d) 32×10^{-12}
964. The equilibrium constant $\text{Br}_2 \rightleftharpoons 2\text{Br}$ at 500 K and 700 K are 10^{-10} and 10^{-5} respectively. The reaction is:
 a) Endothermic b) Exothermic c) Fast d) Slow
965. For the homogenous reaction,

$$4\text{NH}_3 + 5\text{O}_2 \rightleftharpoons 4\text{NO} + 6\text{H}_2\text{O}$$
 the equilibrium constant K_c has the units
 a) conc.^{+10} b) conc.^{+1} c) conc.^{-1} d) It is dimensionless
966. 1 mole of H_2 and 2 moles of I_2 are taken initially in a 2 L vessel. The number of moles of H_2 at equilibrium is 0.2. Then, the number of moles of I_2 and HI at equilibrium are
 a) 1.2, 1.6 b) 1.8, 1.0 c) 0.4, 2.4 d) 0.8, 2.0
967. If the ionic product of water (K_w) is 1.96×10^{-14} at 35°C , what is its value at 10°C ?
 a) 1.96×10^{-14} b) 3.92×10^{-14} c) 2.95×10^{-15} d) 1.96×10^{-13}
968. 0.1 mole of CH_3NH_2 ($K_b = 5 \times 10^{-4}$) is mixed with 0.08 mole of HCl diluted to 1 L. What will be the H^+ concentration in the solution?
 a) $8 \times 10^{-2} \text{ M}$ b) $8 \times 10^{-11} \text{ M}$ c) $1.6 \times 10^{-11} \text{ M}$ d) $8 \times 10^{-5} \text{ M}$

969. The conjugate base of H_3BO_3 is:
 a) $\text{B}(\text{OH})_4^-$ b) H_2BO_3^- c) HBO_3^{2-} d) H_4BO_3^+
970. For the reaction, $\text{CO}(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons \text{COCl}_2(\text{g})$ the value of K_p/K_c is equal to :
 a) 1.0 b) RT c) \sqrt{RT} d) $\frac{1}{RT}$
971. One mole of nitrogen is mixed with 3 mole of hydrogen in a closed 3 litre vessel, 20% of nitrogen is converted into NH_3 . Then K_c for the $\frac{1}{2} \text{N}_2 + \frac{3}{2} \text{H}_2 \rightleftharpoons \text{NH}_3$ is:
 a) 0.36 litre mol^{-1} b) 0.46 litre mol^{-1} c) 0.5 litre mol^{-1} d) 0.2 litre mol^{-1}
972. Which is a reversible reaction?
 a) $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$
 b) $\text{H}_2\text{SO}_4 + \text{Ba}(\text{OH})_2 \rightarrow \text{BaSO}_4 \downarrow + 2\text{H}_2\text{O}$
 c) $\text{NaCl} + \text{AgNO}_3 \rightarrow \text{NaNO}_3 + \text{AgCl} \downarrow$
 d) $2\text{KClO}_3 \rightarrow 2\text{KCl} + 3\text{O}_2 \uparrow$
973. $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$
 The equilibrium constant of the above reaction is 6.4 at 300 K. If 0.25 mole each of H_2 and I_2 are added to the system, the equilibrium constant will be
 a) 6.4 b) 0.8 c) 3.2 d) 1.6
974. Would gaseous HCl be considered as an Arrhenius acid?
 a) Yes b) No
 c) Not known d) Gaseous HCl does not exist
975. Buffer solution is prepared by mixing
 a) Strong acid + its salt of strong base b) Weak acid + its salt of weak base
 c) Strong acid + its salt of weak base d) Weak acid + its salt of strong base
976. Which of the following acids will have lowest value of $\text{p}K_a$?
 a) $\text{CH}_3\text{CH}_2\text{COOH}$ b) $\text{CH}_3\text{CH}(\text{Br})\text{COOH}$
 c) $\text{CH}_3\text{CH}(\text{F})\text{COOH}$ d) $\text{FCH}_2\text{CH}_2\text{COOH}$
977. 2 moles of PCl_5 were heated in a closed vessel of 2 L capacity. At equilibrium 40% of PCl_5 is dissociated into PCl_3 and Cl_2 . The value of equilibrium constant is
 a) 0.266 b) 0.366 c) 2.66 d) 3.66
978. $\text{p}K_a$ of a weak acid is defined as
 a) $\log_{10} K_a$ b) $\frac{1}{\log_{10} K_a}$ c) $\log_{10} \frac{1}{K_a}$ d) $-\log_{10} \frac{1}{K_a}$
979. For a reaction equilibrium, $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$, the concentrations of N_2O_4 and NO_2 at equilibrium are 4.8×10^{-2} and 1.2×10^{-2} mol/L respectively. The value of K_c for the reaction is
 a) 3×10^{-3} mol/L b) 3.3×10^{-3} mol/L c) 3×10^{-1} mol/L d) 3.3×10^{-1} mol/L
980. If α is the degree of ionisation, C the concentration of a weak electrolyte and K_a the acid ionisation constant then the correct relationship between α , C and K_a is
 a) $\alpha^2 = \sqrt{\frac{K_a}{C}}$ b) $\alpha^2 = \sqrt{\frac{C}{K_a}}$ c) $\alpha = \sqrt{\frac{K_a}{C}}$ d) $\alpha = \sqrt{\frac{C}{K_a}}$
981. Which of the following behaves as Lewis acid and not as Bronsted acid?
 a) HCl b) H_2SO_4 c) HSO_3^- d) SO_3
982. If little heat is added to ice \rightleftharpoons liquid equilibrium in a sealed container, then:
 a) Pressure will rise b) Temperature will rise c) Temperature will fall d) No change in P and T

983. An aqueous solution in which the H^+ ion concentration is greater than $10^{-7} M$ is said to be
 a) Acidic b) Alkaline c) Neutral d) None of these
984. The conjugate base of H_2SO_4 in the following reaction is:
 $H_2SO_4 + H_2O \rightleftharpoons H_3O^+ + HSO_4^-$
 a) H_2O b) HSO_4^- c) H_3O^+ d) SO_4^{2-}
985. For the reaction, $H_2 + I_2 \rightleftharpoons 2HI$, the equilibrium concentration of H_2 , I_2 and HI are 8.0, 3.0 and 28.0 mol/L respectively. The equilibrium constant is
 a) 28.34 b) 32.66 c) 34.78 d) 38.88
986. $HClO$ is a weak acid. The concentrations of $[H^+]$ ions in 0.1 M solution of $HClO$ ($K_a = 5 \times 10^{-8}$) will be equal to:
 a) $7.07 \times 10^{-5} M$ b) $5 \times 10^{-7} M$ c) $6 \times 10^{-7} M$ d) $7 \times 10^{-4} M$
987. At a certain temperature, $2HI \rightleftharpoons H_2 + I_2$ only 50% HI is dissociated at equilibrium. The equilibrium constant is :
 a) 1.0 b) 3.0 c) 0.5 d) 0.25
988. Aqueous solution of CH_3COOH contains:
 a) CH_3COOH, H^+ b) $CH_3COO^-, H_3O^+, CH_3CO$ c) CH_3COO^-, H_3O^+, H^+ d) CH_3COOH, CH_3COO^-, H^+
989. In the manufacture of ammonia by Haber's process,
 $N_2(g) + 3H_2 \rightleftharpoons 2NH_3(g) + 92.3 kJ$
 Which of the following condition is unfavourable?
 a) Increasing the temperature
 b) Increasing the pressure
 c) Reducing the temperature
 d) Removing ammonia as it is formed
990. If $CuSO_4 \cdot 5H_2O(s) \rightleftharpoons CuSO_4 \cdot 3H_2O(s) + 2H_2O(v)$ $K_p = 1.086 \times 10^{-4} atm^2$ at $25^\circ C$. The efflorescent nature of $CuSO_4 \cdot 5H_2O$ can be noticed when vapour pressure of H_2O in atmosphere is :
 a) $> 7.92 mm$ b) $< 7.92 mm$ c) $\geq 7.92 mm$ d) None of these
991. Conjugate acid-base pair differs by a/an:
 a) Electron b) Electron pair c) Proton d) Neutron
992. The hydrogen ion concentration in a solution of weak acid of dissociation constant K_a and concentration c is nearly equal to:
 a) $\sqrt{\frac{K_a}{c}}$ b) $\frac{c}{K_a}$ c) $K_a c$ d) $\sqrt{K_a c}$
993. For the liquefaction of gas, the favourable conditions are:
 a) Low T and high P
 b) Low T and low P
 c) Low T and high P and a catalyst
 d) Low T and catalyst
994. 0.5 M ammonium benzoate is hydrolysed to 0.25 percent, hence, its hydrolysis constant is
 a) 2.5×10^{-5} b) 1.5×10^{-4} c) 3.125×10^{-6} d) 6.25×10^{-6}
995. For the equilibrium, $CaCO_3(s) \rightleftharpoons CaO(s) + CO_2(g)$, which of the following expression is correct?
 a) $\frac{K_p}{[CaO][CO_2]/[CaCO_3]}$ b) $\frac{K_p}{P_{CaCO_3}} = (p_{CaO} + p_{CO_2})$ c) $K_p = p_{CO_2}$ d) $\frac{K_p}{(p_{CaO} + p_{CO_2}/p_{CaCO_3})}$
996. When $NaNO_3$ is heated in a closed vessel, O_2 is liberated and $NaNO_2$ is left behind. At equilibrium,
 (i) Addition of $NaNO_3$ favours forward reaction
 (ii) Addition of $NaNO_2$ favours backward reaction
 (iii) Increasing pressure favours reverse reaction
 (iv) Increasing temperature favours forward reaction

Correct option is

- a) (i), (ii), (iii) b) (ii), (iii), (iv) c) (i), (iii), (iv) d) (i), (ii) (iii), (iv)

997. Given pH of a solution *A* is 3 and it is mixed with another solution *B* having pH 2. After mixing are resultant pH of the solution will be

- a) 3.2 b) 1.9 c) 3.4 d) 3.5

998. To neutralise completely 20 mL of 0.1 *M* aqueous solution of phosphorus acid H_3PO_3 , the volume of 0.1 *M* KOH solution required is:

- a) 60 mL b) 20 mL c) 40 mL d) 10 mL

999. The relation between equilibrium constant K_p and K_c is

- a) $K_p = K_c(RT)^{\Delta n_g}$ b) $K_c = K_p(RT)^{\Delta n_g}$ c) $K_p = \left(\frac{K_c}{RT}\right)^{\Delta n_g}$ d) $K_p - K_c = (RT)^{\Delta n_g}$

100. On mixing equal volumes of two buffer solutions of pH value 3 and 5, the pH of the resultant solution will be

- a) 3.3 b) 4.0 c) 4.7 d) 5.3

