

GPLUS EDUCATION

Date :
Time :
Marks :

CHEMISTRY

HYDROGEN

Single Correct Answer Type

- The percentage of para hydrogen in ordinary hydrogen increases when:
a) Temperature is lowered
b) Temperature is increased
c) Pressure is increased and temperature is decreased.
d) None of the above
- Manufacture of H_2 is made by:
a) Lane's process b) Bosch process c) From natural gas d) All of these
- H_2O_2 on treatment with chlorine gives:
a) H_2 b) Oxygen c) Hypochlorous acid d) ClO_2
- Radioactive isotope of hydrogen is
a) Tritium b) Deuterium c) Para hydrogen d) Ortho hydrogen
- K_a of H_2O_2 is of the order of
a) 10^{-12} b) 10^{-14} c) 10^{-16} d) 10^{-10}
- The hardness of water is estimated by
a) EDTA method b) Titrimetic method c) Conductivity method d) Distillation method
- The H – O – O bond angle in H_2O_2 is
a) 107.28° b) 97° c) 104.5° d) 109.28°
- Hydrogen loses its electron to form H^+ ion. In this respect it resembles to:
a) Transition metals b) Alkali metals c) Halogens d) Noble gases
- $TiH_{1.73}$ is an example of :
a) Ionic hydride b) Covalent hydride c) Metallic hydride d) Polymeric hydride
- The decomposition of H_2O_2 can be slowed down by the addition of small amount of phosphoric acid which acts as:
a) Stopper b) Detainer c) Inhibitor d) promoter
- The ortho and para hydrogen possess:
a) Same physical properties but different chemical properties
b) Different physical properties but same chemical properties
c) Same chemical and physical properties
d) Different, physical and chemical properties
- The volume strength of 1.5 N H_2O_2 solution is
a) 4.8 b) 8.4 c) 4.2 d) 2.4
- Which of the following is correct about heavy water?
a) Water at $4^\circ C$ having maximum density is known as heavy water
b) It is heavier than water(H_2O)
c) It is formed by the combination of heavier isotope of hydrogen and oxygen
d) None of the above
- Which is not present in clear hard water?
a) $Mg(HCO_3)_2$ b) $CaCl_2$ c) $MgSO_4$ d) $MgCO_3$
- Which of the following is not correct regarding the electrolytic preparation of H_2O_2 ?
a) Lead is used as cathode b) 50% H_2SO_4 is used
c) Hydrogen is liberated at anode d) Sulphuric acid undergoes oxidation

16. Electrolysis of X gives Y at anode. Vacuum distillation of Y gives H_2O_2 . The number of peroxy (O - O) bonds present in X and Y respectively are
 a) 1.1 b) 1.2 c) Zero, 1 d) Zero, zero
17. When H_2O_2 is added to acidic ferrous sulphate solutions:
 a) Electrons are gained by Fe^{2+}
 b) Electrons are lost by Fe^{2+}
 c) There is no loss or gain of electrons
 d) Iron hydroxide precipitates
18. Which of the following reactions produces hydrogen?
 a) $H_2S_4O_8 + H_2O$ b) $BaO + HCl$ c) $Mg + H_2O$ d) $Na_2O_2 + 2HCl$
19. H_2O_2 is formed by which of the following compounds?
 a) Na_2O_2 b) $NaOH$ c) Na_2O d) KO_2
20. Which of the following acts as both reducing and oxidising agents?
 a) H_2SO_4 b) H_2O_2 c) KOH d) $KMnO_4$
21. The sum of protons, electrons and neutrons in the heaviest isotope of hydrogen is
 a) 3 b) 5 c) 4 d) 6
22. On shaking H_2O_2 with acidified potassium dichromate and ether, ethereal layer becomes
 a) Green b) Red c) Blue d) Brown
23. The acidified solution of $FeCl_3$ is reduced by passing:
 a) ordinary H_2 b) O_2 c) nascent H d) H_2
24. Hydrogen does not combine with
 a) Helium b) Bismuth c) Antimony d) Sodium
25. H_2 acts as an oxidant in its reaction with:
 a) Br_2 b) Ca c) N_2 d) S
26. Of the two solvents H_2O and D_2O , NaCl dissolves:
 a) Equally in both b) Only in H_2O c) More in D_2O d) More in H_2O
27. What is formed when calcium carbide react with heavy water?
 a) C_2D_2 b) CaD_2 c) CaD_2O d) CD_2
28. When different metals like Zn, Sn, Fe are added to dilute sulphuric acid, same gas, which burns explosively in air, is evolved. The gas is:
 a) O_2 b) N_2 c) Cl_2 d) H_2
29. Heavy water is represented as
 a) $H_2^{18}O$ b) D_2O c) $D_2^{18}O$ d) H_2O at $4^\circ C$
30. Which is not a water softener?
 a) Calgon b) Permutit c) Na_2CO_3 d) Na_2SO_4
31. The boiling point of heavy water is:
 a) $100^\circ C$ b) $101.4^\circ C$ c) $104^\circ C$ d) $102.5^\circ C$
32. The volume of oxygen liberated from 15mL of 20 volume H_2O_2 is
 a) 250mL b) 300mL c) 150mL d) 200mL
33. Decomposition of H_2O_2 is prevented by
 a) KOH b) MnO_2 c) Acetanilide d) Oxalic acid
34. The boiling point of water is high because
 a) Water molecule is linear
 b) Water molecule is not linear
 c) Water molecules possess covalent bond between H and O
 d) Water molecules associate due to H-bonding
35. The volume of '10 vol.' of H_2O_2 required to liberate 500 mL O_2 at NTP is:
 a) 50 mL b) 25 mL c) 100 mL d) 125 mL
36. Which of the following pairs of ions make the water hard?

- a) $\text{NH}_4^+, \text{Cl}^-$ b) $\text{Ca}^+, \text{HCO}_3^-$ c) $\text{Ca}^{2+}, \text{NO}_3^-$ d) $\text{Na}^+, \text{SO}_4^{2-}$
37. Which of the following gas is insoluble in water?
 a) SO_2 b) NH_3 c) H_2 d) CO_2
38. Which will produce hard water?
 a) Saturation of water with CaSO_4
 b) Addition of Na_2SO_4 to water
 c) Saturation of water with CaCO_3
 d) Saturation of water with MgCO_3
39. In Bosch's process which gas is utilised for the production of hydrogen gas?
 a) Producer gas b) Water gas c) Coal gas d) None of these
40. Triple point of water is
 a) 203 K b) 193 K c) 273 K d) 373 K
41. The hybridization of the orbitals of oxygen in H_2O_2 is:
 a) sp^3d b) sp c) sp^2 d) sp^3
42. Which of the following pairs will not produce dihydrogen gas?
 a) $\text{Cu} + \text{HCl}$ (dil.) b) $\text{Fe} + \text{H}_2\text{SO}_4$ c) $\text{Mg} + \text{steam}$ d) $\text{Na} + \text{alcohol}$
43. Calgon used as water softner is
 a) $\text{Na}_2[\text{Na}_4(\text{PO}_3)_6]$ b) $\text{Na}_4[\text{Na}_2(\text{PO}_3)_6]$ c) $\text{Na}_2[\text{Na}_4(\text{PO}_4)_5]$ d) None of these
44. Permutit is:
 a) Hydrated sodium aluminium silicate
 b) Sodium hexa meta-phosphate
 c) Sodium silicate
 d) Sodium meta-aluminate
45. The rubber foam is produced by passing oxygen through rubber foaming material. This oxygen is released from:
 a) Nitric oxide b) Hydrogen peroxide c) Water d) CO_2
46. Which is the poorest reducing agent?
 a) Atomic hydrogen b) Nascent hydrogen
 c) Dihydrogen d) All have same reducing strength
47. In context with the industrial preparation of hydrogen from water gas ($\text{CO} + \text{H}_2$) which of the following is the correct statement.
 a) CO and H_2 are fractionally separated using differences in their densities
 b) CO is removed by absorption in aqueous Cu_2Cl_2 solution
 c) H_2 is removed through occlusion with Pd
 d) CO is oxidised to CO_2 with steam in the presence of a catalyst followed by absorption of CO_2 in alkali
48. The number of radioactive isotopes of hydrogen is:
 a) 1 b) 2 c) 3 d) None of these
49. The oxidation number of oxygen in hydrogen peroxide is
 a) +1 b) -1 c) +2 d) -2
50. The normality of 30 volume H_2O_2 is
 a) 2.678 N b) 5.336 N c) 8.034 N d) 6.685 N
51. Acidified solution of chromic acid on treatment with H_2O_2 yields:
 a) $\text{CrO}_3 + \text{H}_2\text{O} + \text{O}_2$ b) $\text{Cr}_2\text{O}_2 + \text{H}_2\text{O} + \text{O}_2$ c) $\text{CrO}_5 + \text{H}_2\text{O} + \text{K}_2\text{SO}_4$ d) $\text{H}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{O} + \text{O}_2$
52. The hair dyes available in the market generally contain two bottles, one containing the dye and the other hydrogen peroxide. Before applying the dye, the two solutions are mixed. The hydrogen peroxide.
 a) Is added to dilute the solution of the dye
 b) Oxidises the dye to give the desired colour
 c) Reduces the dye to give the desired colour
 d) Acidifies the solution of the dye

53. In periodic table tritium is placed in group:
a) I b) II c) III d) IV
54. The *ortho* and *para* hydrogen differ in respect of which of the following?
a) In the molecular weight b) In the nature of spin of protons
c) In the nature of spin of electrons d) In the number of protons
55. The bond energy of covalent O—H bond in water is:
a) Equal to bond energy of hydrogen bond
b) Greater than bond energy of hydrogen bond
c) Lesser than bond energy of hydrogen bond
d) None of the above
56. Water acts as excellent solvent due to:
a) Hydrogen bonding
b) Neutral nature
c) High dielectric constant
d) None of the above
57. $\text{TiH}_{1.73}$ is an example of which type of the hydride?
a) Metallic b) Ionic c) Covalent d) Polymeric
58. An aqueous solution of hydrogen peroxide is
a) Alkaline b) Neutral c) Strongly acidic d) Weakly acidic
59. The O—O bond length in H_2O_2 is:
a) 1.54 \AA b) 1.48 \AA c) 1.34 \AA d) 1.01 \AA
60. Moist hydrogen peroxide cannot be dried over conc. H_2SO_4 because
a) It can catch fire b) It is reduced by H_2SO_4
c) It is oxidised by H_2SO_4 d) It is decomposed by H_2SO_4
61. The strength in volumes of a solution containing 30.36g/L of H_2O_2 is
a) 10 volume b) 20 volume c) 5 volume d) None of these
62. Tritium emits:
a) α -particles b) β -particles c) γ -rays d) Neutrons
63. The ratio of electron, proton and neutron in tritium is:
a) 1 : 1 : 1 b) 1 : 1 : 2 c) 2 : 1 : 1 d) 1 : 2 : 1
64. Hydrogen directly combines with
a) Cu b) Au c) Ca d) Ni
65. In which of the following reactions, H_2O_2 is acting as a reducing agent?
a) $\text{SO}_2 + \text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{SO}_4$ b) $2\text{KI} + \text{H}_2\text{O}_2 \rightarrow 2\text{KOH} + \text{I}_2$
c) $\text{PbS} + 4\text{H}_2\text{O}_2 \rightarrow \text{PbSO}_4 + 4\text{H}_2\text{O}$ d) $\text{AgO}_2 + \text{H}_2\text{O}_2 \rightarrow 2\text{Ag} + \text{H}_2\text{O} + \text{O}_2$
66. Permutit is the technical name given to
a) Aluminates of calcium and sodium b) Hydrated silicate of aluminium and sodium
c) Silicates of calcium and magnesium d) Silicates of calcium and sodium
67. The best method to test whether a clear liquid is water, is to:
a) Taste the liquid
b) Smell the liquid
c) Add litmus paper
d) Add few drops on anhydrous copper sulphate and look for colour change
68. An inorganic compound liberates O_2 when heated, turns an acid solution of KI brown and reduces acidified KMnO_4 . The substance is:
a) H_2O_2 b) D_2O c) KNO_3 d) $\text{Pb}(\text{NO}_3)_2$
69. Heavy water is qualified as heavy because it is:
a) A heavy liquid
b) An oxide of a heavier isotope of oxygen

- c) An oxide of deuterium
d) Denser than water
70. Permanent hardness can be removed by adding
a) Cl_2 b) Na_2CO_3 c) Ca(OH)Cl d) K_2CO_3
71. The ionization energy of hydrogen is:
a) Lower than alkali metals
b) Lower than halogens
c) Closer to alkali metals
d) Closer to halogens
72. Which one of the following reactions represents the oxidising property of H_2O_2 ?
a) $2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 + 5\text{H}_2\text{O}_2 \rightarrow \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 8\text{H}_2\text{O} + 5\text{O}_2$
b) $2\text{K}_3[\text{Fe(CN)}_6] + 2\text{KOH} + \text{H}_2\text{O}_2 \rightarrow 2\text{K}_4[\text{Fe(CN)}_6] + 2\text{H}_2\text{O} + \text{O}_2$
c) $\text{Pb}_2 + \text{H}_2\text{O}_2 \rightarrow \text{PbO} + \text{H}_2\text{O} + \text{O}_2$
d) $2\text{KI} + \text{H}_2\text{SO}_4 + \text{H}_2\text{O}_2 \rightarrow \text{K}_2\text{SO}_4 + \text{I}_2 + 2\text{H}_2\text{O}$
73. Hydrogen peroxide is prepared in the laboratory by
a) Passing CO_2 into BaO_2 b) Adding MnO_2 to dil. H_2SO_4
c) Adding Na_2O_2 to cold water d) Adding PbO_2 into KMnO_4
74. Heavy water is
a) Water at 0°C
b) Water containing Fe, Cr, Mn
c) D_2O
d) Water obtained after a number of distillations
75. *Ortho* and *para* hydrogen differ in
a) Nuclear charge b) Nuclear reaction c) Electron spin d) Proton spin
76. Hydrogen peroxide is manufactured by the auto-oxidation of:
a) 2-ethylanthraquinol b) Anthraquinone c) Naphthalene d) Anthracene
77. What is the product of the reaction of H_2O_2 WITH Cl_2 ?
a) $\text{O}_2 + \text{HOCl}$ b) $\text{HCl} + \text{O}_2$ c) $\text{H}_2\text{O} + \text{HCl}$ d) $\text{HCl} + \text{H}_2$
78. One mole of calcium phosphide on reaction with excess water gives:
a) One mole of phosphene
b) Two moles of phosphoric acid
c) Two moles of phosphine
d) One mole of phosphorus pentaoxide
79. Hydrogen may be prepared by heating a solution of caustic soda with:
a) Mg b) Zn c) Fe d) Ag
80. H_2O_2 is manufactured these days
a) By the action of H_2O_2 on BaO_2 b) By the action of H_2SO_4 on Na_2O_2
c) By electrolysis of 50% H_2SO_4 d) By burning hydrogen in excess of oxygen
81. On bubbling CO_2 through a solution of barium peroxide in water:
a) O_2 is formed b) H_2CO_3 is formed c) H_2O_2 is formed d) H_2 is formed
82. The most reactive state of hydrogen is:
a) Atomic hydrogen b) Heavy hydrogen c) Molecular hydrogen d) Nascent hydrogen
83. The number of protons, electrons and neutrons respectively in a molecule of heavy water is:
a) 10, 10, 10 b) 8, 10, 11 c) 10, 11, 10 d) 11, 10, 10
84. Ordinary hydrogen is a mixture at:
a) 75% ortho H_2 + 25% para H_2
b) 25% ortho H_2 + 75% para H_2
c) 50% ortho H_2 + 50% para H_2
d) 99% para H_2 + 1%ortho H_2

85. Heavy water freezes at:
 a) -3.8°C b) 3.8°C c) 0°C d) 3.2°C
86. The electronic configuration of deuterium is:
 a) $1s^2$ b) $1s^1, 2s^2$ c) $1s^2, 2s^1$ d) $1s^1$
87. Smell of H_2O_2 resembles:
 a) Alcohol b) Alkali c) Nitric acid d) Chloroform
88. Hydrogen produced in contact with substance which is to be reduced is:
 a) Ortho H_2 b) Para H_2 c) Active H d) Nascent H
89. H_2O_2 acts as an oxidizing agent in:
 a) Neutral medium
 b) Acidic medium
 c) Alkaline medium
 d) acidic as well as in alkaline medium
90. The concentration of H_2O_2 solution of '10 volume' is
 a) 30% b) 3% c) 1% d) 10%
91. Water possesses a high dielectric constant, therefore
 a) It always contains ions b) It is universal solvent
 c) Can dissolve covalent compounds d) Can conduct electricity
92. Tailing of mercury is a laboratory test for:
 a) O_3 b) Hg c) Cl_2 d) O_2
93. Which method cannot be used to remove hardness of water?
 a) Clark's method
 b) By adding washing soda
 c) Calgon process
 d) Filtration
94. Which of the following could act as a propellant for rockets?
 a) Liq. $\text{H}_2 + \text{Liq. O}_2$ b) Liq. $\text{N}_2 + \text{Liq. O}_2$ c) Liq. $\text{H}_2 + \text{Liq. N}_2$ d) Liq. $\text{O}_2 + \text{Liq. Ar}$
95. When electric current is passed through an ionic hydride in the molten state,
 a) Hydrogen is liberated at the cathode
 b) Hydrogen is liberated at the anode
 c) Hydride ion migrates towards cathode
 d) No reaction takes place
96. Deuterium was discovered by:
 a) Urey b) Aston c) Rutherford d) Chadwick
97. The percentage by weight of hydrogen in H_2O_2 is:
 a) 50 b) 25 c) 6.25 d) 5.88
98. Ortho and para-hydrogen differ in the:
 a) Number of protons b) Molecular weight c) Nature of spins of protons d) Nature of spins of electrons
99. Decomposition of H_2O_2 is retarded by:
 a) Acetanilide b) Alcohol c) H_3PO_4 d) All of these
100. Heavy water possesses:
 a) Insoluble impurities like silica
 b) Impurities like carbonates and bicarbonates of calcium and magnesium
 c) High density and different physical properties than those of water
 d) The capacity to expedite the rate of nuclear reactions
101. Which element forms maximum compound in chemistry?
 a) O b) H c) Si d) C
102. The bleaching properties of H_2O_2 are due to its:

- a) Reducing properties b) Oxidizing properties c) Unstable nature d) Acidic nature
103. Which one of the following is called amphoteric solvent?
 a) Ammonium hydroxide b) Chloroform
 c) Benzene d) Water
104. The colour of hydrogen is
 a) Yellow b) Orange c) Black d) Colourless
105. The amount of H_2O_2 present in 1 L of 1.5 N H_2O_2 solution is:
 a) 2.5 g b) 25.5 g c) 3.0 g d) 8.0 g
106. H_2O_2 is prepared in the laboratory when:
 a) MnO_2 is added to dilute cold H_2SO_4
 b) BaO_2 is added to CO_2 bubbling through cold water
 c) PbO_2 is added to an acidified solution of $KMnO_4$
 d) Na_2O_2 is added to boiling water
107. Decolourisation of acidified potassium permanganate occurs when H_2O_2 is added to it. This is due to:
 a) Oxidation of $KMnO_4$
 b) Reduction of $KMnO_4$
 c) Both oxidation and reduction of $KMnO_4$
 d) None of the above
108. Which hydride is neutral?
 a) H_2S b) H_2O c) H_2Se d) H_2Te
109. Hydrogen burns with:
 a) Smoky flame b) Yellow flame c) Blue flame d) Pale yellow flame
110. Zeolites are extensively used in:
 a) Softening of water and catalyst b) Preparing heavy water c) Increasing the hardness of water d) Mond's process
111. Deuterium, an isotope of hydrogen is:
 a) Radioactive b) Non-radioactive c) Heaviest d) Lightest
112. Which is the lightest gas?
 a) Nitrogen b) Hydrogen c) Helium d) Oxygen
113. Temporary hardness is caused due to the presence of:
 a) $CaSO_4$ b) $CaCl_2$ c) $CaCO_3$ d) $Ca(HCO_3)_2$
114. H_2O_2 is:
 a) Diamagnetic b) Paramagnetic c) Ferromagnetic d) None of these
115. Commercial 11.2 volume H_2O_2 solution has a molarity of
 a) 1.0 b) 0.5 c) 11.2 d) 1.12
116. The life period of atomic hydrogen is:
 a) Only five minute
 b) Only one third of a second
 c) Only two hour
 d) 10 second
117. There is a sample of 20 volume of hydrogen peroxide solution. Calculate its strength
 a) 6.07% b) 3.035% c) 2.509% d) 4.045%
118. When the same amount of zinc is treated separately with excess of sulphuric acid and excess of sodium hydroxide, the ratio of volumes of hydrogen evolved is:
 a) 1 : 1 b) 1 : 2 c) 2 : 1 d) 9 : 4
119. Atomic hydrogen is obtained by:
 a) Electrolysis of heavy water
 b) Reaction of water with heavy metals
 c) Thermal decomposition of water

- d) Passing silent electric discharge through hydrogen at low pressure
120. Which loses weight on exposure to the atmosphere?
- Concentrated H_2SO_4
 - Solid NaOH
 - A saturated solution of CO_2
 - Anhydrous sodium carbonate
121. Which can adsorb large volumes of hydrogen gas?
- Colloidal solution of palladium
 - Finely divided nickel
 - Colloidal ferric hydroxide
 - Finely divided platinum
122. In the hydrogen peroxide molecule:
- Two hydrogen atoms are connected to one of the oxygen
 - All the four atoms are in the same plane
 - The four atoms are arranged in a non-linear and non-planar manner
 - O—H bonds are polar but molecule is non-polar
123. Fluorine reacts with water to form:
- Fluorine water
 - Oxygen
 - Ozone
 - Oxygen, ozone
124. The hardness of water sample containing 0.002 mole of magnesium sulphate dissolved in a litre of water is expressed as
- 20ppm
 - 200ppm
 - 2000ppm
 - 120ppm
125. Adsorbed hydrogen by palladium is known as
- Nascent
 - Atomic
 - Heavy
 - Occluded
126. When hydrogen peroxide is added to acidified potassium dichromate, a blue colour is produced due to formation of
- CrO_3
 - Cr_2O_3
 - CrO_5
 - CrO_4^{2-}
127. Which is false about H_2O_2 ?
- Act as both oxidising and reducing agent
 - Two OH bonds lie in the same plane
 - Pale blue liquid
 - Can be oxidised by ozone
128. The reaction of $\text{H}_2\text{S} + \text{H}_2\text{O}_2 \rightarrow \text{S} + 2\text{H}_2\text{O}$ manifests
- Reducing action of H_2O_2
 - Oxidising nature of H_2O_2
 - Alkaline nature of H_2O_2
 - Acidic nature of H_2O_2
129. The reagent commonly used to determine hardness of water titrimetrically is
- Oxalic acid
 - Sodium thiosulphate
 - Sodium citrate
 - Disodium salt of EDTA
130. Ordinary hydrogen has preponderance of:
- Hydrogen atoms
 - Deuterium atoms
 - Tritium atoms
 - The above three are in equal proportions
131. Benzene is oxidized by H_2O_2 in presence of FeSO_4 to :
- Phenol
 - Cyclohexane
 - Benzaldehyde
 - Benzoic acid
132. Which of the following is an example of interstitial hydride?
- NH_3
 - CH_4
 - ZnH_2
 - H_2O
133. If water is boiled for sometime it becomes free from:
- Permanent hardness
 - Temporary hardness

- c) Suspended matter
d) Temporary hardness and dissolved gases
134. Polyphosphates are used as water softening agents because they
a) Form soluble complexes with anionic species
b) Precipitate anionic species
c) Precipitate cationic species
d) Form soluble complexes with cationic species
135. When two ice cubes are pressed over each other they unite to form one cube. Which of the following forces are responsible to hold them together?
a) Ionic interaction
b) Van der Waals' forces
c) Covalent interaction
d) Hydrogen bond formation
136. The pH of a solution of H_2O_2 is 6.0. Some chloride gas is bubbled into this solution. Which of the following is correct?
a) The pH of resultant solution becomes 8.0
b) Hydrogen gas is liberated from resultant solution
c) The pH of resultant solution becomes less than 6.0 and oxygen gas is liberated
d) Cl_2O is formed in the resultant solution
137. Permanent hardness of water can be removed by adding Calgon (NaPO_3)_n. This is an example of:
a) Adsorption b) Exchange of ion c) Precipitation d) None of these
138. Hydrogen molecules are:
a) Monoatomic and form X_2^{2-} ions
b) Diatomic and form X_2^{2-} ions
c) Diatomic and form X^- ions
d) Monoatomic and form X^- ions
139. Hydrogen reacts with even in the dark.
a) Br_2 b) F_2 c) I_2 d) Cl_2
140. 1000 g aqueous solution of CaCO_3 contains 10 g of calcium carbonate. Hardness of the solution is:
a) 10 ppm b) 100 ppm c) 1000 ppm d) 10000 ppm
141. Metal which does not react with cold water but evolves H_2 with steam is:
a) Na b) K c) Pt d) Fe
142. The pair that yields the same gaseous product on reaction with water:
a) K and KO_2 b) Ca and CaH_2 c) Na and Na_2O_2 d) Ba and BaO_2
143. The heaviest among the following is:
a) Deuterium b) Helium c) Tritium d) Hydrogen
144. The molarity of a 100 mL solution containing 5.1 g of hydrogen peroxide is:
a) 0.15 M b) 1.5 M c) 3.0 M d) 50.0 M
145. The metal that does not displace hydrogen from an acid is:
a) Hg b) Zn c) Al d) Ca
146. Deionised water is obtained by passing hard water through
a) Anion exchanger b) Zeolite
c) Cation exchanger d) Both anion and cation exchanger
147. The strength in volumes of a solution containing 30.36 g/L of H_2O_2 is
a) 10 V b) 5 V c) 20 V d) None of these
148. Hydrogen was discovered by:
a) Scheele b) Berzelius c) Cavendish d) Priestley
149. Hard water becomes free from ions when passed through ion exchange resin containing RCOOH groups.




- a) Cl^- b) SO_4^{2-} c) H_3O^+ d) Ca^{2+}
150. The sum of number of neutrons and protons in one of the isotopes of hydrogen is:
 a) 3 b) 4 c) 5 d) 6
151. Water contracts on heating:
 a) To 100°C b) From 0°C to 4°C c) To 273 K d) From 10°C to 20°C
152. Hydrogen combines directly with:
 a) Ca b) Cu c) Zn d) Fe
153. H_2O_2 restores the colour of old lead paintings, blackened by the action of H_2S gas, by:
 a) Converting PbO_2 to Pb
 b) Oxidising PbS to PbSO_4
 c) Converting PbCO_3 to Pb
 d) Oxidising PbSO_3 to PbSO_4
154. 10 volumes of H_2O_2 has a strength of approximately:
 a) 3% b) 30% c) 10% d) 5%
155. Ammonium persulphate solution on heating under reduced pressure gives:
 a) H_2O_2 b) O_2 c) H_2 d) $(\text{NH}_4)_2\text{SO}_4$
156. Which statement about zeolite is false?
 a) They are used as cation exchanger
 b) They have open structure which enables them to take up small molecules
 c) Zeolites are aluminosilicates having three dimensional network
 d) Some of the SiO_4^{4-} units are replaced by AlO_4^{5-} and AlO_6^{9-} ions in zeolites
157. Which of the following metal evolves hydrogen on reacting with cold dilute HNO_3 ?
 a) Fe b) Cu c) Al d) Mg
158. The reaction of water with sodium and potassium is
 a) Endothermic b) Reversible
 c) Exothermic d) Irreversible and endothermic
159. High boiling point of water is due to:
 a) Its high specific heat
 b) Hydrogen bonding
 c) High dielectric constant
 d) Low dissociation constant
160. Ozone reacts with H_2O_2 to give oxygen. One volume of ozone gives:
 a) One volume of oxygen
 b) Half volume of oxygen
 c) 1.5 volume of oxygen
 d) Two volumes of oxygen
161. Which of the following statements do not define the characteristic property of water "water is a universal solvent".?
 a) It has high liquid range
 b) It has very low dielectric constant
 c) It can dissolve maximum number of compounds
 d) None of the above
162. Sodium zeolite is:
 a) $\text{NaAlSi}_2\text{O}_6$ b) $\text{Na}_2\text{Al}_2\text{Si}_2\text{O}_3$ c) $\text{Na}_2\text{Al}_2\text{Si}_2\text{O}_8$ d) $\text{NaAl}_2\text{Si}_2\text{O}_8$
163. Acidified KMnO_4 is decolourised by:
 a) Oxygen b) Hydrogen c) Nitric oxide d) Nascent hydrogen
164. The oxidizing property of H_2O_2 is best explained by assuming that two oxygen atoms in its molecule are:
 a) Bonded differently
 b) Bonded similarly


- c) Bonded covalently
d) Bonded by hydrogen bonds
165. H_2O_2 is stored in:
a) Iron container after addition of stabilizer
b) Glass container after addition of stabilizer
c) Plastic container after addition of stabilizer
d) None of the above
166. Hydrogen is not used for:
a) Manufacture of vegetable ghee
b) Production of high temperature
c) As rocket fuel with kerosene
d) As a reducing agent
167. H_2O_2 is concentrated by:
a) Steam distillation
b) Fractional distillation
c) Freezing in freezing mixture
d) Distillation under reduced pressure
168. Pure H_2 is obtained by the action of:
a) Al over KOH
b) NaH over H_2O
c) Electrolysis of warm solution of $\text{Ba}(\text{OH})_2$ using Ni electrodes
d) All of the above
169. Heavy water is manufactured in India at:
a) Delhi b) Trombay c) Bhilai d) None of these
170. What is formed when calcium carbide reacts with heavy water?
a) C_2D_2 b) CaD_2 c) $\text{Ca}_2\text{D}_2\text{O}$ d) CD_2
171. The ionization of hydrogen atom gives:
a) Hydride ion b) Hydronium ion c) Proton d) Hydroxyl ion
172. Which is not true in case of H_2O_2 ?
a) It is more stable in basic solution
b) It acts as strong oxidizing agent in acid and basic solutions
c) It is decomposed by MnO_2
d) It behaves as reducing agent towards KMnO_4
173. Which one of the following is a true peroxide?
a) SO_2 b) MnO_2 c) NO_2 d) BaO_2
174. What is the volume of "20 volume H_2O_2 " required to get 5000 cm^3 of oxygen at STP?
a) 250 cm^3 b) 20 cm^3 c) 100 cm^3 d) 125 cm^3
175. The melting points of most of the solid substances increase with an increase of pressure. However, ice melts at a temperature lower than its usual melting point when the pressure is increased. This is because:
a) Ice is less denser than water
b) Pressure generates heat
c) The chemical bonds break under pressure
d) Ice is not a true solid
176. Heavy water was discovered by:
a) Nernst b) Haber c) Urey and Washburn d) Aston
177. The maximum possible number of hydrogen bonds a water molecule can form is:
a) 1 b) 2 c) 3 d) 4
178. H_2O_2 acts as antiseptic due to its:

- a) Reducing property b) Oxidizing property c) Bleaching property d) Acidic property
179. Hydrogen gas will not reduce:
 a) Heated cupric oxide
 b) Heated ferric oxide
 c) Heated stannic oxide
 d) Heated aluminium oxide
180. Which pair does not show hydrogen isotopes?
 a) *Ortho* and *para* hydrogen b) Protium and deuterium
 c) Deuterium and tritium d) Tritium and protium
181. The hardness of water is due to Metal ions.
 a) Ca^{2+} and Na^+ b) Mg^{2+} and K^+ c) Ca^{2+} and Mg^{2+} d) Zn^{2+} and Ba^{2+}
182. Under what conditions of temperature and pressure, the formation of atomic hydrogen from molecular hydrogen will be favoured more?
 a) High temperature and low pressure
 b) Low temperature and low pressure
 c) High temperature and high pressure
 d) Low temperature and high pressure
183. Heavy hydrogen is used:
 a) In filling the balloons
 b) In studying reaction mechanism
 c) In calculating heat of formation
 d) Iron hydroxide precipitates
184. The reaction, $\text{H}_2\text{S} + \text{H}_2\text{O}_2 \rightarrow \text{S} + 2\text{H}_2\text{O}$ manifests:
 a) Acidic nature of H_2O_2
 b) Alkaline nature of H_2O_2
 c) Oxidizing nature of H_2O_2
 d) Reducing nature of H_2O_2
185. Decomposition of H_2O_2 is accompanied by:
 a) Decrease in free energy
 b) Increase in free energy
 c) No change in free energy
 d) Evolution of heat
186. Which of the following statements is correct? Dielectric constant of H_2O_2
 a) Increases with dilution b) Decreases with dilution
 c) Is unaffected on dilution d) None of the above
187. Heavy water is not used for drinking because:
 a) It is poisonous
 b) It is costly
 c) Its physiological action is different from ordinary water
 d) Its chemical properties are different from ordinary water
188. Maximum density of heavy water is at:
 a) 0°C b) 4°C c) 11.6°C d) 3.8°C
189. The catalyst used in Bosch process of manufacture of H_2 is:
 a) Finely divided Ni b) V_2O_5 c) Pd d) $\text{Fe}_2\text{O}_3 + \text{Cr}_2\text{O}_3$
190. In which of the following reactions, H_2O_2 behaves as a reducing agent?
 a) $\text{Na}_2\text{SO}_3(\text{aq}) + \text{H}_2\text{O}_2(\text{aq}) \rightarrow \text{Na}_2\text{SO}_4(\text{aq}) + \text{H}_2\text{O}(\text{l})$
 b) $\text{PbO}_2(\text{s}) + \text{H}_2\text{O}_2(\text{aq}) \rightarrow \text{PbO}(\text{s}) + \text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$
 c) $2\text{KI}(\text{aq}) + \text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{KOH}(\text{aq}) + \text{I}_2(\text{s})$
 d) $\text{KNO}_2(\text{aq}) + \text{H}_2\text{O}_2(\text{aq}) \rightarrow \text{KNO}_3(\text{aq}) + \text{H}_2\text{O}(\text{l})$

191. Among CaH_2 , NH_3 , NaH and B_2H_6 , which are covalent hydride?
 a) NH_3 and B_2H_6 b) NaH and CaH_2 c) NaH and NH_3 d) CaH_2 and B_2H_6
192. In which reaction hydrogen is not formed?
 a) Copper and hydrochloric acid
 b) Iron and sulphuric acid
 c) Magnesium and steam
 d) Sodium and alcohol
193. The adsorption of hydrogen by metals is called
 a) Adsorption b) Occlusion c) Hydrogenation d) Dehydrogenation
194. A molten ionic hydride on electrolysis gives:
 a) H^+ ion moving towards the cathode
 b) H^+ ion moving towards the anode
 c) H_2 is liberated at anode
 d) H_2 is liberated at cathode
195. Moist hydrogen cannot be dried over concentrated H_2SO_4 because:
 a) It can catch fire
 b) It is reduced by H_2SO_4
 c) It is oxidized by H_2SO_4
 d) It decomposes H_2SO_4
196. Both temporary and permanent hardness are removed on boiling water with:
 a) $\text{Ca}(\text{OH})_2$ b) Na_2CO_3 c) CaCO_3 d) CaO
197. The weight percentage of deuterium in heavy water is:
 a) 22 b) 11.11 c) 4 d) 20
198. Very pure hydrogen (99.9%) can be made by which of the following processes?
 a) Mixing natural hydrocarbons of high molecular weight
 b) Electrolysis of water
 c) Reaction of salt like hydrides with water
 d) Reaction of methane with steam
199. Density of water is maximum at:
 a) 0°C b) 100°C c) 4°C d) 0 K
200. The most reactive isotope of H is:
 a) ${}_1\text{H}^1$
 b) ${}_1\text{H}^2$
 c) ${}_1\text{H}^3$
 d) All the same reactivity
201. Heavy water is used in atomic reactor as
 a) Moderator b) Coolant
 c) Both moderator and coolant d) Neither coolant nor moderator
202. The exhausted Permutit is generally regenerated by percolating through it a solution of:
 a) Sodium chloride b) Calcium chloride c) Magnesium chloride d) Potassium chloride
203. The best explanations for not placing hydrogen with the group of alkali metals or halogens is:
 a) Hydrogen can form compounds with all other elements
 b) Hydrogen is much lighter element than the alkali metals or the halogens
 c) The ionization energy of hydrogen is too high for group of alkali metals but too low for halogen group
 d) None of the above
204. Hydrogen molecule differs from chlorine molecule in the following respect.
 a) Hydrogen molecule is non-polar but chlorine molecule is polar
 b) Hydrogen molecule is polar while chlorine molecule is non-polar
 c) Hydrogen molecule can form intermolecular hydrogen bonds but chlorine molecule does not

- d) Hydrogen molecule cannot participate in coordinate bond formation but chlorine molecule can
205. The geometry of water molecule is same as that of:
 a) CO_2 b) C_2H_4 c) Chlorine oxide d) Boron trifluoride
206. Hydrogen peroxide does not:
 a) Liberate iodine from KI
 b) Turn the titanium salt yellow
 c) Give silver peroxide with moist silver oxide
 d) Turn the mixture of aniline, KClO_3 and dil. H_2SO_4 violet
207. The most dangerous method of preparing hydrogen would be by the action of dil. HCl and:
 a) Zn b) Fe c) K d) Al
208. When zeolite which is hydrated sodium aluminium silicate is treated with hard water, the sodium ions are exchanged with
 a) H^+ ions b) Mg^{2+} ion c) Ca^{2+} ion d) both Ca^{2+} and Mg^{2+}
209. Hydrolysis of one mole of peroxodisulphuric acid produces:
 a) Two moles of sulphuric acid
 b) Two moles of peroxomonosulphuric acid
 c) One mole of sulphuric acid and one mole of peroxomonosulphuric acid
 d) One mole of sulphuric acid, one mole of peroxomonosulphuric acid and one mole of hydrogen peroxide
210. During hydrogenation of oil the catalyst commonly used is:
 a) Pd on CuCl_2 b) Ni c) Fe d) V_2O_5
211. Oxygen and hydrogen react to form water. This discovery was made by:
 a) Priestley b) Cavendish c) Scheele d) Newton
212. Which one of the following processes will produce hard water?
 a) Saturation of water with CaCO_3 b) Saturation of water with MgCO_3
 c) Saturation of water with CaSO_4 d) Addition of Na_2SO_4 to water
213. The oxygen atom of H_2O_2 used for oxidation is bound by:
 a) Electrovalent bond b) Co-ordinate bond c) Covalent bond d) None of these
214. Which reaction shows oxidizing nature of H_2O_2 ?
 a) $\text{H}_2\text{O}_2 + 2\text{KI} \rightarrow 2\text{KOH} + \text{I}_2$
 b) $\text{Cl}_2 + \text{H}_2\text{O}_2 \rightarrow 2\text{HCl} + \text{O}_2$
 c) $\text{H}_2\text{O}_2 + \text{Ag}_2\text{O} \rightarrow 2\text{Ag} + \text{H}_2\text{O} + \text{O}_2$
 d) $\text{NaClO} + \text{H}_2\text{O}_2 \rightarrow \text{NaCl} + \text{H}_2\text{O} + \text{O}_2$
215. H_2O_2 is manufactured these days
 a) By burning hydrogen in excess of oxygen
 b) By the action of H_2O_2 on BaO_2
 c) By the action of H_2SO_4 on Na_2O_2
 d) By electrolysis of 50% H_2SO_4
216. MnO_2 liberates oxygen from a solution of H_2O_2 (the action being catalytic) only if the solution is:
 a) Basic b) Acidic c) Neutral d) None of these
217. Ionic hydrides react with water to give
 a) Hydride ions b) Acidic solutions c) Protons d) Basic solutions
218. Hydrogen is evolved by the action of cold dilute HNO_3 on:
 a) Fe b) Mg or Mn c) Cu d) Al
219. Hydrogen peroxide for the first time was prepared by:
 a) Priestley b) Thenard c) Gay-Lussac d) Bernard
220. Which pair does not show hydrogen isotopes?
 a) *Ortho* hydrogen and *para* hydrogen b) Protium and deuterium
 c) Deuterium and tritium d) Tritium and protium
221. The strength of 10 volume of H_2O_2 solution is


- a) 10 b) 68 c) 60.70 d) 30.36
222. The conversion of atomic hydrogen into ordinary hydrogen is:
 a) Exothermic change
 b) Endothermic change
 c) Nuclear change
 d) Photochemical change
223. Para hydrogen is:
 a) Less stable than ortho hydrogen
 b) More stable than ortho hydrogen
 c) As stable as ortho hydrogen
 d) None of the above
224. Some statements about heavy water are given below:
 (i) Heavy water is used as a moderator in nuclear reactors
 (ii) Heavy water is more associated than ordinary water
 (iii) Heavy water is more effective solvent than ordinary water
 Which of the above statements are correct?
 a) (i) and (ii) b) (i), (ii) and (iii) c) (ii) and (iii) d) (i) and (iii)
225. H_2O_2 acts as a reducing agent in its:
 a) Reaction with a ferrous salt
 b) Reaction with iodides
 c) Reaction with lead sulphide
 d) Reaction with KMnO_4 in acidic medium
226. When hydrolith is treated with water it yields:
 a) H_2 b) H_2O_2 c) N_2 d) NaH
227. Atomic hydrogen produces formaldehyde when it reacts with:
 a) CO_2 b) CO c) O_2 d) C_2H_2
228. K_a for H_2O_2 is of the order of:
 a) 10^{-12} b) 10^{-14} c) 10^{-16} d) 10^{-10}
229. Which one of the following reaction does not form gaseous product?
 a) $\text{PbO}_2 + \text{H}_2\text{O}_2 \rightarrow$ b) Acidified $\text{KMnO}_4 + \text{H}_2\text{O}_2 \rightarrow$
 c) $\text{PbS} + \text{H}_2\text{O}_2 \rightarrow$ d) $\text{Cl}_2 + \text{H}_2\text{O}_2 \rightarrow$
230. The structure of H_2O_2 is:
 a)  b)  c) $\text{H}-\text{O}-\text{O}-\text{H}$ d) 
231. Which cannot be oxidised by H_2O_2 ?
 a) Na_2SO_3 b) PbS c) KI d) O_3
232. A mixture of hydrazine and 40 to 60 per cent of H_2O_2 solution is:
 a) Antiseptic b) Rocket fuel c) Germicide d) Insecticide
233. Hydrogen peroxide is now generally prepared on industrial scale by the:
 a) Action of H_2SO_4 on barium peroxide
 b) Action of H_2SO_4 on sodium peroxide
 c) Electrolysis of 50% H_2SO_4
 d) Burning hydrogen in excess of oxygen
234. The equilibrium molecular structure of hydrogen peroxide is
 a) Planar as given below b) Linear






- c) Tetrahedral
d) Non-planar
235. A given solution of H_2O_2 is 30 volume. Its concentration in terms of molarity is:
a) 9.1 M b) 2.68 M c) 2.5 M d) 26.8 M
236. H_2O_2 turns an acidified solution of to orange red.
a) BaO_2 b) PbO_2 c) Na_2O_2 d) TiO_2
237. Tritium is obtained by:
a) Nuclear reactions
b) Passing steam over heated C
c) Action of NaOH on Al
d) Action of H_2SO_4 on Zn
238. In the case of H_2O_2 , the angle between the planes containing the hydrogen atom is:
a) 100° b) 90° c) $109^\circ 28'$ d) 180°
239. In laboratory, H_2O_2 is prepared by
a) Cold $\text{H}_2\text{SO}_4 + \text{BaO}_2$
b) $\text{HCl} + \text{BaO}_2$
c) conc $\text{H}_2\text{SO}_4 + \text{Na}_2\text{O}_2$
d) $\text{H}_2 + \text{O}_2$
240. The formula of heavy water is:
a) H_2O^{18} b) D_2O c) T_2O d) H_2O^{17}
241. Hydrogen resembles in many of its properties with:
a) Alkali metals b) Halogens c) Both (a) and (b) d) None of these
242. Hydrogen is not obtained when zinc reacts with
a) Cold water b) hot NaOH solution c) dil. H_2SO_4 d) dil. HCl
243. The H-O-H angle in water molecule is about
a) 105° b) 102° c) 180° d) 90°
244. Hydrogen adsorbed on palladium is known as:
a) Atomic H b) Nascent H c) Occluded H d) Heavy H
245. Hydrogen molecule differs from chlorine molecule in the following respect
a) Hydrogen molecule is non-polar but chlorine molecule is polar
b) Hydrogen molecule is polar while chlorine molecule is non-polar
c) Hydrogen molecule can form intermolecular hydrogen bonds but chlorine molecule does not
d) Hydrogen molecule cannot participate in coordination bond formation but chlorine molecule can
246. Decomposition of H_2O_2 is accelerated by:
a) Traces of acids b) Finely divided metals c) Acetanilide d) Alcohol
247. Which one of the following is used for reviving the exhausted 'permutit'?
a) HCl solution b) 10% CaCl_2 solution c) 10% MgCl_2 solution d) 10% NaCl solution
248. The volume strength of 1.5 N H_2O_2 solution is
a) 16.8 L b) 8.4 L c) 4.2 L d) 5.2 L
249. Nascent hydrogen consists of:
a) Hydrogen atoms with excess energy
b) Hydrogen molecules with excess energy
c) Hydrogen ions in excited state
d) Solvated protons
250. At absolute zero:
a) Only para hydrogen exists

- b) Only ortho hydrogen exists
 c) Both para and ortho hydrogen exist
 d) None of the above
251. Hydrogen peroxide works as:
 a) An oxidant only
 b) A reductant only
 c) An acid only
 d) An oxidant, a reductant and an acid
252. Which of the following will not give H_2O_2 on hydrolysis?
 a) HClO_4 b) $\text{H}_2\text{S}_2\text{O}_8$ c) H_2SO_5 d) HNO_4 (pernitric acid)
253. The n/p ratio for ${}_1\text{H}^1$ is:
 a) 1 b) 2 c) 3 d) Zero
254. The percentage by weight of hydrogen in H_2O_2 is:
 a) 5.88 b) 6.25 c) 25 d) 50
255. Exhausted permutit does not containion.
 a) Na^+ b) Mg^{2+} c) Al^{3+} d) Si^{4+}
256. The molarity of pure water at 4°C is:
 a) 1 M b) 2.5 M c) 5 M d) 55.5 M
257. The gas used in the hydrogenation of oils in presence of nickel as a catalyst is:
 a) Methane b) Ethane c) ozone d) Hydrogen
258. The volume of oxygen liberated from 0.68 g of H_2O_2 is
 a) 112mL b) 224mL c) 56mL d) 336mL
259. Which hydride is an ionic hydride?
 a) NH_3 b) H_2S c) $\text{TiH}_{1.73}$ d) NaH
260. H_2O_2 reduces $\text{K}_3\text{Fe}(\text{CN})_6$ in:
 a) Neutral solution b) Acidic solution c) Alkaline solution d) Non-polar medium
261. Point out the incorrect statement.
 a) Hardness of water depends upon its soap consuming power
 b) Temporary hardness is due to bicarbonates of calcium and magnesium
 c) Permanent hardness is due to soluble sulphates, chlorides and nitrates of Ca and Mg
 d) Permanent hardness can be removed by boiling water
262. H_2O_2 converts potassium ferrocyanide to ferricyanide. The change observed in the oxidation state of iron is:
 a) $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$ b) $\text{Fe} \rightarrow \text{Fe}^{2+}$ c) $\text{Fe}^{3+} \rightarrow \text{Fe}^{2+}$ d) $\text{Fe}^{2+} \rightarrow \text{Fe}^+$
263. Which of the following is correct about heavy water?
 a) Water at 4°C having maximum density is known as heavy water
 b) It is formed by the combination of heavier isotope of hydrogen and oxygen
 c) It is heavier than water
 d) None of the above
264. Hydrogen is prepared on large scale for industrial use
 a) by $\text{Zn} + \text{H}_2\text{SO}_4$ b) by $\text{Al} + \text{NaOH}$ c) by $\text{Na} + \text{C}_2\text{H}_5\text{OH}$ d) From water gas
265. Hydrogen is obtained by the action of an alloy of silicon and iron with NaOH. The process is called:
 a) Wood process b) Bosch process c) Haber process d) Silicol process
266. In transforming 0.01 mole of PbS to PbSO_4 , the volume of 10 volume H_2O_2 required will be
 a) 11.2mL b) 22.4mL c) 33.6mL d) 44.8mL
267. Hydrogen peroxide when added to a solution of potassium permanganate acidified with sulphuric acid
 a) Forms water only
 b) Acts as an oxidising agent

- c) Acts as a reducing agent
d) Reduces sulphuric acid
268. Water is oxidised to oxygen by
a) ClO_2 b) KMnO_4 c) H_2O_2 d) Fluorine
269. The most abundant element in the universe is thought to be
a) Carbon b) Oxygen c) Hydrogen d) Nitrogen
270. In the preparation of hydrogenated oil the chemical reaction involving hydrogen is called:
a) Hydrogenation b) Reduction c) Dehydrogenation d) Oxidation
271. The most abundant isotope of hydrogen is:
a) Tritium b) Deuterium c) Protium d) Para-hydrogen
272. Which statement is not correct for hydrogen peroxide?
a) Pure H_2O_2 is fairly stable
b) It sometimes acts as a reducing agent
c) It acts as an oxidizing agent
d) Aqueous solution of H_2O_2 is weakly basic
273. Which one is correct for perhydrol?
a) It is 30% H_2O_2 or 100 vol. H_2O_2
b) Its molarity is 8.8 M
c) It is used as antiseptic and germicide
d) All of the above
274. Hydrogen has a tendency to gain one electron in order to acquire helium configuration. It thus, resembles:
a) Alkali metals b) Noble gases c) Halogens d) Alkaline earth metals
275. Calgon is an industrial name given to:
a) Normal sodium phosphate
b) Sodium meta-aluminate
c) Sodium hexa meta-phosphate
d) Hydrated sodium aluminium silicate
276. For the bleaching of hair, the substance used is:
a) SO_2 b) Bleaching powder c) H_2O_2 d) O_3
277. In solid hydrogen, the intermolecular bonding is:
a) Ionic b) Van der Waals' c) Metallic d) Covalent
278. The species that does not contains peroxide ions is:
a) PbO_2 b) H_2O_2 c) SrO_2 d) BaO_2
279. The critical temperature of water is higher than that of O_2 because H_2O molecule has:
a) Fewer electrons than oxygen
b) Two covalent bonds
c) V-shape
d) Dipole moment
280. Pure H_2O_2 is:
a) Colourless liquid
b) A gas
c) Blue syrupy liquid
d) Pale blue syrupy liquid
281. When silicon is boiled with caustic soda solution, the gas evolved is:
a) O_2 b) SiH_4 c) H_2 d) None of these
282. In which of the following reactions hydrogen peroxide is a reducing agent?
a) $\text{H}_2\text{SO}_3 + \text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{SO}_4 + \text{H}_2\text{O}$
b) $2\text{HI} + \text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{I}_2$
c) $2\text{FeCl}_2 + 2\text{HCl} + \text{H}_2\text{O}_2 \rightarrow 2\text{FeCl}_3 + 2\text{H}_2\text{O}$



283. Which does not react with cold water?



284. Deuterium resembles hydrogen in chemical properties but reacts:

a) Slower than hydrogen

b) Faster than hydrogen

c) More vigorously than hydrogen

d) Just as hydrogen

