

GPLUS EDUCATION


Date :
Time :
Marks :

PHYSICS


MOVING CHARGES AND MAGNETISM

Single Correct Answer Type

1. A particle moving in a magnetic field increases its velocity then its radius of the circle
a) Decreases b) Increases c) Remains the same d) Becomes half
2. A wire carrying current I and other carrying $2I$ in the same direction produces a magnetic field B at the mid point. What will be the field when $2I$ wire is switched off
a) $B/2$ b) $2B$ c) B d) $4B$
3. Two long parallel copper wires carry currents of $5A$ each in opposite directions. If the wires are separated by a distance of $0.5m$, then the force between the two wires is
a) $10^{-5}N$, attractive b) $10^{-5}N$, repulsive
c) $2 \times 10^{-5}N$, attractive d) $2 \times 10^{-5}N$, repulsive
4. An infinitely long straight conductor AB is fixed and a current is passed through it. Another movable straight wire CD of finite length and carrying current is held perpendicular to it and released. Neglect weight of the wire







- a) The rod CD will move upwards parallel to itself
b) The rod CD will move downward parallel to itself
c) The rod CD will move upward and turn clockwise at the same time
d) The rod CD will move upward and turn anti-clockwise at the same time
5. Two similar coils of radius R are lying concentrically with their planes at right angles to each other. The currents flowing in them are I and $2I$, respectively. The resultant magnetic field induction at the centre will be
a) $\frac{\sqrt{5}\mu_0 I}{2R}$ b) $\frac{3\mu_0 I}{2R}$ c) $\frac{\mu_0 I}{2R}$ d) $\frac{\mu_0 I}{R}$
6. A current (i) carrying circular wire of radius R is placed in a magnetic field B perpendicular to its plane. The tension T along the circumference of wire is



- a) BiR b) $2\pi BiR$ c) πBiR d) $2BiR$
7. A current of i ampere flows in a circular area of wire which subtends an angle of $(3\pi/2)$ radian at its centre, whose radius is R . The magnetic induction B at the centre is
a) $\mu_0 i/R$ b) $\mu_0 i/2R$ c) $2\mu_0 i/R$ d) $3\mu_0 i/8R$
8. A wire in the form of a circular loop of one turn carrying a current produces a magnetic field B at the centre. If the same wire is looped into a coil of two turns and carries the same current, the new value of magnetic induction at the centre is
a) $3B$ b) $5B$ c) $4B$ d) $2B$

9. In which orientation the resultant magnetic moment of two magnets, will be zero, if magnetic moment of each magnets is M in the following figures?



10. A square frame of side 1 m carries a current i , produces a magnetic field B at its centre. The same current is passed through a circular coil having the same perimeter as the square. The magnetic field at the centre of the circular coil is B' . The ratio B/B' is

a) $\frac{8}{\pi^2}$

b) $\frac{8\sqrt{2}}{\pi^2}$

c) $\frac{16}{\pi^2}$

d) $\frac{16}{\sqrt{2}\pi^2}$

11. Two charged particles are projected into a region in which a magnetic field is perpendicular to their velocities. After they enter the magnetic field, you can conclude that

- a) The charges are deflected in opposite directions
b) The charges continue to move in a straight line
c) The charges move in circular paths
d) The charges move in circular paths but in opposite directions

12. Through two parallel wires A and B , 10A and 2A of currents are passed respectively in opposite directions. If the wire A is infinitely long and the length of the wire B is 2m, then force on the conductor B , which is situated at 10 cm distance from A , will be

a) 8×10^{-7} N

b) 8×10^{-5} N

c) 4×10^{-7} N

d) 4×10^{-5} N

13. A circular loop carrying a current is replaced by an equivalent magnetic dipole. A point on the axis of the loop is

- a) An end-on position
b) A broad side-on position
c) Both (a) and (b)
d) Neither (a) nor (b)

14. A horizontal rod of mass 10 gm and length 10 cm is placed on a smooth plane inclined at an angle of 60° with the horizontal, with the length of the rod parallel to the edge of the inclined plane. A uniform magnetic field of induction B is applied vertically downwards. If the current through the rod is 1.73 ampere, then the value of B for which the rod remains stationary on the inclined plane is

a) 1.73 tesla

b) $\frac{1}{1.73}$ tesla

c) 1 tesla

d) None of the above

15. An electron revolves in a circle of radius 0.4 \AA with a speed of 10^5 ms^{-1} . The magnitude of the magnetic field, produced at the center of the circular path due to the motion of the electron, in weber metre^{-2} is


a) 0.01

b) 10.0

c) 1.0

d) 0.005

16. A 100 turns coil shown in figure carries a current of 2 amp in a magnetic field $B = 0.2 \text{ Wb/m}^2$. The torque acting on the coil is



- a) 0.32 Nm tending to rotated the side AD out of the page
b) 0.32 Nm tending to rotated the side AD into the page
c) 0.0032 Nm tending to rotated the side AD out of the page
d) 0.0032 Nm tending to rotated the side AD into the page

17. Two concentric coils each of radius equal to $2\pi \text{ cm}$ are placed at right angles to each other. 3 A and 4 A are the currents flowing in each coil respectively. The magnetic induction in Wbm^{-2} at the centre of the coils will be ($\mu_0 = 4\pi \times 10^{-7} \text{ WbAm}^{-1}$)

a) 12×10^{-5}

b) 10^{-5}

c) 5×10^{-5}

d) 7×10^{-5}

18. A thin circular wire carrying a current I has a magnetic moment M . The shape of the wire is changed to a

square and it carries the same current. It will have a magnetic moment

- a) M b) $\frac{4}{\pi^2}M$ c) $\frac{4}{\pi}M$ d) $\frac{\pi}{4}M$


19. Two parallel long wires carry currents i_1 and i_2 with $i_1 > i_2$. When the currents are in the same direction, the magnetic field midway between the wires is $10\mu T$. When the direction of i_2 is reversed, it becomes $40\mu T$. The ratio i_1/i_2 is
a) 3 : 4 b) 11 : 7 c) 7 : 11 d) 5 : 3
20. A straight wire of length (π^2) metre is carrying a current of $2A$ and the magnetic field due to it is measured at a point distance 1 cm from it. If the wire is to be bent into a circle and is to carry the same current as before, the ratio of the magnetic field at its centre to that obtained in the first case would be
a) 50 : 1 b) 1 : 50 c) 100 : 1 d) 1 : 100
21. A proton of energy 8 eV is moving in a circular path in a uniform magnetic field. The energy of an alpha particle moving in the same magnetic field and along the same path will be
a) 4 eV b) 2 eV c) 8 eV d) 6 eV
22. An electron and proton having same kinetic energy enter into magnetic field perpendicular to it. Then
a) The path of electron is less curved b) The path of proton is less curved
c) Both have equal curved paths d) Both have straight line paths
23. A coil of n number of turns is wound tightly in the form of a spiral with inner and outer radii a and b respectively. When a current of strength I is passed through the coil, the magnetic field at its centre is
a) $\frac{\mu_0 n I}{(b-a)} \log_e \frac{a}{b}$ b) $\frac{\mu_0 n I}{2(b-a)}$ c) $\frac{2\mu_0 n I}{b}$ d) $\frac{\mu_0 n I}{2(b-a)} \log_e \frac{b}{a}$
24. A beam of protons is moving parallel to a beam of electrons. Both the beams will tend to
a) Repel each other b) Come closer c) Move more apart d) Either (b) or (c)
25. A fixed horizontal wire carries a current of 200 A . Another wire having a mass per unit length 10^{-2} kg/m is placed below the first wire at a distance of 2 cm and parallel to it. How much current must be passed through the second wire if it floats in air without any support? What should be the direction of current in it
a) 25 A (direction of current is same to first wire)
b) 25 A (direction of current is opposite to first wire)
c) 49 A (direction of current is same to first wire)
d) 49 A (direction of current is opposite to first wire)
26. A long solenoid of length L has a mean diameter D . It has n layers of winding of N turns each. If it carries a current I , the magnetic field at its centre will be
a) Proportional to D b) Inversely proportional to D
c) Independent of D d) Proportional to L
27. Two galvanometer A and B require 3 mA and 5 mA respectively to produce the same deflection of 10 divisions. Then
a) A is more sensitive than B b) B is more sensitive than A
c) A and B are equally sensitive d) Sensitiveness of B is $5/3$ times that of A
28. A circular current carrying coil has a radius R . The distance from the centre of the coil on the axis of the coil, where the magnetic induction is $\frac{1}{8}$ th of its value at the centre of the coil is
a) $\sqrt{3}R$ b) $R/\sqrt{3}$ c) $\left(\frac{2}{\sqrt{3}}\right)R$ d) $\frac{R}{2\sqrt{3}}$
29. The coil of a galvanometer consists of 100 turns and effective area of 1 square cm . The restoring couple is $10^{-8}\text{ N-m/radian}$. The magnetic field between the pole pieces is 5 T . The current sensitivity of this galvanometer will be
a) $5 \times 10^{-4}\text{ rad}/\mu\text{ amp}$ b) $5 \times 10^{-6}\text{ per amp}$ c) $2 \times 10^{-7}\text{ per amp}$ d) $5\text{ rad}/\mu\text{ amp}$
30. A charged particle of mass m and charge q travels in a circular path of radius r that is perpendicular to a magnetic field B . The time taken by the particle to complete one revolution is

- a) $\frac{2\pi B}{m}$ b) $\frac{2\pi m}{qB}$ c) $\frac{2\pi mq}{B}$ d) $\frac{2\pi q^2 B}{m}$

31. Two circular coils are made of two identical wires of same length. If the number of turns of two coils are 4 and 2, then the ratio of magnetic induction at centres will be zero


- a) 4 : 1 b) 2 : 1 c) 1 : 2 d) 1 : 1

32. A long, straight wire is turned into a loop of radius 10 cm (see figure). If a current of 8 amperes is passed through the loop, then the value of the magnetic field and its direction as the centre C of the loop shall be close to



- a) 5.0×10^{-5} Newton/(amp-meter), upward
b) 3.4×10^{-5} Newton/(amp-meter), upward
c) 1.6×10^{-5} Newton/(amp-meter), downward
d) 1.6×10^{-5} Newton/(amp-meter), upward

33. Net magnetic field at the centre of the circle O due to a current through a loop as shown in figure ($\theta < 180^\circ$)



- a) zero b) Perpendicular to paper inwards
c) Perpendicular to paper outwards Perpendicular to paper inwards if $\theta \leq 90^\circ$ and
d) perpendicular to paper outwards if $90^\circ \leq \theta < 180^\circ$

34. Which of the following while in motion cannot be deflected by magnetic field?

- a) Protons b) Cathode rays c) Alpha particles d) Neutrons


35. A charged particle of mass m and charge q describes circular motion of radius r in a uniform magnetic field of strength B . The frequency of revolution is

- a) $\frac{Bq}{2\pi m}$ b) $\frac{Bq}{2\pi r m}$ c) $\frac{2\pi m}{Bq}$ d) $\frac{Bm}{2\pi q}$

36. A current of 2 amp, flows in a long, straight wire of radius 2 mm. The intensity of magnetic field on the axis of the wire is

- a) $\left(\frac{\mu_0}{\pi}\right) \times 10^3 \text{ tesla}$ b) $\left(\frac{\mu_0}{2\pi}\right) \times 10^3 \text{ tesla}$ c) $\left(\frac{2\mu_0}{\pi}\right) \times 10^3 \text{ tesla}$ d) Zero

37. A metallic block carrying current I is subjected to a uniform magnetic induction \vec{B} as shown in the figure. The moving charges experience a force F given by which results in the lowering of the potential of the face Assume the speed of the carriers to be v



- a) $eVB\hat{k}$, ABCD b) $eVB\hat{k}$, EFGH c) $-eVB\hat{k}$, ABCD d) $-eVB\hat{k}$, EFGH

38. The magnetic field existing in a region is given by $\vec{B} = B_0 \left[1 + \frac{x}{l}\right] \hat{k}$. A square loop of edge l and carrying

current i is placed with its edges parallel to $x - y$ axis. The magnitude of the net magnetic force experienced by the loop is

- a) $2B_0il$ b) B_0i_0l c) B_0il d) Bil

39. A straight conductor carrying current I . If the magnetic field at a distance r is 0.4 T, then magnetic field at a distance $2r$ will be

- a) 0.4 T b) 0.1 T c) 0.8 T d) 0.2 T

40. The proton is energy 1 MeV describes a circular path in plane at right angles to a uniform magnetic field of 6.28×10^{-4} T. The mass of the proton is 1.7×10^{-27} kg. The cyclotron frequency of the proton is very nearly equal to

- a) 10^7 Hz b) 10^5 Hz c) 10^6 Hz d) 10^4 Hz


41. Current i is carried in a wire of length L . If the wire is turned into a circular coil, the maximum magnitude of torque in a given magnetic field B will be

- a) $\frac{LiB^2}{2}$ b) $\frac{Li^2B}{2}$ c) $\frac{L^2iB}{4\pi}$ d) $\frac{Li^2B}{4\pi}$


42. A long solenoid has 800 turns per metre length of solenoid. A current of 1.6 A flows through it. The magnetic induction at the end of the solenoid on its axis is

- a) 16×10^{-4} T b) 8×10^{-4} T c) 32×10^{-4} T d) 4×10^{-4} T

43. Which of the field pattern given in the figure is valid for electric field as well as for magnetic field?



44. AB and CD are long straight conductors, distance d apart, carrying a current I . The magnetic field at the midpoint of BC is



- a) $\frac{-\mu_0 I}{2\pi d} \hat{k}$ b) $\frac{-\mu_0 I}{\pi d} \hat{k}$ c) $\frac{-\mu_0 I}{4\pi d} \hat{k}$ d) $\frac{-\mu_0 I}{8\pi d} \hat{k}$

45. The magnetic force acting on a charge particle of charge $-2\mu c$ in a magnetic field of $2T$ act in y direction, when the particle velocity is $(2i + 3j) \times 10^6 \text{ ms}^{-1}$ is

- a) 8 N in $-z$ direction b) 8 N in z direction c) 8 N in y direction d) 8 N in x direction

46. At the centre of a circular coil of radius 5 cm carrying current, magnetic field due to earth is $0.5 \times 10^{-5} \text{ Wbm}^{-2}$. What should be the current flowing through the coil so that it annuals the earth's magnetic field?

- a) 40 A b) 4 A c) 0.4 A d) 0.2 A


47. An electron enters into a region of uniform magnetic field of strength 10 webers/ m^2 with a speed of $3 \times 10^7 \text{ m/s}$ which of the following is not possible

- a) The electron may or may not experience an acceleration
b) The electron may experience an acceleration but can continue to move with same speed
c) The electron may experience an acceleration and continue to move with same velocity
d) The kinetic energy of the electron remains unchanged


48. Magnetic field at the centre of a circular loop of area A is B . The magnetic moment of the loop will be

- a) $\frac{BA^2}{\mu_0\pi}$ b) $\frac{BA^{3/2}}{\mu_0\pi}$ c) $\frac{BA^{3/2}}{\mu_0\pi^{1/2}}$ d) $\frac{2BA^{3/2}}{\mu_0\pi^{1/2}}$

49. A wire oriented in the east-west direction carries a current eastward. Direction of the magnetic field at a point to the south of the wire is
 a) Vertically down b) Vertically up c) North-east d) South-east
50. Magnetic field induction at the centre O of a square loop of side ' a ' carrying current I as shown in figure is



- a) $\frac{\mu_0 I}{\sqrt{2}\pi a}$ b) $2\sqrt{2}\frac{\mu_0 I}{\pi a}$ c) $\frac{2\mu_0 I}{\pi a}$ d)
51. An electron is projected along the axis of a circular conductor carrying some current. Electron will experience force
 a) Along the axis b) Perpendicular to the axis
 c) At an angle of 4° with axis d) No force experienced
52. If the current is doubled, the deflection is also doubled in
 a) A tangent galvanometer b) A moving coil galvanometer
 c) Both (a) and (b) d) None of these
53. In the adjacent figure is shown a closed path P . A long straight conductor carrying a current I passes through O and perpendicular to the plane of the paper. Then which of the following holds good?



- a) $\int_P \mathbf{B} \cdot d\mathbf{l} = 0$ b) $\int_P \mathbf{B} \cdot d\mathbf{l} = \mu_0 I$ c) $\int_P \mathbf{B} \cdot d\mathbf{l} > \mu_0 I$ d) None of these
54. A charge moves in a circle perpendicular to a magnetic field. The time period of revolution is independent of
 a) Magnetic field b) Charge
 c) Mass of the particle d) Velocity of the particle
55. A stream of electrons is projected horizontally to the right. A straight conductor carrying a current is supported parallel to electron stream and above it. If the current in the conductor is from left to right, then what will be the effect on electron stream?
 a) The electron stream will be speeded up towards the right b) The electron stream will be retarded
 c) The electron stream will be pulled upward d) The electron stream will be pulled downward
56. A proton of mass 1.67×10^{-27} kg and charge 1.6×10^{-19} C is projected with a speed of 2×10^6 ms $^{-1}$ at an angle of 60° to the X -axis. If a uniform magnetic field of 0.104 T is applied along Y -axis, the path of proton is
 a) A circle of radius = 0.2 m and time period = $2\pi \times 10^{-7}$ s b) A circle of radius = 0.1 m and time period = $2\pi \times 10^{-7}$ s
 c) A helix of radius 0.1 m and time period = $2\pi \times 10^{-7}$ s d) A helix of radius 0.2 m and time period = $2\pi \times 10^{-7}$ s
57. Rate of change of torque τ with deflection θ is maximum for a magnet suspended freely in a uniform magnetic field of induction B , when
 a) $\theta = 0^\circ$ b) $\theta = 45^\circ$ c) $\theta = 60^\circ$ d) $\theta = 90^\circ$
58. The magnetic field at the centre of a circular current carrying conductor of radius r is B_c . The magnetic

field on its axis at a distance r from the centre is B_a . The value of $B_c : B_a$ will be

- a) $1 : \sqrt{2}$ b) $1 : 2\sqrt{2}$ c) $2\sqrt{2} : 1$ d) $\sqrt{2} : 1$


59. A coil in the shape of an equilateral triangle of side l is suspended between the pole pieces of a permanent magnet such that \vec{B} is in plane of the coil. If due to a current i in the triangle a torque τ acts on it, the side l of the triangle is

- a) $\frac{2}{\sqrt{3}} \left(\frac{\tau}{Bi} \right)^{\frac{1}{2}}$ b) $\frac{2}{3} \left(\frac{\tau}{Bi} \right)$ c) $2 \left(\frac{\tau}{\sqrt{3}Bi} \right)^{\frac{1}{2}}$ d) $\frac{1}{\sqrt{3}} \frac{\tau}{Bi}$

60. A thin circular disk of radius R is uniformly charged with density $\sigma > 0$ per unit area. The disk rotates about its axis with a uniform angular speed ω . The magnetic moment of the disk is


- a) $2\pi R^4 \sigma \omega$ b) $\pi R^4 \sigma \omega$ c) $\frac{\pi R^4}{2} \sigma \omega$ d) $\frac{\pi R^4}{4} \sigma \omega$

61. A conducting loop carrying a current I is placed in a uniform magnetic field pointing into the plane of the paper as shown. The loop will have a tendency to



- a) Contract
b) Expand
c) Move towards +ve x - axis
d) Move towards -ve x - axis

62. A and B are two conductors carrying a current i in the same direction. x and y are two electron beams moving in the same direction




- a) There will be repulsion between A and B attraction between x and y
b) There will be attraction between A and B repulsion between x and y
c) There will be repulsion between A and B and also x and y
d) There will be attraction between A and B and also x and y
63. An electron ($q = 1.6 \times 10^{-19} \text{ C}$) is moving at right angle to the uniform magnetic field $3.534 \times 10^{-5} \text{ T}$. The time taken by the electron to complete a circular orbit is
- a) $2\mu\text{s}$ b) $4\mu\text{s}$ c) $3\mu\text{s}$ d) $1\mu\text{s}$
64. A particle having a mass of 10^{-2} kg carries a charge of $5 \times 10^{-8} \text{ C}$. The particle is given an initial horizontal velocity of 10^5 ms^{-1} in the presence of electric field \vec{E} and magnetic field \vec{B} . To keep the particle moving in a horizontal direction, it is necessary that
- (1) \vec{B} should be perpendicular to the direction of velocity and \vec{E} should be along the direction of velocity
(2) Both \vec{B} and \vec{E} should be along the direction of velocity
(3) Both \vec{B} and \vec{E} are mutually perpendicular and perpendicular to the direction of velocity
(4) \vec{B} should be along the direction of velocity and \vec{E} should be perpendicular to the direction of velocity
- Which of the following pairs of statements is possible
- a) (1) and (3) b) (3) and (4) c) (2) and (3) d) (2) and (4)
65. Which of the following gives the value of magnetic field according to Biot-Savart' law

- a) $\frac{i\Delta l \sin \theta}{r^2}$ b) $\frac{\mu_0}{4\pi} \frac{i\Delta l \sin \theta}{r}$ c) $\frac{\mu_0}{4\pi} \frac{i\Delta l \sin \theta}{r^2}$ d) $\frac{\mu_0}{4\pi} i\Delta l \sin \theta$

66. Two ions having masses in the ratio 1 : 1 and charges 1 : 2 are projected into uniform magnetic field perpendicular to the field with speeds in the ratio 2 : 3. The ratio of the ratio of circular paths along which the two particles move is

- a) 4 : 3 b) 2 : 3 c) 3 : 1 d) 1 : 4

67. Equal current i flows in two segments of a circular loop in the direction shown in figure. Radius of the loop is r . The magnitude of magnitude field induction at the centre of the loop is




- a) zero b) $\frac{\mu_0 i \theta}{4\pi r}$ c) $\frac{\mu_0 i}{2\pi r} (\pi - \theta)$ d) $\frac{\mu_0 i}{2\pi r} (2\pi - \theta)$

68. An electron beam travels with a velocity of $1.6 \times 10^7 \text{ ms}^{-1}$ perpendicularly to magnetic field of intensity 0.1 T. The radius of the path of the electron beam ($m_e = 9 \times 10^{-31} \text{ kg}$)

- a) $9 \times 10^{-5} \text{ m}$ b) $9 \times 10^{-2} \text{ m}$ c) $9 \times 10^{-4} \text{ m}$ d) $9 \times 10^{-3} \text{ m}$

69. A wire carrying current i is shaped as shown. Section AB is a quarter circle of radius r . The magnetic field is directed




- a) At an angle $\pi/4$ to the plane of the paper
b) Perpendicular to the plane of the paper and directed in to the paper
c) Along the bisector of the angle ACB towards AB
d) Along the bisector of the angle ACB away from AB

70. If same current I passing through two parallel wires separated by a distance b , then force per unit length will be


- a) $\frac{\mu_0 2I^2}{4\pi b}$ b) $\frac{\mu_0 I}{4\pi b^2}$ c) $\frac{\mu_0 I^2}{4\pi b^2}$ d) $\frac{\mu_0 I^2}{4\pi b}$

71. PQ and RS are long parallel conductors separated by certain distance. M is the mid-point between them (see the figure). The net magnetic field at M is B. Now, the current 2 A is switched off. The field at M now becomes




- a) $2B$ b) B c) $\frac{B}{2}$ d) $3B$


72. The magnetic field induction at a point 4 cm from a long current carrying wire is 10^{-3} T. The magnetic field induction at a distance of 1.0 cm from the same current wire will be
a) 2×10^{-4} T b) 3×10^{-4} T c) 4×10^{-3} T d) 1.11×10^{-4} T
73. A circular loop has a radius of 5 cm and it is carrying a current of 0.1 amp. Its magnetic moment is
a) $1.32 \times 10^{-4} \text{ amp} \cdot \text{m}^2$ b) $2.62 \times 10^{-4} \text{ amp} \cdot \text{m}^2$ c) $5.25 \times 10^{-4} \text{ amp} \cdot \text{m}^2$ d) $7.85 \times 10^{-4} \text{ amp} \cdot \text{m}^2$
74. A coil of 100 turns and area $2 \times 10^{-2} \text{ m}^2$, pivoted about a vertical diameter in a uniform magnetic field carries a current of 5A. When the coil is held with its plane in North-South direction, it experience a torque of 0.3 Nm. When the plane is in East-West direction the torque is 0.4 Nm. The value of magnetic induction is (Neglect earth's magnetic field)
a) 0.2 T b) 0.3 T c) 0.4 T d) 0.05 T
75. Two charged particles M and N enter a space of uniform magnetic field, with velocities perpendicular to the magnetic field. The paths are as shown in figure. The possible reason (s) is/are?



- a) The charge of M is greater than that of N b) The momentum of M is greater than that of N
c) Specific charge of M is greater than that of N d) The speed of M is greater than that of N
76. A particle of charge q and mass m moving with a velocity v along the x -axis enters the region $x > 0$ with uniform magnetic field B along the \hat{k} direction. The particle will penetrate in this region in the x -direction upto a distance d equal to
a) Zero b) $\frac{mv}{qB}$ c) $\frac{2mv}{qB}$ d) Infinity
77. A and B are two concentric circular conductors of centre O and carrying currents i_1 and i_2 as shown in the adjacent figure. If ratio of their radii is 1 : 2 and ratio of the flux densities at O due to A and B is 1 : 3, then the value of i_1/i_2 is



- a) 1/6 b) 1/4 c) 1/3 d) 1/2
78. A battery is connected between two points A and B on the circumference of a uniform conducting ring of radius r and resistance R . One of the arcs AB of the ring subtends an angle θ at the centre. Magnetic field due to current at the centre of ring is
a) Zero, only if $\theta = 180^\circ$ b) Zero for all values of θ c) Proportional to $\frac{2(180^\circ - \theta)}{2}$ d) Inversely proportional to r
79. An electron with mass m , velocity v and charge e describes half a revolution in a circle of radius r in a magnetic field B , will acquire energy equal to
a) $1/2 mv^2$ b) $1/4 mv^2$ c) $\pi r B e v$ d) zero
80. A horizontal metal wire is carrying an electric current from the north to the south. Using a uniform magnetic field, it is to be prevented from falling under gravity. The direction of this magnetic field should be towards the
a) North b) South c) East d) West
81. What is the net force on the square coil



- a) $25 \times 10^{-7} N$ moving towards wire
 b) $25 \times 10^{-7} N$ moving away from wire
 c) $35 \times 10^{-7} N$ moving towards wire
 d) $35 \times 10^{-7} N$ moving away from wire
82. The pole pieces of the magnet used in a pivoted coil galvanometer are
 a) Plane surface of a bar magnet
 b) Plane surface of a horse-shoe magnet
 c) Cylindrical surfaces of a bar magnet
 d) Cylindrical surfaces of a horse-shoe magnet
83. A cable carrying a direct current is buried in a wall which stands in a north-south plane. A horizontal compass needle on the west side of the wall is found to point towards south instead of north. The coil is laid
 a) Vertically upwards and the current is also flowing upwards
 b) Vertically upwards and the current is flowing downwards
 c) Horizontal with current from south to north
 d) Horizontal with current from north to south
84. The magnetic moment of a current (i) carrying circular coil of radius (r) and number of turns (n) varies as
 a) $1/r^2$
 b) $1/r$
 c) r
 d) r^2
85. An infinitely long wire carrying current i is along Y-axis such that its one end is at point $(0, b)$ while the wire extends upto ∞ . The magnitude of magnetic field strength at point $P(a, 0)$ is
-
- a) $\frac{\mu_0 i}{4\pi a} \left(1 + \frac{b}{\sqrt{a^2 + b^2}}\right)$
 b) $\frac{\mu_0 i}{4\pi a} \left(1 - \frac{b}{\sqrt{a^2 + b^2}}\right)$
 c) $\frac{\mu_0 i}{4\pi a} \left(1 - \frac{a}{\sqrt{a^2 + b^2}}\right)$
 d) $\frac{\mu_0 i}{4\pi a} \left(\frac{b}{\sqrt{a^2 + b^2}}\right)$
86. If the strength of the magnetic field produced 10cm away from a infinitely long straight conductor is $10^{-5} \text{ weber/m}^2$, the value of the current flowing in the conductor will be
 a) 5 ampere
 b) 10 ampere
 c) 500 ampere
 d) 1000 ampere
87. Two concentric circuit coils of ten turn each are situated in the same plane. Their radii are 20 cm and 40 cm and they carry respectively 0.2 A and 0.3 A current in opposite direction. The magnetic field in tesla at the centre is
 a) $35 \mu_0/4 \text{ T}$
 b) $\mu_0/80 \text{ T}$
 c) $7 \mu_0/80 \text{ T}$
 d) $5 \mu_0/4 \text{ T}$
88. A direct current I flows along the length of an infinitely long straight thin walled pipe, then the magnetic field is
 a) Uniform throughout the pipe but not zero
 b) Zero only along the axis of the pipe
 c) Zero at any point inside the pipe
 d) Maximum at the centre and minimum at the edge
89. In ballistic galvanometer, the frame on which the coil is wound is non-metallic to
 a) Avoid the production of induced e.m.f.
 b) Avoid the production of eddy currents
 c) Increase the production of eddy currents
 d) Increase the production of induced e.m.f.
90. An electron and a proton enter a magnetic field perpendicularly. Both have same kinetic energy. Which of the following is true?
 a) Trajectory of electron is less curved
 b) Trajectory of proton is less curved
 c) Both trajectories are equally curved
 d) Both move on straight line path

91. An electron and a proton of equal linear momentum enter in the direction perpendicular to uniform magnetic field. If the radii of their circular paths be r_e and r_p respectively, then $\frac{r_e}{r_p}$ is equal to $m_e =$ mass of electron, $m_p =$ mass of proton.

a) $\left(\frac{m_p}{m_e}\right)^{1/2}$ b) $\frac{m_p}{m_e}$ c) $\left(\frac{m_e}{m_p}\right)^{1/2}$ d) 1


92. An electron of mass m and charge q is travelling with a speed v along a circular path of radius r at right angles to a uniform magnetic field B . If speed of the electron is doubled and the magnetic field is halved, then resulting path would have a radius of

a) $\frac{r}{4}$ b) $\frac{r}{2}$ c) $2r$ d) $4r$

93. A charged particle enters in a magnetic field whose direction is parallel to velocity of the particle, then the speed of this particle

a) In straight line b) In coiled path c) In circular path d) In ellipse path

94. Two circular concentric loops of radii $r_1 = 20$ cm and $r_2 = 30$ cm are placed in the $X - Y$ plane as shown in the figure. A current $I = 7$ A is flowing through them. The magnetic moment of this loop system is



a) $+0.4 \hat{k} (\text{Am}^2)$ b) $-1.5 \hat{k} (\text{Am}^2)$ c) $+1.1 \hat{k} (\text{Am}^2)$ d) $+1.3 \hat{j} (\text{Am}^2)$

95. A charged particle is moving in a magnetic field of strength B perpendicular to the direction of the field. q and m denote the charge and mass of the particle respectively, then the frequency of rotation of the particle is

a) $f = \frac{qB}{2\pi m}$ b) $f = \frac{qB}{2\pi m^2}$ c) $f = \frac{2\pi^2 m}{qB}$ d) $f = \frac{2\pi m}{qB}$

96. If two parallel wires carry current in opposite directions

a) The wires attract each other b) The wires repel each other
c) The wires experience neither attraction nor repulsion d) The forces of attraction or repulsion do not depend on current direction

97. An electron is shot in steady electric and magnetic fields such that its velocity v , electric field E and magnetic field B are mutually perpendicular. The magnitude of B are mutually perpendicular. The magnitude of E is 1 Vcm^{-1} and that of B is 2 T . Now if it so happens that the Lorentz (magnetic) force cancels the electrostatic force on the electron, then the velocity of the electron is

a) 50 ms^{-1} b) 2 cms^{-1} c) 0.5 cms^{-1} d) 200 cms^{-1}

98. Two parallel conductors A and B of equal lengths carry currents I and $10 I$, respectively, in the same direction. Then

a) A and B will repel each other with same force
b) A and B will attract each other with same force
c) A will attract B , but B will repel A
d) A and B will attract each other with different forces

99. An electron ($q = 1.6 \times 10^{-19} \text{ C}$) is moving at right angle to the uniform magnetic field $3.534 \times 10^{-5} \text{ T}$. The time taken by the electron to complete a circular orbit is

a) $2\mu\text{s}$ b) $4\mu\text{s}$ c) $3\mu\text{s}$ d) $1\mu\text{s}$

100. Two long wires are hanging freely. They are joined first in parallel and then in series and then are connected with a battery. In both cases, which type of force acts between the two wires


a) Attraction force when in parallel and repulsion force when in series
b) Repulsion force when in parallel and attraction force when in series

- c) Repulsion force in both cases
d) Attraction force in both cases

101. An electron having charge 1.6×10^{-19} C and mass 9×10^{-31} kg is moving with speed 4×10^6 ms⁻¹ in a magnetic field of 2×10^{-1} T in a circular orbit. The force acting on electron and the radius of the circular orbit will be


- a) 18.8×10^{-13} N, 1.1×10^{-4} m
b) 12.8×10^{-14} N, 1.1×10^{-3} m
c) 12.8×10^{-13} N, 1.1×10^{-3} m
d) 1.28×10^{-13} N, 1.1×10^{-4} m

102. The unit vectors \hat{i} , \hat{j} and \hat{k} are as shown below. What will be the magnetic field at O in the following figure




- a) $\frac{\mu_0 i}{4\pi a} \left(2 - \frac{\pi}{2}\right) \hat{j}$
b) $\frac{\mu_0 i}{4\pi a} \left(2 + \frac{\pi}{2}\right) \hat{j}$
c) $\frac{\mu_0 i}{4\pi a} \left(2 + \frac{\pi}{2}\right) \hat{i}$
d) $\frac{\mu_0 i}{4\pi a} \left(2 + \frac{\pi}{2}\right) \hat{k}$

103. The figure shows certain wire segments joined together to form a coplanar loop. The loop is placed in a perpendicular magnetic field in the direction going into the plane of the figure. The magnitude of the field increases with time. I_1 and I_2 are the currents in segment **ab** and **cd**. Then,



- a) $I_1 > I_2$
b) $I_1 < I_2$
c) I_1 is in the direction **ba** and I_2 is in the direction **cd**
d) I_1 is in the direction **ab** and I_2 is in the direction **dc**

104. Three long, straight parallel wires, carrying current, are arranged as shown in figure. The force experienced by a 25 cm length of wire C is



- a) 10^{-3} N
b) 2.5×10^{-3} N
c) Zero
d) 1.5×10^{-3}

105.


Work done on an electron moving in a solenoid along its axis is equal to

- a) Zero
b) $-evB$
c) i/B
d) None of the above


106. The magnetic field at the point of intersection of diagonals of a square loop of side L carrying a current I is

- a) $\frac{\mu_0 I}{\pi L}$
b) $\frac{2\mu_0 I}{\pi L}$
c) $\frac{\sqrt{2}\mu_0 I}{\pi L}$
d) $\frac{2\sqrt{2}\mu_0 I}{\pi L}$


107. A long wire AB is placed on a table. Another wire PQ of mass 1.0 g and length 50 cm is set to slide on two rails PS and QR . A current of 50 A is passed through the wires. At what distance above AB , will the wire PQ be in equilibrium




- a) 25 mm b) 50 mm c) 75 mm d) 100 mm
108. A straight section PQ of a circuit lies along the X -axis from $x = \frac{-a}{2}$ to $x = \frac{a}{2}$ and carries a steady current i . The magnetic field due to the section PQ at a distance $x = +a$ will be
 a) Proportional to a b) Proportional to $1/a$ c) Proportional to a^2 d) Zero
109. A current i is passing through a straight conductor of infinite length. The magnetic field at a point situated at a distance R from the conductor is
 a) $\frac{\mu_0}{2\pi} i$ b) $\frac{\mu_0}{2\pi} \frac{i}{R^2}$ c) $\frac{\mu_0}{2\pi} \frac{i}{R^3}$ d) $\frac{\mu_0}{2\pi} \frac{i}{R}$
110. A current of 1 ampere is passed through a straight wire of length 2.0 metres. The magnetic field at a point in air at a distance of 3 metres from either end of wire and lying on the axis of wire will be
 a) $\frac{\mu_0}{2\pi}$ b) $\frac{\mu_0}{4\pi}$ c) $\frac{\mu_0}{8\pi}$ d) Zero
111. A current I enters a circular coil of radius R , branches into two parts and then recombines as shown in the circuit diagram




- The resultant magnetic field at the centre of the coil is
 a) Zero b) $\frac{\mu_0 I}{2R}$ c) $\frac{3}{4} \left(\frac{\mu_0 I}{2R} \right)$ d) $\frac{1}{4} \left(\frac{\mu_0 I}{2R} \right)$
112. The figure shows three situations when an electron with velocity \mathbf{v} travels through a uniform magnetic field \mathbf{B} . In each case, what is the direction of magnetic force on the electron?



- a) +ve z - axis, -ve x - axis, +ve y - axis
 b) -ve z - axis, -ve x - axis and zero
 c) +ve z - axis, +ve y - axis and zero
 d) -ve z - axis, +ve y - axis and zero
113. An electron having mass $(9.1 \times 10^{-31} \text{ kg})$ and charge $(1.6 \times 10^{-19} \text{ C})$ moves in a circular path of radius 0.5 m with a velocity 10^6 ms^{-1} in a magnetic field. Strength of magnetic field is
 a) $1.13 \times 10^{-5} \text{ T}$ b) $5.6 \times 10^{-6} \text{ T}$ c) $2.8 \times 10^{-6} \text{ T}$ d) None of these
114. Due to the flow of current in a circular loop of radius R , the magnetic induction produced at the centre of the loop is B . The magnetic moment of the loop is
 (μ_0 = permeability constant)

- a) $BR^3/2\pi\mu_0$ b) $2\pi BR^3/\mu_0$ c) $BR^2/2\pi\mu_0$ d) $2\pi BR^2/\mu_0$
115. An alternating electric field, of frequency ν , is applied across the dees (radius = R) of a cyclotron that is being used to accelerated protons (mass = m). The operating magnetic field (B) used in the cyclotron and the kinetic energy (K) of the proton beam, produced by it, are given by
- a) $B = \frac{mv}{e}$ and $K = 2m\pi^2\nu^2R^2$ b) $B = \frac{2\pi mv}{e}$ and $K = m^2\pi\nu R^2$
 c) $B = \frac{2\pi mv}{e}$ and $K = 2m\pi^2\nu^2R^2$ d) $B = \frac{mv}{e}$ and $K = m^2\pi\nu R^2$
116. A steady electric current is flowing through a cylindrical conductor
- a) The magnetic field in the vicinity of the conductor is zero
 b) The electric field in the vicinity of the conductor is non-zero
 c) The magnetic field at the axis of the conductor is zero
 d) The electric field at the axis of the conductor is zero
117. Two long parallel wires carry currents i_1 and i_2 such that $i_1 > i_2$. When the currents are in the same direction, the magnetic field at a point midway between the wires is 6×10^{-6} T. If the direction of i_2 is reversed, the field becomes 3×10^{-5} T. The ratio $\frac{i_1}{i_2}$ is
- a) $\frac{1}{2}$ b) 2 c) $\frac{2}{3}$ d) $\frac{3}{2}$
118. A wire of length l metre carrying a current i ampere is bent in the form of a coil having two turns. Its magnitude of magnetic moment will be
- a) $il/4\pi$ b) $i^2l^2/4\pi$ c) $i^2l/8\pi$ d) $il^2/8\pi$
119. Find magnetic field at O
- 
- a) $\frac{5\mu_0 i \theta}{24\pi r}$ b) $\frac{\mu_0 i \theta}{24\pi r}$ c) $\frac{11\mu_0 i \theta}{24\pi r}$ d) Zero
120. A long solenoid has 200 turns/cm and carries a current i . The magnetic field at its centre is 6.28×10^{-2} Wb/m². Another long solenoid has 100 turns/cm and it carries a current $i/3$. The value of the magnetic field at its centre is
- a) 1.05×10^{-2} Wbm⁻² b) 1.05×10^{-5} Wbm⁻² c) 1.05×10^{-3} Wbm⁻² d) 1.05×10^{-4} Wbm⁻²
121. A charge + Q is moving upwards vertically. It enters a magnetic field directed to north. The force on the charge will be towards
- a) North b) South c) East d) West
122. A length of wire carries a steady current. It is bent first to form a circular coil of one turn. The same length is now bent more sharply to give a double loop of smaller radius. The magnetic field at the centre caused by the same current is
- a) Double of its first value b) Quarter of its first value
 c) Four times of its first value d) Same as the first value
123. A particle of mass m and charge q moves with a constant velocity v along the positive x direction. It enters a region containing a uniform magnetic field B directed along the negative z direction, extending from $x = a$ to $x = b$. The minimum value of v required so that the particle can just enter the region $x > b$ is
- a) $qb B/m$ b) $q(b - a)B/m$ c) $qa B/m$ d) $q(b + a)B/2m$
124. A, B and C are parallel conductors of equal length carrying currents I, I and $2I$ respectively. Distance between A and B is x . Distance between B and C is also x . F_1 is the force exerted by B on A and F_2 is the force exerted by C on A . Choose the correct answer



- a) $F_1 = 2F_2$ b) $F_2 = 2F_1$ c) $F_1 = F_2$ d) $F_1 = -F_2$

125. When a positively charged particle enters a uniform magnetic field with uniform velocity, its trajectory can be

- (1) a straight line
(2) a circle
(3) a helix

- a) (1) only b) (1) or (2)
c) (1) or (3) d) Any one of (1), (2) and (3)


126. A current I flow in an infinitely long wire cross-section in the form of a semi-circular ring of radius R . The magnitude of the magnetic induction along its axis is

- a) $\frac{\mu_0 I}{2\pi^2 R}$ b) $\frac{\mu_0 I}{2\pi R}$ c) $\frac{\mu_0 I}{4\pi R}$ d) $\frac{\mu_0 I}{\pi^2 R}$

127. An electron is moving in an orbit of radius R with a time

period T as shown in the figure. The magnetic moment produced may be given by

$|e|$ represents the magnitude of the electron charge.



- a) $\mathbf{M} = \frac{2\pi|e|\mathbf{A}}{T}$ b) $\mathbf{M} = -\frac{2\pi|e|\mathbf{A}}{T}$ c) $\mathbf{M} = \frac{|e|\mathbf{A}}{T}$ d) $\mathbf{M} = -\frac{|e|\mathbf{A}}{T}$

128. 1A current flows through an infinitely long straight wire. The magnetic field produced at a point 1 metre away from it is

- a) 2×10^{-3} tesla b) $\frac{2}{10}$ tesla c) 2×10^{-7} tesla d) $2\pi \times 10^{-6}$ tesla

129. A magnetic needle lying parallel to a magnetic field required W units of work to turn it through 60° . The torque required to maintain the needle in this position will be

- a) $\sqrt{3} W$ b) W c) $\sqrt{3} \frac{W}{2}$ d) $\frac{2 W}{2}$

130. Which of the following particles will describe the smallest circle when projected with the same velocity perpendicular to the magnetic field?

- a) Electron b) Proton c) α – particle d) Deuteron


131. On connecting a battery to the two corners of a diagonal of a square conductor frame of side a , the magnitude of the magnetic field at the centre will be

- a) Zero b) $\frac{\mu_0}{\pi a}$ c) $\frac{2\mu_0}{\pi a}$ d) $\frac{4\mu_0 i}{\pi a}$


132. A solenoid consists of 100 turns of wire and has a length of 10.0 cm. The magnetic field inside the solenoid when it carries a current of 0.500 A will be

- a) 6.28×10^{-4} T b) 6.28×10^{-5} T c) 3.14×10^{-4} T d) None of these

133. A rectangular loop carrying a current i is situated near a long straight wire such that the wire is parallel to the one of the sides of the loop and is in the plane of the loop. If a steady current I is established in wire as shown in figure, the loop will




133. The rectangular loop PQRS is placed to the right of the wire. The magnetic field on the axis of a long solenoid having n turns per unit length and carrying a current i is
- a) Rotate about an axis parallel to the wire b) Move away from the wire to towards right
c) Move towards the wire d) Remain stationary
134. The magnetic field on the axis of a long solenoid having n turns per unit length and carrying a current i is
a) $\mu_0 n i$ b) $\mu_0 n^2 i$ c) $\mu_0 n i^2$ d) None of these
135. In case of Hall effect for a strip having charge Q and area of cross-section A , the Lorentz force is
a) Directly proportional to Q b) Inversely proportional to Q
c) Inversely proportional to A d) Directly proportional to A
136. Circular loop of a wire and long straight wire carry currents I_c and I_e , respectively as shown in figure. Assuming that these are placed in the same plane. The magnetic fields will be zero at the centre of the loop when the separation H is



- a) $\frac{I_e R}{I_c \pi}$ b) $\frac{I_c R}{I_e \pi}$ c) $\frac{\pi I_c}{I_e R}$ d) $\frac{I_e \pi}{I_c R}$
137. An electron enters the space between the plates of a charged capacitor as shown. The charge density on the plate is σ . Electric intensity in the space between the plates is E . A uniform magnetic field B also exists in that space perpendicular to the direction of E . The electron moves perpendicular to both \vec{E} and \vec{B} without any change in direction. The time taken by the electron to travel a distance l in that space is
- a) $\frac{\sigma l}{\epsilon_0 B}$ b) $\frac{\sigma B}{\epsilon_0 l}$ c) $\frac{\epsilon_0 l B}{\sigma}$ d) $\frac{\epsilon_0 l}{\sigma B}$
138. An electron is moving in the north direction. It experiences a force in vertically upward direction. The magnetic field at the position of the electron is in the direction of
a) East b) West c) North d) South
139. In hydrogen atom, the electron is making $6.6 \times 10^{15} \text{ rev/sec}$ around the nucleus in an orbit of radius 0.528 \AA . The magnetic moment ($A - m^2$) will be
a) 1×10^{-15} b) 1×10^{-10} c) 1×10^{-23} d) 1×10^{-27}
140. The magnetic moment of a circular coil carrying current is
a) Directly proportional to the length of the wire in the coil
b) Inversely proportional to the length of the wire in the coil
c) Directly proportional to the square of the length of the wire in the coil
d) Inversely proportional to the square of the length of the wire in the coil
141. A doubly ionized helium ion and a H_2 ion are accelerated through the same potential. The ratio of the speed of helium and H_2 ion is
a) 2 : 1 b) 1 : 2 c) 1 : $\sqrt{2}$ d) $\sqrt{2}$: 1


142. What is the shape of magnet used in moving coil galvanometer to make the magnetic fields radial
 a) Concave b) Horse shoe magnet c) Convex d) None of these
143. A circular coil carrying a current has a radius R . The ratio of magnetic induction at the centre of the coil and at a distance equal to $\sqrt{3} R$ from the centre of the coil on the axis is
 a) 1 : 1 b) 1 : 2 c) 2 : 1 d) 8 : 1
144. A galvanometer of resistance 100Ω gives a full scale deflection for a current of 10^{-5} A. To convert it into a ammeter capable of measuring upto 1A, we should connect a resistance of
 a) 1Ω in parallel b) $10^{-3} \Omega$ in parallel c) $10^5 \Omega$ in series d) 100Ω in series
145. Two wires of same length are shaped into a square and a circle. If they carry same current, ratio of the magnetic moment is
 a) $2 : \pi$ b) $\pi : 2$ c) $\pi : 4$ d) $4 : \pi$
146. If an electron is going in the direction of magnetic field \vec{B} with the velocity of \vec{v} then the force on electron is
 a) Zero b) $e(\vec{v} \cdot \vec{B})$ c) $e(\vec{v} \times \vec{B})$ d) None of these
147. A charge of $2.0 \mu C$ moves with a speed of $3.0 \times 10^6 \text{ ms}^{-1}$ along +ve X -axis. A magnetic field of strength $\vec{B} = -0.2 \hat{k} \text{ tesla}$ exists in space. What is the magnetic force (\vec{F}_m) on the charge
 a) $F_m = 1.2 \text{ N}$ along +ve x – direction b) $F_m = 1.2 \text{ N}$ along –ve x – direction
 c) $F_m = 1.2 \text{ N}$ along +ve y – direction d) $F_m = 1.2 \text{ N}$ along –ve y – direction
148. An arc of a circle of radius R subtends an angle $\pi/2$ at the centre. It carries a current i . The magnetic field at the centre will be
 a) $\frac{\mu_0 i}{2R}$ b) $\frac{\mu_0 i}{8R}$ c) $\frac{\mu_0 i}{4R}$ d) $\frac{2\mu_0 i}{5R}$
149. Lorentz force can be calculated by using the formula
 a) $\vec{F} = q(\vec{E} + \vec{v} \times \vec{B})$ b) $\vec{F} = q(\vec{E} - \vec{v} \times \vec{B})$ c) $\vec{F} = q(\vec{E} + \vec{v} \cdot \vec{B})$ d) $\vec{F} = q(\vec{E} \times \vec{B} + \vec{v})$
150. A current flows in a conductor from east to west. The direction of the magnetic field at a point above the conductor is
 a) Towards east b) Towards west c) Towards north d) Towards south
151. Consider two straight parallel conductors A and B separated by a distance x and carrying individual currents i_A and i_B respectively. If the two conductors attract each other, it indicates that
 a) The two currents are parallel in direction
 b) The two currents are anti-parallel in direction
 c) The magnetic lines induction are parallel
 d) The magnetic lines of induction are parallel to length of conductors
152. The earth's magnetic induction at certain point is $7 \times 10^{-5} \text{ Wb/m}^2$. This is to be annulled by the magnetic induction at the centre of a circular conducting loop of radius 5 cm . The required current in the loop is
 a) 0.56 A b) 5.6 A c) 0.28 A d) 2.8 A
153. Positively charged particles are projected into a magnetic field. If the direction of the magnetic field is along the direction of motion of the charge particles, the particles get
 a) Accelerated b) Decelerated c) Deflected d) No change in velocity
154. The deflection in moving coil galvanometer is reduced to half, when it is shunted with a 40Ω coil. The resistance of the galvanometer is
 a) 60Ω b) 10Ω c) 40Ω d) 20Ω
155. A coil having N turns carry a current I as shown in the figure. The magnetic field intensity at point P is




- a) $\frac{\mu_0 N I R^2}{2(R^2 + x^2)^{3/2}}$ b) $\frac{\mu_0 N I}{2R}$ c) $\frac{\mu_0 N I R^2}{(R + x)^2}$ d) Zero

156. In a cyclotron, the angular frequency of a charged particle is independent of
 a) Mass b) Speed c) Charge d) Magnetic field
157. A square current carrying loop is suspended in a uniform magnetic field acting in the plane of the loop. If the force on one arm of the loop is \vec{F} , the net force on the remaining three arms of the loop is
 a) \vec{F} b) $3\vec{F}$ c) $-\vec{F}$ d) $-3\vec{F}$
158. A solenoid of 1.5 m length and 4.0 cm diameter has 10 turns per cm. A current of 5 A is flowing through it. The magnetic field induction at axis inside the solenoid is
 a) $2\pi \times 10^{-4}$ T b) $2\pi \times 10^{-5}$ T c) 20π G d) 2π G
159. If a copper rod carries a direct current, the magnetic field associated with the current will be
 a) Only inside the rod b) Only outside the rod
 c) Both inside and outside the rod d) Neither inside nor outside the rod
160. An electron and a proton are projected at right angles to a uniform magnetic field with the same kinetic energy. Then
 a) The electron trajectory will be less curved than proton trajectory b) The electron trajectory will be more curved than proton trajectory
 c) Both the trajectories will be equally curved d) Both particles continue to move along a straight line

161. An electron and a proton with equal momentum enter perpendicularly into a uniform magnetic field, then
 a) The path of proton shall be more curved than that of electron
 b) The path of proton shall be less curved than that of electron
 c) Both are equally curved
 d) Path of both will be straight line
162. In the given diagram two long parallel wires carry equal currents in opposite direction. Point O is situated midway between the wires and the XY - plane contains the two wires and the positive Z -axis comes normally out of the plane of paper. The magnetic field B at O is non-zero along



- a) X, Y and Z -axes b) X -axis c) Y -axis d) Z -axis
163. A circular coil is in y - z plane with centre at origin. The coil is carrying a constant current. Assuming direction of magnetic field at $x = -25$ cm to be positive direction of magnetic field, which of the following graphs shows variation of magnetic field along x -axis




164. The unit of electric current "ampere" is the current which when flowing through each of two parallel wires spaced 1 m apart in vacuum and of infinite length will give rise to a force between them equal to
 a) 1 N/m b) 2×10^{-7} N/m c) 1×10^{-2} N/m d) $4\pi \times 10^{-7}$ N/m
165. A straight conductor carries a current of 5A. An electron travelling with a speed 5×10^6 ms⁻¹ parallel to the wire at a distance of 0.1m from the conductor, experiences a force of
 a) 8×10^{-20} N b) 3.2×10^{-19} N c) 8×10^{-18} N d) 1.6×10^{-19} N
166. A deuteron of kinetic energy 50 keV is describing a circular orbit of radius 0.5 m in a plane perpendicular to

magnetic field \vec{B} . The kinetic energy of the proton that describes a circular orbit of radius 0.5 m in the same plane with the same \vec{B} is

- a) 200 keV b) 100 keV c) 50 keV d) 25 keV

167. The resultant force on the current loop PQRS due to a long current carrying conductor will be



- a) 10^{-4} N b) 3.6×10^{-4} N c) 1.8×10^{-4} N d) 5×10^{-4} N


168. An electron moving in a circular orbit of radius r makes n rotations per second. The magnetic field produced at the centre has a magnitude of

- a) $\frac{\mu_0 n e}{2r}$ b) $\frac{\mu_0 n^2 e}{2r}$ c) $\frac{\mu_0 n e}{2\pi r}$ d) Zero


169. A current i flows along the length of an infinitely long, straight, thin-walled pipe. Then

- a) The magnetic field at all points inside the pipe is the same, but not zero b) The magnetic field at any point inside the pipe is zero
c) The magnetic field is zero only on the axis of the pipe d) The magnetic field is different at different points inside the pipe

170. The magnetic field lines due to a bar magnet are correctly shown in



171. In a square loop PQRS made with a wire of cross-section current i enters from point P and leaves from point S. The magnitude of magnetic field induction at the centre O of the square is



- a) $\frac{\mu_0}{4\pi} \frac{2\sqrt{2}i}{a}$ b) $\frac{\mu_0}{4\pi} \frac{4\sqrt{2}i}{a}$ c) $\frac{\mu_0}{4} \frac{2\sqrt{2}i}{a}$ d) zero

172. A cell is connected between two points of a uniformly thick circular conductor. The magnetic field at the centre of the loop will be

(Here i_1 and i_2 are the currents flowing in the two parts of the circular conductor of radius ' a ' and μ_0 has the usual meaning)

- a) Zero b) $\frac{\mu_0}{2a} (i_1 - i_2)$ c) $\frac{\mu_0}{2a} (i_1 + i_2)$ d) $\frac{\mu_0}{a} (i_1 + i_2)$

173. A electron moving with kinetic energy 6×10^{-16} J enters a field of magnetic induction 6×10^{-3} Wbm⁻² at right angle to its motion. The radius of its path is

- a) 3.42 cm b) 4.23 cm c) 6.17 cm d) 7.7 cm

174. A straight wire of diameter 0.5 mm carrying a current of 1 A is repaced by another wire of 1 mm diameter

carrying the same current. The strength of magnetic field far away is

- a) Twice the earlier value b) Half of the earlier value
c) Quarter of its earlier value d) Unchanged

175. The expression for the torque acting on a coil having area of cross-section A , number of turns n , placed in a magnetic field of strength B , making an angle θ with the normal to the plane of the coil, when a current i is flowing in it, will be

- a) $ni AB \tan \theta$ b) $ni AB \cos \theta$ c) $ni AB \sin \theta$ d) $ni AB$

176. An α -particle with a specific charge of $2.5 \times 10^7 \text{ C kg}^{-1}$ moves with a speed of $2 \times 10^5 \text{ ms}^{-1}$ in a perpendicular magnetic field of 0.05 T. Then the radius of the circular path described by it is

- a) 8 cm b) 4 cm c) 16 cm d) 2 cm

177. A proton (mass m and charge $+e$) and an α -particle (mass $4m$ and charge $+2e$) are projected with the same kinetic energy at right angles to the uniform magnetic field. Which one of the following statements will be true

- a) The α -particle will be bent in a circular path with a small radius that for the proton
b) The radius of the path of the α -particle will be greater than that of the proton
c) The α -particle and the proton will be bent in a circular path with the same radius
d) The α -particle and the proton will go through the field in a straight line

178. A vertical circular coil of radius 0.1 m and having 10 turns carries a steady current. When the plane of the coil is normal to the magnetic meridian, a neutral point is observed at the centre of the coil. If $B_H = 0.314 \times 10^{-4}$ the current in the coil is

- a) 0.5 A b) 0.25 A c) 2 A d) 1 A


179. Two particles A and B having equal charges $+6C$, after being accelerated through the same potential difference, enter a region of uniform magnetic field and describe circular paths of radii 2cm and 3cm respectively. The ratio of mass of A to that of B is

- a) $\frac{4}{9}$ b) $\frac{9}{5}$ c) $\frac{1}{2}$ d) $\frac{1}{3}$

180. The magnetic force on a charged particle moving in the field does not work, because

- a) Kinetic energy of the charged particle does not change
b) The charge of the particle remains same
c) The magnetic force is parallel to velocity of the particle
d) The magnetic force is parallel to magnetic field

181. An infinitely long hollow conducting cylinder with inner radius $R/2$ and outer radius R carries a uniform current density along its length. The magnitude of the magnetic field $|\vec{B}|$ as a function of the radial distance r from the axis is best represented by



182. A uniform magnetic field $\vec{B} = B_0 \hat{j}$ exists in space. A particle of mass m and charge q is projected towards x -axis with speed v from a point $(a, 0, 0)$. The maximum value of v for which the particle does not hit the y - z plane is

- a) $\frac{B q a}{m}$ b) $\frac{B q a}{2 m}$ c) $\frac{B q}{a m}$ d) $\frac{B q}{2 a m}$


183. Force acting on a magnetic pole of $7.5 \times 10^{-2} \text{ A-m}$ is 1.5 N. Magnetic field at the point is

- a) 20 Wbm^{-2} b) 50 Wbm^{-2} c) 112.5 T d) 2.0 T


184. A winding wire which is used to frame a solenoid can bear a maximum 10 A current. If length of solenoid is 80 cm and it's cross sectional radius is 3 cm then required length of winding wire is ($B = 0.2 \text{ T}$)

- a) $1.2 \times 10^2 \text{ m}$ b) $4.8 \times 10^2 \text{ m}$ c) $2.4 \times 10^3 \text{ m}$ d) $6 \times 10^3 \text{ m}$

185. A moving coil galvanometer has N number of turns in a coil of effective area A , it carries a current I . The magnetic field B is radial. The torque acting on the coil is
 a) NA^2B^2I b) $NAB I^2$ c) N^2ABI d) $NABI$
186. When a magnetic field is applied in a direction perpendicular to the direction of cathode rays, then their
 a) Energy decreases b) Energy increases
 c) Momentum increases d) Momentum and energy remain unchanged
187. A wire of length l is bent in the form of a circular coil of some turns. A current i flows through the coil. The coil is placed in a uniform magnetic field B . The maximum torque on the coil can be
 a) $\frac{iBl^2}{2\pi}$ b) $\frac{iBl^2}{4\pi}$ c) $\frac{iBl^2}{\pi}$ d) $\frac{2iBl^2}{\pi}$
188. A wire in the form of a square of side ' a ' carries a current i . Then the magnetic induction at the centre of the square wire is (Magnetic permeability of free space = μ_0)
 a) $\frac{\mu_0 i}{2\pi a}$ b) $\frac{\mu_0 i\sqrt{2}}{\pi a}$ c) $\frac{2\sqrt{2}\mu_0 i}{\pi a}$ d) $\frac{\mu_0 i}{\sqrt{2}\pi a}$
189. A charged particle is moving with velocity v in a magnetic field of induction B . The force on the particle will be maximum when
 a) v and B are in the same direction b) v and B are in opposite directions
 c) v and B are perpendicular d) v and B are at an angle of 45°
190. A particle of mass m and charge q is placed at a rest in a uniform electric field E and then released. The kinetic energy attained by the particle after moving a distance y is
 a) qEy^2 b) qE^2y c) qEy d) q^2Ey
191. The magnetic induction at the centre O in the figure shown is




- a) $\frac{\mu_0 i}{4} \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$ b) $\frac{\mu_0 i}{4} \left(\frac{1}{R_1} + \frac{1}{R_2} \right)$ c) $\frac{\mu_0 i}{4} (R_1 - R_2)$ d) $\frac{\mu_0 i}{4} (R_1 + R_2)$
192. A small coil of N turns has area A and a current I flows through it. The magnetic dipole moment of this coil will be
 a) NI/A b) NI^2A c) N^2AI d) NIA
193. The ratio of the magnitude field at the centre of a current carrying coil of the radius a and at a distance ' a ' from centre of the coil and perpendicular to the axis of coil is
 a) $\frac{1}{\sqrt{2}}$ b) $\sqrt{2}$ c) $\frac{1}{2\sqrt{2}}$ d) $2\sqrt{2}$
194. The figure shows the cross-section of a long cylindrical conductor of radius a carrying a uniformly distributed current i . The magnetic field due to current at P is



- a) $\mu_0 ir/(2\pi a^2)$ b) $\mu_0 ir^2/(2\pi a)$ c) $\mu_0 ia/(2\pi r^2)$ d) $\mu_0 ia^2/(\pi r^2)$
195. Identify the correct statement from the following
 a) Cyclotron frequency is dependent on speed of the charged particle
 b) Kinetic energy of charged particle in cyclotron does not depend on its mass
 c) Cyclotron frequency does not depend on speed of charged particle

- d) Kinetic energy of charged particle in cyclotron is independent of its charge
196. Two parallel beam of electrons moving in the same direction produce a mutual force
 a) Of attraction in plane of paper b) Of repulsion in plane of paper
 c) Upwards perpendicular to plane of paper d) Downward perpendicular to plane of paper
197. A thin disc having radius r and charge q distributed uniformly over the disc is rotated n rotations per second about its axis. The magnetic field at the centre of the disc is
 a) $\frac{\mu_0 qn}{2r}$ b) $\frac{\mu_0 qn}{r}$ c) $\frac{\mu_0 qn}{4r}$ d) $\frac{3\mu_0 qn}{4r}$
198. A one metre long wire is lying at right angles to the magnetic field. A force of 1 kg wt. is acting on it in a magnetic field of 0.98 tesla . The current flowing in it will be
 a) 100 A b) 10 A c) 1 A d) Zero
199. A long solenoid carrying a current produces a magnetic field B along its axis. If the current is doubled and the number of turns per cm is halved, the new value of the magnetic field is
 a) B b) $2B$ c) $4B$ d) $B/2$
200. A proton, a deuteron and an α – particle with the same kinetic energy enter a region of uniform magnetic field moving at right angles to B . What is the ratio of the radii of their circular paths?
 a) $1 : \sqrt{2} : \sqrt{2}$ b) $1 : \sqrt{2} : 1$ c) $\sqrt{2} : 1 : 1$ d) $\sqrt{2} : \sqrt{2} : 1$
201. A metallic loop is placed in a magnetic field. If a current is passed through it, then
 a) The ring will feel a force of attraction
 b) The ring will feel a force of repulsion
 c) It will move to and from about its centre of gravity
 d) None of these
202. A charge is fired through a magnetic field. The force acting on the charge is maximum when the angle between the direction of motion of charge and the magnetic field is
 a) Zero b) $\frac{\pi}{4}$ c) π d) $\frac{\pi}{2}$
203. The field due to a long straight wire carrying a current I is proportional to
 a) I b) I^3 c) \sqrt{I} d) $1/I$
204. The forces existing between two parallel current carrying conductors is F . If the current in each conductor is doubled, then the value of force will be
 a) $2F$ b) $4F$ c) $5F$ d) $F/2$
205. The force between two parallel current carrying wires is independent of
 a) Their distance of separation b) The length of the wires
 c) The magnitude of currents d) The radii of the wires
206. An electron (mass = $9.1 \times 10^{-31} \text{ kg}$, charge = $1.6 \times 10^{-18} \text{ C}$) experience no deflection if subjected to an electric field of $3.2 \times 10^5 \text{ V/m}$, and a magnetic field of $2.0 \times 10^{-3} \text{ Wb/m}^2$. Both the fields are normal to the path of electron and to each other. If the electric field is removed, then the electron will revolve in an orbit of radius
 a) 45 m b) 4.5 m c) 0.45 m d) 0.045
207. If the direction of the initial velocity of the charged particle is neither along nor perpendicular to that of the magnetic field, then the orbit will be
 a) A straight line b) An ellipses c) A circle d) A helix
208. An electron and a proton have equal kinetic energies. They enter in a magnetic field perpendicular to B , then
 a) Both will follow a circular path with same radius
 b) Both will follow a helical path
 c) Both will follow a parabolic path
 d) All the statements are false
209. A current carrying loop is placed in a uniform magnetic field in four different orientations, I, II, III & IV. Arrange them in the decreasing order of potential energy



- a) $I > III > II > IV$ b) $I > II > III > IV$ c) $I > IV > II > III$ d) $III > IV > I > II$


210. The ratio of magnetism potentials due to magnetic dipole in the end on position to that in broad side on position for the same distance from it is

- a) Zero b) ∞ c) 1 d) 2

211. A long copper tube of inner radius R carries a current i . The magnetic field B inside the tube is

- a) $\frac{\mu_0 i}{2\pi R}$ b) $\frac{\mu_0 i}{4\pi R}$ c) $\frac{\mu_0 i}{2R}$ d) Zero

212. Three infinite straight wires A , B and C carry currents as shown. The net force on the wire B is directed



- a) Toward A b) Toward C
c) Normal to plane of paper d) zero

213. A wire of length l is bent into a circular loop of radius R and carries a current I . The magnetic field at the centre of the loop is B . The same wire is now bent into a double loop of equal radii. If both loop carry the same current I and it is in the same direction, the magnetic field at the centre of the double loop will be

- a) Zero b) $2B$ c) $4B$ d) $8B$

214. Gauss is unit of which quantity

- a) H b) B c) ϕ d) I


215. Due to 10 ampere of current flowing in a circular coil of 10 cm radius, the magnetic field produced at its centre is $3.14 \times 10^{-3} \text{ weber/m}^2$. The number of turns in the coil will be

- a) 5000 b) 100 c) 50 d) 25

216. A beam of electrons and protons move parallel to each other in the same direction, then they

- a) Attract each other b) Repel each other
c) No relation d) Neither attract nor repel

217. The $(\tau - \theta)$ graph for a coil is




218. A bar magnet is cut into two equal halves by a plane parallel to the magnetic axis. Of the following physical quantities, the one which remains unchanged is

- a) Pole strengths b) Magnetic moment
c) Intensity of magnetization d) Moment of inertia

219. A particle of charge q and mass m starts moving from the origin under the action of an electric field $\vec{E} = E_0 \hat{i}$ and $\vec{B} = B_0 \hat{i}$ with a velocity $\vec{V} = v_0 \hat{j}$. The speed of the particle will become $\frac{\sqrt{5}}{2} v_0$ after a time

- a) $\frac{mv_0}{qE}$ b) $\frac{m v_0}{2q E}$ c) $\frac{\sqrt{3} m v_0}{2q E}$ d) $\frac{\sqrt{5} m v_0}{2q E}$


220. Two insulated rings, one of slightly smaller diameter than the other are suspended along the common diameter as shown. Initially the planes of the rings are mutually perpendicular. When a steady current is set up in each of them,



- a) The two rings rotate into a common plane b) The inner ring oscillates about its initial position
c) The inner rings stays stationary while the outer one moves into plane of the inner ring d) The outer ring stays stationary while the inner one moves into plane of the outer ring
221. Current i_0 is passed through a solenoid of length l having number of turns N when it is connected to a DC source. A charged particle with charge q is projected along the axis of the solenoid with a speed v_0 . The velocity of the particle in the solenoid
- a) Increases b) Decreases c) Remain same d) Becomes zero
222. An electric current passes through a long straight copper wire. At a distance 5 cm from the straight wire, the magnetic field is B . The magnetic field at 20 cm from the straight wire would be
- a) $\frac{B}{6}$ b) $\frac{B}{4}$ c) $\frac{B}{3}$ d) $\frac{B}{2}$
223. The areas of cross-section of three magnets of same length area A , $2A$ and $6A$ respectively. The ratio of their magnetic moments will be
- a) $6 : 2 : 1$ b) $1 : 2 : 6$ c) $2 : 6 : 1$ d) $1 : 1 : 1$
224. A long wire carries a steady current. It is bent into a circle of one turn and the magnetic field at the centre of the coil is B . It is then bent into a circular loop of n turns. The magnetic field at the centre of the coil will be
- a) nB b) n^2B c) $2nB$ d) $2n^2B$
225. A proton a mass m and charge $+e$ is moving in a circular orbit in a magnetic field with energy 1 MeV . What should be the energy of α -particle (mass = $4m$ and charge = $+2e$), so that it can revolve in the path of same radius
- a) 1 MeV b) 4 MeV c) 2 MeV d) 0.5 MeV
226. Four wires each of length 2.0 m are bent into four P , Q , R and S and then suspended into a uniform magnetic field. Same current is passed in each loop
-
- a) Couple on loop P will be maximum b) Couple on loop Q will be maximum
c) Couple on loop R will be maximum d) Couple on loop S be maximum
227. A long straight wire carries a current of $\pi \text{ amp}$. The magnetic field due to it will be $5 \times 10^{-5} \text{ weber/m}^2$ at what distance from the wire [μ_0 = permeability of air]
- a) $10^4 \mu_0 \text{ metre}$ b) $\frac{10^4}{\mu_0} \text{ metre}$ c) $10^6 \mu_0 \text{ metre}$ d) $\frac{10^6}{\mu_0} \text{ metre}$
228. An ammeter has resistance R_0 and range I . What resistance should be connected in parallel with it to increase its range by nI ?
- a) $R_0/(n - 1)$ b) $R_0/(n + 1)$ c) R_0/n d) None of these
229. An electron (mass = $9.0 \times 10^{-31} \text{ kg}$ and charge ($1.6 \times 10^{-19} \text{ C}$) is moving in a circular orbit in a magnetic field of $1.0 \times 10^{-4} \text{ Wbm}^{-2}$. Its period of revolution is
- a) $2.1 \times 10^{-6} \text{ s}$ b) $1.05 \times 10^{-6} \text{ s}$ c) $7 \times 10^{-7} \text{ s}$ d) $3.5 \times 10^{-7} \text{ s}$
230. A charged particle (charge q) is moving in a circle of radius R with uniform speed v . The associated magnetic moment μ is given by

- a) $\frac{qvR}{2}$ b) qvR^2 c) $\frac{qvR^2}{2}$ d) qvR

231. A current I flowing through the loop as shown in figure. The magnetic field at centre O is



- a) $\frac{7\mu_0 I}{16R} \otimes$ b) $\frac{7\mu_0 I}{16R} \odot$ c) $\frac{7\mu_0 I}{16R} \odot$ d) $\frac{5\mu_0 I}{16R} \odot$


232. A man carrying suitable instruments for measuring electric and magnetic field passes by a stationary electron with velocity V . Then these instruments will note

- a) Electric field b) Magnetic field c) Both a and b d) None of these

233. The magnetic field B with in the solenoid having n turns per metre length and carrying a current of i ampere is given by


- a) $\frac{\mu_0 n i}{e}$ b) $\mu_0 n i$ c) $4\pi\mu_0 n i$ d) $n i$

234. Two magnets of equal magnetic moments M each are placed as shown in figure. The resultant magnetic moment is




- a) M b) $\sqrt{3}M$ c) $\sqrt{2}M$ d) $M/2$

235. The correct curve between the magnetic induction (B) along the axis of a long solenoid due to current flow i in it and distance x from one end is



236. Two wires PQ and QR , carry equal currents i as shown in figure. One end of both the wires extends to infinity $\angle PQR = \theta$. The magnitude of the magnetic field at O on the bisector angle of these two wires at a distance r from point Q is




- a) $\frac{\mu_0 i}{4\pi r} \sin\left(\frac{\theta}{2}\right)$ b) $\frac{\mu_0 i}{4\pi r} \cot\left(\frac{\theta}{2}\right)$ c) $\frac{\mu_0 i}{4\pi r} \tan\left(\frac{\theta}{2}\right)$ d) $\frac{\mu_0 i}{4\pi r} \frac{(1 + \cos\theta/2)}{(\sin\theta/2)}$

237. The expression for magnetic induction inside a solenoid of length L carrying a current I and having N number of turns is



- a) $\frac{\mu_0 N}{4\pi L I}$ b) $\mu_0 N I$ c) $\frac{\mu_0}{4\pi} N L I$ d) $\mu_0 \frac{N}{L} I$

238. Wires 1 and 2 carrying currents i_1 and i_2 respectively are inclined at an angle θ to each other. What is the force on a small element dl of wire 2 at a distance of r from wire 1 (as shown in figure) due to the magnetic field of wire 1



- a) $\frac{\mu_0}{2\pi r} i_1 i_2 dl \tan \theta$ b) $\frac{\mu_0}{2\pi r} i_1 i_2 dl \sin \theta$ c) $\frac{\mu_0}{2\pi r} i_1 i_2 dl \cos \theta$ d) $\frac{\mu_0}{4\pi r} i_1 i_2 dl \sin \theta$


239. A magnetic field $\vec{B} = B_0 \hat{j}$ exists in the region $a < x < 2a$ and $\vec{B} = -B_0 \hat{j}$, in the region $2a < x < 3a$, where B_0 is a positive constant. A positive point charge moving with a velocity $\vec{V} = V_0 \hat{i}$, where V_0 is a positive constant, enters the magnetic field at $x = a$. The trajectory of the charge in this region can be like



240. If two streams of protons move parallel to each other in the same direction, then they

- a) Do not exert any force on each other b) Repel each other
c) Attract each other d) Get rotated to be perpendicular to each other

241. A particle having a charge of $10.0 \mu\text{C}$ and mass $1 \mu\text{g}$ moves in a circle of radius 10 cm under the influence of a magnetic field of induction 0.1 T . When the particle is at a point P , a uniform electric field is switched on so that the particle starts moving along the tangent with a uniform velocity. The electric field is




- a) 0.1 V/m b) 1.0 V/m c) 10.0 V/m d) 100 V/m

242. Two very long, straight, parallel wires carry steady current i and $-i$ respectively. The distance between the wires is d . At a certain instant of time, a point charge q is at a point equidistant from the two wires, in the plane of the wires. Its instantaneous magnitude of the force due to the magnetic field acting on the charge at this instant is

- a) $\frac{\mu_0 i q v}{2\pi d}$ b) $\frac{\mu_0 i q v}{\pi d}$ c) $\frac{2\mu_0 i q v}{\pi d}$ d) zero

243. The magnetic field due to a straight conductor of uniform cross section of radius a and carrying a steady current is represented by



244. The magnetic field at the point of intersection of diagonals of a square wire loop of side L carrying a current I is

- a) $\frac{\mu_0 I}{\pi L}$ b) $\frac{2\mu_0 I}{\pi L}$ c) $\frac{\sqrt{2}\mu_0 I}{\pi L}$ d) $\frac{2\sqrt{2}\mu_0 I}{\pi L}$

245. A proton and a deuteron both having the same kinetic energy, enter perpendicularly into a uniform

magnetic field B . For motion of proton and deuteron on circular path of radius R_p and R_d respectively, the correct statement is

- a) $R_d = \sqrt{2} R_p$ b) $R_d = R_p/\sqrt{2}$ c) $R_d = R_p$ d) $R_d = 2R_p$

246. Magnetic dipole moment of a rectangular loop is

- a) Inversely proportional to current in loop
b) Inversely proportional to area of loop
c) Parallel to plane of loop and proportional to area of loop
d) Perpendicular to plane of loop and proportional to area of loop

247. A cylindrical conductor of radius ' R ' carries a current ' i '. The value of magnetic field at a point which is $R/4$ distance inside from the surface is $10T$. Find the value of magnetic field at point which is $4R$ distance outside from the surface

- a) $4/3 T$ b) $8/3 T$ c) $40/3 T$ d) $80/3 T$


248. A short bar magnet placed with its axis at 30° , with a uniform external magnetic field of $0.25 T$ experiences a torque of $4.5 \times 10^{-2} \text{ N-m}$. Magnetic moment of the magnet is

- a) 0.36 J T^{-1} b) 0.72 J T^{-1} c) 0.18 J T^{-1} d) Zero

249. A circular conductor of radius 5 cm produces a magnetic field of $7 \times 10^{-6} T$. The current flowing through the conductor is


- a) $3.0 A$ b) $2.25 A$ c) $4.5 A$ d) $0.56 A$

250. A neutron, a proton, an electron and an α – particle enter a region of uniform magnetic field with the same velocities. The magnetic field is perpendicular and directed into the plane of the paper. The tracks of the particles are labeled in the figure. The electron follows the track



- a) D b) C c) B d) A

251. A wire shown in figure carries a current of $40 A$. If $r = 3.14 \text{ cm}$, the magnetic field at point P will be



- a) $1.6 \times 10^{-3} T$ b) $3.2 \times 10^{-2} T$ c) $4.8 \times 10^{-3} T$ d) $6.0 \times 10^{-4} T$





252. Two parallel wires carrying currents in the same direction attract each other because of


- a) Potential difference between them b) Mutual inductance between them
c) Electric force between them d) Magnetic force between them

253. A proton (or charged particle) moving with velocity v is acted upon by electric field E and magnetic field B . The proton will move undeflected if

- a) E is perpendicular to B
b) E is parallel to v and perpendicular to B
c) E, B and v are mutually perpendicular and $v = \frac{E}{B}$
d) E and B both are parallel to v





254. A neutral point is obtained at the centre of a vertical circular coil carrying current. The angle between the plane of the coil and the magnetic meridian is

- a) 0 b) 45° c) 60° d) 90°
255. An electron enters a region where magnetic field (\vec{B}) and electric field (\vec{E}) are mutually perpendicular to one another then
- a) It will always move in the direction of \vec{B} b) It will always move in the direction of \vec{E}
 c) It always possesses circular motion d) It can go undeflected also
256. The deflection in a galvanometer falls from 50 division to 20 when a $12\ \Omega$ shunt is applied. The galvanometer resistance is
- a) $18\ \Omega$ b) $36\ \Omega$ c) $24\ \Omega$ d) $30\ \Omega$
257. A length l of wire carries a steady current i . It is bent first to form a circular plane coil of one turn. The same length is now bent more sharply to give three loops of smaller radius. The magnetic field at the centre caused by the same current is
- a) One-third of its first value b) Unaltered
 c) Three times of its initial value d) Nine times of its initial value
258. A long thin hollow metallic cylinder of radius ' R ' has a current i ampere. The magnetic induction ' B '-away from the axis at a distance r from the axis varies as shown in
- a)  b)  c)  d) 
259. A circular coil of radius R carries an electric current. The magnetic field due to the coil at a point on the axis of the coil located at a distance r from the centre of the coil, such that $r \gg R$, varies as
- a) $\frac{1}{r}$ b) $\frac{1}{r^{3/2}}$ c) $\frac{1}{r^2}$ d) $\frac{1}{r^3}$
260. A wire of length L is bent in the form of a circular coil and current i is passed through it. If this coil is placed in a magnetic field then the torque acting on the coil will be maximum when the number of turns is
- a) As large as possible b) Any number c) 2 d) 1
261. A charge moving with velocity v in X -direction is subjected to a field of magnetic induction in the negative X -direction. As a result, the charge will
- a) Remain unaffected b) Start moving in a circular path in Y - Z plane
 c) Retard along X -axis d) Move along a helical path around X -axis
262. A uniform magnetic field $B = 1.2\text{ mT}$ is directed vertically upward throughout the volume of a laboratory chamber. A proton ($m_p = 1.67 \times 10^{-27}\text{ kg}$) enters the laboratory horizontally from south to north. Calculate the magnitude of centripetal acceleration of the proton if its speed is $3 \times 10^7\text{ m/s}$
- a) $3.45 \times 10^{12}\text{ m/s}^2$ b) $1.67 \times 10^{12}\text{ m/s}^2$ c) $5.25 \times 10^{12}\text{ m/s}^2$ d) $2.75 \times 10^{12}\text{ m/s}^2$
263. A loop carrying current I lies in the x - y plane as shown in the figure. The unit vector \hat{k} is coming out of the plane of the paper. The magnetic moment of the current loop is




- a) $a^2 I \hat{k}$ b) $\left(\frac{\pi}{2} + 1\right) a^2 I \hat{k}$ c) $-\left(\frac{\pi}{2} + 1\right) a^2 I \hat{k}$ d) $(2\pi + 1) a^2 U \hat{k}$
264. Two parallel beams of protons and electrons, carrying equal currents are fixed at a separation d . The protons and electrons move in opposite directions. P is a point on a line joining the beams, at a distance x from any one beam. The magnetic field at P is B . If B is plotted against x , which of the following best

represents the resulting curve




265. A long insulated copper wire is closely wound as a spiral of N turns. The spiral has inner radius a and outer radius b . The spiral lies in the $X - Y$ plane and a steady current I flows through the wire. The Z -component of the magnetic field at the centre of the spiral is



- a) $\frac{\mu_0 NI}{2(b-a)} \ln\left(\frac{b}{a}\right)$ b) $\frac{\mu_0 NI}{2(b-a)} \ln\left(\frac{b+a}{b-a}\right)$ c) $\frac{\mu_0 NI}{2b} \ln\left(\frac{b}{a}\right)$ d) $\frac{\mu_0 NI}{2b} \ln\left(\frac{b+a}{b-a}\right)$

266. Two wires with currents 2 A and 1 A are enclosed in a circular loop. Another wire with current 3 A is situated outside the loop as shown. The $\oint \vec{B} \cdot d\vec{l}$ around the loop is



- a) μ_0 b) $3\mu_0$ c) $6\mu_0$ d) $2\mu_0$

267. The distance at which the magnetic field on axis as compared to the magnetic field at the center of the coil carrying current I and radius R is $\frac{1}{8}$, would be

- a) R b) $\sqrt{2}R$ c) $2R$ d) $\sqrt{3}R$


268. A particle of mass M and charge Q moving with velocity \vec{v} describes a circular path of radius R when subjected to a uniform transverse magnetic field of induction B . The work done by the field when the particle completes one full circle is

- a) $BQv2\pi R$ b) $\left(\frac{Mv^2}{R}\right) 2\pi R$ c) Zero d) $BQ2\pi R$

269. The coil of a moving coil galvanometer is wound over a metal frame in order to

- a) Reduce hysteresis b) Provide electromagnetic damping
c) Increase the moment of inertia d) Increase the sensitivity


270. Current I is flowing in conductor shaped as shown in the figure. The radius of the curved part is r and the length of straight portion is very large. The value of the magnetic field at the centre O will be




- a) $\frac{\mu_0 I}{4\pi r} \left(\frac{3\pi}{2} + 1\right)$ b) $\frac{\mu_0 I}{4\pi r} \left(\frac{3\pi}{2} - 1\right)$ c) $\frac{\mu_0 I}{4\pi r} \left(\frac{\pi}{2} + 1\right)$ d) $\frac{\mu_0 I}{4\pi r} \left(\frac{\pi}{2} - 1\right)$

271. The helium nucleus makes a full rotation in a circle of radius 0.8 m in 2 s. The value of the magnetic field induction B in tesla at the centre of circle will be

- a) $2 \times 10^{-19} \mu_0$ b) $10^{-19}/\mu_0$ c) $10^{-19} \mu_0$ d) $2 \times 10^{-20}/\mu_0$
272. An electron rotates about a proton, the induced magnetic field is $14T$ at the centre, find out the angular velocity of electron if radius of rotation is 0.5 nm
 a) $4.4 \times 10^{17} \text{ rad/sec}$ b) $4.4 \times 10^{12} \text{ rad/sec}$ c) $3.14 \times 10^{-15} \text{ rad/sec}$ d) $4.2 \times 10^{10} \text{ rad/sec}$
273. The electrons in the beam of television tube move horizontally from south to north. The vertical component of the earth's magnetic field points down. The electron is deflected towards
 a) West b) No deflection c) East d) North to south
274. A circular coil of 5 turns and of 10 cm mean diameter is connected to a voltage source. If the resistance of the coil is 10Ω , the voltage of the source so is to nullify the horizontal component of earth's magnetic field of 30 A turn m^{-1} at centre of the coil should be
 a) 6 V , plane of the coil normal to magnetic meridian
 b) 2 V , plane of the coil normal to magnetic meridian
 c) 6 V , plane of the coil along the magnetic meridian
 d) 2 V , plane of the coil along the magnetic meridian
275. The magnetic field at the centre of a current carrying circular loop is B . If the radius of the loop is doubled, keeping the current same, the magnetic field at the centre of the loop would be
 a) $\frac{B}{4}$ b) $\frac{B}{2}$ c) $2B$ d) $4B$
276. A current loop consists of two identical semicircular parts each of radius R , one lying in the $x - y$ plane and the other in $x - z$ plane. If the current in the loop is i . The resultant magnetic field due to the two semicircular part at their common centre is
 a) $\frac{\mu_0 i}{2\sqrt{2}R}$ b) $\frac{\mu_0 i}{2R}$ c) $\frac{\mu_0 i}{4R}$ d) $\frac{\mu_0 i}{\sqrt{2}R}$
277. $PQRS$ is a square loop made of uniform wire. If the current enters the loop at P and leaves at S , then the magnetic field will be




- a) Maximum at the centre of the loop b) Zero at the centre of loop
 c) Zero at all points inside the loop d) Zero at all points outside of the loop
278. An electron (charge q coulomb) enters a magnetic field of $H \text{ weber/m}^2$ with a velocity of vm/s in the same direction as that of the field. The force on the electron is
 a) Hqv newtons in the direction of the magnetic field
 b) Hqv dynes in the direction of the magnetic field
 c) Hqv newtons at right angles to the direction of the magnetic field
 d) Zero
279. Two long parallel conductors carry currents in opposite directions as shown. One conductor carries a current of 10 A and the distance between the wires is $d = 10 \text{ cm}$. Current I is adjusted, so that the magnetic field at P is zero. P is at a distance of 5 cm to the right of the 10 A current. Value of I is



- a) 40 A b) 30 A c) 20 A d) 10 A
280. A current carrying loop is free to turn in a uniform magnetic field. The loop will then come into equilibrium when its plane is inclined at


- a) 0° to the direction of the field
 b) 45° to the direction of the field
 c) 90° to the direction of the field
 d) 135° to the direction of the field

281. Two identical bar magnets are fixed with their centres at a distance d apart. A stationary charge Q is placed at P in between the gap of the two magnets at a distance D from the centre O as shown in the figure



The force on the charge Q is

- a) Zero
 b) Directed along OP
 c) Directed along PO
 d) Directed perpendicular to the plane of paper
282. A circular coil having N turns is made from a wire of length L meter. If a current I ampere is passed through it and is placed in a magnetic field of B tesla, the maximum torque on it is
- a) Directly proportional to N
 b) Inversely proportional to N
 c) Inversely proportional to N^2
 d) Independent of N
283. The strength of the magnetic field around a long straight wire, carrying current, is
- a) Same everywhere around the wire at any distance
 b) Inversely proportional to the distance from the wire
 c) Inversely proportional to the square of the distance from the wire
 d) Directly proportional to the square of the distance from the wire
284. A positive charge is moving towards an observer. The direction of magnetic induction is
- a) Clockwise
 b) Anticlockwise
 c) Right
 d) Left
285. Motion of a moving electron is not affected by
- a) An electric field applied in the direction of motion
 b) Magnetic field applied in the direction of motion
 c) Electric field applied perpendicular to the direction of motion
 d) Magnetic field applied perpendicular to the direction of motion
286. An electron and a proton travel with equal speed in the same direction at 90° to a uniform magnetic field as this is switched on. They experience forces which are initially
- a) Identical
 b) Equal but in opposite direction
 c) In the same direction and differing by a factor of about 1840
 d) In opposite direction and differing by a factor of about 1840
287. At a distance of 10 cm from a long straight wire carrying current, the magnetic field is 0.04 T . At the distance of 40 cm , the magnetic field will be
- a) 0.01 T
 b) 0.02 T
 c) 0.08 T
 d) 0.16 T
288. If a current is passed in a spring, it
- a) Gets compressed
 b) Get expanded
 c) Oscillates
 d) Remains unchanged
289. A uniform magnetic field acts at right angles to the direction of motion of electrons. As a result, the electron moves in a circular path of radius 2 cm . If the speed of the electrons is doubled, then the radius of the circular path will be
- a) 2.0 cm
 b) 0.5 cm
 c) 4.0 cm
 d) 1.0 cm
290. A straight rod of mass m and length L is suspended from the identical springs as shown in figure. The spring is stretched a distance x_0 due to the weight of the wire. The circuit has total resistance R . When the magnetic field perpendicular to the plane of paper is switched on, springs are observed to extend further by the same distance. The magnetic field strength is



- a) $\frac{2mgR}{LE}$ b) $\frac{mgR}{LE}$ c) $\frac{mgR}{2LE}$ d) $\frac{mgR}{E}$

291. If the direction of the initial velocity of the charged particle is perpendicular to the magnetic field, then the orbit will be

Or

The path executed by a charged particle whose motion is perpendicular to magnetic field is

- a) A straight line b) An ellipse c) A circle d) A helix

292. A metal wire of mass m slides without friction on two rails placed at a distance l apart. The track lies in a uniform vertical magnetic field B . A constant current i flows along the rails across the wire and back down the other rail. The acceleration of the wire is

- a) $\frac{B m i}{l}$ b) $m B i l$ c) $\frac{B i l}{m}$ d) $\frac{m i l}{B}$

293. If a proton, deuteron and α -particle on being accelerated by the same potential difference enters perpendicular to the magnetic field, then the ratio of their kinetic energies is

- a) 1 : 2 : 2 b) 2 : 2 : 1 c) 1 : 2 : 1 d) 1 : 1 : 2

294. Proton and α -particle are projected perpendicularly in a magnetic field, if both move in a circular path with same speed. Then ratio of their radii is

- a) 1 : 2 b) 2 : 1 c) 1 : 4 d) 1 : 1


295. Ampere's circuital law is equivalent to

- a) Biot-Savart law b) Coulomb's law c) Faraday's law d) Kirchhoff's law

296. We have a galvanometer of resistance 25Ω . It is shunted by a 2.5Ω wire. The part of total current i_0 that flows through the galvanometer is given as


- a) $(i/i_0) = (1/11)$ b) $(i/i_0) = (1/10)$ c) $(i/i_0) = (1/9)$ d) $(i/i_0) = (2/11)$

297. In the given figure and magnetic field at O will be



- a) $\frac{2\mu_0 i}{3\pi a} \sqrt{4 - \pi^2}$ b) $\frac{\mu_0 i}{3\pi a} \sqrt{4 + \pi^2}$ c) $\frac{2\mu_0 i}{3\pi a^2} \sqrt{4 + \pi^2}$ d) $\frac{\mu_0 i}{3\pi a} \sqrt{4 - \pi^2}$





298. For a positively charged particle moving in a x - y plane initially along the x -axis, there is a sudden change in its path due to the presence of electric and/or magnetic fields beyond P. The curved path is shown in the x - y plane and is found to be non-circular. Which one of the following combination is possible




- a) $\vec{E} + 0; \vec{B} = b\hat{i} + c\hat{k}$ b) $\vec{E} + a\hat{i}; \vec{B} = c\hat{k} + a\hat{i}$ c) $\vec{E} + 0; \vec{B} = c\hat{j} + b\hat{k}$ d) $\vec{E} + a\hat{i}; \vec{B} = c\hat{k} + b\hat{j}$


299. Four charged particles are projected perpendicularly into the magnetic field with equal. Which will have minimum frequency?


- a) Proton b) Electron c) Li^+ d) He^+

300. A particle of mass 0.6 g and having charge of 25 nC is moving horizontally with a uniform velocity $1.2 \times 10^4\text{ ms}^{-1}$ in a uniform magnetic field, then the value of the magnetic induction is ($g = 10\text{ ms}^{-2}$)
 a) Zero b) 10 T c) 20 T d) 200 T
301. An electron and a proton enter region of uniform magnetic field in a direction at right angles to the field with the same kinetic energy. They describe circular paths of radius r_e and r_p respectively. Then
 a) $r_e = r_p$
 b) $r_e < r_p$
 c) $r_e > r_p$
 d) r_e may be less than or greater than r_p depending on the direction of the magnetic field
302. A moving coil galvanometer has N number of turns in a coil of effective area A , it carries a current I . The magnetic field B is radial. The torque acting on the coil is
 a) A moving coil galvanometer has N number of turns in a coil of effective area A , it carries a current I . The magnetic field B is radial. The torque acting on the coil is
 b) A moving coil galvanometer has N number of turns in a coil of effective area A , it carries a current I . The magnetic field B is radial. The torque acting on the coil is
 c) A moving coil galvanometer has N number of turns in a coil of effective area A , it carries a current I . The magnetic field B is radial. The torque acting on the coil is
 d) A moving coil galvanometer has N number of turns in a coil of effective area A , it carries a current I . The magnetic field B is radial. The torque acting on the coil is
303. The magnetic dipole moment of a current loop is independent of
 a) Magnetic field in which it is lying b) Number of turns
 c) Area of the loop d) Current in the loop
304. A pulsar is a neutron star having magnetic field is 10^{12} G at its surface. The maximum magnetic force experienced by an electron moving with velocity 0.9 c is
 a) 43.2 N b) $4.32 \times 10^{-3}\text{ N}$ c) $4.32 \times 10^3\text{ N}$ d) zero
305. In hydrogen atom, the electron is making $6.6 \times 10^{15}\text{ rev s}^{-1}$ around the nucleus of radius of 53 \AA . The magnetic field produced at the centre of the orbit is nearly
 a) 0.14 Wbm^{-2} b) 1.4 Wbm^{-2} c) 14 Wbm^{-2} d) 140 Wbm^{-2}
306. A particle of charge q and mass m is moving along the x -axis with a velocity v and enters a region of electric field E and magnetic field B as shown in figure below. For which figure the net force on the charge may be zero
 a)  b) 
 c)  d) 
307. Two infinite length wires carry currents 8 A and 6 A respectively and are placed along X and Y -axis. Magnetic field at a point $P(0,0,d)\text{ m}$ will be
 a) $\frac{7\mu_0}{\pi l}$ b) $\frac{10\mu_0}{\pi l}$ c) $\frac{14\mu_0}{\pi l}$ d) $\frac{5\mu_0}{\pi l}$
308. A closely wound flat circular coil of 25 turns of wire has diameter of 10 cm and carries a current of 4 ampere . Determine the flux density at the centre of a coil
 a) $1.679 \times 10^{-5}\text{ tesla}$ b) $2.028 \times 10^{-4}\text{ tesla}$ c) $1.257 \times 10^{-3}\text{ tesla}$ d) $1.512 \times 10^{-6}\text{ tesla}$
309. A long wire A carries a current of 10 amp . Another long wire B , which is parallel to A and separated by 0.1 m from A , carries a current of 5 amp , in the opposite direction to that in A . What is the magnitude and nature of the force experience per unit length of B
 ($\mu_0 = 4\pi \times 10^{-7}\text{ weber/amp} - \text{m}$)
 a) Repulsive force of 10^{-4} N/m b) Attractive force of 10^{-4} N/m


- c) Repulsive force of $2\pi \times 10^{-5} N/m$ d) Attractive force of $2\pi \times 10^{-5} N/m$
310. The current in the windings on a toroid is 2.0A. There are 400 turns and the mean circumferential length is 40cm. If the inside magnetic field is 1.0T, the relative permeability is near to
 a) 100 b) 200 c) 300 d) 400
311. Magnetic field at the centre of a circular coil of radius R due to i flowing through it is B . The magnetic field at a point along the axis at distance R from the centre is
 a) $\frac{B}{2}$ b) $\frac{B}{4}$ c) $\frac{B}{\sqrt{8}}$ d) $\sqrt{8}B$
312. A candidate connects a moving coil ammeter A and a moving coil voltmeter V and a resistance R as shown in figure



- If the voltmeter reads 20 V and the ammeter reads 4 A, then R is
 a) Equal to 5 Ω b) Greater than 5 Ω
 c) Less than 5 Ω d) Greater or less than 5 Ω depending upon its material
313. A bar magnet of length 3 cm has a point A and B along axis at a distance of 24 cm and 48 cm on the opposite ends. Ratio of magnetic fields at these points will be

 a) 8 b) 3 c) 4 d) $1/2\sqrt{2}$
314. Figure shows the cross-sectional view of the hollow cylindrical conductor with inner radius ' R ' and outer radius ' $2R$ '. Cylinder is carrying uniformly distributed current along its axis. The magnetic induction at point ' P ' at a distance $\frac{3R}{2}$ from the axis of the cylinder will be



- a) Zero b) $\frac{5\mu_0 i}{72\pi R}$ c) $\frac{7\mu_0 i}{18\pi R}$ d) $\frac{5\mu_0 i}{36\pi R}$
315. The radius of circular path of an electron when subjected to a perpendicular magnetic field is
 a) $\frac{mv}{Be}$ b) $\frac{me}{Be}$ c) $\frac{mE}{Be}$ d) $\frac{Be}{mv}$
316. A current carrying coil is subjected to a uniform magnetic field. The coil will orient so that its plane becomes
 a) Inclined at 45° to the magnetic field b) Inclined at any arbitrary angle to the magnetic field
 c) Parallel to the magnetic field d) Perpendicular to the magnetic field
317. A power line lies along the east-west direction and carries a current of 10 ampere. The force per metre due to the earth's magnetic field of 10^{-4} tesla is
 a) $10^{-5} N$ b) $10^{-4} N$ c) $10^{-3} N$ d) $10^{-2} N$
318. A current carrying square loop is placed near an infinitely long current carrying wire as shown in figure. The torque acting on the loop is




- a) $\frac{\mu_0}{2\pi} i_1 i_2 r$ b) $\frac{\mu_0}{2\pi} i_1 i_2 \log_e 2$ c) $\frac{\mu_0}{2\pi} \frac{i_1 i_2 r}{2}$ d) zero

319. A steady current I goes through a wire loop PQR having shape of a right angle triangle with $PQ = 3x$, $PR = 4x$ and $QR = 5x$. If the magnitude of the magnetic field at P due to this loop is $k \left(\frac{\mu_0 I}{48\pi x} \right)$, find the value of k


- a) 8 b) 3 c) 7 d) None of these

320. A particle with charge q , moving with a momentum p , enters a uniform magnetic field normally. The magnetic field has magnitude B and is confined to a region of width d , where $d < \frac{p}{Bq}$. If the particle is deflected by an angle θ in crossing the field, then



- a) $\sin \theta = \frac{Bqd}{p}$ b) $\sin \theta = \frac{p}{Bqd}$ c) $\sin \theta = \frac{Bp}{qd}$ d) $\sin \theta = \frac{pd}{Bq}$

321. A candidate connects a moving coil voltmeter V and a moving coil ammeter A and resistor R as shown in figure?



If the voltmeter reads 10 V and the ammeter reads 2 A and then R is

- a) Equal to 5Ω b) Greater than 5Ω
c) Less than 5Ω d) Greater or less than 5Ω depending upon its material

322. A circular coil A of radius r carries current i . Another circular coil B of radius $2r$ carries current of i . The magnetic fields at the centres of the circular coils are in the ratio of

- a) 3 : 1 b) 4 : 1 c) 1 : 1 d) 2 : 1


323. A charge q is moving in a magnetic field then the magnetic force does not depend upon

- a) Charge b) Mass c) Velocity d) Magnetic field


324. A current carrying circular loop is freely suspended by a long thread. The plane of the loop will point in the direction

- a) Wherever left free b) North-south
c) East-west d) At 45° with the east-west direction


325. A current is flowing through a thin cylindrical shell of radius R . If energy density in the medium, due to magnetic field, at a distance r from axis of the shell is equal to U then which of the following graphs is correct



326. The figure shows three long straight wires P , Q and R carrying currents normal to the plane of the paper. All three currents have the same magnitude. Which arrow best shows the direction of the resultant force on the wire P




- a) A b) B c) C d) D
327. Magnetic induction at the centre of a circular loop of area $\pi \text{ m}^2$ is 0.1 T . The magnetic moment of the loop is ($\mu_0 =$ permeability of air)
- a) $\frac{0.1\pi}{\mu_0}$ b) $\frac{0.2\pi}{\mu_0}$ c) $\frac{0.3\pi}{\mu_0}$ d) $\frac{0.4\pi}{\mu_0}$
328. A long straight wire of radius a carries a steady current i . The current is uniformly distributed across its cross-section. The ratio of the magnetic field at $\frac{a}{2}$ and $2a$ is
- a) $\frac{1}{4}$ b) 4 c) 1 d) $\frac{1}{2}$
329. Figure shows a straight wire of length l carrying current i . The magnitude of a magnetic field produced by the current at point P is




- a) $\frac{\sqrt{2}\mu_0 i}{\pi l}$ b) $\frac{\mu_0 i}{4\pi l}$ c) $\frac{\sqrt{2}\mu_0 i}{8\pi l}$ d) $\frac{\mu_0 i}{2\sqrt{2}\pi l}$
330. The magnetic field near a current carrying conductor is given by
- a) Coulomb's law b) Lenz's law c) Biot-Savart's law d) Kirchhoff's law
331. An electron of charge e moves with a constant speed v along a circle of radius r . Its magnetic moment will be
- a) evr b) $evr/2$ c) $\pi r^2 ev$ d) $2\pi rev$
332. Uniform magnetic B is directed vertically upwards and 3 wires of equal length L , carrying equal current i are lying in a horizontal plane such that the first one is along north, second one along north-east and the third one at 60° north of east. Force exerted by magnetic field B on them is
- a) Zero on the first b) $\frac{BiL}{\sqrt{2}}$ on the second c) $\sqrt{3}\frac{BiL}{2}$ on the third d) BiL on all of them
333. There are 50 turns of a wire in every cm length of a long solenoid. If 4 ampere current is flowing in the solenoid, the approximate value of magnetic field along its axis at an internal point and at one end will be respectively
- a) $12.6 \times 10^{-3} \text{ weber/m}^2, 6.3 \times 10^{-3} \text{ weber/m}^2$
 b) $12.6 \times 10^{-3} \text{ weber/m}^2, 25.1 \times 10^{-3} \text{ weber/m}^2$
 c) $25.1 \times 10^{-3} \text{ weber/m}^2, 6.3 \times 10^{-3} \text{ weber/m}^2$
 d) $25.1 \times 10^{-5} \text{ weber/m}^2, 6.3 \times 10^{-5} \text{ weber/m}^2$

334. A current of 5 *ampere* is flowing in a wire of length 1.5 *metres*. A force of 7.5 *N* acts on it when it is placed in a uniform magnetic field of 2 *tesla*. The angle between the magnetic field and the direction of the current is
 a) 30° b) 45° c) 60° d) 90°
335. The torque required to hold a small circular coil of 10 turns, $2 \times 10^{-4} \text{ m}^2$ area and carrying 0.5 A current in the middle of a long solenoid of 10^3 turnsm^{-1} carrying 3 A current. With its axis perpendicular to the axis of the solenoid is
 a) $12\pi \times 10^{-7} \text{ Nm}$ b) $6\pi \times 10^{-7} \text{ Nm}$ c) $4\pi \times 10^{-7} \text{ Nm}$ d) $2\pi \times 10^{-7} \text{ Nm}$
336. A particle of charge e and mass m moves with a velocity v in a magnetic field B applied perpendicular to the motion of the particle. The radius r of its path in the field is
 a) $\frac{Bv}{em}$ b) $\frac{ev}{Bm}$ c) $\frac{Be}{mv}$ d) $\frac{mv}{Be}$
337. A current i is flowing through the loop. The direction of the current and the shape of the loop are as shown in the figure.
 The magnetic field at the centre of the loop is $\frac{\mu_0 i}{R}$ times.
 ($MA = R, MB = 2R, \angle DMA = 90^\circ$)



- a) $\frac{5}{16}$, but out of the plane of the paper b) $\frac{5}{16}$, but into the plane of the paper
 c) $\frac{7}{16}$, but out of the plane of the paper d) $\frac{7}{16}$, but into the plane of the paper
338. A wire of length l is bent into a circular coil of one turn of radius R_1 . Another wire of the same material and same area of cross-section and same lengths is bent into a circular coil of two turns of radius R_2 . When the same current flows, through the two coils, the ratio of magnetic induction at the centres of the two coils is
 a) 1 : 2 b) 1 : 1 c) 1 : 4 d) 3 : 1
339. Two free parallel wires carrying currents in the opposite directions
 a) attract each other b) repel each other
 c) do not effect each other d) get rotated to be perpendicular to each other
340. A conduction rod of 1 m length and 1 kg mass is suspended by two vertical wires through its ends. An external magnetic field of 2 T is applied normal to the rod. Now the current to be passed through the rod so as to make the tension in the wires zero is
 [Take $g = 10 \text{ ms}^{-2}$]
 a) 0.5 A b) 15 A c) 5 A d) 1.5 A
341. In a hydrogen atom, the electron is making 6.6×10^{15} rps in a circular path of radius 0.53 Å. What is the magnetic induction produced at the centre of orbit?
 a) 6.3 T b) 12.6 T c) 18.9 T d) 25.2 T
342. A closed loop PQRS carrying a current is placed in a uniform magnetic field. If the magnetic forces on segment PS, SR and RQ are F_1 , F_2 and F_3 respectively and are in the plane of the paper and along the directions shown, the force on the segment QP is



- a) $\sqrt{(F_3 - F_1)^2 - F_2^2}$ b) $F_3 + F_1 - F_2$ c) $F_3 - F_1 + F_2$ d) $\sqrt{(F_3 - F_1)^2 + F_2^2}$

343. When current is passed through a circular wire prepared from a long conducting wire, the magnetic field produced at its centre is B . Now a loop having two turns is prepared from the same wire and the same current is passed through it. The magnetic field at its centre will be

- a) $4B$ b) $\frac{B}{4}$ c) $\frac{B}{2}$ d) $16B$

344. A straight wire of length 0.5 metre and carrying a current of 1.2 ampere is placed in a uniform magnetic field of induction 2 tesla. The magnetic field is perpendicular to the length of the wire. The force on the wire is

- a) 2.4 N b) 1.2 N c) 3.0 N d) 2.0 N

345. A proton of energy 200 MeV enters the magnetic field of 5 T. If direction of field is from south to north and motion is upward, the force acting on it will be

- a) Zero b) $1.6 \times 10^{-10} \text{ N}$ c) $3.2 \times 10^{-8} \text{ N}$ d) $1.6 \times 10^{-6} \text{ N}$

346. An electron, moving in a uniform magnetic field of induction of intensity \vec{B} , has its radius directly proportional to

- a) Its charge b) Magnetic field c) Speed d) None of these

347. A coil carrying a heavy current and having large number of turns is mounted in a N-S vertical plane and a current flows in clockwise direction. A small magnetic needle at its centre will have its north pole in

- a) East-north direction b) West-north direction c) East-south direction d) West-south direction

348. A moving charge will gain energy due to the application of

- a) Electric field b) Magnetic field c) Both of these d) None of these

349. A charged particle with charge q enters a region of constant, uniform and mutually orthogonal fields \vec{E} and \vec{B} with a velocity \vec{v} perpendicular to both \vec{E} and \vec{B} , and comes out without any change in magnitude or direction of \vec{v} . Then

- a) $\vec{v} = \vec{E} \times \vec{B}/B^2$ b) $\vec{v} = \vec{E} \times \vec{B}/B^2$ c) $\vec{v} = \vec{E} \times \vec{B}/E^2$ d) $\vec{v} = \vec{B} \times \vec{E}/E^2$

350. The magnetic induction due to an infinitely long straight wire carrying a current i at a distance r from wire is given by

- a) $|\vec{B}| = \left(\frac{\mu_0}{4\pi}\right) \frac{2i}{r}$ b) $|\vec{B}| = \left(\frac{\mu_0}{4\pi}\right) \frac{r}{2i}$ c) $|\vec{B}| = \left(\frac{4\pi}{\mu_0}\right) \frac{2i}{r}$ d) $|\vec{B}| = \left(\frac{4\pi}{\mu_0}\right) \frac{r}{2i}$


351. A conductor in the form of a right angle ABC with $AB = 3 \text{ cm}$ and $BC = 4 \text{ cm}$ carries a current of 10 A. There is a uniform magnetic field of 5 T perpendicular to the plane of the conductor. The force on the conductor will be

- a) 1.5 N b) 2.0 N c) 2.5 N d) 3.5 N

352. Two concentric coils of 10 turns each are placed in the same plane. Their radii are 20 cm and 40 cm and carry 0.2 A and 0.3 A current respectively in opposite directions. The magnetic induction (in tesla) at the centre is


- a) $\frac{3}{4} \mu_0$ b) $\frac{5}{4} \mu_0$ c) $\frac{7}{4} \mu_0$ d) $\frac{9}{4} \mu_0$

353. Two long straight wires are set parallel to each other. Each carries a current i in the opposite direction and the separation between them is $2R$. The intensity of the magnetic field midway between them is




- a) Zero b) $\frac{\mu_0 i}{4\pi R}$ c) $\frac{\mu_0 i}{2\pi R}$ d) $\frac{\mu_0 i}{\pi R}$


354. A rectangular loop carrying current is placed near a long straight fixed wire carrying strong current such that long sides are parallel to wire. If the current in the nearer long side of loop is parallel to current in the wire. Then the loop



- a) Experiences no force b) Experiences a force towards wire
c) Experiences a force away from wire d) Experiences a torque but no force
355. A long, straight, solid metal wire of radius 2 mm carries a current uniformly distributed over its circular cross-section. The magnetic field induction at a distance 2 mm from its axis is B . Then the magnetic field induction at distance 1 mm from axis will be
a) B b) $B/2$ c) $2B$ d) B
356. Two particles of masses m_a and m_b and same charge are projected in a perpendicular magnetic field. They travel along circular paths of radius r_a and r_b such that $r_a > r_b$. Then which is true?
a) $m_a v_a > m_b v_b$ b) $m_a > m_b$ and $v_a > v_b$
c) $m_a = m_b$ and $v_a = v_b$ d) $m_b v_b > m_a v_a$
357. The relation between voltage sensitivity (σ_v) and current sensitivity (σ_i) of a moving coil galvanometer is (Resistance of galvanometer = G)
a) $\frac{\sigma_i}{G} = \sigma_v$ b) $\frac{\sigma_v}{G} = \sigma_i$ c) $\frac{G}{\sigma_v} = \sigma_i$ d) $\frac{G}{\sigma_i} = \sigma_v$
358. A straight wire of mass 200 g and length 1.5 m carries a current of 2 A. It is suspended in mid-air by a uniform horizontal magnetic field B . The magnitude of B (in tesla) is
a) 2 b) 1.5 c) 0.55 d) 0.65
359. A long solenoid has n turns per meter and current I A is flowing through it. The magnetic field at the ends of the solenoid is
a) $\frac{\mu_0 n I}{2}$ b) $\mu_0 n I$ c) Zero d) $2\mu_0 n I$
360. The maximum velocity to which a proton can be accelerated in a cyclotron of 10 MHz frequency and radius 50 cm is
a) 6.28×10^8 m/s b) 3.14×10^8 m/s c) 6.28×10^7 m/s d) 3.14×10^7 m/s
361. Two straight parallel wires, both carrying 10 ampere in the same direction attract each other with a force of 1×10^{-3} N. If both currents are doubled, the force of attraction will be
a) 1×10^{-3} N b) 2×10^{-3} N c) 4×10^{-3} N d) 0.25×10^{-3} N
362. A circular coil of radius 4 cm and of 20 turns carries a current of 3 amperes. It is placed in a magnetic field of intensity of 0.5 weber/m². The magnetic dipole moment of the coil is
a) 0.15 ampere – m² b) 0.3 ampere – m² c) 0.45 ampere – m² d) 0.6 ampere – m²
363. A circular loop of radius 0.0175m carries a current of 2.0 amp. The magnetic field at the centre of the loop is
($\mu_0 = 4\pi \times 10^{-7}$ weber/amp – m)
a) 1.57×10^{-5} weber/m² b) 8.0×10^{-5} weber/m²
c) 2.5×10^{-5} weber/m² d) 3.14×10^{-5} weber/m²
364. Two thick wires and two thin wires, all of same material and same length, form a square in three different ways P , Q and R as shown in the figure. With correct connections shown, the magnetic field due to the current flow, at the centre of the loop will be zero in case of



- a) Q and R b) P only c) P and Q d) P and R
365. In a mass spectrometer used to measuring the masses of ions, the ions are initially accelerated by an electric potential V and then made to describe semicircular paths of radius R using a magnetic field B . If V and B are kept constant, the ratio $\left(\frac{\text{charge on the ion}}{\text{mass of the ion}}\right)$ will be proportional
- a) $1/R$ b) $1/R^2$ c) R^2 d) R
366. If cathode rays are projected at right angles to a magnetic field, their, trajectory is
- a) Ellipse b) Circle c) Parabola d) None of these
367. A current carrying closed loop in the form of a right angle isosceles triangle ABC is placed in a uniform magnetic field acting along AB . If the magnetic force on the arm BC is \vec{F} , the force on the arm AC is
-
- a) $\sqrt{2}\vec{F}$ b) $-\sqrt{2}\vec{F}$ c) $-\vec{F}$ d) \vec{F}
368. A charged particle enters a magnetic field H with its initial velocity making an angle of 45° with H . The path of the particle will be
- a) A straight line b) A circle c) An ellipse d) A helix
369. In a hydrogen atom, an electron moves in a circular orbit of radius $5.2 \times 10^{-11} \text{ m}$ and produces a magnetic induction of 12.56 T at its nucleus. The current produced by the motion of the electron will be (Given $\mu_0 = 4\pi \times 10^{-7} \text{ Wb/A} - \text{m}$)
- a) $6.53 \times 10^{-3} \text{ ampere}$ b) $13.25 \times 10^{-10} \text{ ampere}$
c) $9.6 \times 10^6 \text{ ampere}$ d) $1.04 \times 10^{-3} \text{ ampere}$
370. The cyclotron frequency of an electron grating in a magnetic field of 1 T is approximately
- a) 28 MHz b) 280 MHz c) 2.8 GHz d) 28 GHz
371. A coil of 100 turns and area $2 \times 10^{-2} \text{ m}^2$, pivoted about a vertical diameter in a uniform magnetic field carries a current of 5 A . When the coil is held with its plane in North-South direction, it experiences a torque of 0.3 Nm . When the plane is in East-West direction the torque is 0.4 Nm . The value of magnetic induction is (Neglect earth's magnetic field)
- a) 0.2 T b) 0.3 T c) 0.4 T d) 0.5 T
372. Which is a vector quantity
- a) Density b) Magnetic flux
c) Intensity of magnetic field d) Magnetic potential
373. The number of lines of force passing through a unit area placed perpendicularly to the magnetic lines of force is termed as
- a) Magnetic induction b) Magnetic flux density
c) Intensity of magnetic field d) All of the above
374. A coil $PQRS$ carrying a current i ampere is placed in a powerful horse shoe magnet NS of uniform magnetic field \vec{B} figure. If A is the area of the coil and θ is the inclination of the plane of the coil with the magnetic field in equilibrium, then the deflecting couple will be



- a) $BiA \cot \theta$ b) $BiA \cos \theta$ c) $BiA \operatorname{cosec} \theta$ d) $BiA \sin \theta$


375. When a certain length of wire is turned into one circular loop, the magnetic induction at the centre of coil due to some current flowing is B_0 . If the same wire is turned into three loops to make a circular coil, the magnetic induction at the center of this coil for the same current will be

- a) B_0 b) $9 B_0$ c) $3 B_0$ d) $27 B_0$

376. Two straight long conductors AOB and COD are perpendicular to each other and carry currents i_1 and i_2 . The magnitude of the magnetic induction at a point P at a distance a from the point O in a direction perpendicular to the plane $ABCD$ is

- a) $\frac{\mu_0}{2\pi a} (i_1 + i_2)$ b) $\frac{\mu_0}{2\pi a} (i_1 - i_2)$ c) $\frac{\mu_0}{2\pi a} (i_1^2 + i_2^2)^{1/2}$ d) $\frac{\mu_0}{2\pi a} \frac{i_1 i_2}{(i_1 + i_2)}$

377. For the arrangement as shown in the figure, the magnetic induction at the centre is



- a) $\frac{3\mu_0 i \pi}{4a}$ b) $\frac{\mu_0 i}{4\pi a} (1 + \pi)$ c) $\frac{\mu_0 i}{4\pi a}$ d) $\frac{3\mu_0 i}{8a}$

378. The magnetic field $d\vec{B}$ due to a small current element $d\vec{l}$ at a distance \vec{r} and element carrying current i is
Or

Vector form of Bio-savart's law is

- a) $d\vec{B} = \frac{\mu_0}{4\pi} i \left(\frac{d\vec{l} \times \vec{r}}{r} \right)$ b) $d\vec{B} = \frac{\mu_0}{4\pi} i^2 \left(\frac{d\vec{l} \times \vec{r}}{r} \right)$
c) $d\vec{B} = \frac{\mu_0}{4\pi} i^2 \left(\frac{d\vec{l} \times \vec{r}}{r^2} \right)$ d) $d\vec{B} = \frac{\mu_0}{4\pi} i \left(\frac{d\vec{l} \times \vec{r}}{r^3} \right)$


379. A proton and a deuteron with the same initial kinetic energy enter a magnetic field in a direction perpendicular to the direction of the field. The ration of the radii of the circular trajectories described by them is

- a) $1 : 4$ b) $1 : \sqrt{2}$ c) $1 : 1$ d) $1 : 2$

380. The magnetic field at the centre of a circular coil of radius r carrying current I is B_1 . The field at the centre of another coil of radius $2r$ carrying same current I is B_2 . The ratio $\frac{B_1}{B_2}$ is

- a) $1/2$ b) 1 c) 2 d) 4

381. A very long straight wire carries a current I . At the instant when a charge $+Q$ at point P has velocity \vec{V} , as shown, the force on the charge is




- a) Opposite OX b) Along OX c) Opposite OY d) Along OY

382. Two concentric circular lops of radii R and $2R$ carry currents of $2I$ and I respectively in opposite sense

(i.e., clockwise in one coil and counter-clockwise in the other coil). The resultant magnetic field at their common centre is

- a) $\mu_0 \frac{I}{4R}$ b) $\mu_0 \frac{5I}{4R}$ c) $\mu_0 \frac{3I}{4R}$ d) $\mu_0 \frac{I}{2R}$

383. A proton accelerated by a potential difference 500 KV moves through a transverse magnetic field of 0.51 T as shown in figure. The angle θ through which the proton deviates from the initial direction of its motion is



- a) 15° b) 30° c) 45° d) 60°


384. A proton, a deuteron and an α - particle having the same kinetic energy are moving in circular trajectories in a constant magnetic field. If r_p , r_d and r_α denote respectively the radii of the trajectories of these particles, then

- a) $r_\alpha = r_d > r_p$ b) $r_\alpha = r_d = r_p$ c) $r_\alpha < r_d < r_p$ d) $r_\alpha = r_p < r_d$

385. The magnetic field at the centre of coil of n turns, bent in the form of a square of side $2l$, carrying i , is

- a) $\frac{\sqrt{2}\mu_0 ni}{\pi l}$ b) $\frac{\sqrt{2}\mu_0 ni}{2\pi l}$ c) $\frac{\sqrt{2}\mu_0 ni}{4\pi l}$ d) $\frac{2\mu_0 ni}{\pi l}$

386. The magnetic field induction at the centre O , in the arrangement shown in figure is



- a) $\frac{\mu_0 i}{4\pi r} (4 + \pi)$ b) $\frac{\mu_0 i}{4\pi r} (3 + \pi)$ c) $\frac{\mu_0 i}{4\pi r} (2 + \pi)$ d) $\frac{\mu_0 i}{4\pi r} (1 + \pi)$

387. A circular coil of wire consisting of 100 turns, each of radius 8.0 cm carries a current of 0.40 A. What is the magnitude of the magnetic field B at the centre of the coil?

- a) $\pi \times 10^{-3}$ T b) $2\pi \times 10^{-4}$ T c) $\pi \times 10^{-4}$ T d) Zero

388. One Tesla is equal to

- a) 10^7 gauss b) 10^{-4} gauss c) 10^4 gauss d) 10^{-8} gauss


389. Two wires A and B are of lengths 40 cm and 30 cm. A is bent into a circle of radius r and B into an arc of radius r . A current i_1 is passed through A and i_2 through B . To have the same magnetic inductions at the centre, the ratio of $i_1 : i_2$ is

- a) 3 : 4 b) 3 : 5 c) 2 : 3 d) 4 : 3

390. A homogenous electric field \vec{E} and a uniform magnetic field \vec{B} are pointing in the same direction. A proton is projected with its velocity parallel to \vec{E} . It will

- a) Go on moving in the same direction with increasing velocity
b) Go on moving in the same direction with constant velocity
c) Turn to its right
d) Turn to its left

391. Which of the following graph represents the variation of magnetic flux density B with distance r for a straight long wire carrying an electric current?




392. A vertical straight conductor carries a current vertically upwards. A point P lies to the east of it at a small distance and another point Q lies to the west at the same distance. The magnetic field at P is

- a) Greater than at Q
- b) Same as at Q
- c) Less than at Q
- d) Greater or less than at Q depending upon the strength of the current

393. Magnetic field due to a ring having n turns at a distance x on its axis is proportional to (if r = radius of ring)

- a) $\frac{r}{(x^2 + r^2)}$
- b) $\frac{r^2}{(x^2 + r^2)^{3/2}}$
- c) $\frac{nr^2}{(x^2 + r^2)^{3/2}}$
- d) $\frac{n^2r^2}{(x^2 + r^2)^{3/2}}$

394. A wire PQR is bent as shown in figure and is placed in a region of uniform magnetic field B . The length of $PQ = QR = l$. A current I ampere flows through the wire as shown. The magnitude of the force on PQ and QR will be



- a) $Bil, 0$
- b) $2Bil, 0$
- c) $0, Bil$
- d) $0, 0$

395. A small circular flexible loop of wire of radius r carries a current I . It is placed in a uniform magnetic field B . The tension in the loop will be doubled if

- a) I is doubled
- b) B is halved
- c) r is doubled
- d) Both B and I are doubled

396. Magnetic fields at two points on the axis of a circular coil at a distance of 0.05 m and 0.2 m from the centre are in the ratio $8:1$. The radius of the coil is

- a) 1.0 m
- b) 0.1 m
- c) 0.15 m
- d) 0.2 m

397. If a particle of charge 10^{-12} C moving along the x -direction with a velocity of 10^5 ms^{-1} experience a force of 10^{-10} N in y -direction due to magnetic field, then the minimum value of magnetic field is

- a) $6.25 \times 10^3\text{ T}$ in z -direction
- b) 10^{-15} T in z -direction
- c) $6.25 \times 10^{-3}\text{ T}$ in z -direction
- d) 10^{-3} T in z -direction

398. A current carrying rectangular coil is placed in a uniform magnetic field. In which orientation, the coil will not tend to rotate

- a) The magnetic field is parallel to the plane of the coil
- b) The magnetic field is perpendicular to the plane of the coil
- c) The magnetic field is at 45° with the plane of the coil
- d) Always in any orientation

399. The magnetic induction at a point P which is at a distance 4 cm from a long current carrying wire is 10^{-8} tesla . The field of induction at a distance 12 cm from the same current would be

- a) $3.33 \times 10^{-9}\text{ tesla}$
- b) $1.11 \times 10^{-4}\text{ tesla}$
- c) $3 \times 10^{-3}\text{ tesla}$
- d) $9 \times 10^{-2}\text{ tesla}$

400. A triangular loop of side l carries a current I . It is placed in a magnetic field B such that the plane of the loop is in the direction of B . The torque on the loop is

- a) Zero b) IBl c) $\frac{\sqrt{3}}{2} Il^2 B^2$ d) $\frac{\sqrt{3}}{4} IBl^2$

401. Two circular coils mounted parallel to each other on the same axis carry steady currents. If an observer between the coils reports that one coil is carrying a clockwise current i_1 , while the other is carrying a counter clockwise current i_2 , between the two coils, then there is

- a) A steady repulsive force b) Zero force
c) A repulsive force d) A steady attractive force

402. An electron is revolving around a proton in a circular path of diameter 0.1 nm. It produces a magnetic field 14 T at a proton. Then the angular speed of the electron is

- a) $8.8 \times 10^6 \text{ rad s}^{-1}$ b) $4.4 \times 10^{16} \text{ rad s}^{-1}$ c) $2.2 \times 10^{16} \text{ rad s}^{-1}$ d) $1.1 \times 10^{16} \text{ rad s}^{-1}$

403. Under the influence of a uniform magnetic field a charged particle is moving in a circle of radius R with constant speed v . The time period of the motion

- a) Depends on v and not on R b) Depends on both R and v
c) Is independent of both R and v d) Depends on R and not on v

404. A particle mass m and charge q enters a magnetic field B perpendicularly with a velocity v . The radius of the circular path described by it will be

- a) Bq/mv b) mq/Bv c) mB/qv d) mv/Bq


405. Two short bar magnets with magnetic moments 400 ab-amp cm² and 800 ab-amp cm² are placed with their axis in the same straight line with similar poles facing each other and with their centers at 20 cm from each other. Then the force of repulsion is

- a) 12 dyne b) 6 dyne c) 800 dyne d) 150 dyne

406. A charge q coulomb moves in a circle at n revolutions per second and the radius of the circle is r metre. Then the magnetic field at the centre of the circle is

- a) $\frac{2\pi q}{nr} \times 10^{-7} \text{ NA}^{-1} \text{ m}^{-1}$ b) $\frac{2\pi q}{r} \times 10^{-7} \text{ NA}^{-1} \text{ m}^{-1}$
c) $\frac{2\pi nq}{r} \times 10^{-7} \text{ NA}^{-1} \text{ m}^{-1}$ d) $\frac{2\pi q}{r} \times 10^{-6} \text{ NA}^{-1} \text{ m}^{-1}$

407. An equilateral triangle of side l is formed from a piece of wire of uniform resistance. The current i is fed as shown in the figure. The magnitude of the magnetic field at its centre O is



- a) $\frac{\sqrt{3}\mu_0 i}{2\pi l}$ b) $\frac{3\sqrt{2}\mu_0 i}{2\pi l}$ c) $\frac{\mu_0 i}{2\pi l}$ d) zero

408. A uniform electric field and a uniform magnetic field exist in a region in the same direction. An electron is projected with a velocity pointed in the same direction. Then the electron will

- a) Be deflected to the left without increase in speed
b) Be deflected to the right without increase in speed
c) Not be deflected but its speed will decrease
d) Not be deflected but its speed will increase

409. A solenoid 1.5 m long and 0.4 cm in diameter possesses 10 turns/cm length. A current of 5 A flows through it. The magnetic field at the axis inside the solenoid is

- a) $2\pi \times 10^{-3} \text{ T}$ b) $2\pi \times 10^{-5} \text{ T}$ c) $4\pi \times 10^{-2} \text{ T}$ d) $4\pi \times 10^{-3} \text{ T}$

410. A particle of mass m , charge Q and kinetic energy T enters a transverse uniform magnetic field of induction \vec{B} . After 3 seconds the kinetic energy of the particle will be

- a) T b) $4T$ c) $3T$ d) $2T$

411. A horizontal straight wire 10 m long extending from east to west falling with a speed of 5.0 m/s, at right

angles to the horizontal component of the earth's magnetic field of strength $0.30 \times 10^{-4} \text{ Wb/m}^2$. The instantaneous value of the induced potential gradient in the wire, from west to east is

- a) $+1.5 \times 10^{-3} \text{ V/m}$ b) $-1.5 \times 10^{-3} \text{ V/m}$ c) $+1.5 \times 10^{-4} \text{ V/m}$ d) $-1.5 \times 10^{-4} \text{ V/m}$


412. The radius of curvature of the path of the charged particle in a uniform magnetic field is directly proportional to

- a) The charge on the particle b) The momentum of the particle
c) The energy of the particle d) The intensity of the field

413. An electron moving towards the east enters a magnetic field directed towards the north. The force on the electron will be directed

- a) Vertically upward b) Vertically downward c) Towards the west d) Towards the south

414. Three long, straight and parallel wires carrying currents are arranged as shown in the figure. The wire *C* which carries a current of 5.0 amp is so placed that it experiences no force. The distance of wire *C* from wire *D* is then



- a) 9 cm b) 7 cm c) 5 cm d) 3 cm


415. A very high magnetic field is applied to a stationary charge. Then the charge experiences

- a) A force in the direction of magnetic field
b) A force perpendicular to the magnetic field
c) A force in an arbitrary direction
d) No force

416. A straight wire carrying current *i* is turned into a circular loop. If the magnitude of magnetic moment associated with it in MKS unit is *M*, the length of wire will be

- a) $\frac{4\pi}{M}$ b) $\sqrt{\frac{4\pi M}{i}}$ c) $\sqrt{\frac{r\pi i}{M}}$ d) $\frac{M\pi}{4i}$

417. A pair of stationary and infinite long bent wires are placed in the *x* – *y* plane. The wires carrying currents of 10 A each as shown in figure. The segments *L* and *M* are parallel to *x*-axis. The segments *P* and *Q* are parallel to *y*-axis, such that *OS* = *OR* = 0.02 m. The magnetic field induction at the origin *O* is



- a) 10^{-3} T b) $4 \times 10^{-3} \text{ T}$ c) $2 \times 10^{-4} \text{ T}$ d) 10^{-4} T

418. In a current carrying long solenoid, the field produced does not depend upon

- a) Number of turns per unit length b) Current flowing
c) Radius of the solenoid d) All of the above

419. The magnetic induction at the centre of a current carrying circular of radius *r*, is

- a) Directly proportional to *r* b) Inversely proportional to *r*
c) Directly proportional to r^2 d) Inversely proportional to r^2

420. Maximum kinetic energy of the positive ion in the cyclotron is

- a) $\frac{q^2 B r_0}{2m}$ b) $\frac{q B^2 r_0}{2m}$ c) $\frac{q^2 B^2 r_0^2}{2m}$ d) $\frac{q B r_0}{2m^2}$

421. The ratio of magnetic field and magnetic moment at the centre of a current carrying circular loop is *x*. When both the current and radius is doubled then the ratio will be

- a) $\frac{x}{8}$ b) $\frac{x}{4}$ c) $\frac{x}{2}$ d) $2x$

422. Consider the following statements regarding a charged particle in a magnetic field
 (i) starting with zero velocity, it accelerates in a direction perpendicular to the magnetic field
 (ii) While deflecting in the magnetic field, its energy gradually increases
 (iii) Only the component of magnetic field perpendicular to the direction of motion of the charged particle is effective in deflecting it
 (iv) Direction of deflecting force on the moving charged particle is perpendicular to its velocity. Of these statements
 a) (ii) and (iii) are correct b) (iii) and (iv) are correct
 c) (ii) . (iii) and (iv) are correct d) (i). (ii) and (iii) are correct
423. Toroid is
 a) Ring shaped closed solenoid b) Rectangular shaped solenoid
 c) Ring shaped open solenoid d) Square shaped solenoid
424. A steady current i flows in a small square loop of wire of side l in a horizontal plane. The loop is now folded about its middle such that half of it lies in a vertical plane. Let \hat{M}_1 and \hat{M}_2 respectively denote the magnetic moments due to current loop before and after folding. Then
 a) $\hat{M}_2 = 0$ b) \hat{M}_1 and \hat{M}_2 are in the same direction
 c) $M_1/M_2 = \sqrt{2}$ d) $M_1/M_2 = 1/\sqrt{2}$
425. A proton and an electron both moving with the same velocity v enter into a region of magnetic field directed perpendicular to the velocity of the particles. They will now move in circular orbits such that
 a) Their time periods will be same b) The time period for proton will be higher
 c) The time period for electron will be higher d) The orbital radii will be the same
426. Two similar coils are kept mutually perpendicular such that their centres coincide. At the centre, find the ratio of the magnetic field due to one coil and the resultant magnetic field by both coils, if the same current is flown
 a) $1 : \sqrt{2}$ b) $1 : 2$ c) $2 : 1$ d) $\sqrt{3} : 1$
427. A charge $+q$ is moving upwards vertically. It enters a magnetic field directed to the north. The force on the charged will be towards
 a) North b) South c) West d) East
428. A ring of radius R , made of an insulating material carries a charge Q uniformly distributed on it. If the ring rotates about the axis passing through its centre and normal to plane of the ring with constant angular speed ω , then the magnitude of the magnetic moment of the ring is
 a) $Q\omega R^2$ b) $1/2 Q\omega R^2$ c) $Q\omega^2 R$ d) $1/2 Q\omega^2 R$
429. The velocity of two α -particles A and B in a uniform magnetic field is in the ratio of $1 : 3$. They move in different circular orbits in the magnetic field. The ratio of radius of curvatures of their paths is
 a) $1 : 2$ b) $1 : 3$ c) $3 : 1$ d) $2 : 1$
430. The strength of the magnetic field at a point r near a long straight current carrying wire is B . The field at a distance $\frac{r}{2}$ will be
 a) $\frac{B}{2}$ b) $\frac{B}{4}$ c) $2B$ d) $4B$
431. A current carrying loop is placed in a uniform magnetic field. The torque acting on it does not depend upon
 a) Shape of the loop b) Area of the loop c) Value of the current d) Magnetic field
432. An ammeter has a resistance of $G \Omega$ and a range of i ampere. The value of resistance used in parallel to convert it into an ammeter of range in ampere is
 a) nG b) $(n-1)G$ c) $\frac{G}{n}$ d) $\frac{G}{n-1}$
433. A particle with 10^{-11} coulomb of charge and 10^{-7} kg mass is moving with a velocity of 10^8 m/s along the y -axis. A uniform static magnetic field $B = 0.5 \text{ tesla}$ is acting along the x -direction. The force on the particle

is

- a) $5 \times 10^{-11} \text{ N}$ along \hat{i} b) $5 \times 10^3 \text{ N}$ along \hat{k} c) $5 \times 10^{-11} \text{ N}$ along $-\hat{j}$ d) $5 \times 10^{-4} \text{ N}$ along $-\hat{k}$


434. A current of 3A is flowing in a linear conductor having a length of 40 cm. The conductor is placed in a magnetic field of strength 500 G and makes an angle of 30° with the direction of the field. It experiences a force of magnitude

- a) $3 \times 10^{-4} \text{ N}$ b) $3 \times 10^{-2} \text{ N}$ c) $3 \times 10^2 \text{ N}$ d) $3 \times 10^4 \text{ N}$

435. The direction of induced magnetic field \mathbf{dB} due to current element $i \mathbf{dL}$, at a point of distance r from it, when a current i passes through a long conductor is in the direction

- a) Of position vector \mathbf{r} of the point b) Of current element \mathbf{dL}
c) Perpendicular to both \mathbf{dL} and \mathbf{r} d) Perpendicular to \mathbf{dL} only

436. In the figure shown, the magnetic field induction at the point O will be



- a) $\frac{\mu_0 i}{2\pi r}$ b) $\left(\frac{\mu_0}{4\pi}\right)\left(\frac{i}{r}\right)(\pi + 2)$ c) $\left(\frac{\mu_0}{4\pi}\right)\left(\frac{i}{r}\right)(\pi + 1)$ d) $\frac{\mu_0 i}{4\pi r}(\pi - 2)$

437. Two thin, long, parallel wires, separated by a distance d carry a current of i ampere in the same direction. They will

- a) Attract each other with a force of $\frac{\mu_0 i^2}{(2\pi d)}$ b) Repel each other with a force of $\frac{\mu_0 i^2}{(2\pi d)}$
c) Attract each other with a force of $\frac{\mu_0 i^2}{(2\pi d^2)}$ d) Repel each other with a force of $\frac{\mu_0 i^2}{(2\pi d^2)}$

438. The current is flowing in south direction along a power line. The direction of magnetic field above the power line (neglecting earth's field) is

- a) South b) East c) North d) West

439. The magnitude of the magnetic field required to accelerate protons (mass = $1.67 \times 10^{-27} \text{ kg}$) in a cyclotron that is operated at an oscillator frequency 12 MHz is approximately

- a) 0.8 T b) 1.6 T c) 2.0 T d) 3.2 T

440. A current carrying small loop behaves like a small magnet. If A be its area and M its magnetic moment, the current in the loop will be

- a) M/A b) A/M c) MA d) $A^2 M$

441. An electron is accelerated by a potential difference of 12000 volts. It then enters a uniform magnetic field of 10^{-3} T applied perpendicular to the path of electron. Find the radius of path. Given mass of electron = $9 \times 10^{-31} \text{ kg}$ and charge on electron = $1.6 \times 10^{-19} \text{ C}$

- a) 36.7 m b) 36.7 cm c) 3.67 m d) 3.67 cm


442. A charged particle is moving in a circular orbit of radius 6 cm with a uniform speed of $3 \times 10^6 \text{ m/s}$ under the action of a uniform magnetic field $2 \times 10^{-4} \text{ wb/m}^2$ at right angles to the plane of the orbit. The charge to mass ratio of the particle is

- a) $5 \times 10^9 \text{ C/kg}$ b) $2.5 \times 10^{11} \text{ C/kg}$ c) $5 \times 10^{11} \text{ C/kg}$ d) $5 \times 10^{12} \text{ C/kg}$

443. Which of the following statements is true

- a) The presence of a large magnetic flux through a coil maintains a current in the coil if the circuit is continuous
b) A coil of a metal wire kept stationary in a non-uniform magnetic field has an e.m.f. induced in it
c) A charged particle enters a region of uniform magnetic field at an angle of 85° to the magnetic line of force; the path of the particle is a circle
d) There is no change in the energy of a charged particle moving in a magnetic field although a magnetic force is acting on it


444. A uniform magnetic field B and a uniform electric field E , act in a common region. An electron is entering this region of space. The correct arrangement for it to escape undeviated is



445. An electron is revolving round a proton, producing a magnetic field of 16 weber/m^2 in a circular orbit of radius 1\AA . It's angular velocity will be

- a) 10^{17} rad/sec b) $1/2\pi \times 10^{12} \text{ rad/sec}$ c) $2\pi \times 10^{12} \text{ rad/sec}$ d) $4\pi \times 10^{12} \text{ rad/sec}$

446. A current carrying wire LN is bent in the form shown below. If wire carries a current of 10 A and it is placed in a magnetic field of 5 T which acts perpendicular to the paper outward then it will experience a force



- a) Zero b) 5 N c) 30 N d) 20 N

447. A proton and an α -particle are projected normally into a magnetic field. What will be the ratio of radii of the trajectories of the proton and α -particle?

- a) $2 : 1$ b) $1 : 2$ c) $4 : 1$ d) $1 : 4$

448. A uniform magnetic field B is acting from south to north and is of magnitude 1.5 wb/m^2 . If a proton having mass $= 1.7 \times 10^{-27} \text{ kg}$ and charge $= 1.6 \times 10^{-19} \text{ C}$ moves in this field vertically downward with energy 5 MeV , then the force acting on it will be

- a) $7.4 \times 10^{12} \text{ N}$ b) $7.4 \times 10^{-12} \text{ N}$ c) $7.4 \times 10^{19} \text{ N}$ d) $7.4 \times 10^{-19} \text{ N}$


449. Field inside a solenoid is

- a) Directly proportional to its length b) Directly proportional to current
c) Inversely proportional to total number of turns d) Inversely proportional to current

450. Two long parallel wires carry equal current i flowing in the same direction and are at a distance $2d$ apart. The magnetic field B at a point lying on the perpendicular line joining the wires and at a distance x from the midpoint is

- a) $\frac{\mu_0 i d}{\pi(d^2 + x^2)}$ b) $\frac{\mu_0 i x}{\pi(d^2 - x^2)}$ c) $\frac{\mu_0 i x}{(d^2 + x^2)}$ d) $\frac{\mu_0 i d}{(d^2 - x^2)}$

451. A conducting rod of length l and mass m is moving down a smooth inclined plane of inclination θ with constant speed v . A vertically upward magnetic field \vec{B} exists in space there. The magnitude of magnetic field \vec{B} is



- a) $\frac{mg}{il} \sin \theta$ b) $\frac{mg}{il} \cos \theta$ c) $\frac{mg}{il} \tan \theta$ d) $\frac{mg}{il \sin \theta}$


452. Two parallel wires in free space are 10 cm apart and each carries a current of 10 A in the same direction. The force one wire exerts on the other per metre of length is

- a) $2 \times 10^{-4} \text{ N}$, attractive b) $2 \times 10^{-4} \text{ N}$, repulsive
c) $2 \times 10^{-7} \text{ N}$, attractive d) $2 \times 10^{-7} \text{ N}$, repulsive

453. Two very long straight parallel wires carry current i and $2i$ in opposite directions. The distance between the wires is r . At a certain instant of time a point charge q is at a point equidistant from the two wires in the plane of the wires. Its instantaneous velocity \vec{v} is perpendicular to this plane. The magnitude of the force due to the magnetic field acting on the charge at this instant is

- a) zero b) $\frac{3\mu_0 i q v}{2\pi r}$ c) $\frac{\mu_0 i q v}{\pi r}$ d) $\frac{\mu_0 i q v}{2\pi r}$

454. A thin flexible wire of length L is connected to two adjacent fixed points and carries a current I in the clockwise direction, as shown in the figure. When the system is put in a uniform magnetic field of strength B going into the plane of the paper, the wire takes the shape of a circle. The tension in the wire is



- a) IBL b) $\frac{IBL}{\pi}$ c) $\frac{IBL}{2\pi}$ d) $\frac{IBL}{4\pi}$

455. Two concentric coplanar circular loops of radii r_1 and r_2 carry currents of respectively i_1 and i_2 in opposite directions (one clockwise and the other anticlockwise.) The magnetic induction at the centre of the loops is half that due to i_1 alone at the centre. If $r_2 = 2r_1$. The value of i_2/i_1 is

- a) 2 b) $1/2$ c) $1/4$ d) 1

456. A cyclotron is used to accelerate protons, deuterons α – particle etc. If the energy attained, after acceleration, by the protons is E , the energy attained by α – particles shall be

- a) $4E$ b) $2E$ c) E d) $E/4$


457. A positively charged particle moving due east enters a region of uniform magnetic field directed vertically upwards. The particle will

- a) Get deflected vertically upwards
b) Move in a circular orbit with its speed increased
c) Move in a circular orbit with its speed unchanged
d) Continue to move due east

458. Positively charged particles are projected into a magnetic field. If the direction of the magnetic field is along the direction of motion of the charge particles, the particles get

- a) Accelerated b) Decelerated
c) Deflected d) No change in velocity

459. Figure shows a square loop $ABCD$ with edge length a . The resistance of the wire ABC is r and that of ADC is $2r$. The value of magnetic field at the centre of the loop assuming uniform wire is



- a) $\frac{\sqrt{2}\mu_0 i}{3\pi a} \odot$ b) $\frac{\sqrt{2}\mu_0 i}{3\pi a} \otimes$ c) $\frac{\sqrt{2}\mu_0 i}{\pi a} \odot$ d) $\frac{\sqrt{2}\mu_0 i}{\pi a} \otimes$

460. Two long conductors, separated by a distance d carry currents I_1 and I_2 in the same direction. They exert a force F on each other. Now the current in one of them is increased to two times and its direction is

reversed. The distance is also increased to $3d$. The new value of the force between them is

- a) $-2F$ b) $F/3$ c) $-2F/3$ d) $-F/3$

461. A long straight wire along the z -axis carries a current I in the negative z direction. The magnetic vector field \vec{B} at a point having coordinates (x, y) in the $z = 0$ plane is

- a) $\frac{\mu_0 I (y\hat{i} - x\hat{j})}{2\pi(x^2 + y^2)}$ b) $\frac{\mu_0 I (x\hat{i} + y\hat{j})}{2\pi(x^2 + y^2)}$ c) $\frac{\mu_0 I (x\hat{i} - y\hat{j})}{2\pi(x^2 + y^2)}$ d) $\frac{\mu_0 I (x\hat{i} - y\hat{j})}{2\pi(x^2 + y^2)}$

462. A straight conductor of length l carrying a current I , is bent in the form of a semicircle. The magnetic field (in tesla) at the centre of the semicircle is

- a) $\frac{\pi^2 I}{l} \times 10^{-7}$ b) $\frac{\pi I}{l} \times 10^{-7}$ c) $\frac{\pi I}{l^2} \times 10^{-7}$ d) $\frac{\pi I^2}{l} \times 10^{-7}$

463. A charged particle moving in a uniform magnetic field penetrates layer of lead and there by loss one-half of its kinetic energy. How does the radius of curvature of its path change?

- a) The radius reduces to $r\sqrt{2}$ b) The radius reduces to $\frac{r}{\sqrt{2}}$
c) The radius remains the same d) The radius becomes $r/2$

464. Magnetic effect of current was discovered by

- a) Faraday b) Oersted c) Ampere d) Bohr

465. A current I flows along the length of an infinitely long, straight, thin walled pipe. Then

- a) The magnetic field is zero only on the axis of the pipe
b) The magnetic field is different at different points inside the pipe
c) The magnetic field at any point inside the pipe is zero
d) The magnetic field at all points inside the pipe is the same, but not zero


466. The deflection in a moving coil galvanometer is

- a) Directly proportional to the torsional constant b) Directly proportional to the number of turns in the coil
c) Inversely proportional to the area of the coil d) Inversely proportional to the current flowing

467. Two thin long parallel wires separated by a distance b are carrying currents of i amp each, the magnitude of the force per unit length exerted by one wire over the other is


- a) $\frac{\mu_0 i^2}{b^2}$ b) $\frac{\mu_0 i^2}{2\pi b}$ c) $\frac{\mu_0 i}{2\pi b}$ d) $\frac{\mu_0 i}{2\pi b^2}$

468. A current i_1 carrying wire AB is placed near an another long wire CD carrying current i_2 as shown in figure. If free to move, wire AB will have







- a) Rotational motion only b) Translational motion only
c) Rotational as well as translational motion d) Neither rotational nor translational motion

469. A cell is connected between the points A and C of a circular conductor $ABCD$ with O as centre and angle $AOC = 60^\circ$. If B_1 and B_2 are the magnitudes of the magnetic fields at O due to the currents in ABC and ADC respectively, then ratio $\frac{B_1}{B_2}$ is




- a) 1 b) 2 c) 5 d) 6
470. A beam of well collimated cathode rays travelling with a speed of $5 \times 10^6 \text{ ms}^{-1}$ enter a region of mutually perpendicular electric and magnetic fields and emerge undeviated from this region. If $|B| = 0.02 \text{ T}$, the magnitude of the electric field is
- a) 10^5 Vm^{-1} b) $2.5 \times 10^8 \text{ Vm}^{-1}$ c) $1.25 \times 10^{-10} \text{ Vm}^{-1}$ d) $2 \times 10^3 \text{ Vm}^{-1}$
471. Two infinitely long parallel wires carry equal current in same direction. The magnetic field at a mid point in between the two wires is
- a) Twice the magnetic field produced due to each of the wires
b) Half of the magnetic field produced due to each of the wires
c) Square of the magnetic field produced due to each of the wires
d) Zero
472. A beam of electrons passes undeflected through mutually perpendicular electric and magnetic fields. If the electric field is switched off and the same magnetic field is maintained the electrons move
- a) In an elliptical orbit b) In a circular orbit
c) Along a parabolic path d) Along a straight line
473. An electron ($e = 1.6 \times 10^{-19} \text{ C}$) moves in a circular orbit of radius 1.42 cm with a speed of 10^5 ms^{-1} in presence of magnetic field of $4 \times 10^{-2} \text{ T}$. If the mass of electron is $9.1 \times 10^{-31} \text{ kg}$ the energy gained by the electron in going one round the circular orbit is
- a) zero b) $4.54 \times 10^{-28} \text{ J}$ c) $9.08 \times 10^{-28} \text{ J}$ d) $28.55 \times 10^{-28} \text{ J}$
474. For the magnetic field to be maximum due to a small element of current carrying conductor at a point, the angle between the element and the line joining the element to the given point must be
- a) 0° b) 90° c) 180° d) 45°
475. The magnetic potential due to a magnetic dipole at a point on its axis distant 40 cm from its center is found to be $2.4 \times 10^{-5} \text{ JA}^1 \text{ m}^{-1}$. The magnetic moment of the dipole will be
- a) 28.6 Am^2 b) 32.2 Am^2 c) 38.4 Am^2 d) None of these
476. A charged particle enters a uniform magnetic field with a certain speed at right angles to it. In the magnetic field a change could occur in its
- a) Kinetic energy b) Angular momentum c) Linear momentum d) Speed
477. At a specific instant emission of radioactive compound is deflected in a magnetic field. The compound can emit
- (i) Electrons (ii) Circle
(iii) He^{2+} (iv) Neutrons
- The emission at the instant can be
- a) i, ii, iii b) i, ii, iii, iv c) iv d) ii, iii
478. If current flowing through shell of previous objective is equal to i , then energy density at a point distance $2R$ from axis of the shell varies according to the graph



479. An electron moves in a circular orbit with a uniform speed v . It produces a magnetic field B at the centre of the circle. The radius of the circle is proportional to

- a) $\frac{B}{v}$ b) $\frac{v}{B}$ c) $\sqrt{\frac{v}{B}}$ d) $\sqrt{\frac{B}{v}}$

480. In the following figure a wire bent in the form of a regular polygon of n sides is inscribed in a circle of radius a . Net magnetic field at centre will be



- a) $\frac{\mu_0 i}{2\pi a} \tan \frac{\pi}{n}$ b) $\frac{\mu_0 n i}{2\pi a} \tan \frac{\pi}{n}$ c) $\frac{2 n i}{\pi a} \mu_0 \tan \frac{\pi}{n}$ d) $\frac{n i}{2a} \mu_0 \tan \frac{\pi}{n}$

481. A closely wound solenoid of 2000 turns and area of cross-section $1.5 \times 10^{-4} \text{ m}^2$ carries a current of 2.0 A. It is suspended through its centre and perpendicular to its length, allowing it to turn in a horizontal plane in a uniform magnetic field $5 \times 10^{-2} \text{ tesla}$ making an angle of 30° with the axis of the solenoid. The torque on the solenoid will be

- a) $3 \times 10^{-3} \text{ N.m}$ b) $1.5 \times 10^{-3} \text{ N.m}$ c) $1.5 \times 10^{-2} \text{ N.m}$ d) $3 \times 10^{-2} \text{ N.m}$

482. A current of i ampere flows along an infinitely long straight thin walled tube, then the magnetic induction at any point inside the tube is

- a) Infinite b) zero c) $\frac{\mu_0 2i}{4\pi r} \text{ T}$ d) $\frac{\mu_0 i_0}{2r} \text{ T}$

483. A circular disc of radius 0.2 m is placed in a uniform magnetic field of induction $\frac{1}{\pi} (\text{Wb/m}^2)$ in such a way that its axis makes an angle of 60° with vector \vec{B} . The magnetic flux linked with the disc is

- a) 0.08 Wb b) 0.01 Wb c) 0.02 Wb d) 0.06 Wb


484. An electron enters a magnetic field whose direction is perpendicular to the velocity of the electron. Then

- a) The speed of the electron will increase b) The speed of the electron will decrease
c) The speed of the electron will remain the same d) The velocity of the electron will remain the same

485. A beam of ions with velocity $2 \times 10^5 \text{ m/s}$ enters normally into a uniform magnetic field of $4 \times 10^{-2} \text{ tesla}$. If the specific charge of the ion is $5 \times 10^7 \text{ C/kg}$, then the radius of the circular path described will be

- a) 0.10 m b) 0.16 m c) 0.20 m d) 0.25 m

486. A vertical wire kept in $Z\text{-}X$ plane carries a current from Q to P (see figure). The magnetic field due to current will have the direction at the origin O along



- a) OX b) OX' c) OY d) OY'

487. A charged particle moves with velocity \vec{v} in a uniform magnetic field \vec{B} . The magnetic force experienced by the particle is

- a) Always zero b) Never zero
c) Zero, if \vec{B} and \vec{v} are perpendicular d) Zero, if \vec{B} and \vec{v} are parallel

488. A charged particle moving in a magnetic field experiences a resultant force


- a) In the direction of field
b) In the direction opposite to the field
c) In the direction perpendicular to both the field and its velocity

d) None of the above

489. A long solenoid has a radius a and number of turns per unit length n . If it carries a current i , then the magnetic field on its axis is directly proportional to

- a) ani b) ni c) $\frac{ni}{a}$ d) n^2i

490. Two thin metallic strips, carrying current in the direction shown, cross each other perpendicularly without touching but being close to each other, as shown in the figure. The regions which contain some points of zero magnetic induction are



- a) I and II b) I and III c) I and IV d) II and III

491. An electron is moving with a speed of 10^8 ms^{-1} perpendicular to a uniform magnetic field of induction B . Suddenly induction of magnetic field is reduced to $B/2$. The radius of the path becomes from the original value of r

- a) No change b) Reduces to $r/2$ c) Increases to $2r$ d) Stops moving

492. In the above question, the magnetic induction at O due to the whole length of the conductor is

- a) $\frac{\mu_0 i}{r}$ b) $\frac{\mu_0 i}{2r}$ c) $\frac{\mu_0 i}{4r}$ d) Zero

493. $3A$ of current is flowing in a linear conductor having a length of 40 cm . The conductor is placed in a magnetic field of strength 500 gauss and makes an angle of 30° with direction of the field. It experiences a force of magnitude

- a) $3 \times 10^4 \text{ N}$ b) $3 \times 10^2 \text{ N}$ c) $3 \times 10^{-2} \text{ N}$ d) $3 \times 10^{-4} \text{ N}$

494. A strong magnetic field is applied on a stationary electron, then

- a) The electron moves in the direction of the field
b) The electron moves in an opposite direction
c) The electron remains stationary
d) The electron starts spinning

495. A tangent galvanometer is connected directly to an ideal battery. If the number of turns in the coil is doubled, the deflection will

- a) Increase b) Decrease
c) Remain unchanged d) Either increase or decrease

496. A magnetic field can be produced by

- a) A moving charge b) A changing electric field
c) None of these d) Both of these



497. Three long straight wires are connected parallel to each other across a battery of negligible internal resistance. The ratio of their resistances are $3 : 4 : 5$. What is the ratio of distances of middle wire from the others if the net force experienced by it is zero

- a) $4 : 3$ b) $3 : 1$ c) $5 : 3$ d) $2 : 3$


498. If a long hollow copper pipe carries a direct current, the magnetic field associated with the current will be

- a) Only inside the pipe b) Only outside the pipe
c) Neither inside nor outside the pipe d) Both inside and outside the pipe

499. Graph of force per unit length between two long parallel currents carrying conductor and the distance between them is


- a) Straight line
c) Ellipse
- b) Parabola
d) Rectangular hyperbola
500. A coil carrying electric current is placed in uniform magnetic field, then
a) Torque is formed
b) E.M.f. is induced
c) Both (a) and (b) are correct
d) None of these
501. An electron moving with a uniform velocity along the positive x -direction enters a magnetic field directed along the positive y -direction. The force on the electron is directed along
a) Positive y -direction
b) Negative y -direction
c) Positive z -direction
d) Negative z -direction
502. An element $d\vec{l} = dx\hat{i}$ (where $dx = 1$ cm) is placed at the origin and carries a large current $i = 10$ A. What is the magnetic field on the y -axis at a distance f 0.5 m?
a) $2 \times 10^{-8}\hat{k}$ T
b) $4 \times 10^{-8}\hat{k}$ T
c) $-2 \times 10^{-8}\hat{k}$ T
d) $-4 \times 10^{-8}\hat{k}$ T
503. A small coil of N turns has an effective area A and carries a current I . It is suspended in a horizontal magnetic field \vec{B} such that its plane is perpendicular to \vec{B} . The work done in rotating it by 180° about the vertical axis is
a) $NAIB$
b) $2NAIB$
c) $2\pi NAIB$
d) $4\pi NAIB$
504. Potential energy of a bar magnet of magnetic moment M placed in a magnetic field of induction B such that it makes an angle θ with the direction of B is
a) $MB \sin \theta$
b) $-MB \cos \theta$
c) $MB(1 - \cos \theta)$
d) $MB(1 + \cos \theta)$
505. A wire along x -axis carries a current 3.5 A. Find the force in newton on a 1 cm section of the wire exerted by a magnetic field $\vec{B} = (0.74\hat{j} + 0.36\hat{k})$ T.
a) $(1.26\hat{k} - 2.59\hat{j})10^{-2}$ N
b) $(-1.26\hat{k} + 2.59\hat{j}) \times 10^{-2}$ N
c) $(-2.59\hat{k} + 1.26\hat{j}) \times 10^{-2}$ N
d) $(2.59\hat{k} - 1.26\hat{j}) \times 10^{-2}$ N
506. In the figure shown the magnetic induction at the centre of the arc due to the current in portion AB will be

a) $\frac{\mu_0 i}{r}$
b) $\frac{\mu_0 i}{2r}$
c) $\frac{\mu_0 i}{4r}$
d) Zero
507. Two parallel long straight conductors are placed at right angle to the meter scale at the 2 cm and 6 cm marks as shown in the figure. If they carry currents i and $3i$ respectively in the same direction, then they will produce zero magnetic field at

a) Zero mark
b) 9 cm mark
c) 3 cm mark
d) 7 cm mark
508. Two magnets have the same length and the same pole strength. But one of the magnets has a small hole at its center. Then
a) Both the equal magnetic moment
b) One with hole has smaller magnetic moment
c) One with hole has large magnetic moment
d) One with hole has loses magnetism through the hole
509. A rectangular coil $20\text{ cm} \times 20\text{ cm}$ has 100 turns and carries a current of 1 A. It is placed in a uniform magnetic field $B = 0.5\text{ T}$ with the direction of magnetic field parallel to the plane of the coil. The magnitude of the torque required to hold this coil in this position is
a) Zero
b) 200 N-m
c) 2 N-m
d) 10 N-m
510. A vertical wire carrying a current in the upward direction is placed in horizontal magnetic field directed towards north. The wire will experience a force directed towards
a) North
b) South
c) East
d) West
511. An elastic circular wire of length l carries a current I . It is placed in a uniform magnetic field \vec{B} (out of

paper) such that its plane is perpendicular to the direction of \vec{B} . The wire will experience



- a) No force b) A stretching force c) A compressive force d) A torque

512. The square loop $ABCD$, carrying a current i , is placed in uniform magnetic field B , as shown. The loop can rotate about the axis XX' . The plane of the loop makes an angle θ ($\theta < 90^\circ$) with the direction of B . Through what angle will the loop rotate by itself before the torque on it becomes zero



- a) θ b) $90^\circ - \theta$ c) $90^\circ + \theta$ d) $180^\circ - \theta$

513. The oscillating frequency of a cyclotron is 10 MHz . If the radius of its Dees is 0.5 m , then kinetic energy of a proton, which is accelerated by the cyclotron is

- a) 10.2 MeV b) 2.55 MeV c) 20.4 MeV d) 5.1 MeV

514. If \vec{m} is magnetic moment and B is the magnetic field, then the torque is given by

- a) $\vec{m} \cdot \vec{B}$ b) $\frac{|\vec{m}|}{|\vec{B}|}$ c) $\vec{m} \times \vec{B}$ d) $|\vec{m}| \cdot |\vec{B}|$

515. An α particle and a proton travel with same velocity in a magnetic field perpendicular to the direction of their velocities. Find the ratio of the radii of their circular path

- a) $4 : 1$ b) $1 : 4$ c) $2 : 1$ d) $1 : 2$

516. A circular coil of 20 turns and radius 10 cm is placed in uniform magnetic field of 0.10 T normal to the plane of the coil. If the current in coil is 5 A , then the torque acting on the coil will be

- a) 31.4 Nm
b) 3.14 Nm
c) 0.314 Nm
d) zero

517. In Nebraska the horizontal component of earth's field is 0.2 G . If a vertical wire carries a current of 30 A upward there. What is the magnitude and direction of the force on 1 m of wire?


($1 \text{ G} = 10^{-4} \text{ T}$)

- a) 6 E to W b) $6 \times 10^{-3} \text{ E to W}$ c) $6 \times 10^{-3} \text{ E to W}$ d) $6 \times 10^{-4} \text{ E to W}$

518. A rectangular loop carrying a current i is placed in a uniform magnetic field B . The area enclosed by the loop is A . If there are n turns in the loop, the torque acting on the loop is given by

- a) $ni \vec{A} \times \vec{B}$ b) $ni \vec{A} \cdot \vec{B}$ c) $\frac{1}{n}(i \vec{A} \times \vec{B})$ d) $\frac{1}{n}(i \vec{A} \cdot \vec{B})$


519. A uniform conducting wire ABC has a mass of 10 g . A current of 2 A flows through it. The wire is kept in a uniform magnetic field $B = 2 \text{ T}$. The acceleration of the wire will be



- a) Zero b) 12 ms^{-2} along y -axis

- c) $1.2 \times 10^{-3} \text{ms}^{-2}$ along y-axis d) $0.6 \times 10^{-3} \text{ms}^{-2}$ along y-axis
520. A charged particle is moving in a uniform magnetic field in a circular path. Radius of circular path is R . When energy of particle is doubled, then new radius will be
 a) $R\sqrt{2}$ b) $R\sqrt{3}$ c) $2R$ d) $3R$
521. A particle of mass m and charge q released from the origin in a region occupied by electric field E and magnetic field B ,
 $B = -B_0 \hat{j}$; $E = E_0 \hat{i}$
 The velocity of the particle will be
 a) $\sqrt{\frac{2qE_0}{m}}$ b) $\sqrt{\frac{qE_0}{m}}$ c) $\sqrt{\frac{qE_0}{2m}}$ d) None of these
522. Imaging that an electron revolves round a circle of radius $5.3 \times 10^{-11} \text{m}$ with a linear velocity of $7.5 \times 10^4 \text{ms}^{-1}$ in a hydrogen atom. The magnetic field produced at the centre of the circle due to the electron is
 a) 43Wbm^{-2} b) $43 \times 10^2 \text{Wbm}^{-2}$ c) 0.43Wbm^{-2} d) $43 \times 10^{-4} \text{Wbm}^{-2}$
523. A charged particle is projected in a plane perpendicular to a uniform magnetic field. The area bounded by the path described by the particle is proportional to
 a) The velocity b) The momentum c) The kinetic energy d) None of these
524. When a charged particle moving with velocity \vec{v} is subjected to a magnetic field of induction \vec{B} , the force on it is non-zero. This implies that
 a) Angles between \vec{v} and \vec{B} can have any value other than zero and 180°
 b) Angle between \vec{v} and \vec{B} is either zero or 180°
 c) Angle between \vec{v} and \vec{B} is necessarily 90°
 d) Angle between \vec{v} and \vec{B} can have any value other than 90°
525. A circular loop and a square loop are formed from the same wire and the same current is passed through them. Find the ratio of their dipole moments.
 a) 4π b) $\frac{4}{\pi}$ c) $\frac{2}{\pi}$ d) 2π
526. A proton, a deuteron and an alpha particle with the same kinetic energy enter a region of uniform magnetic field B at right angles to the field. The ratio of the radii of their circular paths is
 a) $1 : 1 : 1$ b) $1 : \sqrt{2} : \sqrt{2}$ c) $\sqrt{2} : 1 : 1$ d) $\sqrt{2} : \sqrt{2} : 1$
527. A proton with energy of 2 MeV enters a uniform magnetic field of 2.5 T normally. The magnetic force on the proton is
 (Take mass of proton to be $1.6 \times 10^{-27} \text{kg}$)
 a) $3 \times 10^{-12} \text{N}$ b) $8 \times 10^{-10} \text{N}$ c) $8 \times 10^{-12} \text{N}$ d) $2 \times 10^{-10} \text{N}$
528. A current carrying conductor produces
 a) Only electric field b) Only magnetic field
 c) Both electric and magnetic fields d) Neither electric nor magnetic field
529. A current carrying straight wire is kept along the axis of a circular loop carrying a current. The straight wire
 a) Will exert an inward force on the circular loop
 b) Will exert an outward force on the circular loop
 c) Will exert a force on the circular loop parallel to itself
 d) Will not exert any force on the circular loop
530. A deuteron of kinetic energy 50 keV is describing a circular orbit of radius 0.5m, is plane perpendicular to magnetic field \vec{B} . The kinetic energy of a proton that describes a circular orbit of radius 0.5 m in the same plane with the same magnetic field \vec{B} is
 a) 200 keV b) 50 keV c) 100 keV d) 25 keV
531. A current of 10 A is passing through a long wire which has semicircular loop of the radius 20 cm as shown

in the figure. Magnetic field produced at the centre of the loop is



- a) $10 \pi \mu$ tesla b) $5 \pi \mu$ tesla c) $4 \pi \mu$ tesla d) $2 \pi \mu$ tesla
532. The direction of magnetic lines of force produced by passing a direct current in a conductor is given by
 a) Lenz's law b) Fleming's left hand rule
 c) Right hand palm rule d) Maxwell's law
533. A long hollow copper tube carries a current I . Then which of the following will be true?
 a) The magnetic field B will be zero at all points inside the tube
 b) The magnetic field B will be zero only at points on the axis of the tube
 c) The magnetic field B will be maximum at points on the axis of the tube
 d) The magnetic field will be zero at any point outside the tube
534. An arbitrary shaped closed coil is made of a wire of length L and a current I ampere is flowing in it. If the plane of the coil is perpendicular to magnetic field \vec{B} , the force on the coil is
 a) Zero b) IBL c) $2IBL$ d) $\frac{1}{2}IBL$
535. A proton (mass = $1.67 \times 10^{-27} \text{ kg}$ and charge = $1.6 \times 10^{-19} \text{ C}$) enters perpendicular to a magnetic field of intensity 2 weber/m^2 with a velocity $3.4 \times 10^7 \text{ m/sec}$. The acceleration of the proton should be
 a) $6.5 \times 10^{15} \text{ m/sec}^2$ b) $6.5 \times 10^{13} \text{ m/sec}^2$ c) $6.5 \times 10^{11} \text{ m/sec}^2$ d) $6.5 \times 10^9 \text{ m/sec}^2$
536. "On flowing current in a conducting wire the magnetic field produces around it." It is a law of
 a) Lenz b) Ampere c) Ohm d) Maxwell
537. A straight wire carrying a current 10 A is bent into a semicircular arc of radius 5 cm . The magnitude of magnetic field at the centre is
 a) $1.5 \times 10^{-5} \text{ T}$ b) $3.14 \times 10^{-5} \text{ T}$ c) $6.28 \times 10^{-5} \text{ T}$ d) $19.6 \times 10^{-5} \text{ T}$
538. The magnetic field normal to the plane of a wire of n turns and radius r which carries a current i is measured on the axis of the coil at a small distance h from the centre of the coil. This is smaller than the magnetic field at the centre by the fraction
 a) $(2/3)r^2/h^2$ b) $(3/2)r^2/h^2$ c) $(2/3)h^2/r^2$ d) $(3/2)h^2/r^2$
539. An electron is moving on a circular path of radius r with speed v in a transverse magnetic field B . e/m for it will be
 a) $\frac{v}{Br}$ b) $\frac{B}{rv}$ c) Bvr d) $\frac{vr}{B}$
540. A proton moving with a velocity $2.5 \times 10^7 \text{ m/s}$, enters a magnetic field of intensity 2.5 T making an angle 30° with the magnetic field. The force on the proton is
 a) $3 \times 10^{-12} \text{ N}$ b) $5 \times 10^{-12} \text{ N}$ c) $6 \times 10^{-12} \text{ N}$ d) $9 \times 10^{-12} \text{ N}$
541. The magnetic flux density B at a distance r from a long straight rod carrying a steady current varies with r as shown in figure.
- a)

b)

c)

d)
542. A long straight wire carrying a current of 30 A is placed in an external uniform magnetic field of induction $4 \times 10^{-4} \text{ T}$. The magnetic field is acting parallel to the direction of current. The magnitude of the resultant magnetic induction in tesla at a point 2.0 cm away from the wire is ($\mu_0 = 4\pi \times 10^{-7} \text{ H/m}$)
 a) 10^{-4} b) 3×10^{-4} c) 5×10^{-4} d) 6×10^{-4}
543. A $2 \mu\text{C}$ charge moving around a circle with a frequency of $6.25 \times 10^{12} \text{ Hz}$ produces a magnetic field 6.28

tesla at the centre of the circle. The radius of the circle is

- a) 2.25 m b) 0.25 m c) 13.0 m d) 1.25 m

544. A charge particle of mass m and charge q enters a region of uniform magnetic field \mathbf{B} perpendicular of its velocity \mathbf{v} . The particle initially at rest was accelerated by a potential difference V (volts) before it entered the region of magnetic field. What is the diameter of the circular path followed by the charged particle in the region of magnetic field?

- a) $\frac{2}{B} \sqrt{\frac{mV}{q}}$ b) $\frac{2}{B} \sqrt{\frac{2mV}{q}}$ c) $B \sqrt{\frac{2mV}{q}}$ d) $\frac{B}{q} \sqrt{\frac{2mV}{B}}$

545. The magnetic induction at a distance r from the axis of an infinitely straight conductor which carries current i is

- a) $\frac{\mu_0 i}{2\pi r}$ b) $\frac{\mu_0 i}{2r}$ c) ∞ d) Zero

546. A proton and an alpha particle are separately projected in a region where a uniform magnetic field exists. Their initial velocities are perpendicular to direction of magnetic field. If both the particles move around magnetic field in circles of equal radii, the ratio of momentum of proton to alpha particle $\left(\frac{P_p}{P_\alpha}\right)$ is

- a) 1 b) 1/2 c) 2 d) 1/4

547. Two parallel long wires A and B carry currents i_1 and i_2 ($< i_1$). When i_1 and i_2 are in the same direction, the magnetic field at a point mid way between the wires is $10 \mu\text{ T}$. If i_2 is reversed, the field becomes $30 \mu\text{ T}$. The ratio i_1/i_2 is

- a) 1 b) 2 c) 3 d) 4

548. A proton, a deuteron and a α - particle enter a magnetic field perpendicular to field with same velocity. What is the ratio of the radii of circular paths?

- a) 1 : 2 : 2 b) 2 : 1 : 1 c) 1 : 1 : 2 d) 1 : 2 : 1

549. A current i flows in a circular coil of radius r . If the coil is placed in a uniform magnetic field B with its plane parallel to the field, magnitude of the torque that acts on the coil is

- a) Zero b) $2\pi r i B$ c) $\pi r^2 i B$ d) $2\pi r^2 i B$

550. If a current is passed in a spring, it

- a) Gets compressed b) Gets expanded c) Oscillates d) Remains unchanged


551. The radius of a circular loop is r and a current i is flowing in it. The equivalent magnetic moment will be

- a) ir b) $2\pi ir$ c) $i\pi r^2$ d) $\frac{1}{r^2}$

552. A circular coil carrying a certain current produces a magnetic field B_0 at its centre. The coil is now rewound so as to have 3 turns and the same current is passed through it. The new magnetic field at the centre is

- a) $\frac{B_0}{9}$ b) $9B_0$ c) $\frac{B_0}{3}$ d) $3B_0$

553. Current through ABC and $A'B'C'$ is I . What is the magnetic field at P ? $BP = PB' = r$ (Here $C'B'PBC$ are collinear)



- a) $B = \frac{1}{4\pi} \frac{2I}{r}$ b) $B = \frac{\mu_0}{4\pi} \left(\frac{2I}{r} \right)$ c) $B = \frac{\mu_0}{4\pi} \left(\frac{I}{r} \right)$ d) Zero

554. A voltmeter with a resistance $50 \times 10^3 \Omega$ is used to measure voltage in a circuit. To increase its range to 3 times, the additional resistance to be put in series is

- a) $9 \times 10^6 \Omega$ b) $10^5 \Omega$ c) $1.5 \times 10^5 \Omega$ d) $9 \times 10^5 \Omega$

555. A galvanometer has a resistance G and a current i_g flowing in it produces full scale deflection. S_1 is the value of the shunt which converts it into an ammeter of range 0 to i and S_2 is the value of the shunt for the range 0 to $2i$. The ratio $\frac{S_1}{S_2}$ is

- a) $\left(\frac{2i - i_g}{i - i_g} \right)$ b) $\frac{1}{2} \left(\frac{i - i_g}{2i - i_g} \right)$ c) 2 d) 1

556. Energy in a current carrying coil is stored in the form of

- a) Electrical energy b) Magnetic field c) Heat d) None of these


557. Two particles X and Y having equal charges, after being accelerated through the same potential difference, enter a region of uniform magnetic field and describe circular paths of radius R_1 and R_2 respectively. The ratio of mass of X to that of Y is

- a) $(R_1/R_2)^{1/2}$ b) R_2/R_1 c) $(R_1/R_2)^2$ d) R_1/R_2

558. A straight wire of mass 200 g and length 1.5 m carries a current of 2 A. It is suspended in mid-air by a uniform horizontal magnetic field B . The magnitude of B (in tesla) is

- a) 2 b) 1.5 c) 0.55 d) 0.65

559. A wire carrying a current i is placed in a uniform magnetic field in the form of the curve $y = a \sin\left(\frac{\pi x}{L}\right)$, $0 \leq x \leq 2L$. The force acting on the wire is



- a) $\frac{iBL}{\pi}$ b) $iBL\pi$ c) $2iBL$ d) Zero


560. A long hollow copper tube carries a current I . Then which of the following will be true?

- a) The magnetic field B will be zero at all points inside the tube
 b) The magnetic field B will be zero only at points on the axis of the tube
 c) The magnetic field B will be maximum at points on the axis of the tube
 d) The magnetic field will be zero at any point outside the tube

561. Cyclotron is used to accelerate

- a) Electrons b) Neutrons c) Positive ions d) Negative ions
562. When deuterium and helium are subjected to an accelerating field simultaneously then
- a) Both acquire same energy b) Deuterium accelerates faster
- c) Helium accelerates faster d) Neither of them is accelerated

563. A and B are two infinitely long straight parallel conductors. C is another straight conductor of length 1 m kept parallel to A and B as shown in the figure. Then the force experienced by C is.



- a) Towards A equal to $0.6 \times 10^{-5} \text{ N}$ b) Towards B equal to $5.4 \times 10^{-5} \text{ N}$
- c) Towards A equal to $5.4 \times 10^{-5} \text{ N}$ d) Towards B equal to $0.6 \times 10^{-5} \text{ N}$
564. An electron enters a region where electrostatic field is 20 N/C and magnetic field is 5 T . If electron passes undeflected through the region, the velocity of electron will be
- a) 0.25 ms^{-1} b) 2 ms^{-1} c) 4 ms^{-1} d) 8 ms^{-1}
565. A current carrying wire in the neighborhood produces
- a) No field b) Electric field only
- c) Magnetic field only d) Electric and magnetic field
566. Magnetic field intensity at the centre of coil of 50 turns, radius 0.5 m and carrying a current of 2 A is
- a) $0.5 \times 10^{-5} \text{ T}$ b) $1.25 \times 10^{-4} \text{ T}$ c) $3 \times 10^{-5} \text{ T}$ d) $4 \times 10^{-5} \text{ T}$
567. An electric current is passed through a circuit containing two wires of the same material, connected in parallel. If the lengths and radii of the wires are in the ratio of $4/3$ and $2/3$, then the ratio of the currents passing through the wire will be
- a) 3 b) $1/3$ c) $8/9$ d) 2
568. To make the field radial in a moving coil galvanometer
- a) The number of turns in the coil is increased
- b) Magnet is taken in the form of horse-shoe
- c) Poles are cylindrically cut
- d) Coil is wound on aluminium frame
569. When a charged particle enters a uniform magnetic field, its kinetic energy
- a) Remains constant b) Increases c) Decreases d) Becomes zero
570. The force on a charged particle moving with a velocity v in a magnetic field B is not
- a) Perpendicular to both v and B b) Maximum if v is perpendicular to B
- c) Maximum, if v is parallel to B d) Zero if v is parallel to B
571. A particle of charge $-16 \times 10^{-18} \text{ coulomb}$ moving with velocity 10 ms^{-1} along the x -axis enters a region where a magnetic field of induction B is along the y -axis, and an electric field of magnitude 10^4 V/m is along the negative z -axis. If the charged particle continues moving along the x -axis, the magnitude of B is
- a) 10^{-3} Wb/m^2 b) 10^3 Wb/m^2 c) 10^5 Wb/m^2 d) 10^{16} Wb/m^2
572. When the current flowing in a circular coil is doubled and the number of turns of the coil in it is halved, the magnetic field at its centre will become
- a) Four times b) Same c) Half d) Double
573. The magnetic field due to a current carrying circular loop of radius 3 cm at a point on the axis at a distance of 4 cm from the centre is $54 \mu\text{T}$. What will be its value at the centre of the loop?
- a) $250 \mu\text{T}$ b) $150 \mu\text{T}$ c) $125 \mu\text{T}$ d) $75 \mu\text{T}$
574. The dimension of the magnetic field intensity B is
- a) $\text{MLT}^{-2}\text{A}^{-1}$ b) $\text{MT}^{-2}\text{A}^{-1}$ c) $\text{ML}^2\text{TA}^{-2}$ d) $\text{M}^2\text{LT}^{-2}\text{A}^{-1}$


575. If in a circular coil A of radius R , current i is flowing and in another coil B of radius $2R$ a current $2i$ is flowing, then the ratio of the magnetic fields, B_A and B_B produced by them will be

- a) 1 b) 2 c) $\frac{1}{2}$ d) 4

576. Biot-Savart's law may be represented in vector form as

- a) $d\mathbf{B} = \frac{\mu_0}{4\pi} i \frac{d\mathbf{l} \times \mathbf{r}}{r^3}$ b) $d\mathbf{B} = \frac{\mu_0}{4\pi} i d\mathbf{l} \times \mathbf{r}$ c) $d\mathbf{B} = \frac{\mu_0}{4\pi} i \frac{d\mathbf{l} \times \mathbf{r}}{r^2}$ d) $d\mathbf{B} = \frac{\mu_0}{4\pi} i \frac{d\mathbf{l} \times \mathbf{r}}{r}$

577. What will be the resultant magnetic field at origin due to four infinite length wires if each wire produces magnetic field ' B ' at origin




- a) $4B$ b) $\sqrt{2}B$ c) $2\sqrt{2}B$ d) Zero

578. A long solenoid is formed by winding 20 turns/cm. The current necessary to produce a magnetic field of 20 millitesla inside the solenoid will be approximately

$$\left(\frac{\mu_0}{4\pi} = 10^{-7} \text{ tesla} - \text{metre/ampere}\right)$$


- a) 8.0 A b) 4.0 A c) 2.0 A d) 1.0 A

579. Two wires A and B carry currents as shown in figure. The magnetic interactions




- a) Push i_2 away from i_1 b) Pull i_2 closer to i_1
c) Turn i_2 clockwise d) Turn i_2 counterclockwise

580. Which of the following graphs represent variation of magnetic field B with distance r for a straight long wire carrying current?



581. Following figure shows the path of an electron that passes through two regions containing uniform magnetic fields of magnitudes B_1 and B_2 . It's path in each region is a half circle, choose the correct option




- a) B_1 is into the page and it is stronger than B_2
b) B_1 is into the page and it is weaker than B_2
c) B_1 is out of the page and it is weaker than B_2
d) B_1 is out of the page and it is stronger than B_2

582. The radius of the path of an electron moving at a speed of 3×10^7 m/s perpendicular to a magnetic field 5×10^{-4} T is nearly

- a) 15 cm b) 45 cm c) 27 cm d) 34 cm

583. In the given figure, the electron enters into the magnetic field. It deflects in direction



- a) +ve X direction b) -ve X direction c) +ve Y direction d) -ve Y direction

584. An electron, a proton, a deuteron and an alpha particle, each having the same speed are in a region of constant magnetic field perpendicular to the direction of the velocities of the particles. The radius of the circular orbits of these particles are respectively R_e, R_p, R_d and R_α . It follows that

- a) $R_e = R_p$ b) $R_p = R_d$ c) $R_d = R_\alpha$ d) $R_p = R_\alpha$

585. A small cylindrical soft iron piece is kept in a galvanometer so that

- a) A radial uniform magnetic field is produced b) A uniform magnetic field is produced
c) There is a steady deflection of the coil d) All of these


586. A particle is moving in a uniform magnetic field, then

- a) Its momentum changes but total energy remains the same
b) Both momentum and total energy remain the same
c) Both will change
d) Total energy changes but momentum remains the same

587. A large magnet is broken into two pieces so that their lengths are in the 2:1. The pole strengths of the two pieces will have ratio



- a) 2 : 1 b) 1 : 2 c) 4 : 1 d) 1 : 1



588. In the figure shown there are two semicircles of radii r_1 and r_2 in which a current i is flowing. The magnetic induction at the centre O will be



- a) $\frac{\mu_0 i}{r} (r_1 + r_2)$ b) $\frac{\mu_0 i}{4} (r_1 - r_2)$ c) $\frac{\mu_0 i}{4} \left(\frac{r_1 + r_2}{r_1 r_2} \right)$ d) $\frac{\mu_0 i}{4} \left(\frac{r_2 - r_1}{r_1 r_2} \right)$

589. Two long parallel wires are at a distance $2d$ apart. They carry steady equal current flowing out of the plane of the paper as shown. The variation of the magnetic field along the line XX' is given by






590. A voltmeter has resistance of $2000\ \Omega$ and it can measure upto 2V. If we want to increase its range by 8V, then required resistance in series will be
 a) $4000\ \Omega$ b) $6000\ \Omega$ c) $7000\ \Omega$ d) $8000\ \Omega$
591. The direction of magnetic lines of forces close to a straight conductor carrying current will be
 a) Along the length of the conductor b) Radially outward
 c) Circular in a plane perpendicular to the conductor d) Helical
592. The magnetic field at the centre of current carrying coil is
 a) $\frac{\mu_0 ni}{2r}$ b) $\frac{\mu_0 ni}{2\pi r}$ c) $\frac{\mu_0 ni}{4r}$ d) $\mu_0 ni$
593. A microammeter has a resistance of $100\ \Omega$ and full scale range of $50\ \mu\text{A}$. It can be used as a voltmeter of as a higher range ammeter provided a resistance is added to it. Pick the correct range and resistance combinations
 a) 50 V range with $10\ \text{k}\Omega$ resistance in series b) 10 V range with $200\ \text{k}\Omega$ resistance in series
 c) 10 mA range with $1\ \Omega$ resistance in parallel d) 10 mA range with $0.1\ \Omega$ resistance in parallel
594. Two parallel wires of length $9\ \text{m}$ each are separated by a distance $0.15\ \text{m}$. If they carry equal currents in the same direction and exert a total force of $30 \times 10^{-7}\ \text{N}$ on each other, then the value of current must be
 a) 2.5 amp b) 3.5 amp c) 1.5 amp d) 0.5 amp
595. A charge Q is uniformly distributed over the surface of non-conducting disc of radius R . The disc rotates about an axis perpendicular to its plane and passing through its centre with an angular velocity ω . As a result of this rotation a magnetic field of induction B is obtained at the centre of the disc. If we keep both the amount of charge placed on the disc and its angular velocity to be constant and vary the radius of the disc then the variation of the magnetic induction at the centre of the disc will be represented by the figure
 a) b) c) d)
596. If a particle of charge $10^{-12}\ \text{C}$ moving along the x -direction with a velocity of $10^5\ \text{m/s}$ experience a force of $10^{-10}\ \text{N}$ in y -direction due to magnetic field, then the minimum value of magnetic field is
 a) $6.25 \times 10^3\ \text{T}$ in z - direction b) $10^{-15}\ \text{T}$ in z - direction
 c) $6.25 \times 10^{-3}\ \text{T}$ in z - direction d) $10^{-3}\ \text{T}$ in z - direction
597. An electron has a circular path of radius $0.01\ \text{m}$ in a perpendicular magnetic induction $10^{-3}\ \text{T}$. The speed of the electron is nearly
 a) $1.76 \times 10^4\ \text{ms}^{-1}$ b) $1.76 \times 10^6\ \text{ms}^{-1}$ c) $3.52 \times 10^6\ \text{ms}^{-1}$ d) $7.04 \times 10^6\ \text{ms}^{-1}$
598. A proton, a deuteron and an α -particle with the same KE enter a region of uniform magnetic field, moving at right angle to B . What is the ratio of the radius of their circular paths ?
 a) $1 : \sqrt{2} : 1$ b) $1 : \sqrt{2} : \sqrt{2}$ c) $\sqrt{2} : 1 : 1$ d) $\sqrt{2} : \sqrt{2} : 1$
599. A particle carrying a charge equal to 100 times the charge on an electron is rotating per second in a circular path of radius $0.8\ \text{metre}$. The value of the magnetic field produced at the centre will be ($\mu_0 =$ permeability for vacuum)

- a) $\frac{10^{-7}}{\mu_0}$ b) $10^{-17}\mu_0$ c) $10^{-6}\mu_0$ d) $10^{-7}\mu_0$

600. An electron is travelling along the x -direction. It encounters a magnetic field in the y -direction. Its subsequent motion will be

- a) Straight line along the x -direction b) A circle in the xz -plane
c) A circle in the yz -plane d) A circle in the xy -plane

601. A square loop of side ' a ' hangs from an insulating hanger of spring balance. The magnetic field of strength B occurs only at the lower edge. It carries a current I . Find the change in the reading of the spring balance if the direction of current is reversed



- a) IaB b) $2IaB$ c) $\frac{IaB}{2}$ d) $\frac{3}{2}IaB$


602. An electron moving around the nucleus with an angular momentum l has a magnetic moment

- a) $\frac{e}{m}l$ b) $\frac{e}{2m}l$ c) $\frac{2e}{m}l$ d) $\frac{e}{2\pi m}l$

603. A uniform electric field and a uniform magnetic field are produced, pointing in the same direction. If an electron is projected with its velocity pointing in the same direction


- a) The electron will turn to its right
b) The electron will turn to its left
c) The electron velocity will increase in magnitude
d) The electron velocity will decrease in magnitude

604. A part of a long wire carrying a current i is bent into a circle of radius r as shown in figure. The net magnetic field at the centre O of the circular loop is



- a) $\frac{\mu_0 i}{4r}$ b) $\frac{\mu_0 i}{2r}$ c) $\frac{\mu_0 i}{2\pi r}(\pi + 1)$ d) $\frac{\mu_0 i}{2\pi r}(\pi - 1)$

605. In the figure, what is the magnetic field at the point O



- a) $\frac{\mu_0 I}{4\pi r}$ b) $\frac{\mu_0 I}{4\pi r} + \frac{\mu_0 I}{2\pi r}$ c) $\frac{\mu_0 I}{4r} + \frac{\mu_0 I}{4\pi r}$ d) $\frac{\mu_0 I}{4r} + \frac{\mu_0 I}{4\pi r}$

606. An electron having kinetic energy E is moving in a circular orbit of radius R perpendicular to a uniform magnetic field induction \vec{B} . If kinetic energy is doubled and magnetic field induction is tripled, the radius

will become

- a) $R\sqrt{9/4}$ b) $R\sqrt{3/2}$ c) $R\sqrt{2/9}$ d) $R\sqrt{4/3}$

607. Two long parallel wires P and Q are both perpendicular to the plane of the paper with distance 5 m between them. If P and Q carry current of 2.5 amp and 5 amp respectively in the same direction, then the magnetic field at a point half way between the wires is

- a) $\frac{\sqrt{3}\mu_0}{2\pi}$ b) $\frac{\mu_0}{\pi}$ c) $\frac{3\mu_0}{2\pi}$ d) $\frac{\mu_0}{2\pi}$


608. Electrons move at right angles to a magnetic field of $1.5 \times 10^{-2}\text{ tesla}$ with a speed of $6 \times 10^7\text{ m/s}$. If the specific charge of the electron is $1.7 \times 10^{11}\text{ C/kg}$, the radius of the circular path will be

- a) 2.9 cm b) 3.9 cm c) 2.35 cm d) 3 cm

609. A moving coil galvanometer gives full scale deflection, when a current of 0.005 A is passed through its coil. It is converted into a voltmeter reading upto 5 V by using an external resistance of $975\ \Omega$. What is the resistance of the galvanometer coil?

- a) $30\ \Omega$ b) $25\ \Omega$ c) $50\ \Omega$ d) $40\ \Omega$

610. Three long straight wires A , B and C are carrying currents as shown in figure. Then the resultant force on B is directed



- a) perpendicular to the plane of paper and outward
b) perpendicular to the plane of paper and inward
c) towards A
d) towards B

611. A horizontal overhead powerline is at a height of 4 m from the ground and carries a current of 100 A from east to west. The magnetic field directly below it on the ground is

($\mu_0 = 4\pi \times 10^{-7}\text{ TmA}^{-1}$)

- a) $2.5 \times 10^{-7}\text{ T}$, southward b) $5.0 \times 10^{-6}\text{ T}$, northward
c) $5.0 \times 10^{-6}\text{ T}$, southward d) $2.5 \times 10^{-7}\text{ T}$, northward

612. In order to increase the sensitivity of a moving coil galvanometer, one should decrease

- a) The strength of its magnet b) The torsional constant of its suspension
c) The number of turns in its coil d) The area of its coil

613. A current $i\text{ A}$ flows along an infinitely long straight thin walled tube, then the magnetic induction at any point inside the tube is

- a) Infinite b) Zero c) $\frac{\mu_0}{4\pi} \cdot \frac{2i}{r}\text{ T}$ d) $\frac{2i}{r}\text{ T}$

614. A wire of length 2 m carrying a current of 1 A is bent to form a circle, the magnetic moment of the coil is

- a) $2\pi\text{ Am}^2$ b) $\frac{1}{\pi}\text{ Am}^2$ c) $\pi\text{ Am}^2$ d) $\frac{2}{\pi}\text{ Am}^2$

615. A loosely wound helix made of stiff wire is mounted vertically with the lower end just touching a dish of mercury. When a current from a battery is started in the coil through the mercury

- a) The wire oscillates b) The wire continues making contact
c) The wire breaks contact just as current is passed d) The mercury will expand by heating due to passage of current

616. The resultant magnetic moment of neon atom will be

- a) Infinity b) μ_B c) Zero d) $\mu_B/2$

617. Two particles of equal charges after being accelerated through the same potential difference enter a uniform transverse magnetic field and describe circular path of radii R_1 and R_2 respectively. Then the ratio of their masses (M_1/M_2) is

a) $\frac{R_1}{R_2}$

b) $\left(\frac{R_1}{R_2}\right)^2$

c) $\frac{R_2}{R_1}$

d) $\left(\frac{R_2}{R_1}\right)^2$

