

# GPLUS EDUCATION

Date :  
Time :  
Marks :

CHEMISTRY


## CHEMICAL KINETICS

### Single Correct Answer Type


- The activation energies of two reactions are  $E_1$  and  $E_2$  ( $E_1 > E_2$ ). If the temperature of the system is increased from  $T_1$  to  $T_2$ , the rate constant of the reactions changes from  $k_1$  to  $k_1'$  in the first reaction and  $k_2$  and  $k_2'$  in the second reaction. Predict which of the following expression is correct?  
a)  $\frac{k_1'}{k_1} = \frac{k_2'}{k_2}$       b)  $\frac{k_1'}{k_1} > \frac{k_2'}{k_2}$       c)  $\frac{k_1'}{k_1} < \frac{k_2'}{k_2}$       d)  $\frac{k_1'}{k_1} = \frac{k_2'}{k_2} = 1$
- Effective collisions are those in which molecules must:  
a) Have energy equal to or greater than the threshold energy  
b) Have proper orientation  
c) Acquire the energy of activation  
d) All of the above
- Consider the following statements,  
The rate law for the acid catalysed hydrolysis of an ester being given as  $Rate = k[H^+][ester] = k'[ester]$ .  
If the acid concentration is doubled at constant ester concentration  
1. The second order rate constant,  $k$  is doubled.  
2. The pseudo first order rate constant,  $k'$  is double.  
3. The rate of the reaction is doubled.  
Which of the above statements are correct?  
a) 1 and 2      b) 2 and 3      c) 1 and 3      d) 1, 2 and 3
- Half-life of two samples is 0.1 and 0.8 s. Their respective concentration is 400 and 50 respectively. The order of reaction is  
a) 0      b) 2      c) 1      d) 4
- The units of rate of reaction are  
a)  $L mol^{-1} s^{-1}$       b)  $mol L^{-1} s^{-1}$       c)  $mol s^{-1}$       d) None of these
- Units of rate constant of first and zero order reactions in terms of molarity  $M$  unit are respectively  
a)  $s^{-1}, M s^{-1}$       b)  $s^{-1}, M$       c)  $M s^{-1}, s^{-1}$       d)  $M, s^{-1}$
- The half time of a second order reaction is:  
a) Inversely proportional to the square of the initial concentration of the reactants  
b) Inversely proportional to the initial concentration of the reactants  
c) Proportional to the initial concentration of reactants  
d) Independent of the initial concentration of reactants
- $\frac{1}{[A]^2}$  vs times are a straight line. Order of reaction is  
a) First      b) Second      c) Zero      d) Third
- For an endothermic reaction where,  $\Delta H$  represents the enthalpy of the reaction in kJ/mol, the minimum value for the energy of activation will be  
a) Less than  $\Delta H$       b) Zero      c) More than  $\Delta H$       d) Equal to  $\Delta H$
- The unit of rate constant for a zero order reaction  
a)  $L s^{-1}$       b)  $L mol^{-1} s^{-1}$       c)  $mol L^{-1} s^{-1}$       d)  $mol s^{-1}$
- What is the formula to find value of  $t_{1/2}$  for a zero order reaction?

- a)  $\frac{k}{[R]_0}$                       b)  $\frac{2k}{[R]_0}$                       c)  $\frac{[R]_0}{2k}$                       d)  $\frac{0.693}{k}$

12. For the reaction,  $A + B \rightarrow C + D$ . The variation of the concentration of the products is given by the curve:




- a) X                                      b) Y                                      c) Z                                      d) W
13. Acid hydrolysis of sucrose is a
- a) Pseudo first order reaction                      b) Zero order reaction
- c) Second order reaction                              d) Unimolecular reaction
14. For a first order reaction the graph  $\log [A]$  vs  $t$  is given below



$x$  is equal to

- a)  $\frac{0.693}{k}$                                       b)  $\frac{k}{2.303}$                                       c)  $-\frac{k}{2.303}$                                       d)  $\log [A]_0$


15. The rate constant of a first order reaction is  $4 \times 10^{-3} \text{sec}^{-1}$ . At a reactant concentration of  $0.02 \text{ M}$ , the rate of reaction would be:
- a)  $8 \times 10^{-5} \text{ M sec}^{-1}$                       b)  $4 \times 10^{-3} \text{ M sec}^{-1}$                       c)  $2 \times 10^{-1} \text{ M sec}^{-1}$                       d)  $4 \times 10^{-1} \text{ M sec}^{-1}$
16. The rate constant for the reaction,  $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$  is  $3.0 \times 10^{-5} \text{ s}^{-1}$ . If the rate is  $2.4 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$  then the concentration of  $\text{N}_2\text{O}_5$  (in  $\text{mol L}^{-1}$ ) is
- a) 0.04                                      b) 0.8                                      c) 0.07                                      d) 1.4
17. Activation energy of a reaction is:
- a) The energy released during the reaction
- b) The energy evolved when activated complex is formed
- c) Minimum amount of energy needed to overcome the potential barrier of reaction
- d) The energy needed to form one mole of the product
18. The activation energy for a reaction is  $9.0 \text{ Kcal/mol}$ . The increase in the rate constant when its temperature is increased from  $298 \text{ K}$  to  $308 \text{ K}$  is:
- a) 10%                                      b) 100%                                      c) 50%                                      d) 63%
19. The rate of first order reaction,  $A \rightarrow \text{Products}$ , is  $7.5 \times 10^{-4} \text{ mol litre}^{-1} \text{ sec}^{-1}$ . If the concentration of  $A$  is  $0.5 \text{ mol litre}^{-1}$  the rate constant is:
- a)  $3.75 \times 10^{-4} \text{ sec}^{-1}$                       b)  $2.5 \times 10^{-5} \text{ sec}^{-1}$                       c)  $1.5 \times 10^{-3} \text{ sec}^{-1}$                       d)  $8.0 \times 10^{-4} \text{ sec}^{-1}$
20.  $2\text{N}_2\text{O}_5 \rightleftharpoons 4\text{NO}_2 + \text{O}_2$
- For the above reaction which of the following is not correct above rates of reaction?
- a)  $\frac{-d[\text{N}_2\text{O}_5]}{dt} = 2 \frac{d[\text{O}_2]}{dt}$                                       b)  $\frac{-2d[\text{N}_2\text{O}_5]}{dt} = \frac{d[\text{NO}_2]}{dt}$
- c)  $\frac{d[\text{NO}_2]}{dt} = 4 \frac{d[\text{O}_2]}{dt}$                                       d)  $\frac{-2d[\text{N}_2\text{O}_5]}{dt} = 4 \frac{d[\text{NO}_2]}{dt} = \frac{d[\text{O}_2]}{dt}$
21. A substance undergoes first order decomposition. The decomposition follows to parallel first order reactions as:





The percentage distribution of B and C are:

- a) 80% B and 20% C  
 b) 76.83% B and 23.17% C  
 c) 90% B and 10% C  
 d) 60% B and 40% C
22. In Arrhenius plot intercept is equal to  
 a)  $-E_a/R$                       b)  $\ln A$                       c)  $\ln k$                       d)  $\log_{10} a$
23. Half-life period of a first order reaction is 1386 seconds. The specific rate constant of the reaction is:  
 a)  $5.0 \times 10^{-2} \text{ s}^{-1}$               b)  $5.0 \times 10^{-3} \text{ s}^{-1}$               c)  $0.5 \times 10^{-2} \text{ s}^{-1}$               d)  $0.5 \times 10^{-3} \text{ s}^{-1}$
24. On addition of  $\text{AgNO}_3$  to  $\text{NaCl}$ , white ppt. occurs:  
 a) Instantaneously  
 b) With a measurable speed  
 c) Slowly  
 d) None of these
25. Which is correct about zero order reaction?  
 a) Rate of reaction depends on decay constant.  
 b) Rate of reaction is independent of concentration.  
 c) Unit of rate constant is  $\text{conc}^{-1}$   
 d) Unit of rate constant is  $\text{conc}^{-1} \text{ time}^{-1}$
26. The half-life period of a first order reaction is 1 min 40 s. Calculate its rate constant.  
 a)  $6.93 \times 10^{-3} \text{ min}^{-1}$               b)  $6.93 \times 10^{-3} \text{ s}^{-1}$               c)  $6.93 \times 10^{-3} \text{ s}$               d)  $6.93 \times 10^3 \text{ s}$
27. The reaction  $2A + B + C \rightarrow D + E$  is found to be first order in A, second in B and zero order in C. What is the effect on the rate of increasing concentration of A, B and C two times?  
 a) 72 times                      b) 8 times                      c) 24 times                      d) 36 times
28. In a reaction, the threshold energy is equal to:  
 a) Activation energy + normal energy of reactants  
 b) Activation energy - normal energy of reactants  
 c) Activation energy  
 d) Normal energy of reactants
29. Which one is not correct?  
 a) Rate of zero order reaction depends upon initial concentration of reactant  
 b) Rate of zero order reaction does not depend upon initial concentration of reactant  
 c)  $t_{1/2}$  of first order reaction is independent of initial concentration of reaction  
 d)  $t_{1/2}$  of zero order reaction is dependent of initial concentration of reaction
30. A reaction proceeds by first order, 75% of this reaction was completed in 32 min. the time required for 50% completion is  
 a) 8 min                      b) 16 min                      c) 20 min                      d) 24 min
31. The rate of the reaction  $\text{CCl}_3\text{CHO} + \text{NO} \rightarrow \text{CHCl}_3 + \text{NO} + \text{CO}$  is equal to rate  $k[\text{CCl}_3\text{CHO}][\text{NO}]$ . If concentration is expressed in mol/L. The unit of  $k$  is  
 a)  $\text{L mol}^{-1} \text{ s}^{-1}$                       b)  $\text{mol L}^{-1} \text{ s}^{-1}$                       c)  $\text{L}^2 \text{ mol}^{-2} \text{ s}^{-1}$                       d)  $\text{s}^{-1}$
32. Observe the following reaction,  
 $2A + B \rightarrow C$   
 The rate of formation of C is  $2.2 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$ .  
 What is the value of  $-\frac{d[A]}{dt}$  ( $\text{mol L}^{-1} \text{ min}^{-1}$ ) ?

- a)  $2.2 \times 10^{-3}$                       b)  $1.1 \times 10^{-3}$                       c)  $4.4 \times 10^{-3}$                       d)  $5.5 \times 10^{-3}$
33. The unit of rate constant of a third order chemical reaction is  
 a)  $\text{mol}^{-1} \text{L s}^{-1}$     b)  $\text{mol}^{-1} \text{s}^{-2}$   
 c)  $\text{mol L}$     d)  $\text{s}^{-1} \text{mol}^{-2} \text{L}^2$
34.  $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{H}_2\text{O} \xrightarrow{\text{H}^+} \text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH}$  is an example of ..... order.  
 a) Zero    b) Second    c) Third    d) Pseudo first order
35. Collision theory is applicable to  
 a) First order reactions    b) Zero order reactions  
 c) Bimolecular reactions    d) Intra-molecular reactions
36. The efficiency of an enzyme in catalyzing a reaction is due to its capacity  
 a) To form a strong enzyme substrate complex                      b) To decrease the bond energy of all substrate molecules  
 c) To change the shape of the substrate molecule                      d) To lower the activation energy of the reaction
37. The reaction  
 $2A + B \rightarrow 3C + D$   
 Which of the following does not express the reaction rate?  
 a)  $\frac{d[D]}{dt}$     b)  $-\frac{d[A]}{2dt}$     c)  $-\frac{d[C]}{3dt}$     d)  $-\frac{d[B]}{dt}$
38. If  $E_f$  and  $E_r$  are the activation energies of the forward and reverse reactions and the reaction is known to be exothermic then  
 a)  $E_f < E_r$   
 b)  $E_f > E_r$   
 c)  $E_f = E_r$   
 d) No relation can be given between  $E_f$  and  $E_r$  as data are not sufficient
39. Milk turns sour at  $40^\circ\text{C}$  three times as faster as at  $0^\circ\text{C}$ . The energy of activation for souring of milk is:  
 a) 4.693 kcal    b) 2.6 kcal    c) 6.6 kcal    d) None of these
40. Which plots will give the value of activation energy?  
 a)  $K$  vs.  $T$     b)  $1/K$  vs.  $T$     c)  $\ln K$  vs.  $T$     d)  $\ln K$  vs.  $\frac{1}{T}$
41. In a second order reaction when the concentration of both reactant are equal, the reaction is completed in 500 s. How long will it take for the reaction to go to 60% completion?  
 a) 1000 s    b) 300 s    c) 3000 s    d) 2000 s
42. The rate constant ( $K$ ) for the reaction  $2A + B \rightarrow \text{Product}$  was found to be  $2.5 \times 10^{-5} \text{ litre mol}^{-1} \text{sec}^{-1}$  after 15 sec,  $2.60 \times 10^{-5} \text{ litre mol}^{-1} \text{sec}^{-1}$  after 30 sec and  $2.55 \times 10^{-5} \text{ litre mol}^{-1} \text{sec}^{-1}$  after 50 sec. The order of reaction is:  
 a) 2    b) 3    c) Zero    d) 1
43. The differential rate expression for the reaction  $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$  is:  
 a)  $\frac{-d[\text{H}_2]}{dt} = \frac{-d[\text{I}_2]}{dt} = \frac{-d[\text{HI}]}{dt}$   
 b)  $\frac{d[\text{H}_2]}{dt} = \frac{d[\text{I}_2]}{dt} = \frac{d[\text{HI}]}{dt}$   
 c)  $\frac{1}{2} \frac{d[\text{H}_2]}{dt} = \frac{1}{2} \frac{d[\text{I}_2]}{dt} = \frac{d[\text{HI}]}{dt}$   
 d)  $-2 \frac{d[\text{H}_2]}{dt} = -2 \frac{d[\text{I}_2]}{dt} = \frac{d[\text{HI}]}{dt}$
44. For the elementary step,  
 $(\text{CH}_3)_3\text{CBr}(aq) \rightarrow (\text{CH}_3)_3\text{C}^+(aq) + \text{Br}^-(aq)$  the molecularity is:  
 a) Zero    b) 1    c) 2    d) Cannot ascertained
45. A graph plotted between  $\log t_{50\%}$  vs.  $\log a$  concentration is a straight line. What conclusion can you draw from the given graph?



- a)  $n = 1, t_{1/2} = \frac{1}{K \cdot a}$       b)  $n = 2, t_{1/2} = 1/a$       c)  $n = 1, t_{1/2} = \frac{0.693}{K}$       d) None of the above
46. If  $a$  is the initial concentration then time required to decompose half of the substance for  $n$ th order is inversely proportional to:  
 a)  $a^n$       b)  $a^{n-1}$       c)  $a^{1-n}$       d)  $a^{n-2}$
47. The hydrolysis of ethyl acetate,  
 $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{H}_2\text{O} \xrightarrow{\text{H}^+} \text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH}$  is:  
 a) First order      b) Second order      c) Third order      d) Zero order
48. The rate law for a reaction between the substances A and B is given by  
 rate =  $k[A]^n[B]^m$ . On doubling the concentration of A and halving the concentration of B, the ratio of the new rate to the earlier rate of the reaction will be as  
 a)  $\frac{1}{2^{m+n}}$       b)  $(m+n)$       c)  $(n-m)$       d)  $2^{(n-m)}$
49. For the reaction  
 $\text{H}_2(\text{g}) + \text{Br}_2(\text{g}) \rightarrow 2\text{HBr}(\text{g})$   
 The experimental data suggest  
 rate =  $k[\text{H}_2][\text{Br}_2]^{1/2}$   
 the molecularity and order of the reaction are respectively  
 a)  $1, \frac{1}{2}$       b)  $1, 1$       c)  $\frac{3}{2}, \frac{3}{2}$       d)  $2, \frac{3}{2}$
50. The rate of reaction increases with temperature due to  
 a) Decrease in activation energy      b) Increase in activation energy  
 c) Increase in collision frequency      d) Increase in concentration
51. In a first order reaction, the concentration of the reactant is decreased from  $1.0 \text{ M}$  to  $0.25 \text{ M}$  in 20 minute. The rate constant of the reaction would be:  
 a)  $10 \text{ min}^{-1}$       b)  $6.931 \text{ min}^{-1}$       c)  $0.6931 \text{ min}^{-1}$       d)  $0.06931 \text{ min}^{-1}$
52. The reaction obey I order with respect to  $\text{H}_2$  and  $\text{ICl}$  both  
 $\text{H}_2(\text{g}) + 2\text{ICl}(\text{g}) \rightarrow 2\text{HCl}(\text{g}) + \text{I}_2(\text{g})$   
 Which of the following mechanism is in consistent with the given fact?  
 Mechanism A:  $\text{H}_2(\text{g}) + 2\text{ICl} \rightarrow 2\text{HCl}(\text{g}) + \text{I}_2(\text{g})$   
 Mechanism B: (i)  $\text{H}_2(\text{g}) + \text{ICl}(\text{g}) \xrightarrow{\text{slow}} \text{HCl}(\text{g}) + \text{HI}(\text{g})$   
 (ii)  $\text{HI}(\text{g}) + \text{ICl}(\text{g}) \rightarrow \text{HCl}(\text{g}) + \text{I}_2$   
 a) A and B both      b) Neither A nor B      c) A only      d) B only
53. Two reactions  $A \rightarrow$  products and  $B \rightarrow$  products have rate constants  $K_A$  and  $K_B$  at temperature  $T$  and activation energies  $E_A$  and  $E_B$  respectively. If  $K_A > K_B$  and  $E_A < E_B$  and assuming that  $A$  for both the reactions is same, then:  
 a) At higher temperatures  $K_A$  will be greater than  $K_B$   
 b) At lower temperature  $K_A$  and  $K_B$  will differ more and  $K_A > K_B$   
 c) As temperature rises  $K_A$  and  $K_B$  will be close to each other in magnitude  
 d) All of the above
54. The half life for a reaction ... of temperature.  
 a) Independent  
 b) Increased with increase  
 c) Decreased with increase  
 d) Dependent
55. The following mechanism has been proposed for the reaction of  $\text{NO}$  with  $\text{Br}_2$  to form  $\text{NOBr}$




If the second step is the rate determining step, the order of the reaction with respect to NO(g) is


- a) 1                                      b) 0                                      c) 3                                      d) 2
56. The unit and value of rate constant and that of rate of reaction are same for  
a) Zero order                            b) First order                            c) Second order                            d) Third order
57. According to collision theory of reaction rates:  
a) Every collision between reactants leads to chemical reaction  
b) Rate of reaction is proportional to velocity of molecules  
c) All reactions which occur in gaseous phase are zero order reactions  
d) Rate of reaction is directly proportional to collision frequency
58. Half-life of a reaction is found to be inversely proportional to the cube of initial concentration. The order of reaction is  
a) 4                                      b) 3                                      c) 5                                      d) 2
59. A reaction involving two different reactants can never be  
a) Bimolecular reaction                                      b) Second order reaction  
c) First order reaction                                      d) Unimolecular reaction
60. For the non-equilibrium process,  $A + B \rightarrow \text{Products}$ , the rate is first order with respect to  $A$  and second order respect to  $B$ . If 1.0 mole each of  $A$  and  $B$  are introduced into a 1 litre vessel and the initial rate was  $1.0 \times 10^{-2}$  mol/litre-sec. The rate (in mol litre<sup>-1</sup> sec<sup>-1</sup>) when half of the reactants have been used:  
a)  $1.2 \times 10^{-3}$                             b)  $1.2 \times 10^{-2}$                             c)  $2.5 \times 10^{-4}$                             d) None of these
61. The activation energy of a reaction is zero. The rate constant for the reaction  
a) Decreases with decrease of temp                                      b) Increases with increase of temp  
c) Decreases with increase of temp                                      d) Is nearly independent of temp
62. The burning of coal represented by the equation;  $\text{C(s)} + \text{O}_2\text{(g)} \rightarrow \text{CO}_2\text{(g)}$ . The rate of this reaction is increased by:  
a) Decrease in the concentration of oxygen  
b) Powdering the lumps of coal  
c) Decreasing the temperature  
d) Providing inert atmosphere for burning
63. At room temperature, the reaction between NO and O<sub>2</sub> to give NO<sub>2</sub> is fast, while that between CO and O<sub>2</sub> is slow. It is due to:  
a) CO is smaller in size than that of NO  
b) CO is poisonous  
The activation energy for the reaction,  
c)  $2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$  is less than  
 $2\text{CO} + \text{O}_2 \rightarrow 2\text{CO}_2$   
d) None of the above
64. The rate of first order reaction is  $1.5 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$  at 0.5 M concentration of the reactant. The half-life of reaction is  
a) 0.383 min                            b) 23.1 min                            c) 8.73 min                            d) 7.53 min
65. The rate constant of a first order reaction at 27°C is  $10^{-3} \text{ min}^{-1}$ . The temperature coefficient of this reaction is 2. What is the rate constant (in  $\text{min}^{-1}$ ) at 17°C for this reaction?  
a)  $10^{-3}$                                       b)  $5 \times 10^{-4}$                                       c)  $2 \times 10^{-3}$                                       d)  $10^{-2}$
66. The minimum energy required for the reacting molecules to undergo reaction is  
a) Potential energy                                      b) Kinetic energy  
c) Thermal energy                                      d) Activation energy
67. The decomposition of  $\text{N}_2\text{O}_5$  occur as  $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ , and follows 1st order kinetics, hence  
a) The reaction is unimolecular                                      b) The reaction is bimolecular  
c)  $t_{1/2} \propto a^0$                                       d) None of the above

68. The rate of a chemical reaction doubles for every 10°C rise of temperature. If the temperature is raised by 50°C, the rate of the reaction increases by about  
 a) 10 times                      b) 24 times                      c) 32 times                      d) 64 times
69. Which of the following statement is incorrect about the molecularity of a reaction?  
 a) Molecularity of a reaction is the number of molecules of the reactants presents in the balanced equation  
 b) Molecularity of a reaction is the number of molecules in the slowest step  
 c) Molecularity is always a whole number  
 d) There is no difference between order and molecularity of a reaction
70. For a reaction  $A + B \rightarrow \text{Products}$ , the rate of the reaction was doubled when the concentration of  $A$  was doubled. When the concentration of  $A$  and  $B$  were doubled, the rate was again doubled, the order of the reaction with respect to  $A$  and  $B$  are:  
 a) 1, 1                      b) 2, 0                      c) 1, 0                      d) 0, 1

71. An exothermic chemical reaction occurs in two steps as follows



The progress of the reaction can be best represented by



d) All are correct

72. According to the Arrhenius equation a straight line is to be obtained by plotting the logarithm of the rate constant of a chemical reaction ( $\log k$ ) against

- a)  $T$                       b)  $\log T$                       c)  $\frac{1}{T}$                       d)  $\log \frac{1}{T}$

73. The rate constant is numerically the same for three reactions of first, second and third order respectively. Which one is true for rate of three reaction?

- a)  $r_1 = r_2 = r_3$                       b)  $r_1 > r_2 > r_3$                       c)  $r_1 < r_2 < r_3$                       d) All of these

74. Mathematical expression for  $t_{1/4}$  i.e., when (1/4)th reaction is over following first order kinetics can be given by

- a)  $t_{1/2} = \frac{2.303}{k} \log 4$                       b)  $t_{1/2} = \frac{2.303}{k} \log 2$                       c)  $t_{1/2} = \frac{2.303}{k} \log \frac{4}{3}$                       d)  $t_{1/2} = \frac{2.303}{k} \log \frac{3}{4}$

75. The rate of reaction:

$2\text{NO} + \text{Cl}_2 \rightarrow 2\text{NOCl}$  is given by the rate, equation  $\text{rate} = k[\text{NO}]^2[\text{Cl}_2]$ . The value of the rate constant can be increased by:

- a) Increasing the temperature  
 b) Increasing the concentration of NO  
 c) Increasing the concentration of the  $\text{Cl}_2$   
 d) Doing all of these

76. A reaction was observed for 15 days and the percentage of the reactant remaining after the days indicated was recorded in the following table.

Time (days)	% Reactant remaining
0	100
2	50
4	39
6	25
8	21
10	18
12	15


14	12.5
15	10

Which one of following best describes the order and the half-life of the reaction?

**Reaction order      Half-life (days)**

- |           |   |          |   |
|-----------|---|----------|---|
| a) First  | 2 | b) First | 6 |
| c) Second | 2 | d) Zero  | 6 |

77. In the reaction



The rate of appearance of bromine ( $\text{Br}_2$ ) is related to rate of disappearance of bromide ions as following:

- a)  $\frac{d(\text{Br}_2)}{dt} = \frac{3}{5} \frac{d(\text{Br}^-)}{dt}$       b)  $\frac{d(\text{Br}_2)}{dt} = -\frac{3}{5} \frac{d(\text{Br}^-)}{dt}$       c)  $\frac{d(\text{Br}_2)}{dt} = -\frac{5}{3} \frac{d(\text{Br}^-)}{dt}$       d)  $\frac{d(\text{Br}_2)}{dt} = \frac{5}{3} \frac{d(\text{Br}^-)}{dt}$

78. Which one of the following is a second order reaction?

- |   |   |
|---|---|
| a) $\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$                   | b) $\text{NH}_4\text{NO}_3 \rightarrow \text{N}_2 + 3\text{H}_2\text{O}$                              |
| c) $\text{H}_2 + \text{Cl}_2 \xrightarrow{\text{Sunlight}} 2\text{HCl}$ | d) $\text{CH}_3\text{COOCH}_3 + \text{NaOH} \rightarrow \text{CH}_3\text{COONa} + \text{H}_2\text{O}$ |

79. The temperature coefficient of most of the reactions lies between

- |            |            |            |            |
|------------|------------|------------|------------|
| a) 1 and 3 | b) 2 and 3 | c) 1 and 4 | d) 2 and 4 |
|------------|------------|------------|------------|

80. In respect of the equation  $k = Ae^{-E_a/RT}$  in chemical kinetics, which one of the statement is correct?

- |                                  |                                |
|----------------------------------|--------------------------------|
| a) $R$ is Rydberg constant       | b) $K$ is equilibrium constant |
| c) $E_a$ is energy of activation | d) $A$ is adsorption factor    |

81. The rate of chemical reaction (except zero order):

- a) Decreases from moment to moment  
 b) Remains constant throughout  
 c) Is independent of the order of reaction  
 d) None of the above

82. For a zero order reaction

- |                          |                            |                            |                              |
|--------------------------|----------------------------|----------------------------|------------------------------|
| a) $t_{1/2} \propto R_0$ | b) $t_{1/2} \propto 1/R_0$ | c) $t_{1/2} \propto R_0^2$ | d) $t_{1/2} \propto 1/R_0^2$ |
|--------------------------|----------------------------|----------------------------|------------------------------|

83. Effect of temperature on reaction rate is given by

- |                               |                        |
|-------------------------------|------------------------|
| a) Claisen-Clapeyron equation | b) Arrhenius equation  |
| c) Gibbs Helmholtz equation   | d) Kirchoff's equation |

84. The Arrhenius equation expressing the effect of temperature on the rate constant of reaction is:

- |                         |                       |                                |                      |
|-------------------------|-----------------------|--------------------------------|----------------------|
| a) $K = \frac{E_a}{RT}$ | b) $K = Ae^{-E_a/RT}$ | c) $K = \log_e \frac{E_a}{RT}$ | d) $K = e^{-E_a/RT}$ |
|-------------------------|-----------------------|--------------------------------|----------------------|

85. Find the two third life ( $t_{2/3}$ ) of a first order reaction in which  $k = 5.48 \times 10^{-14}$  per second

- |                                   |                                    |                                   |                                     |
|-----------------------------------|------------------------------------|-----------------------------------|-------------------------------------|
| a) $201 \times 10^{13} \text{ s}$ | b) $2.01 \times 10^{13} \text{ s}$ | c) $201 \times 10^{20} \text{ s}$ | d) $0.201 \times 10^{10} \text{ s}$ |
|-----------------------------------|------------------------------------|-----------------------------------|-------------------------------------|

86.  $A + B \rightarrow \text{Product}$

If concentration of A is doubled, rate increases 4 times. If concentration of A and B are doubled, rate increases 8 times. The differential rate equation of the reaction will be

- |                                      |  |  |  |
|--------------------------------------|--|--|--|
| a) $\frac{dC}{dt} = kC_A \times C_B$ | b) $\frac{dC}{dt} = kC_A^2 \times C_B^3$ | c) $\frac{dC}{dt} = kC_A^2 \times C_B$ | d) $\frac{dC}{dt} = kC_A^2 \times C_B^2$ |
|--------------------------------------|--|--|--|


87. For the reaction  $A \rightarrow B$ , the rate expression is  $r = k[A]^n$ . When the concentration of A is doubled, the rate of reaction is quadrupled. The value of  $n$  is

- |      |         |      |      |
|------|---------|------|------|
| a) 1 | b) Zero | c) 3 | d) 2 |
|------|---------|------|------|

88. The rate constant for the first order reaction is  $60 \text{ s}^{-1}$ . How much time will it take to reduce the concentration of the reaction to  $1/16 \text{ M}$  value ?

- |                                   |                                |                                |                                   |
|-----------------------------------|--------------------------------|--------------------------------|-----------------------------------|
| a) $4.6 \times 10^{-2} \text{ s}$ | b) $4.6 \times 10^4 \text{ s}$ | c) $4.6 \times 10^2 \text{ s}$ | d) $4.6 \times 10^{-4} \text{ s}$ |
|-----------------------------------|--------------------------------|--------------------------------|-----------------------------------|

89. In the reaction,




- |            |            |            |            |
|------------|------------|------------|------------|
| a) 490 atm | b) 250 atm | c) 480 atm | d) 420 atm |
|------------|------------|------------|------------|


90. For a chemical reaction, ..... can never to a fraction



- a) Order                                      b) Half life                                      c) Rate constant                                      d) Molecularity
91. The time taken for the completion of  $3/4$  of a first order reaction is  
 a)  $(2.303/k) \log 3/4$                       b)  $(2.303/k) \log 4$                       c)  $(2.303/k) \log 1/4$                       d)  $(2.303/0.75) \log k$
92.  $2N_2O_5(g) \rightarrow 4NO_2(g) + O_2(g)$   
 What is the ratio of the rate of decomposition of  $N_2O_5$  to rate of formation of  $O_2$  ?  
 a) 1:2                                      b) 2:1                                      c) 1:4                                      d) 4:1
93. A first order reaction is 75% complete after 32 min. when was 50% of the reaction completed?  
 a) 16 min                                      b) 8 min                                      c) 4 min                                      d) 32 min
94. For a reaction,  $A + 2B \rightarrow C$ , rate is given by  $+\frac{d[C]}{dt} = k[A][B]$ , hence, the order of the reaction is  
 a) 3                                      b) 2                                      c) 1                                      d) 0
95. The accompanying figure depicts the change in concentration of species X and Y for the reaction  $X \rightarrow Y$ , as a function of time. The point of intersection of the two curves represents:



- a)  $t_{1/2}$   
 b)  $t_{3/4}$   
 c)  $t_{2/3}$   
 d) Data is insufficient to predict
96. The rate constant of a reaction at temperature 200 K is 10 times less than the rate constant at 400 K. What is the activation energy ( $E_a$ ) of the reaction?  
 a)  $1842.4 R$                                       b)  $921.2 R$                                       c)  $460.6 R$                                       d)  $230.3 R$
97. A zero order reaction is one:  
 a) In which reactants do not react  
 b) In which one of the reactants is in large excess  
 c) Whose rate does not change with time  
 d) Whose rate increases with time
98. In a first order reaction the  $a/(a - x)$  was found to be 8 after 10 minute. The rate constant is:  
 a)  $(2.303 \times 3 \log 2)/10$                       b)  $(2.303 \times 2 \log 3)/10$                       c)  $10 \times 2.303 \times 2 \log 3$                       d)  $10 \times 2.303 \times 3 \log 2$
99. If the rate of reaction  $A \rightarrow B$  doubles on increasing the concentration of A by 4 times, the order of the reaction is  
 a) 2                                      b) 1                                      c)  $\frac{1}{2}$                                       d) 4
100. The rate of chemical reaction  
 a) Increase as the reaction proceeds                                      b) Decrease the reaction proceeds  
 c) May increase or decrease during reaction                                      d) Remains constant as the reaction proceeds
101. For zero order reaction, the integrated rate equation is  
 a)  $kt = \frac{[A]}{[A]_0}$                                       b)  $kt = [A] - [A]_0$                                       c)  $[A] = -kt + [A]_0$                                       d)  $[A] = kt - [A]_0$
102. The half-life period of a first order reaction is 69.3 s. what is the rate constant?  
 a)  $0.01s^{-1}$                                       b)  $0.1s^{-1}$                                       c)  $1s^{-1}$                                       d)  $10s^{-1}$
103. A reaction has a rate constant of  $0.5 \text{ mol}^{-1} \text{ dm}^3 \text{ min}^{-1}$ . If initial concentration of the reactant is  $0.2 \text{ mol dm}^{-3}$ , half-life of the reaction  
 a) 1.4 min                                      b) 10 min                                      c) 15 min                                      d) 20 min
104. The bromination of acetone that occurs in acid solution is represented by this equation.



These kinetic data were obtained for given reaction concentrations.

Initial concentrations, $M$			
$[\text{CH}_3\text{COCH}_3]$	$[\text{Br}_2]$	$[\text{H}^+]$	Initial rate, disappearance of $\text{Br}_2, M s^{-1}$
0.30	0.05	0.05	$5.7 \times 10^{-5}$
0.30	0.10	0.05	$5.7 \times 10^{-5}$
0.30	0.10	0.10	$1.2 \times 10^{-4}$
0.40	0.05	0.20	$3.1 \times 10^{-4}$

Based on these data, the rate equation is:

- a) rate =  $k[\text{CH}_3\text{COCH}_3][\text{Br}_2]$
- b) rate =  $k[\text{CH}_3\text{COCH}_3][\text{Br}_2][\text{H}^+]^2$
- c) rate =  $k[\text{CH}_3\text{COCH}_3][\text{Br}_2][\text{H}^+]$
- d) rate =  $k[\text{CH}_3\text{COCH}_3][\text{H}^+]$

105. The rate constant for a chemical reaction has units  $L mol^{-1} s^{-1}$ , order of the reaction will be

- a) 0
- b) 1
- c) 2
- d) 3

106. Activation energy of a chemical reaction can be determined by

- a) Evaluating rate constant at standard temperatures
- b) Evaluating velocities of reaction at two different temperatures
- c) Evaluating rate constants at two different temperatures
- d) Changing concentration of reactants

107. Which statement about molecularity of a reaction is wrong?

- a) It is the number of molecules of the reactants taking part in a single step of reaction
- b) It is calculated from the reaction mechanism
- c) It may be either whole number or fractional
- d) None of the above

108. Arrhenius equation may not be represented as

- a)  $\ln \frac{A}{k} = \frac{E_a}{RT}$
- b)  $\frac{d \ln k}{dT} = \frac{E}{RT^2}$
- c)  $\log A = \log k + \frac{E_a}{2.303 RT}$
- d)  $\log \left[ -\frac{E_a}{RT} \right] = \frac{k}{A}$


109. The reaction  $2\text{SO}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{SO}_3(g)$  is carried out in a  $1 dm^3$  vessel and  $2 dm^3$  vessel separately. The ratio of the reaction velocities will be

- a) 1:8
- b) 1:4
- c) 4:1
- d) 8:1

110. The rate for the reaction,  $\text{RCl} + \text{NaOH}(aq) \rightarrow \text{ROH} + \text{NaCl}$  is given by rate =  $k[\text{RCl}]$ , the freezing point of the reaction is

- a) Unaffected by increasing the temperature of the reaction
- b) Decreased on increasing the temperature of the reaction
- c) Halved on reducing the concentration of RCl to half
- d) Doubled on doubling the concentration of NaOH

111. In the sequence of reaction,



- a)  $A \rightarrow B$
- b)  $B \rightarrow C$
- c)  $C \rightarrow D$
- d)  $A \rightarrow D$

112. A first order reaction is 20% complete in 10 min. What is the rate constant of the reaction?

- a) 0.223
- b) 0.0223
- c) 0.322
- d) 0.0322

113. The activation energy of exothermic reaction  $A \rightarrow B$  is  $80 \text{ kJ mol}^{-1}$ . The heat of reaction is  $200 \text{ kJ mol}^{-1}$ . The activation energy for the reaction  $B \rightarrow A$  (in  $\text{kJ mol}^{-1}$ ) will be

- a) 80
- b) 120
- c) 40
- d) 280

114. An endothermic reaction  $A \rightarrow B$  has an activation energy of 15 kcal/mol and the energy of reaction is 5 kcal/mol. The activation energy for the reaction  $B \rightarrow A$  is  
 a) 20 kcal/mol                      b) 15 kcal/mol                      c) 10 kcal/mol                      d) Zero
115.  $K$  for a zero order reaction is  $2 \times 10^{-2} \text{ mol L}^{-1} \text{ sec}^{-1}$ . If the concentration of the reactant after 25 sec is 0.5 M, the initial concentration must have been:  
 a) 0.5 M                                  b) 1.25 M                                  c) 12.5 M                                  d) 1.0 M
116. Rate constant for a reaction is  $10^{-3} \text{ s}^{-1}$ . The to leave 25% reaction is  
 a) 693 s                                      b) 1386 s                                      c) 6930 s                                      d) 2029 s
117. By increase in temperature by 10 K, the rate of reaction becomes double. How many times the rate of reaction will be if the temperature is increased from 303K to 353 K?  
 a) 4    b) 8    c) 16    d) 32
118. Temperature coefficient of a reaction is 2. When temperature is increased from 30°C to 100°C, rate of the reaction increases by  
 a) 128 times                                  b) 100 times                                  c) 500 times                                  d) 250 times
119. The activation energy of a reaction is 9 kcal/mol. The increase in the rate constant when its temperature is raised from 295 to 300 K is approximately  
 a) 10%    b) 50%    c) 100%    d) 28%
120. For a reaction  $\frac{1}{2}A \rightarrow 2B$ , rate of disappearance of 'A' is related to the rate of appearance of B by the expression  
 a)  $-\frac{d[A]}{dt} = \frac{1}{2} \frac{d[B]}{dt}$                       b)  $-\frac{d[A]}{dt} = \frac{1}{4} \frac{d[B]}{dt}$                       c)  $-\frac{d[A]}{dt} = \frac{d[B]}{dt}$                       d)  $-\frac{d[A]}{dt} = 4 \frac{d[B]}{dt}$
121. For a first order reaction, ( $A \rightarrow$  products), the concentration of A changes from 0.1 M to 0.025 M in 40 minutes. The rate of reaction when the concentration of A is 0.01 M, is:  
 a)  $3.47 \times 10^{-4} \text{ M/min}$                       b)  $3.47 \times 10^{-5} \text{ M/min}$                       c)  $1.73 \times 10^{-4} \text{ M/min}$                       d)  $1.73 \times 10^{-5} \text{ M/min}$
122. In a 1<sup>st</sup> order reaction, reactant concentration C varies with time t as  
 a) 1/C increases linearly with t                      b) Log C decreases linearly with t  
 c) C decreases with 1/t                                  d) Log C decreases with 1/t
123. The rate constant of a zero order reaction is  $0.2 \text{ mol dm}^{-3} \text{ h}^{-1}$ . If the concentration of the reactant after 30 min is  $0.05 \text{ mol dm}^{-3}$ . Then its initial concentration would be  
 a)  $0.01 \text{ mol dm}^{-3}$                                   b)  $0.15 \text{ mol dm}^{-3}$                                   c)  $0.25 \text{ mol dm}^{-3}$                                   d)  $4.00 \text{ mol dm}^{-3}$
124. For a chemical reaction  $2X + Y \rightarrow Z$ , the rate of appearance of Z is  $0.05 \text{ mol L}^{-1}$ . The rate of disappearance of X will be  
 a)  $0.05 \text{ mol L}^{-1} \text{ h}^{-1}$                                   b)  $0.05 \text{ mol L}^{-1} \text{ min}^{-1}$                                   c)  $0.1 \text{ mol L}^{-1} \text{ min}^{-1}$                                   d)  $0.25 \text{ mol L}^{-1} \text{ min}^{-1}$
125. For a reaction,  $x(g) \rightarrow y(g) + z(g)$  the half-life period is 10 min. in what period of time would the concentration of X be reduce to 10% of original concentration?  
 a) 20 min    b) 33 min    c) 15 min    d) 25 min
126. When is the activation energy for the decomposition of  $\text{N}_2\text{O}_5$  as,  

$$\text{N}_2\text{O}_5 \rightleftharpoons 2\text{NO}_2 + \frac{1}{2}\text{O}_2$$
 If the values of rate constant =  $3.45 \times 10^{-5}$  at 27°C and rate constant =  $6.9 \times 10^{-3}$  at 67°C?  
 a) 112.5 kJ    b) 200 kJ    c) 149.5 kJ    d) 11.25 kJ
127. Rate equation for a second order reaction is:  
 a)  $K = \frac{2.303}{t} \log \frac{a}{(a-x)}$                       b)  $K = \frac{1}{t} \log \frac{a}{a(a-x)}$                       c)  $K = \frac{1}{t} \cdot \frac{x}{a(a-x)}$                       d)  $K = \frac{1}{t^2} \cdot \frac{a}{(a-x)}$
128. Which of the following statements are correct?  
 1. Order of a reaction can be known from experimental result and not from the stoichiometry of reaction  
 2. Overall molecularity of a reaction may be determined in a manner similar to overall order of reaction  
 3. Overall order of reaction,  
 $A^m + B^n \rightarrow AB_x$  is  $(m + n)$   
 4. Molecularity of a reaction refers to

(i) Molecularity of each of the elementary steps (slow steps) in a multistep reaction

(ii) Molecularity of that particular step in a single step reaction

Select the correct answer by using the codes given below

- a) 1, 3 and 4                      b) 1, 2 and 3                      c) 2, 3 and 4                      d) 1, 2 and 4

129. For the order reaction with rate constant 'K' and initial concentration 'a', the half-life period given by

- a)  $\frac{1n2}{k}$                                       b)  $\frac{1}{ka}$                                       c)  $\frac{3}{2k.a^2}$                                       d) None of these

130. For the reaction,  $2A + B \rightarrow C + D$ , the order of reaction is

- a) One with respect[B]                                      b) Two with respect to [A]  
c) Three                                      d) Cannot be predicted

131. Which expression is wrong for first order reaction?

- a)  $k = \frac{2.303}{t} \log \left( \frac{A_0}{A_t} \right)$                                       b)  $k = \frac{t}{2.303} \log \left( \frac{A_0}{A_t} \right)$   
c)  $-k = \frac{t}{2.303} \log \left( \frac{A_t}{A_0} \right)$                                       d) Rate=k[A]

132. For a first order reaction, the half-life period is

- a) Dependent on the square of the initial concentration.  
b) Dependent on first power of initial concentration.  
c) Dependent on the square root of initial concentration.  
d) Independent on initial concentration

133. Give relation between half reaction time ( $t_{1/2}$ ) and initial concentration of reactant for  $(n - 1)$  order reaction.

- a)  $t_{1/2} \propto [R]_0$                       b)  $t_{1/2} \propto [R]_0^{2-n}$                       c)  $t_{1/2} \propto [R]_0^{n+1}$                       d)  $t_{1/2} \propto [R]_0^{n-2}$

134. For a first order reaction  $A \rightarrow \text{Product}$ , the initial concentration of A is 0.1 M and after 40 minute it becomes 0.025 M. Calculate the rate of reaction at reactant concentration of 0.01M:

- a)  $3.47 \times 10^{-4} \text{ M min}^{-1}$   
b)  $3.47 \times 10^{-5} \text{ M min}^{-1}$   
c)  $1.735 \times 10^{-6} \text{ M min}^{-1}$   
d)  $1.735 \times 10^{-4} \text{ M min}^{-1}$

135. Rate of reaction

- a) Decreases with increase in temperature  
b) Increases with increase in temperature  
c) May increase or decrease with increase in temperature  
d) Does not depends on temperature

136. For the first order reaction with the rate constant  $k$ , which expression gives the rate half-life period? (Initial conc. =  $a$ )

- a)  $\frac{3}{2}ka^2$                                       b)  $\frac{1}{ka}$                                       c)  $\frac{1}{k}$                                       d)  $\frac{0.693}{k}$

137. The rate constant ( $k'$ ) of one of the reaction is found to be double that of the rate constant ( $k''$ ) of another reaction. Then the relationship between the corresponding activation energies of the two reaction ( $E'_a$  and  $E''_a$ ) can be represented as

- a)  $E'_a > E''_a$                                       b)  $E'_a = 4E''_a$                                       c)  $E'_a = E''_a$                                       d)  $E'_a < E''_a$

138. For a given reaction of first order, it takes 15 minute for the concentration to drop from 0.8 M litre<sup>-1</sup> to 0.4 M litre<sup>-1</sup>. The time required for the concentration to drop from 0.1 M litre<sup>-1</sup> to 0.025 M litre<sup>-1</sup> will be:

- a) 60 minute                                      b) 15 minute                                      c) 7.5 minute                                      d) 30 minute

139. The rate constant for a second order reaction is  $8 \times 10^{-5} \text{ M}^{-1} \text{ min}^{-1}$ . How long will it take a 1M solution to be reduced to 0.5M?

- a)  $8.665 \times 10^3$  minute                      b)  $8 \times 10^{-5}$  minute                      c)  $1.25 \times 10^4$  minute                      d)  $4 \times 10^{-5}$  minute

140. The slope in Arrhenius plot, is equal to

- a)  $-\frac{E_a}{2.303 R}$       b)  $\frac{E_a}{R}$       c)  $-\frac{R}{2.303 E_a}$       d) None of the above

141. The rate constants  $k_1$  and  $k_2$  for two different reactions are  $10^{16}e^{-2000/T}$  and  $10^{15}e^{-1000/T}$ , respectively. The temperature at which  $k_1 = k_2$  is:

- a)  $\frac{2000}{2.303}$  K      b) 2000 K      c)  $\frac{1000}{2.303}$  K      d) 1000 K

142. If the volume of the vessel in which the reaction  $2NO + O_2 \rightarrow 2NO_2$  is occurring is diminished to  $1/3^{\text{rd}}$  of its initial volume. The rate of the reaction will be increased by

- a) 3 times      b) 9 times      c) 27 times      d) 36 times

143. The time for half-life period of a creation reaction  $A \rightarrow \text{products}$  is 1 h. when the initial concentration of the reactant 'A', is  $2.0 \text{ mol L}^{-1}$ , how much time does it take for its concentration to come from  $0.50$  to  $0.25 \text{ mol L}^{-1}$ , if it is a zero order reaction?

- a) 4 h      b) 0.5 h      c) 0.25 h      d) 1 h

144. For a reaction  $A + B \rightarrow C + D$ , if the concentration of A is doubled without altering the concentration of B, the rate gets doubled. If the concentration of B is increased by nine times without altering the concentration of A, the rate gets tripled. The order of the reaction is

- a) 2      b) 1      c)  $3/2$       d)  $4/3$

145. What fraction of a reactant showing first order remains after 40 minute if  $t_{1/2}$  is 20 minute?

- a)  $1/4$       b)  $1/2$       c)  $1/8$       d)  $1/6$

146. At 500 k, the half-life period of a gaseous reaction at an initial pressure of 80 kPa is 350 s. when the pressure is 40 kPa, the half-life period is 175 s. The order of the reaction is

- a) Zero      b) One      c) Two      d) Three


147. Which of the following statements are incorrect?

- a) Rate of the reaction involving conversion of *ortho* hydrogen to *para* hydrogen =  $-\frac{d[H_2]}{dt} = k[H_2]^{3/2}$   
 b) Rate of the reaction involving the thermal decomposition of acetaldehyde =  $k[CH_3CHO]^{1/2}$   
 c) In the formation of phosgene from CO and  $Cl_2$ , the rate of the reaction =  $k[CO][Cl_2]^{1/2}$   
 d) In the decomposition of  $H_2O_2$ , the rate of reaction =  $k[H_2O_2]$

148. At 373 K, a gaseous reaction  $A \rightarrow 2B + C$  is found to be of first order. Starting with pure A, the total pressure at the end of 10 min was 176 mm and after a long time when A was completely dissociated, it was 270 mm. The pressure of A at the end of 10 min was

- a) 94 mm      b) 47 mm      c) 43 mm      d) 90 mm

149. For a reversible reaction,  $A \rightleftharpoons B$ , which one of the following statements is wrong from the given energy profile diagram?



- a) Activation energy of forward reaction is greater than backward reaction  
 b) The forward reaction is endothermic  
 c) The threshold energy is less than that of activation energy  
 d) The energy of activation of forward reaction is equal to the sum of heat of reaction and the energy of activation of backward reaction

150. Which one of the following is wrongly matched?

- a) Saponification of  $CH_3COOC_2H_5$  -second order reaction  
 b) Hydrolysis of  $CH_3COOCH_3$  -pseudo unimolecular

- c) Decomposition of  $H_2O_2$  -first order reaction  
 d) Combination of  $H_2$  and  $Br_2$  to give HBr -first order reaction
151. For the reaction,  
 $2N_2O_5(g) \rightarrow 4NO_2(g) + O_2(g)$   
 If the concentration of  $NO_2$  increase by  $5.2 \times 10^{-3} M$  in 100 s then the rate of the reactions  
 a)  $1.3 \times 10^{-5} Ms^{-1}$       b)  $0.5 \times 10^{-4} Ms^{-1}$       c)  $7.6 \times 10^{-4} Ms^{-1}$       d)  $2 \times 10^{-3} Ms^{-1}$
152. The rate of the reaction  $A \rightarrow$  product, at the initial concentration of  $3.24 \times 10^{-2} M$  is nine times its rate at another initial concentration of  $1.2 \times 10^{-3} M$ . The order of the reaction is  
 a)  $\frac{1}{2}$       b)  $\frac{3}{4}$       c)  $\frac{3}{2}$       d)  $\frac{2}{3}$
153. The half-life period for zero order reaction  $A \rightarrow$  product, is 100 min. How long will it take in 80% completion?  
 a) 80 min      b) 160 min      c) 100 min      d) 200 min
154. Consider the reaction  $2A + B \rightarrow$  product  
 When concentration of B alone was doubled, the half-life did not change. When the concentration of A alone was doubled, the rate increased by two times. The unit of rate constant for this reaction is  
 a)  $L mol^{-1} s^{-1}$       b) No unit      c)  $mol L^{-1} s^{-1}$       d)  $s^{-1}$
155. The expression for rate constant of a first order chemical reaction is  
 a)  $k = \frac{1}{t} \cdot \frac{x}{a(a-x)}$       b)  $k = \frac{2.303}{t} \log_{10} \frac{a}{(a-x)}$   
 c)  $k = \frac{x}{t}$       d)  $k = \frac{1}{2t} \left[ \frac{1}{(a-x)^2} - \frac{1}{a^2} \right]$
156. In gaseous reactions important for the understanding of the upper atmosphere  $H_2O$  and O react bimolecularly to form two OH radicals.  $\Delta H$  for this reaction is 72kJ at 500 K and  $E_a$  is 77 kJ  $mol^{-1}$ , then  $E_a$  for the bimolecular recombination of two OH radicals to form  $H_2O$  and O is:  
 a) 3 kJ  $mol^{-1}$       b) 4 kJ  $mol^{-1}$       c) 5 kJ  $mol^{-1}$       d) 7 kJ  $mol^{-1}$
157. Activation energy of a reaction  
 a) Is independent of temperature  
 b) Increases with temperature  
 c) Gets doubled for every 10 degree rise in temperature  
 d) Decreases with temperature
158. For a I order reaction  $A \rightarrow B$  the reaction rate at reactant concentration 0.01M is found to be  $2.0 \times 10^{-5} Ms^{-1}$ . The half-life period of the reaction is:  
 a) 30s      b) 300s      c) 220s      d) 347s
159. For a zero order reaction, the plot of concentration of reactant vs time is (intercept refers to concentration axis)  
 a) Linear with positive slope and zero intercept  
 b) Linear with negative slope and zero intercept  
 c) Linear with negative slope and non-zero intercept  
 d) Linear with positive slope and non-zero intercept
160. The rate of reaction between two reactants A and B decreases by a factor 4, if the concentration of reactant B is doubled. The order of this reaction with respect to B is:  
 a) -1      b) -2      c) 2      d) 1
161. The velocity constant of a reaction at 290 K was found to be  $3.2 \times 10^{-3}$  at 300 K, it will be  
 a)  $6.4 \times 10^{-3}$       b)  $1.28 \times 10^{-2}$       c)  $9.6 \times 10^{-3}$       d)  $3.2 \times 10^{-4}$
162. The term  $\frac{-dc}{dt}$  in a rate equation refers to  
 a) The decrease in concentration of the reactant with time  
 b) The concentration of the reactant  
 c) The change in concentration of the reactant  
 d) The velocity constant of the reaction

163. In a first order reaction the concentration of reactant decreases from  $800 \text{ mol/dm}^6$  to  $50 \text{ mol/dm}^6$  in  $2 \times 10^4 \text{ s}$ . The rate constant of reaction in  $\text{s}^{-1}$  is  
 a)  $2 \times 10^4$                       b)  $3.45 \times 10^{-5}$                       c)  $1.386 \times 10^{-4}$                       d)  $2 \times 10^{-4}$
164. Consider the chemical reaction,  
 $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$   
 The rate of this reaction can be expressed in terms of time derivative of concentration of  $\text{N}_2(\text{g})$ ,  $\text{H}_2(\text{g})$  and  $\text{NH}_3(\text{g})$ . Identify the correct relationship amongst the rate expressions:  
 a)  $\text{Rate} = -d[\text{N}_2]/dt = -\frac{1}{3}d[\text{H}_2]/dt = \frac{1}{2}d[\text{NH}_3]/dt$   
 b)  $\text{Rate} = -d[\text{N}_2]/dt = -3d[\text{H}_2]/dt = 2d[\text{NH}_3]/dt$   
 c)  $\text{Rate} = d[\text{N}_2]/dt = \frac{1}{3}d[\text{H}_2]/dt = \frac{1}{2}d[\text{NH}_3]/dt$   
 d)  $\text{Rate} = -d[\text{N}_2]/dt = -d[\text{H}_2]/dt = d[\text{NH}_3]/dt$
165. Rate of reaction can be expressed by following rate expression,  $\text{rate} = k[\text{A}]^2[\text{B}]$ , if concentration of A is increased by 3 times and concentration of B is increased by 2 times, how many times rate of reaction increases?  
 a) 9 times                      b) 27 times                      c) 18 times                      d) 8 times
166. As the reaction progresses, the rate of reaction  
 a) Increases                      b) Decreases  
 c) Remains constant                      d) First increases, then decreases
167. The data for the reaction,  $\text{A} + \text{B} \rightarrow \text{C}$
- | Ex | $[\text{A}]_0$ | $[\text{B}]_0$ | Initial rate |
|----|----------------|----------------|--------------|
| 1  | 0.012          | 0.035          | 0.10         |
| 2  | 0.024          | 0.070          | 0.80         |
| 3  | 0.024          | 0.035          | 0.10         |
| 4  | 0.012          | 0.070          | 0.80         |
- The rate law corresponds to the above data is  
 a)  $\text{rate} = k[\text{B}]^3$                       b)  $\text{rate} = k[\text{B}]^4$                       c)  $\text{rate} = k[\text{A}][\text{B}]^3$                       d)  $\text{rate} = k[\text{A}]^2[\text{B}]^2$
168. In a reaction, when the concentration of reactant is increased two times, the increase in rate of reaction was four times. Order of reaction is  
 a) Zero                      b) 1                      c) 2                      d) 3
169. For the reaction  
 $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$   
 The rate of change of concentration for hydrogen is  $0.3 \times 10^{-4} \text{ Ms}^{-1}$   
 The rates of change of concentration of ammonia is  
 a)  $-0.2 \times 10^{-4}$                       b)  $0.2 \times 10^{-4}$                       c)  $0.1 \times 10^{-4}$                       d)  $0.3 \times 10^{-4}$
170. Which of the following statement is in accordance with collision theory?  
 Rate is directly proportional to collision frequency  
 Rate depend upon orientation of atoms  
 Temperature determines the rate  
 a) Only III                      b) Only I and II  
 c) Only II and III                      d) All of these
171.  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g}) + 22 \text{ kcal}$ .  
 The activation energy for the forward reaction is 50 kcal. What is the activation energy for the backward reaction?  
 a) -72 kcal                      b) -28 kcal                      c) +28 kcal                      d) +72 kcal
172. According to collision theory:  
 a) Collisions are sufficiently violent  
 b) All collision are responsible for reaction  
 c) All collisions are effective  
 d) Only highly energetic molecules have enough energy to react

173. The rate constant of a first order reaction whose half-life is 480 s is  
 a)  $1.44 \text{ s}^{-1}$                       b)  $1.44 \times 10^{-3} \text{ s}^{-1}$                       c)  $2.88 \times 10^{-3} \text{ s}^{-1}$                       d)  $0.72 \times 10^{-3} \text{ s}^{-1}$

174.  $2A \rightarrow B + C$ ; It would be a zero order reaction when:  
 a) The rate of reaction is proportional to square of conc. of A  
 b) The rate of reaction remains same at any conc. of A  
 c) The rate remains unchanged at any conc. of B and C  
 d) The rate of reaction doubles if conc. of B is increased to double

175. For a reaction  $A + 2B \rightarrow C$ , rate is given by

$$r = K[A][B]^2$$


The order of reaction is:

- a) 3                                      b) 6                                      c) 5                                      d) 7


176. Rate constant for a reaction is  $\lambda$ . Average life is represent by

- a)  $\frac{1}{\lambda}$                                       b)  $\frac{1n^2}{\lambda}$                                       c)  $\frac{\lambda}{\sqrt{2}}$                                       d)  $\frac{0.693}{\lambda}$


177. The plot between concentration versus time for a zero order reaction is represented by



178. For the decomposition of  $\text{N}_2\text{O}_5(\text{g})$ , it is given that:



Activation energy  $E_a$



Activation energy  $E'_a$

then;

- a)  $E_a = E'_a$                                       b)  $E_a > E'_a$                                       c)  $E_a < E'_a$                                       d)  $E_a = 2E'_a$

179. During the kinetic study of the reaction  $2A + B \rightarrow C + D$  following results were obtained.

Run	[A] in M	[B] in M	Initial rate of formation of D in $\text{Ms}^{-1}$
I	0.1	0.1	$6.0 \times 10^{-3}$
II	0.3	0.2	$7.2 \times 10^{-2}$
III	0.3	0.4	$2.88 \times 10^{-1}$
IV	0.4	0.1	$2.40 \times 10^{-2}$

On the basis of above data which one is correct:

- a)  $r = k[A]^2[B]$                                       b)  $r = k[A][B]$                                       c)  $r = k[A^2][B]^2$                                       d)  $r = k[A][B]^2$


180. If the reaction rate at a given temperature becomes slower then

- a) The free energy of activation is higher  
 b) The free energy of activation is lower  
 c) The entropy changes  
 d) The initial concentration of the reactants remains constant

181. The number of molecules of the reactants taking part in a single step of the reaction tells about:

- a) Molecularity of the reaction  
 b) Mechanism of the reaction  
 c) Order of reaction  
 d) All of the above

182. For the reaction system,






Volume is suddenly reduced to half its value by increasing the pressure on it. If the reaction is of first order with respect to  $O_2$  and second order with respect to  $NO$ ; the rate of reaction will

- a) Diminish to one –fourth of its initial value                      b) Diminish to one –eighth of its initial value  
c) Increase to eight time of its initial value                      d) Increase to four time of its initial value

183. The reaction,



- a) Biomolecular reaction  
b) II order reaction  
c) Both (a) and (b)  
d) None of these


184. Which is correct relation in between  $\frac{dc}{dt}$ ,  $\frac{dn}{dt}$  and  $\frac{dP}{dt}$ , where  $c$ ,  $n$ , and  $P$ , represent concentration, mole and pressure terms for gaseous phase reactant  $A(g) \rightarrow$  product?

- a)  $-\frac{dc}{dt} = -\frac{1}{V} \frac{dn}{dt}$                       b)  $\frac{dc}{dt} = \frac{dn}{dt} = -\frac{dP}{dt}$                       c)  $\frac{dc}{dt} = \frac{RT}{V} \frac{dn}{dt} = -\frac{dP}{dt}$                       d) All of the above  
 $= -\frac{1}{RT} \frac{dP}{dt}$

185. The rate constant of a reaction is found to be  $3 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$ . The order of reaction is

- a) Zero                      b) 1                      c) 2                      d) 1.5

186. A reactant (A) forms two products :



If  $E_{a_2} = 2 E_{a_1}$ , then  $k_1$  and  $k_2$  are related as

- a)  $k_1 = 2k_2 e^{E_{a_2}/RT}$                       b)  $k_1 = k_2 e^{E_{a_1}/RT}$                       c)  $k_2 = k_1 e^{E_{a_2}/RT}$                       d)  $k_1 = Ak_2 e^{E_{a_1}/RT}$



187. For the reaction  $2A + B \rightarrow A_2B$ , the rate Law given is

- a)  $k[2A][B]$                       b)  $k[A]^3[B]$                       c)  $k[A][B]^3$                       d)  $k[A]^2[B]$

188. For producing the effective collisions the colliding molecules must have:

- a) A certain minimum amount of energy  
b) Energy lesser than threshold energy  
c) Improper orientation  
d) Proper orientation and energy equal or greater than threshold energy

189. The chemical reaction  $2O_3 \rightarrow 3O_2$  proceeds as follows



The rate law expression should be

- a)  $r = k[O_3]^2$                       b)  $r = k[O_3]^2 [O_2]^{-1}$                       c)  $r = k[O_3][O_2]$                       d) Unpredictable

190. Two substances A and B are present such that  $[A] = 4[B]$  and half-life of A is 5 minute and of B is 15 minute. If they start decaying at the same time following first order, how much time later will the concentration of both of them would be same?

- a) 15 minute                      b) 10 minute                      c) 5 minute                      d) 12 minute

191. A reaction involving A, B and C as reactants is found to obey the rate law,  $\text{rate} = k[A]^x[B]^y[C]^z$ . When the concentration of A, B and C are doubled separately, the rate is also found to increase two, zero and four times respectively. The overall order of the reaction is

- a) 1                      b) 2                      c) 3                      d) 4

192. The rate constant of  $n$ th order has units:

- a)  $\text{litre}^{1-n} \text{ mol}^{1-n} \text{ sec}^{-1}$                       b)  $\text{mol}^{n-1} \text{ litre}^{n-1} \text{ sec}^{-1}$                       c)  $\text{mol}^{1-n} \text{ litre}^{n-1} \text{ sec}^{-1}$                       d) None of these


193. The reaction;  $N_2O_5$  in  $2NO_2 + 1/2 O_2(g)$  is of first order for  $N_2O_5$  with rate constant  $6.2 \times 10^{-4} \text{ s}^{-1}$ . What is the value of rate of reaction when  $[N_2O_5] = 1.25 \text{ mole L}^{-1}$ ?

- a)  $5.15 \times 10^{-5} \text{ mole L}^{-1} \text{ s}^{-1}$   
b)  $6.35 \times 10^{-3} \text{ mole L}^{-1} \text{ s}^{-1}$




206. In the first order reaction, the concentration of the reactants is reduced to 25% in one hour. The half-life period of the reaction is  
 a) 2h                                      b) 4h                                      c) 1/2h                                      d) 1/4h

207. In the given graph the activation energy,  $E_a$  for the reverse reaction will be



- a) 150 kJ                                      b) 50 kJ                                      c) 200 kJ                                      d) 100 kJ
208. What is the order of a reaction which has a rate expression rate =  $k[A]^{3/2}[B]^{-1}$ ?  
 a)  $\frac{3}{2}$                                       b) Zero                                      c)  $\frac{1}{2}$                                       d) None of these
209. For a first order reaction, the concentration changes from 0.8 to 0.4 in 15 min. The time taken for the concentration to change from 0.1 M to 0.025 M is  
 a) 30 Min                                      b) 15 Min                                      c) 7.5 min                                      d) 60 min

210. Give the hypothetical reaction mechanism



Species formed	Rate of its Formation
B	0.002 mol/h, per mole of A
C	0.030 mol/h, per mole of B
D	0.011 mol/h, per mole of C
E	0.420 mol/h, per mole of D

The rate determining step is

- a) Step I                                      b) Step II                                      c) Step III                                      d) Step IV
211. For  $A + B \rightarrow C + D$ ,  $\Delta H = -20 \text{ kJ mol}^{-1}$  the activation energy of the forward reaction is  $85 \text{ kJ mol}^{-1}$ . The activation energy for backward reaction is ... $\text{kJ mol}^{-1}$ .  
 a) 65                                      b) 105                                      c) 85                                      d) 40
212. For the reaction  $\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$ , the rate  $\frac{d[\text{NH}_3]}{dt} = 2 \times 10^{-4} \text{ M s}^{-1}$ . Therefore, the rate  $-\frac{d[\text{N}_2]}{dt}$  is given as:  
 a)  $10^{-4} \text{ M s}^{-1}$                                       b)  $10^4 \text{ M s}^{-1}$                                       c)  $10^{-2} \text{ M s}^{-1}$                                       d)  $10^{-4} \text{ s M}^{-1}$
213. The reaction  $L \rightarrow M$  is started with 10.0 g of L. After 30 and 90 min, 5.0 g and 1.25 g of L respectively are left. The order of the reaction is  
 a) 0                                      b) 1                                      c) 2                                      d) 3
214. The rate of a reaction is expressed in different ways as follows

$$+\frac{1}{2} \frac{d[C]}{dt} = -\frac{1}{5} \frac{d[D]}{dt} = +\frac{1}{3} \frac{d[A]}{dt} = -\frac{d[B]}{dt}$$

The reaction is

- a)  $4A + B \rightarrow 2C + 3D$                                       b)  $B + 5D \rightarrow 3A + 2C$   
 c)  $4A + 2B \rightarrow 2C + 3D$                                       d)  $B + \frac{1}{2}D \rightarrow 4A + 2C$

215. The rate of elementary reaction,  $A \rightarrow B$ , increases by 100 times when the concentration of  $A$  is increased ten folds. The order of the reaction with respect to  $A$  is:  
 a) 1    b) 2    c) 10    d) 100
216. The differential rate law for the reaction  $4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \rightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{g})$  is  
 a)  $-\frac{d[\text{NH}_3]}{dt} = -\frac{d[\text{O}_2]}{dt} = -\frac{d[\text{NO}]}{dt} = -\frac{d[\text{H}_2\text{O}]}{dt}$     b)  $\frac{d[\text{NH}_3]}{dt} = \frac{d[\text{O}_2]}{dt} = -\frac{1}{4} \frac{d[\text{NO}]}{dt} = -\frac{1}{6} \frac{d[\text{H}_2\text{O}]}{dt}$   
 c)  $\frac{1}{4} \frac{d[\text{NH}_3]}{dt} = \frac{1}{5} \frac{d[\text{O}_2]}{dt} = \frac{1}{4} \frac{d[\text{NO}]}{dt} = \frac{1}{6} \frac{d[\text{H}_2\text{O}]}{dt}$     d)  $-\frac{1}{4} \frac{d[\text{NH}_3]}{dt} = -\frac{1}{5} \frac{d[\text{O}_2]}{dt} = \frac{1}{4} \frac{d[\text{NO}]}{dt} = \frac{1}{6} \frac{d[\text{H}_2\text{O}]}{dt}$
217. For the reaction  $A \rightarrow B$ , when concentration of  $A$  is made 1.5 times, the rate of reaction becomes 1.837 times. The order of reaction is  
 a) 1    b) 1.5    c) 2    d) 2.5
218. For the reaction,  $\text{Cl}_2 + 2\text{I}^- \rightarrow \text{I}_2 + 2\text{Cl}^-$ , the initial concentration of  $\text{I}^-$  was  $0.20 \text{ mol L}^{-1}$  and the concentration after 20 min was  $0.20 \text{ mol L}^{-1}$ . Then the rate of formation of  $\text{I}_2$  in  $\text{mol L}^{-1}$  would be  
 a)  $1 \times 10^{-4}$     b)  $5 \times 10^{-4}$     c)  $1 \times 10^{-3}$     d)  $5 \times 10^{-3}$
219. The energies of activation for forward and reverse reactions for  $\text{A}_2 + \text{B}_2 \rightleftharpoons 2\text{AB}$  are  $180 \text{ kJ mol}^{-1}$  and  $200 \text{ kJ mol}^{-1}$  respectively. The presence of a catalyst lowers the activation energy of both (forward and reverse) reactions by  $100 \text{ kJ mol}^{-1}$ . The enthalpy change of the reaction ( $\text{A}_2 + \text{B}_2 \rightarrow 2\text{AB}$ ) in the presence of catalyst will be (in  $\text{kJ mol}^{-1}$ )  
 a) 300    b) 120    c) 280    d) 20
220. Which statement is correct?  
 a) Reactions with low activation energy are usually exothermic  
 b) The rate law sometimes enable to deduce the mechanism of a reaction  
 c) The rate law for reaction is an algebraic expression relating the forward reaction rate to product concentration  
 d) Increase in the total pressure of a gas phase reaction increase the fraction of collisions effective in producing reactions
221. The temperature coefficient of a reaction is:  
 a) The rate constant at a fixed temperature  
 b) The ratio of rate constant at two temperature  
 c) The ratio of rate constant differing by  $10^\circ$  preferably  $25^\circ$  and  $35^\circ\text{C}$   
 d) None of the above
222. The term  $(-dc/dt)$  in rate equation refers to:  
 a) The concentration of a reactant  
 b) The decrease in concentration of the reactant with time  
 c) The velocity constant of reaction  
 d) None of the above
223. For a first order reaction, the initial concentration of a reactant is  $0.05 \text{ M}$ . After 45 min it is decreased by  $0.015 \text{ M}$ . calculation half reaction time ( $t_{1/2}$ )  
 a) 87.42 min    b) 25.90 min    c) 78.72 min    d) 77.20 min
224. The order of a gaseous phase reaction for which rate becomes half if volume of container having same amount of reactant is doubled is:  
 a) 1    b)  $1/2$     c) 2    d)  $1/3$
225. For the reaction,  $\text{N}_2\text{O}_5 \rightarrow 2\text{NO}_2 + \frac{1}{2}\text{O}_2$   
 Given,  $-\frac{d[\text{N}_2\text{O}_5]}{dt} = K_1[\text{N}_2\text{O}_5]$   
 $\frac{d[\text{NO}_2]}{dt} = K_2[\text{N}_2\text{O}_5]$   
 and  $\frac{d[\text{O}_2]}{dt} = K_3[\text{N}_2\text{O}_5]$   
 The relation in between  $K_1, K_2$  and  $K_3$  is:  
 a)  $2K_1 = K_2 = 4K_3$     b)  $K_1 = K_2 = K_3$     c)  $2K_1 = 4K_2 = K_3$     d) None of these

226. Temperature dependent equation can be written as  
 a)  $\ln k = \ln A - e^{E_a/RT}$  b)  $\ln k = \ln A + e^{E_a/RT}$   
 c)  $\ln k = \ln A - e^{RT/E_a}$  d) None of these
227. A first order reaction is 50% complete in 30 min at 27°C and in 10 min at 47°C. The energy of activation of the reaction is  
 a) 52.8 kJ b) 23.6 kJ c) 29.5 kJ d) 43.8 kJ
228. Increase in the concentration of the reactants leads to the change in  
 a) Activation energy b) Heat of reaction  
 c) Collision frequency d) Threshold energy
229. When a graph is plotted between  $\ln k$  and  $1/T$  for a first order reaction, a straight line is obtained. The slope of the line is equal to  
 a)  $-\frac{E_a}{2.303}$  b)  $-\frac{E_a}{2.303R}$  c)  $-\frac{2.303}{E_a R}$  d)  $-\frac{E_a}{R}$
230. For the two gaseous reactions, following data are given  
 $A \rightarrow B; k_1 = 10^{10} e^{-20,000/T}$   
 $C \rightarrow D; k_2 = 10^{12} e^{-24,606/T}$   
 The temperature at which  $k_1$  becomes equal to  $k_2$  is  
 a) 400 K b) 1000 K c) 800 K d) 1500 K
231. Hydrogenation of vegetable ghee at 25°C reduces pressure of  $H_2$  from 2 atm to 1.2 atm in 50 minute. The rate of reaction in terms of molarity per second is:  
 a)  $1.09 \times 10^{-6}$  b)  $1.09 \times 10^{-5}$  c)  $1.09 \times 10^{-7}$  d)  $1.09 \times 10^{-8}$
232. In the reaction  $A + B \rightarrow \text{products}$ , if  $B$  is taken in excess, then it is an example of  
 a) Second order reaction b) Zero order reaction  
 c) Pseudounimolecular reaction d) First order reaction
233. The rate of a chemical reaction depends upon:  
 a) Time b) Pressure c) Concentration d) All of these
234. The rate constant for the reaction,  
 $2N_2O_5 \rightarrow 4NO_2 + O_2$  is  $3.0 \times 10^{-5} \text{ s}^{-1}$ . If the rate is  $2.40 \times 10^{-5}$  then the concentration of  $N_2O_5$  (in mol/L) is  
 a) 1.4 b) 1.2 c) 0.04 d) 0.8
235. The following data are for the decomposition of ammonium nitrite in aqueous solution.
- | Vol. of $N_2$ in cc | Time (min) |
|---------------------|------------|
| 6.25                | 10         |
| 9.00                | 15         |
| 11.40               | 20         |
| 13.65               | 25         |
| 33.05               | Infinity   |
- The order of reaction is  
 a) Zero b) One c) Two d) Three
236. For a reaction  $A \rightarrow \text{Product}$ , rate law is  $-\frac{d[A]}{dt} = K[A]_0$ . The concentration of  $A$  left after time  $t$  when  $t = \frac{1}{K}$  is:  
 a)  $\frac{[A]_0}{e}$  b)  $[A]_0 \times e$  c)  $\frac{[A]_0}{e^2}$  d)  $\frac{1}{[A]_0}$
237. For a first order reaction  $A \rightarrow P$ , the temperature ( $T$ ) dependent rate constant ( $k$ ) was found to follow the equation.  
 $\log k = -(2000)/T + 6.0$   
 The pre-exponential factor  $A$  and the activation energy  $E_a$ , respectively, are  
 a)  $1.0 \times 10^6 \text{ s}^{-1}$  and  $9.2 \text{ kJ mol}^{-1}$  b)  $6.0 \text{ s}^{-1}$  and  $16.6 \text{ kJ mol}^{-1}$   
 c)  $1.0 \times 10^6 \text{ s}^{-1}$  and  $16.6 \text{ kJ mol}^{-1}$  d)  $1.0 \times 10^6 \text{ s}^{-1}$  and  $38.3 \text{ kJ mol}^{-1}$


238. In Arrhenius equation  $K = Ae^{-E_a/RT}$ , the quantity  $-E_a/RT$  is referred as:  
 a) Boltzmann factor      b) Frequency factor      c) Activation factor      d) None of these
239. Among the following reaction, the fastest one is:  
 a) Burning of coal  
 b) Rusting of iron in moist air  
 c) Conversion of monoclinic sulphur to rhombic sulphur  
 d) Precipitation of silver chloride by mixing silver nitrate and sodium chloride solutions
240. The following homogeneous gaseous reactions were experimentally found to be second order overall.  
 1.  $2\text{NO} \rightarrow \text{N}_2 + \text{O}_2$     2.  $3\text{O}_2 \rightarrow 2\text{O}_3$   
 3.  $\text{N}_2\text{O}_3 \rightarrow \text{NO} + \text{NO}_2$     4.  $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$   
 Which of these are most likely to be elementary reaction that occur in one step?  
 a) 3 only      b) 1 and 3      c) 1 and 4      d) 3 and 4
241. Consider a reaction;  $aG + bH \rightarrow \text{Products}$   
 When concentration of both the reactants G and H is doubled, the rate increases by eight times. However, when concentration of G is doubled keeping the concentration of H fixed, the rate is doubled. The overall of the reaction,  
 a) 0      b) 1      c) 2      d) 3
242. The rate constant for a zero order reaction is  
 a)  $k = \ln \frac{c_0 - c_t}{2t}$       b)  $k = \frac{c_0 - c_t}{t}$       c)  $k = \frac{c_0}{c_t}$       d)  $k = \frac{c_0}{2t}$
243. The ratio of the times for 99.9% of the reaction to complete and half of the reaction to complete is  
 a) 2      b) 4      c) 8      d) 10
244. The activation energy for a simple chemical reaction  $A \rightarrow B$  is  $E_a$  in forward direction. The activation energy for the reverse reaction  
 a) Is negative of  $E_a$       b) Is always less than  $E_a$   
 c) Can be less than or more than  $E_a$       d) Is always double of  $E_a$
245. After how many second will the concentration of the reactant in a first order reaction be halved if the rate constant is  $1.155 \times 10^{-3} \text{ s}^{-1}$ ?  
 a) 600      b) 100      c) 60      d) 10
246. For the reaction  $A + 2B \rightarrow C$ , the rate of reaction at a given instant can be given by  
 a)  $+\frac{d[A]}{dt} = +\frac{1}{2} \frac{d[B]}{dt} = +\frac{d[C]}{dt}$       b)  $\frac{d[A]}{dt} = -\frac{1}{2} \frac{d[B]}{dt} = -\frac{d[C]}{dt}$   
 c)  $-\frac{d[A]}{dt} = -\frac{1}{2} \frac{d[B]}{dt} = +\frac{d[C]}{dt}$       d)  $+\frac{d[A]}{dt} = +\frac{1}{2} \frac{d[B]}{dt} = +\frac{d[C]}{dt}$
247. Which of the following theory is not related to chemical kinetics?  
 a) Collision theory  
 b) Activated complex theory  
 c) Absolute reaction rate theory  
 d) VSPER theory
248. For the chemical change  $A \rightarrow B$  it is found that the rate of reaction doubles when the concentration is increased by 4 times. The order of the reaction is  
 a) One      b) Two      c) Half      d) None of these
249. If a certain reaction is first order with respect to A, second order with respect to B and zero order with respect to C then what is the order of reaction?  
 a) First      b) Second      c) Third      d) Zero
250. If a plot of  $\log_{10} C$  versus t give a straight line for a given reaction, then the reaction is  
 a) Zero order      b) First order      c) Second order      d) Third order
251. The given reaction  
 $2\text{FeCl}_3 + \text{SnCl}_2 \rightarrow 2\text{FeCl}_2 + \text{SnCl}_4$  is an example of  
 a) Third order reaction      b) Second order reaction  
 c) First order reaction      d) None of these

252. The order for the reaction,  $\text{H}_2 + \text{Cl}_2 \xrightarrow{h\nu} 2\text{HCl}$  over water is:  
 a) 0                                      b) 1                                      c) 2                                      d) 3
253. The inversion of cane sugar into glucose and fructose is:  
 a) I order                                      b) II order                                      c) III order                                      d) Zero order
254. For the reaction,  $\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$ , if  $\frac{d[\text{NH}_3]}{dt} = 2 \times 10^{-4} \text{ mol L}^{-1}\text{s}^{-1}$ , the value of  $-\frac{d[\text{H}_2]}{dt}$  would be:  
 a)  $1 \times 10^{-4} \text{ mol L}^{-1}\text{s}^{-1}$     b)  $3 \times 10^{-4} \text{ mol L}^{-1}\text{s}^{-1}$     c)  $4 \times 10^{-4} \text{ mol L}^{-1}\text{s}^{-1}$     d)  $6 \times 10^{-4} \text{ mol L}^{-1}\text{s}^{-1}$
255. For the reaction  $\text{N}_2\text{O}_5 \xrightarrow{(g)} 2\text{NO}_2 + \frac{1}{2}\text{O}_2$ , the rate of disappearance of  $\text{N}_2\text{O}_5$  is  $6.25 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$ . The rate of formation of  $\text{NO}_2$  and  $\text{O}_2$  will be respectively.  
 a)  $6.25 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$  and  $6.25 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$   
 b)  $1.25 \times 10^{-2} \text{ mol L}^{-1}\text{s}^{-1}$  and  $3.125 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$   
 c)  $6.25 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$  and  $3.125 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$   
 d)  $1.25 \times 10^{-2} \text{ mol L}^{-1}\text{s}^{-1}$  and  $6.25 \times 10^{-3} \text{ mol L}^{-1}\text{s}^{-1}$
256. If the concentration units are reduced by  $n$  times, then the value of rate constant of first order will:  
 a) Increase by  $n$  times  
 b) Decrease by factor of  $n$   
 c) Not change  
 d) None of these
257. Unit of frequency factor ( $A$ ) is  
 a) mol/L                                      b) mol/L $\times$  s  
 c) Depends upon order of reaction    d) It does not have any unit
258. The ionic reactions are usually very fast because:  
 a) It does not involve bond breaking  
 b) The energy of activation between charged ions is greater than that between neutral molecules  
 c) Collision frequency is very low  
 d) The reactions are highly exothermic
259. In the first order reaction, 75% of the reactant gets disappeared in 1.386h. The rate constant of the reaction is  
 a)  $3.0 \times 10^{-3} \text{ s}^{-1}$                       b)  $2.8 \times 10^{-4} \text{ s}^{-1}$                       c)  $17.2 \times 10^{-3} \text{ s}^{-1}$                       d)  $1.8 \times 10^{-3} \text{ s}^{-1}$
260. Number of mole of a substance present in 1 litre volume is known as:  
 a) Activity                                      b) Molar concentration    c) Active mass                                      d) None of these
261. The rate of a reaction is doubled when temperature increases by  $10^\circ\text{C}$ . If temperature is increased by  $100^\circ\text{C}$ , then rate of reaction will become  
 a) 64 times                                      b) 256 times                                      c) 512 times                                      d) 1024 times
262. For a hypothetical reaction  
 $A + 2B \rightarrow 3C + D$   
 $d[C]/dt$  is equal to  
 a)  $\frac{3d[A]}{dt}$                                       b)  $-\frac{3}{2} \frac{d[B]}{dt}$                                       c)  $-\frac{d[B]}{dt}$                                       d)  $-\frac{d[A]}{dt}$
263. The half-life of two samples is 0.1 and 0.4 s. Their reactive concentration is 200 and 50 respectively. What is the order of reaction?  
 a) 0                                      b) 2                                      c) 1                                      d) 4
264. Consider following two reaction,  
 $A \rightarrow \text{product} - \frac{d[A]}{dt} = k_1[A]^0$   
 $B \rightarrow \text{product} - \frac{d[B]}{dt} = k_2[B]^0$   
 $k_1$  and  $k_2$  are expressed in term of molarity ( $\text{mol L}^{-1}$ ) and time ( $\text{s}^{-1}$ ) as  
 a)  $\text{s}^{-1}, \text{M s}^{-1}\text{L}^{-1}$                       b)  $\text{Ms}^{-1}, \text{M s}^{-1}$                       c)  $\text{s}^{-1}, \text{M}^{-1}\text{s}^{-1}$                       d)  $\text{Ms}^{-1}, \text{L s}^{-1}$
265. The reaction,  $\text{NO} + (1/2)\text{O}_2 \rightarrow \text{NO}_2$  exhibits:

- a) Small negative temperature coefficient  
 b) Decrease in value of  $K$  with temperature  
 c) Decrease in value of rate with temperature  
 d) All of the above
266. Consider the reaction,  
 $Cl_2(aq) + H_2S(aq) \rightarrow S(s) + 2H^+(aq) + 2Cl^-(aq)$  the rate equation for this reaction is , rate =  $k[Cl_2][H_2S]$   
 Which of these mechanisms is/are consistent with this rate equation?
- I.  $Cl_2 + H_2S(aq) \rightarrow H^+Cl^- + Cl^+ + HS^-(slow)$   
 $Cl^+ + HS^- \rightarrow H^+Cl^- + S(fast)$
- II.  $H_2S \leftrightarrow H^+ + HS^-(fast\ equilibrium)$   
 $Cl_2 + HS^- \rightarrow 2Cl^- + H^+S(slow)$
- a) (B) only                      b) Both (A) and (B)                      c) Neither (A) nor (B)                      d) (A)Only
267. In a reaction  $2A \rightarrow$  Products; the concentration of  $A$  decreases from  $0.5\ mol\ litre^{-1}$  to  $0.4\ mol\ litre^{-1}$  in 10 minute. The rate of reaction during this interval is:  
 a)  $0.05\ M\ min^{-1}$                       b)  $0.005\ M\ min^{-1}$                       c)  $0.5\ M\ min^{-1}$                       d)  $5\ M\ min^{-1}$
268. What is the two third life of a first order reaction having  $k = 5.48 \times 10^{-14}\ s^{-1}$ ?  
 a)  $2.01 \times 10^{13}\ s$                       b)  $2.01 \times 10^{12}\ s$                       c)  $4.02 \times 10^{13}\ s$                       d)  $4.02 \times 10^{26}\ s$
269. In a 1st order reaction the concentration of reactant decreases from  $800\ mol/dm^3$  to  $50\ mol/dm^3$  in  $2 \times 10^4\ s$ . The rate constant of reaction in  $S^{-1}$  is  
 a)  $2 \times 10^4$                       b)  $3.45 \times 10^{-5}$                       c)  $1.386 \times 10^{-4}$                       d)  $2 \times 10^{-4}$
270. The rate constant of a first order reaction is  $6.9 \times 10^{-3}\ s^{-1}$ . How much time will it take to reduce the initial concentration to its  $1/8^{th}$  value ?  
 a) 100s                      b) 200s                      c) 300s                      d) 400s
271. In a reaction, the rate expression is, rate =  $K[A][B]^{2/3}[C]^0$ , the order of reaction is:  
 a) 1                      b) 2                      c)  $5/3$                       d) Zero
272. In the Synthesis of ammonia by Haber process, if 60 moles of ammonia is obtained in one hour, then the rate of disappearance of nitrogen is  
 a) 30 mol/min                      b) 6 mol/min                      c) 0.5 mol/min                      d) 60 mol/min
273. Half-life period of second order reaction is  
 a) Proportional to initial concentration of reactants  
 b) Independent of initial concentration of reactants  
 c) Inversely proportional to initial concentration of reactants  
 d) None of the above
274. A reactant with initial concentration  $1.386\ mol\ litre^{-1}$  showing first order change takes 40 minute to become half. If it shows zero order change taking 20 minute to becomes half under the similar conditions, the ratio,  $\frac{K_1}{K_0}$  for first order and zero order kinetics will be:  
 a)  $0.5\ mol^{-1}\ litre$                       b)  $1.0\ mol/litre$                       c)  $1.5\ mol/litre$                       d)  $2.0\ mol^{-1}\ litre$
275. The reason for almost doubling the rate of reaction on increasing the temperature of the reaction system by  $10^\circ C$  is  
 a) The value of threshold energy increases  
 b) Collision frequency increases  
 c) The fraction of the molecules having energy equal to threshold energy increases  
 d) Activation energy decreases
276. Plot of  $\log(a - x)$  versus time  $t$  is straight line. This indicates that the reaction is of:  
 a) Second order                      b) First order                      c) Zero order                      d) Third order
277. The  $t_{1/2}$  of the first order reaction is  
 a) Dependent of initial concentration                      b) Directly proportional to initial concentration  
 c) indirectly proportional to initial concentration                      d) Independent of initial concentration




278. In a zero-order reaction for every  $10^\circ$  rise of temperature, the rate is doubled. If the temperature is increased from  $10^\circ\text{C}$  to  $100^\circ\text{C}$ , the rate of the reaction will become:
- a) 64 times                      b) 128 times                      c) 256 times                      d) 512 times
279. The time required for 100% completion of a zero order reaction is
- a)  $ak$                               b)  $\frac{a}{2k}$                               c)  $\frac{a}{k}$                               d)  $\frac{2k}{a}$
280. The reaction,  $2\text{N}_2\text{O}_5 \rightleftharpoons 2\text{N}_2\text{O}_4 + \text{O}_2$  is
- a) Bimolecular and second order                      b) Unimolecular and first order  
c) Bimolecular and first order                      d) Bimolecular and zero order
281. The thermal decomposition of a compound is of first order. If a sample of the compound decompose 50% in 120 min. What time will it take to undergo 90% decomposition?
- a) Nearly 400 min                      b) Nearly 45 min                      c) Nearly 480 min                      d) Nearly 240 min
282. Which one of the following statements for the order of a reaction is incorrect?
- a) Order of reaction is always a whole number  
b) Order can be determined only experimentally  
c) Order is not influenced by stoichiometric coefficient of the reactants  
d) Order of reaction is sum of power to the concentration terms of reactants to express the rate of reaction
283. The rate of chemical reaction depends on the nature of chemical reactions, because:
- a) The threshold energy level differs from one reaction to another  
b) Some of the reactants are solid at room temperature  
c) Some of the reactants are coloured  
d) All of the above
284. If the rate of reaction between  $A$  and  $B$  is given by, rate =  $K[A][B]^n$ , then the reaction is:
- a) First order in  $A$   
b)  $n$ th order in  $B$   
c) Overall order is  $(1 + n)$   
d) All are correct
285. In a reaction,  $A + B \rightarrow \text{Product}$ , rate is doubled when the concentration of  $B$  is doubled, and rate increases by a factor of 8 when the concentrations of both the reactants ( $A$  and  $B$ ) are doubled, rate law for the reaction can be written as:
- a) Rate =  $k[A][B]$                       b) Rate =  $k[A]^2[B]$                       c) Rate =  $k[A][B]^2$                       d) Rate =  $k[A]^2[B]^2$
286. Combustion of carbon is exothermic, but coal stored in coal depots does not burn automatically because of:
- a) High threshold energy barrier  
b) Kinetic stability of coal  
c) Higher energy of activation needed for burning  
d) All of the above
287. A drop of a solution (volume = 0.05 mL) contains  $6 \times 10^{-7}$  mol of  $\text{H}^+$ . If the rate of disappearance of  $\text{H}^+$  is  $6.0 \times 10^5$  mol/L  $\times$  s, how long will it take for  $\text{H}^+$  to disappear from the drop
- a)  $8.0 \times 10^{-8}$  s                      b)  $2.0 \times 10^{-8}$  s                      c)  $6.0 \times 10^{-6}$  s                      d)  $2.0 \times 10^{-2}$  s
288. The inversion of cane sugar into glucose and fructose according to the equation  $\text{C}_{12}\text{H}_{22}\text{O}_{11} + \text{H}_2\text{O} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + \text{C}_6\text{H}_{12}\text{O}_6$  is an example of
- a) First order reaction                      b) Third order reaction  
c) Second order reaction                      d) Zero order reaction
289. In the reversible reaction



The rate of disappearance of  $\text{NO}_2$  is equal to

- a)  $\frac{2k_1}{k_2} [\text{NO}_2]^2$                       b)  $2k_1[\text{NO}_2]^2 - 2k_2[\text{N}_2\text{O}_4]$

- c)  $2k_2[\text{NO}_2]^2 - k_2[\text{N}_2\text{O}_4]$  d)  $(2k_1 - k_2)[\text{NO}_2]$
290. For the reaction,  $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$  the rate of change of concentration for hydrogen is  $-0.3 \times 10^{-4} \text{ Ms}^{-1}$ . The rate of change of concentration of ammonia is  
 a)  $-0.2 \times 10^{-4}$  b)  $0.2 \times 10^{-4}$  c)  $0.1 \times 10^{-4}$  d)  $0.3 \times 10^{-4}$
291. In the reaction;  $\text{A} + 2\text{B} \rightarrow 3\text{C} + \text{D}$  which of the following expressions does not describe changes in the concentration of various species as a function of time?  
 a)  $\frac{d[\text{C}]}{dt} = \frac{-3d[\text{A}]}{dt}$  b)  $\frac{3d[\text{D}]}{dt} = \frac{d[\text{C}]}{dt}$  c)  $\frac{3d[\text{B}]}{dt} = \frac{-2d[\text{C}]}{dt}$  d)  $\frac{2d[\text{B}]}{dt} = \frac{d[\text{A}]}{dt}$
292. The order of reaction, with respect to one of the reacting component Y, is zero. It implies that  
 a) The reaction is going on at a constant rate.  
 b) The rate of reaction does not vary with temperature.  
 c) The reaction rate is independent of the concentration of Y.  
 d) The rate of formation of the activated complex is zero.
293. The rate of disappearance of  $\text{SO}_2$  in the reaction;  $2\text{SO}_2 + \text{O}_2 \rightarrow 2\text{SO}_3$  is  $1.28 \times 10^{-3} \text{ g/sec}$ . Then the rate of formation of  $\text{SO}_3$  is:  
 a)  $0.64 \times 10^{-3} \text{ g/sec}$  b)  $0.80 \times 10^{-3} \text{ g/sec}$  c)  $1.28 \times 10^{-1} \text{ g/sec}$  d)  $1.60 \times 10^{-3} \text{ g/sec}$
294. For the first order reaction half-life is 14 s, the time required for the initial concentration to reduce to  $1/8$  of its value is  
 a)  $(14)^3 \text{ s}$  b) 28 s c) 42 s d)  $(14)^2 \text{ s}$
295. Given that  $K$  is the rate constant for some order of any reaction at temp.  $T$  then the value of  $\lim_{T \rightarrow \infty} \log K =$  (where  $A$  is the Arrhenius constant):  
 a)  $A/2.303$  b)  $A$  c)  $2.303 A$  d)  $\log A$
296. The rate constant of a first order reaction is  $3 \times 10^{-6}$  per second and initial concentration is 0.10 M. Then the initial rate of reaction is  
 a)  $3 \times 10^{-6} \text{ ms}^{-1}$  b)  $3 \times 10^{-8} \text{ ms}^{-1}$  c)  $3 \times 10^{-7} \text{ ms}^{-1}$  d)  $3 \times 10^{-9} \text{ ms}^{-1}$
297. With respect to the figure given below which of the following statements is correct?



- a)  $E_a$  for the forward reaction is  $C - B$   
 b)  $E_a$  for the forward reaction is  $B - A$   
 c)  $E_{a(\text{forward})} < E_{a(\text{backward})}$   
 d)  $E_a$  (for reverse reaction) =  $C - A$
298. For a first order reaction,  $\text{A} \rightarrow \text{products}$ , the rate of reaction at  $[\text{A}] = 0.2 \text{ M}$  is  $1.0 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$ . The half-life period for the reaction is  
 a) 476 s b) 496 s c) 832 s d) 242 s
299. From the following data, the activation energy for the reaction (cal/mol):  
 $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$   

$T, K$	$1/T, K^{-1}$	$\log_{10} K$
769	$1.3 \times 10^{-3}$	2.9
667	$1.5 \times 10^{-3}$	1.1

  
 a)  $4 \times 10^4$  b)  $2 \times 10^4$  c)  $8 \times 10^4$  d)  $3 \times 10^4$
300. An elementary reaction is given as  $2\text{P} + \text{Q} \rightarrow \text{products}$ . If concentration of  $\text{Q}$  is kept constant and concentration of  $\text{P}$  is doubled then rate of reaction is:  
 a) Doubled b) Halved c) Quadrupled d) Remains same
301. The hydrolysis of ethyl acetate was carried out separately with 0.05 M HCl and 0.05 M  $\text{H}_2\text{SO}_4$ . The rate constants were found to be  $k_1$  and  $k_2$  respectively then  
 a)  $k_1 < k_2$  b)  $k_1 > k_2$  c)  $k_1 = k_2$  d)  $k_2 = 2k_1$

302. Which one of the following statement for order of reaction is not correct ?  
 a) Order can be determined experimentally  
 b) Order of reaction is equal to sum of the power of concentration terms in differential rate law  
 c) It is not affected with stoichiometric coefficients of the reactants  
 d) Order cannot be fractional
303. For a single step, reaction,  $A + 2B \rightarrow \text{Products}$ , the molecularity is  
 a) 0                                      b) 1                                      c) 2                                      d) 3
304. Which of the following statement is correct for a reaction  $X + 2Y \rightarrow \text{Products}$ ?  
 a) The rate of disappearance of X = twice the rate of disappearance of Y  
 b) The rate of disappearance of X = 1/2 rate of appearance of products  
 c) The rate of appearance of products = 1/2 the rate of disappearance of Y  
 d) The rate of appearance of products = 1/2 the rate of disappearance of X
305. For the reaction  $A \rightarrow B$ , the rate law is; rate =  $K[A]$ . Which of the following statements is incorrect?  
 a) The reaction follows first order kinetics  
 b) The  $t_{1/2}$  of reaction depends upon initial concentration of reactant  
 c)  $K$  is constant for the reaction at a constant temperature  
 d) The rate law provides a simple way of predicting the concentration of reactants and products at any times after the start of the reaction
306. For a first order reaction  $k = 100 \text{ s}^{-1}$ . The time for completion of 50% reaction is  
 a) 1 milli-second                      b) 4 milli-second                      c) 7 mili-second                      d) 10 milli-second
307. Pieces of wood burn faster than a log of wood of the same mass because  
 a) Surface area of log of wood is larger and needs more time to burn  
 b) Pieces of wood have large surface area  
 c) All pieces of wood catch fire at the same time  
 d) Block of wood has higher density than pieces of the same wood
308. Which statement is not correct?  
 a) For endothermic reactions, heat of reaction is lesser than energy of activation  
 b) For exothermic reactions, heat of reaction is more than energy of activation  
 c) For exothermic reactions energy of activation is less in forward reaction than in backward reaction  
 d) For endothermic reactions energy of activation is more in forward reaction than in backward reaction
309. Which statement is true?  
 a) Endothermic reactions have higher activation energies than exothermic reactions  
 b) The specific rate constant for a reaction is independent of the concentration of the reacting species  
 c) There is a single rate determining step in any reaction mechanism  
 d) None of the above
310. If  $k_1 =$  rate contant at temperature  $T_1$  and  $k_2 =$ rate constant at temperature  $T_2$  for a first order reaction, then which of the following relations is correct?  
 ( $E_a$  : activation energy)  
 a)  $\log \frac{k_1}{k_2} = \frac{2.303 E_a}{R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$                                       b)  $\log \frac{k_2}{k_1} = \frac{E_a}{2.303 R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$   
 c)  $\log \frac{k_2}{k_1} = \frac{E_a}{2.303 R} \left( \frac{T_1 T_2}{T_2 + T_1} \right)$                                       d)  $\log \frac{k_1}{k_2} = \frac{E_a}{2.303 R} \left( \frac{T_1 T_2}{T_2 - T_1} \right)$
311. What is the half-life of Na-24 if  $2 \times 10^{-4}$  g sample of it disintegrate at the rate of  $7.0 \times 10^{12}$  atoms per s?  
 a)  $4.97 \times 10^5 \text{ s}$                       b)  $4.97 \times 10^4 \text{ s}$                       c)  $4.97 \times 10^6 \text{ s}$                       d)  $4.97 \times 10^2 \text{ s}$
312. The concentration of a reactant X decreases from 01 M to 0.005 m in 40 min. If the reaction follows first order kinetics, the rate of the reaction when the concentration of X is 0.01 M will be  
 a)  $1.73 \times 10^{-4} \text{ Mmin}^{-1}$                                       b)  $3.47 \times 10^{-4} \text{ Mmin}^{-1}$   
 c)  $3.47 \times 10^{-5} \text{ M min}^{-1}$                                       d)  $7.5 \times 10^{-4} \text{ M min}^{-1}$
313. For zero order reaction the integrated rate equation is

a)  $kt = \frac{[A]}{[A]_0}$       b)  $kt = [A] - [A]_0$       c)  $[A] = -kt + [A]_0$       d)  $[A] = kt - [A]_0$

314. The rate equation for the reaction  $2A + B \rightarrow C$  is found to be

Rate =  $k[A][B]$

The correct statement in relation to this reaction that the

- a) Unit of k must be  $s^{-1}$
- b)  $t_{1/2}$  is constant
- c) Rate of formation of C is twice the rate of disappearance of A
- d) Value of k is independent of the initial concentration of A and B

315. The unit of rate constant of second order reaction

- a)  $time^{-1}$
- b)  $conc^{-1}time^{-1}$
- c)  $conc\ time^{-1}$
- d)  $conc^{-2}time^{-1}$

316. The rate constant for the first order reaction is  $6\ s^{-1}$ . How much time will it take to reduce the concentration of the reactant to  $\frac{1}{16}$  th value?

- a)  $4.6 \times 10^{-2}\ s$
- b)  $4.6 \times 10^4\ s$
- c)  $4.6 \times 10^2\ s$
- d)  $4.6 \times 10^{-4}\ s$

317. Rate constant of a chemical reaction can be kept constant by:

- a) Stirring the compounds
- b) Keeping the temperature constant
- c) Both (a) and (b)
- d) None of the above

318. The rate of a chemical reaction doubled for every  $10^\circ\text{C}$  rise in temperature. If the temperature is increased by  $60^\circ\text{C}$  the rate of reaction increase by

- a) 20 times
- b) 32 times
- c) 64 times
- d) 128 times

319. If 'a' is the initial concentration of the reactant, the half-life period of the reaction of nth order is proportional to

- a)  $a^{n-1}$
- b)  $a^n$
- c)  $a^{1-n}$
- d)  $a^{n+1}$




320. Rate of reaction depends upon

- a) temperature
- b) catalyst
- c) concentration
- d) All of these

321. For a reaction, the rate of reaction was found to increase about 1.8 times when the temperature was increased by  $10^\circ\text{C}$ . The increase in rate is due to:

- a) Increase in number of active molecules
- b) Increase in activation energy of reactants
- c) Decrease in activation energy of reactants
- d) Increase in the number of collisions between reacting molecules

322. A hypothetical reaction  $A_2 + B_2 \rightarrow 2AB$  follows the mechanism as given below,



The order of the overall reaction is

- a) 2
- b) 1
- c)  $1\frac{1}{2}$
- d) 0

323. A chemical reaction proceeds following formula

$k = PZe^{-E_a/RT}$



Which of the following process will increase the rate of reaction?

- a) Lowering of  $E_a$
- b) Lowering of P
- c) Lowering of Z
- d) Independent of all the above factors

324. In the respect of the equation  $k = Ae^{-E_a/RT}$  in chemical kinetics, which one of the following statements is correct?

- a) K is equilibrium constant
- b) A is adsorption factor
- c)  $E_a$  is energy of activation
- d) R is Rydberg constant


325. For the reaction  $2\text{NO}_2 + \text{F}_2 \rightarrow 2\text{NO}_2\text{F}$ , following mechanism has been provided,



Thus, rate expression of the above reaction can be written as:

a)  $r = K[\text{NO}_2]^2[\text{F}_2]$       b)  $r = K[\text{NO}_2][\text{F}_2]$       c)  $r = K[\text{NO}_2]$       d)  $r = K[\text{F}_2]$


326. For the reaction:



Then correct statement is/are:

- a) Rate constant for forward reaction =  $2 \times 10^{-4}$   
 b) Rate constant for backward reaction =  $3 \times 10^5$   
 c) Equilibrium constant for the reaction =  $6.6 \times 10^{-10}$   
 d) All of the above

327. For a reaction between gaseous compounds,



The reaction rate =  $k[\text{A}][\text{B}]$ . If the volume of the container is made  $\frac{1}{4}$  of the initial, then what will be the rate of reaction as compared to the initial rate?

- a) 16 times      b) 4 times      c) 1/8 times      d) 1/16 times

328. The rate constant for a first order reaction whose half-life, is 480 s is

- a)  $2.88 \times 10^{-3} \text{s}^{-1}$       b)  $2.72 \times 10^{-3} \text{s}^{-1}$       c)  $1.44 \times 10^{-3} \text{s}^{-1}$       d)  $1.44 \text{s}^{-1}$

329. If  $X$  is the total number of collisions which a gas molecule register with others per unit time under particular conditions, then the collision frequency of the gas containing  $N$  molecules per unit volume is

- a)  $X/N$       b)  $NX$       c)  $2NX$       d)  $NX/2$

330. For a reaction, the rate constant is  $2.34 \text{s}^{-1}$ . The half-life period for reaction is

- a) 0.30 s      b) 0.60 s      c) 3.3 s      d) Data is insufficient

331. If " $a$ " and " $t_{1/2}$ " are initial concentration of reactant and half-life of a zero order reaction respectively, which of the following is correct ?

- a)  $t_{1/2} \propto \frac{1}{a}$       b)  $t_{1/2} \propto a$       c)  $t_{1/2} \propto \frac{1}{a^2}$       d)  $t_{1/2} \propto a^2$

332. The temperature dependence of rate constant ( $k$ ) of a chemical reaction is written in terms of Arrhenius equation,  $k = Ae^{-E^*/RT}$ . Activation energy ( $E^*$ ) of the reaction can be calculated by plotting

- a)  $\log k$  vs  $\frac{1}{T}$       b)  $\log k$  vs  $\frac{1}{\log T}$       c)  $k$  vs  $T$       d)  $k$  vs  $\frac{1}{\log T}$

333. The rate constant of a reaction is given by  $k = 2.1 \times 10^{10} \exp(-2700 RT)$ .

It means that

- a)  $\log k$  vs  $1/T$  will be a curved line with slope =  $-\frac{2700}{2.303}$   
 b)  $\log k$  vs  $1/T$  will be a straight line with intercept on  $\log k$  axis =  $\log 2.1 \times 10^{10}$   
 c) The number of effective collisions are  $2.1 \times 10^{10} \text{cm}^{-3} \text{s}^{-1}$   
 d) Half-life of the reaction increases with increase of temperature

334. The unit of the rate of a second order reaction



- a)  $\text{time}^{-1}$       b)  $\text{mol L}^{-1}\text{time}^{-1}$       c)  $\text{L mol}^{-1}\text{time}^{-1}$       d)  $\text{L}^2 \text{mol}^{-2} \text{time}^{-1}$

335. Rate of a reaction can be expressed of by following rate expression

Rate =  $k[\text{A}]^2[\text{B}]$ , if concentration of A is reduced by half by what times concentration of B is to be increased to have same rate of reaction?

- a) 4 times      b) 2 times      c)  $\frac{1}{4}$  times      d) 8 times

336. Select the intermediate in the following reaction mechanism:



- a)  $\text{O}_3(\text{g})$       b)  $\text{O}(\text{g})$       c)  $\text{O}_2(\text{g})$       d) None of these


337. For a certain reaction of order 'n' the time for half change  $t_{1/2}$  is given by;  $t_{1/2} = \frac{2-\sqrt{2}}{K} \times c_0^{1/2}$ , where K is rate constant  $c_0$  is initial concentration. The value of n is:  
 a) 1                                      b) 2                                      c) 0                                      d) 0.5

338. For a reaction between A and B, the initial rate of reaction is measured for various initial concentrations A and B. the data provided are

	[A]	[B]	Initial reaction rate
1	0.20 M	0.30 M	$5 \times 10^{-5}$
2	0.20 M	0.10 M	$5 \times 10^{-5}$
3	0.40 M	0.05 M	$1 \times 10^{-5}$


The overall order of the reaction is

- a) One                                      b) Two                                      c) Two and half                                      d) Three
339. Which order of reaction obeys the relation  $t_{1/2} = 1/Ka$ ?  
 a) First                                      b) Second                                      c) Third                                      d) Zero
340. How much faster would a reaction proceed at 25°C than at 0°C if the activation energy is 65 kJ?  
 a) 2 times                                      b) 16 times                                      c) 11 times                                      d) 6 times
341. The activation energy of a reaction at a given temperature is found to be  $2.303RT \text{ J mol}^{-1}$ . The ratio of rate constant to the Arrhenius factor is  
 a) 0.01                                      b) 0.1                                      c) 0.02                                      d) 0.001
342. Consider an endothermic reaction  $X \rightarrow Y$  with the activation energies  $E_b$  and  $E_f$  for the backward and forward reactions respectively. In general  
 a) There is no definite relation between  $E_b$  and  $E_f$     b)  $E_b = E_f$   
 c)  $E_b > E_f$                                       d)  $E_b < E_f$
343. For the reaction  $A \rightarrow nB$ , at the point of intersection of two curves show, the [B] is can be given by:



- a)  $\frac{nA_0}{2}$                                       b)  $\frac{A_0}{n-1}$                                       c)  $\frac{nA_0}{n+1}$                                       d)  $\left[\frac{n-1}{n+1}\right] A_0$
344. The elementary step of the reaction,  $2\text{Na} + \text{Cl}_2 = 2\text{NaCl}$  is found to follow III order kinetics, its molecularity is:  
 a) 1                                      b) 2                                      c) 3                                      d) 4
345. Following mechanism has been proposed for a reaction,  
 $2A + B \rightarrow D + E$   
 $A + B \rightarrow C + D$                       ... (Slow)  
 $A + C \rightarrow E$                       ... (fast)  
 The rate law expression for the reaction is:  
 a)  $r = K[A]^2[B]$                       b)  $r = K[A][B]$                       c)  $r = K[A]^2$                       d)  $r = K[A][C]$
346. Two first order reaction have half-lives in the ratio 8 : 1. Calculate the ratio of time intervals  $t_1 : t_2$ . The time  $t_1$  and  $t_2$  are the time period for (1/4)th and (3/4)th completion  
 a) 1 : 0.602                      b) 2 : 301                      c) 0.256 : 0.603                      d) 0.2 : 0.301
347. Order of a reaction can be  
 a) Fractional                      b) Zero                      c) Integer                      d) All of these
348. The half-life period for a zero order reaction is equal to

- a)  $\frac{2k}{[A]_0}$       b)  $\frac{[A]_0}{2k}$       c)  $\frac{0.693}{k}$       d)  $\frac{0.693}{k[A]_0}$
349. In a reaction  $A + B \rightarrow C$ , the rate expression is  $R = k[A][B]^2$ . If the concentration of both the reaction is doubled at constant volume then the rate of the reaction will be  
 a) Eight time      b) Double      c) Quadruple      d) Triple
350. For a gaseous reaction, the units of rate of reaction are  
 a)  $L \text{ atm s}^{-1}$       b)  $\text{atm s}^{-1}$       c)  $\text{atm mol}^{-1} \text{ s}^{-1}$       d)  $\text{mol s}^{-1}$
351. The rate constant is given by the equation  $K = Ae^{-E_a/RT}$  which factor should register a decrease for the reaction to proceed more rapidly?  
 a)  $T$       b)  $Z$       c)  $A$       d)  $E_a$
352. The activation energy for most of the reactions is approximately  $50 \text{ kJ mol}^{-1}$ . The value of temperature coefficient for such reactions is  
 a)  $> 2$       b)  $> 3$       c)  $< 1$       d)  $> 4$
353. The half-life period for a first order reaction is 693 s. The rate constant of this reaction would be  
 a)  $0.1 \text{ s}^{-1}$       b)  $0.01 \text{ s}^{-1}$       c)  $0.001 \text{ s}^{-1}$       d)  $0.0001 \text{ s}^{-1}$
354. For the reaction  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$  under certain condition of temperature and partial pressure of the reactants, the rate of formation of  $\text{NH}_3$  is  $10^{-3} \text{ kg hr}^{-1}$ . The rate of conversion of  $\text{H}_2$  under same condition is:  
 a)  $1.5 \times 10^{-3} \text{ kg hr}^{-1}$       b)  $1.76 \times 10^{-4} \text{ kg hr}^{-1}$       c)  $2 \times 10^{-3} \text{ kg hr}^{-1}$       d)  $3 \times 10^{-3} \text{ kg hr}^{-1}$
355. In a first order reaction  $A \rightarrow B$ , if  $K$  is the rate constant initial concentration of the reactant is  $0.5 \text{ M}$ , then half-life is:  
 a)  $\frac{\ln 2}{K}$       b)  $\frac{\ln 2}{K\sqrt{0.5}}$       c)  $\frac{\log_{10} 2}{K}$       d)  $\frac{0.693}{0.5 K}$
356. A reaction  $A \rightarrow B$  follows a second order kinetics. Doubling the concentration of A will increase the rate of formation of B by a factor of  
 a)  $\frac{1}{4}$       b) 4      c)  $\frac{1}{2}$       d) 2
357. With increase in temperature, rate of reaction  
 a) increases      b) decreases  
 c) Remains same      d) May increase or decrease
358. Which of the following statement is not correct?  
 a) In zero order reaction the rate of the reaction remains constant throughout.  
 b) A second order reaction would become a pseudo first order reaction when one of the reactant is taken in large excess.  
 c) The value of first order rate constant depends on the units of the concentration term used.  
 d) In a first order reaction the plot of  $\log(a-x)$  vs time gives a straight line.
359. A drop of solution (volume  $0.05 \text{ mL}$ ) contains  $3.0 \times 10^{-6}$  mole of  $\text{H}^+$ . If the rate constant of disappearance of  $\text{H}^+$  is  $1.0 \times 10^{-7} \text{ mol litre}^{-1} \text{ sec}^{-1}$ . How long would it take for  $\text{H}^+$  in drop to disappear?  
 a)  $6 \times 10^{-8} \text{ sec}$       b)  $6 \times 10^{-7} \text{ sec}$       c)  $6 \times 10^{-9} \text{ sec}$       d)  $6 \times 10^{-10} \text{ sec}$
360. For the reaction,  $2A + B \rightarrow \text{products}$ , the active mass of B is kept constant, and that of A is doubled. The rate of reaction will be then  
 a) Decrease 4 times      b) Decrease 2 times      c) Increase 4 times      d) Increase 2 times
361. Which of the following expression is correct for second order reaction. ( $C_0$  refers to initial concentration of reactant)?  
 a)  $t_{1/2} \propto C_0$       b)  $t_{1/2} \propto C_0^0$       c)  $t_{1/2} \propto C_0^{-1}$       d)  $t_{1/2} \propto C_0^{-2}$
362. A first order reaction is 10% complete in 20 min. The time taken for 19% completion is  
 a) 30 min      b) 40 Min      c) 50 min      d) 38 min
363. A graph plotted between  $\log k$  versus  $\frac{1}{T}$  for calculating activation energy is shown by




364. In the following reaction  $A \rightarrow B + C$ , rate constant is  $0.001 \text{ Ms}^{-1}$ . Half-life and completion time of the given reaction are  
 a) 500 s, 1000 s                      b) 500 s, 750 s                      c) 250 s, 500 s                      d) 300 s, 600 s
365. A reaction was found to order with respect to the concentration of carbon monoxide. If the concentration of carbon monoxide is doubled, with everything else kept the same, the rate of reaction will  
 a) Remain unchanged                      b) Triple  
 c) Increases by factor of four                      d) Double
366. The following data were obtained the first order decomposition of  $2 A(g) \rightarrow B(g) + C(S)$  at a constant volume and at a particular temperature

S N	Time	Total pressure in Pascal
1	At the end of 10 min	300
2	After completion	200

The rate constant in  $\text{min}^{-1}$  is

- a) 0.0693                      b) 69.3                      c) 6.93                      d)  $6.93 \times 10^{-4}$
367. According to Arrhenius equation, the rate constant ( $k$ ) is related to temperature ( $T$ ) as  
 a)  $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$                       b)  $\ln \frac{k_2}{k_1} = -\frac{E_a}{R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$   
 c)  $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left[ \frac{1}{T_1} + \frac{1}{T_2} \right]$                       d)  $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left[ \frac{1}{T_1} + \frac{1}{T_2} \right]$
368. Inversion of cane-sugar in dilute acid is a  
 a) Bimolecular reaction                      b) Pseudo-unimolecular reaction  
 c) Unimolecular reaction                      d) Trimolecular reaction

369. Consider the reaction




The equality relationship between  $\frac{d[\text{NH}_3]}{dt}$  and  $-\frac{d[\text{H}_2]}{dt}$  is:

- a)  $+\frac{d[\text{NH}_3]}{dt} = -\frac{3}{2} \frac{d[\text{H}_2]}{dt}$   
 b)  $\frac{d[\text{NH}_3]}{dt} = -\frac{d[\text{H}_2]}{dt}$   
 c)  $\frac{d[\text{NH}_3]}{dt} = -\frac{1}{3} \frac{d[\text{H}_2]}{dt}$   
 d)  $+\frac{d[\text{NH}_3]}{dt} = -\frac{2}{3} \frac{d[\text{H}_2]}{dt}$
370. For a reaction  $aA \rightarrow bB$  when  $[A] = 2.2 \text{ mM}$ , the rate was found to be  $2.4 \text{ mM s}^{-1}$ . On reducing concentration of  $[A]$  to half, the rate changes to  $0.6 \text{ mM s}^{-1}$ . The order of reaction with respect to  $A$  is:  
 a) 1.5                      b) 2.0                      c) 2.5                      d) 3.0
371. The units of the rate constant of a second order reaction are  
 a)  $\text{mol}^{-1} \text{L}^{-1} \text{s}^{-1}$                       b)  $\text{mol}^{-1} \text{L s}^{-1}$                       c)  $\text{mol}^{-1} \text{L s}$                       d)  $\text{mol L}^{-1} \text{s}^{-1}$
372. A follows first order reaction,  $A \rightarrow \text{product}$  Concentration of  $A$ , change from  $0.1 \text{ M}$  to  $0.025 \text{ M}$  in  $40 \text{ min}$ . find the rate of reaction of  $A$  when concentration of  $A$  is  $0.01 \text{ M}$ .  
 a)  $3.47 \times 10^{-4} \text{ M min}^{-1}$   
 b)  $3.47 \times 10^{-5} \text{ M min}^{-1}$   
 c)  $1.73 \times 10^{-4} \text{ M min}^{-1}$



- d)  $1.73 \times 10^{-5} M \text{ min}^{-1}$
373. In the reaction  $2A + B \rightarrow A_2B$ , if the concentration of  $A$  is doubled and of  $B$  is halved, then the rate of the reaction will
- Increase by two times
  - Decrease by two times
  - Increase by four times
  - Remain the same
374. Energy of activation of an exothermic reaction is
- Negative
  - Positive
  - Zero
  - Can't be predict
375. For a reaction, the rate constant is  $2.34s^{-1}$ . The half-life period for the reaction is
- 0.30 s
  - 0.60 s
  - 3.3 s
  - Data is insufficient
376. The rate of a reaction get doubles when the temperature changes from  $7^\circ\text{C}$  to  $17^\circ\text{C}$ . By what factor will it change for the temperature change from  $17^\circ\text{C}$  to  $27^\circ\text{C}$ ?
- 1.81
  - 1.71
  - 1.91
  - 1.76
377. Arrhenius equation is
- $\Delta H = \Delta E + \Delta n_g RT$
  - $\Delta G = \Delta H - T \cdot \Delta S$
  - $k = Ae^{-E_a/RT}$
  - None of these
378. Which rate expression suggests an over all order of 0.5 for the reaction involving substances  $X, Y, Z$ ?
- Rate =  $K[X][Y][Z]$
  - Rate =  $K[X]^{0.5}[Y]^{0.5}[Z]^{0.5}$
  - Rate =  $K[X]^{1.5}[Y]^{-1}[Z]^0$
  - Rate =  $K[X][Y]^0/[Z]^2$
379. Rate of a reaction can be expressed by Arrhenius equation as  
 $k = Ae^{-E/RT}$   
 In this equation,  $E$  represents
- The energy above which all the colliding molecules will react
  - The energy below which colliding molecules will not react
  - The total energy of the reacting molecules at a temperature,  $T$
  - The fraction of molecules with energy greater than the activation energy of the reaction
380. The minimum energy required for a molecule to take part in a reaction is called
- Threshold energy
  - Nuclear energy
  - Potential energy
  - Kinetic energy
381. The rate of reaction becomes 2 times for every  $10^\circ\text{C}$  rise in temperature. How the rate of reaction will increase when temperature is increased from  $30^\circ\text{C}$  to  $80^\circ\text{C}$ ?
- 16
  - 32
  - 64
  - 128
382. In a gaseous phase reaction:  
 $A_2(g) \rightarrow B(g) + (1/2)C(g)$ , the increase in pressure from 100 mm to 120 mm is noticed in 5 minute. The rate of disappearance of  $A_2$  in  $\text{mm min}^{-1}$  is:
- 4
  - 8
  - 16
  - 2
383. The unit of rate constant of second order reaction is
- $\text{Mol/Ls}$
  - $\text{L / Mol s}$
  - $\text{L}^2 / \text{mol}^2$
  - Per second
384. The given reaction,  
 $2\text{FeCl}_3 + \text{SnCl}_2 \rightarrow \text{SnCl}_4 + 2\text{FeCl}_2$  is an example of:
- First order reaction
  - Third order reaction
  - Second order reaction
  - None of these
385. For the reaction  $A + B \rightarrow C$ , it is found that doubling the concentration of  $A$  increases the rate by four times and doubling the concentration of  $B$  doubles the reaction rate. What is the overall order of the reaction?
- 4
  - $3/2$
  - 3
  - 1
386. The rate constant  $K_1$  of a reaction is found to be double that of rate constant  $K_2$  of another reaction. The relationship between corresponding activation energies of the two reactions at same temperature ( $E_1$  and  $E_2$ ) can be represented as:

- a)  $E_1 > E_2$                       b)  $E_1 < E_2$                       c)  $E_1 = E_2$                       d) None of these
387. For the reaction,  $H_2 + I_2 \rightarrow 2HI$ , the differential rate law is
- a)  $-\frac{d[H_2]}{dt} = -\frac{d[I_2]}{dt} = \frac{1}{2} \frac{d[HI]}{dt}$                       b)  $-\frac{d[H_2]}{dt} = -2 \frac{d[I_2]}{dt} = \frac{1}{2} \frac{d[HI]}{dt}$
- c)  $-\frac{d[H_2]}{dt} = -\frac{d[I_2]}{dt} = \frac{d[HI]}{dt}$                       d)  $-\frac{d[H_2]}{dt} = -\frac{d[I_2]}{dt} = -\frac{d[HI]}{dt}$
388. The time for half-life of a first order reaction is 1 hr. What is the time taken for 87.5% completion of the reaction?
- a) 1 hour                      b) 2 hour                      c) 3 hour                      d) 4 hour
389. DDT on exposure to water decomposes. Half-life is 10 yr. How much time it will take for its decomposition to 99%?
- a) 50 yr                      b) 70 yr                      c) 500 yr                      d) 700 yr
390. In Arrhenius equation,  $k = Ae^{-E_a/RT}$ ;  $A$  may be called the rate constant at
- a) Very low temperature                      b) Zero activation energy
- c) The boiling temperature of reaction mixture                      d) All of the above
391. The phenomenon of emission of visible light as a result of chemical change is known as
- a) Chemiluminescence                      b) Fluorescence
- c) Phosphorescence                      d) Photosensitization
392. Chemical reactions with very high  $E_a$  values are generally
- a) Very fast                      b) Very slow                      c) Moderately fast                      d) Spontaneous
393. In the reaction  $3A \rightarrow 2B$ , rate of reaction  $+\frac{d(B)}{dt}$  is equal to
- a)  $-\frac{1}{3} \frac{d[A]}{dt}$                       b)  $-\frac{2}{3} \frac{d[A]}{dt}$                       c)  $+\frac{2d[A]}{dt}$                       d)  $-\frac{3}{2} \frac{d[A]}{dt}$
394. If the half-time for a particular reaction is found to be constant and independent of the initial concentration of the reactants, then the ratio is of
- a) First order                      b) Zero order                      c) Second order                      d) None of these
395. Under the same reaction condition, initial concentration of  $1.386 \text{ mol dm}^{-3}$  of a substance become half in 40 s and 20 s through first order and zero order kinetics respectively. Ratio  $\left(\frac{k_1}{k_0}\right)$  of the rate constants for first order ( $k_1$ ) and zero order ( $k_0$ ) of the reaction is
- a)  $0.5 \text{ mol}^{-1} \text{ dm}^{-3}$                       b)  $1.0 \text{ mol}^{-1} \text{ dm}^{-3}$                       c)  $1.5 \text{ mol}^{-1} \text{ dm}^{-3}$                       d)  $2 \text{ mol}^{-1} \text{ dm}^{-3}$
396. The order of a reaction with rate equal to  $kC_A^{3/2} C_B^{-1/2}$  is
- a) 1                      b)  $-\frac{1}{2}$                       c)  $-\frac{3}{2}$                       d) 2
397. For reaction  $A \rightarrow B$ , the rate constant  $k_1 = A_1 e^{-E_{a1}/RT}$  and for the reaction  $X \rightarrow Y$ , the rate constant  $k_2 = A_2 e^{-E_{a2}/RT}$ . If  $A_1 = 10^8$ ,  $A_2 = 10^{10}$  and  $E_{a1} = 600 \text{ cal/mol}$ ,  $E_{a2} = 1800 \text{ cal/mol}$ , then the temperature at which  $k_1 = k_2$  is ( $R = 2 \text{ cal/k mol}$ )
- a)  $\frac{200}{3.506} \text{ K}$                       b)  $\frac{300}{2.303} \text{ K}$                       c)  $\frac{400}{6.506} \text{ K}$                       d)  $\frac{200}{5.204} \text{ K}$
398. Which of these does not influence the rate of reaction?
- a) Nature of the reactants                      b) Concentration of the reactants
- c) Temperature of the reaction                      d) Molecularity of the reaction
399. The concentration of R in the reaction  $R \rightarrow P$  was measured as a function of time and the following data is obtained
- |            |      |      |      |      |
|------------|------|------|------|------|
| [R](molar) | 1.00 | 0.75 | 0.40 | 0.10 |
| t (min)    | 0.00 | 0.05 | 0.12 | 0.18 |
- The order of the reaction is
- a) Zero                      b) First                      c) Second                      d) Third
400. A graph plotted between concentration of reactant consumed at any time ( $x$ ) and time  $t$  is found to be a straight line passing through the origin. Thus, reaction is of:

- a) First order                      b) Zero order                      c) Third order                      d) Second order
401. Rate constant of a reaction depends upon  
 a) Speed of reaction                      b) Concentration of the reactants  
 c) Pressure of the surrounding                      d) Temperature
402. An endothermic reaction  $A \rightarrow B$  has an activation energy as  $x \text{ kg mol}^{-1}$  of  $A$ . If energy change of the reaction is  $y \text{ kJ}$ , the activation energy of the reverse reactions is  
 a)  $-x$                       b)  $y - x$                       c)  $x + y$                       d)  $x - y$
403. Consider the following statement in respect of zero order reaction.  
 III. The rate of the reaction is independent of reactant concentration.  
 IV. The rate of the reaction is independent of temperature.  
 V. The rate constant of the reaction is independent of temperature.  
 VI. The rate constant of reaction is independent of reactant cogeneration.  
 Choose the correct statements/s  
 a) I only                      b) I and II only                      c) III and IV only                      d) I and IV only
404. If concentration of reactants is increased by 'X', the rate constant  $K$  becomes:  
 a)  $e^{K/X}$                       b)  $K/X$                       c)  $K$                       d)  $X/K$
405. The correct expression for the rate of reaction of elementary reaction,  $A + B \rightarrow C$  is:  
 a)  $\frac{d[C]}{dt} = K[A]$                       b)  $\frac{d[C]}{dt} = K[B]$                       c)  $\frac{-d[A]}{dt} = K[A][B]$                       d)  $\frac{-d[A]}{dt} = K[A]$
406. The threshold energy of a chemical reaction depends upon:  
 a) Nature of reacting species  
 b) Temperature  
 c) Concentration of species  
 d) Number of collisions per unit time or collision frequency
407. A first order reaction has a half-life period of 69.3 sec. At  $0.10 \text{ mol litre}^{-1}$  reactant concentration, rate will be:  
 a)  $10^{-4} \text{ M sec}^{-1}$                       b)  $10^{-3} \text{ M sec}^{-1}$                       c)  $10^{-1} \text{ M sec}^{-1}$                       d)  $6.93 \times 10^{-1} \text{ M sec}^{-1}$
408. The rate constant for the reaction  $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$  is  $3.0 \times 10^{-5} \text{ s}^{-1}$ . If the rate is  $2.40 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$ , then the concentration of  $\text{N}_2\text{O}_5$  is  $\text{mol L}^{-1}$  is  
 a) 0.4                      b) 0.8                      c) 1.2                      d) 3.2
409. The enzyme catalysed reaction is faster than metal catalysed reaction because its activation energy is:  
 a) Greater                      b) Lower                      c) Same                      d) None of these
410. Plots showing the variation of the rate constant ( $k$ ) with temperature ( $T$ ) are given below. The plot that follows Arrhenius equation is  

411. For the following homogeneous reaction, the unit of rate constant is  $A + B \xrightarrow{k} C$   
 a)  $\text{mol}^{-1} \text{L s}^{-1}$                       b)  $\text{s}^{-1}$                       c)  $\text{S}$                       d)  $\text{s}^{-1} \text{mol L}^{-1}$
412. If 'a' is the initial concentration of a substance which reacts according to zero order kinetics and  $K$  is rate constant, the time for the reaction to go to completion is:  
 a)  $a/K$                       b)  $2/Ka$                       c)  $K/a$                       d)  $2K/a$
413. What is the two third life of a first order reaction having  $= 5.48 \times 10^{-14} \text{ s}^{-1}$  ?  
 a)  $2.01 \times 10^{11} \text{ s}$                       b)  $2.01 \times 10^{13} \text{ s}$                       c)  $8.08 \times 10^{13} \text{ s}$                       d)  $16.04 \times 10^{11} \text{ s}$
414. The velocity constant of a reaction at 290 K was found to be  $3.2 \times 10^{-3} \text{ s}^{-1}$ . When the temperature is raised to 310 K, it will be about

- a)  $6.4 \times 10^{-3}$                       b)  $3.2 \times 10^{-4}$                       c)  $9.6 \times 10^{-3}$                       d)  $1.28 \times 10^{-2}$
415. For fourth order reaction, what is the unit of K?  
 a)  $\left(\frac{\text{mol}}{\text{L}}\right)^{-3} \text{ s}^{-1}$                       b)  $\left(\frac{\text{mol}}{\text{L}}\right)^{+3} \text{ s}^{-1}$                       c)  $\left(\frac{\text{mol}}{\text{L}}\right)^{-3} \text{ s}$                       d)  $\left(\frac{\text{mol}}{\text{L}}\right)^{-3}$
416. The branch of chemistry which deals with the reaction rates and reaction mechanism is called:  
 a) Thermochemistry                      b) Photochemistry                      c) Analytical chemistry                      d) Chemical kinetics
417. For the reaction;  $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ , rate and rate constant are  $1.02 \times 10^{-4} \text{ M sec}^{-1}$  and  $3.4 \times 10^{-5} \text{ sec}^{-1} \text{ M}$  respectively, then concentration of  $\text{N}_2\text{O}_5$ , at that time will be:  
 a)  $1.732 \text{ M}$                       b)  $3 \text{ M}$                       c)  $1.02 \times 10^{-4} \text{ M}$                       d)  $3.5 \times 10^5 \text{ M}$
418. In the following first order competing reactions:  
 $A + \text{Reagent} \rightarrow \text{Product}$   
 $B + \text{Reagent} \rightarrow \text{Product}$   
 Find the ratio of  $K_1/K_2$  if only 50% of B will have been reacted when 94% of A has been reacted in same time is:  
 a) 4.06                      b) 0.246                      c) 2.06                      d) 0.06
419. For which order half-life period is independent of initial concentration?  
 a) Zero                      b) First                      c) Second                      d) Third
420. For a given reaction, pressure of catalyst reduces the energy of activation by 2 kcal at  $27^\circ\text{C}$ . The rate of reaction will be increased by:  
 a) 20 times                      b) 14 times                      c) 28 times                      d) 2 times
421. For a reversible reaction  $A \xrightleftharpoons[K_2]{K_1} B$ ,  
 1st order in both the directions, the rate of reaction is given by:  
 a)  $K_1[A]$   
 b)  $-K_2[B]$   
 c)  $K_1[A] + K_2[B]$   
 d)  $K_1[A] - K_2[B]$
422. A first order reaction is 20% complete in 10 min. Calculate the time for 75% completion of the reaction  
 a) 0.233 min                      b) 62.18 min                      c) 112.12 min                      d) 36.18 min
423. Order of radioactive disintegration reaction is  
 a) Zero                      b) First                      c) Second                      d) Third
424. Ethylene is produced by cyclobutane as:  
 $\text{C}_4\text{H}_8 \xrightarrow{\Delta} 2\text{C}_2\text{H}_4$   
 The rate constant is  $2.48 \times 10^{-4} \text{ sec}^{-1}$ . In what time will the molar ratio of the ethylene to cyclobutane in reaction mixture attain the value 1?  
 a) 27.25 minute                      b) 28.25 minute                      c) 25 minute                      d) 20 minute
425. For a reaction for which the activation energies of forward and reverse reactions are equal?  
 a)  $\Delta H = 0$                       b)  $\Delta S = 0$                       c) The order is zero                      d) There is no catalyst
426. The half-life period of a first order chemical reaction is 6.93 min. the time required for the completion of 99% of the chemical reaction will be ( $\log 2 = 0.302$ )  
 a) 230.3 min                      b) 23.03 min                      c) 46.06 min                      d) 460.6 min
427. The rate of the elementary reaction,  $2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$ , when the volume of the reaction vessel is doubled:  
 a) Will grow eight times of its initial rate  
 b) Reduce to one-eight of its initial rate  
 c) Will grow four times of its initial rate  
 d) Reduce to one-fourth of its initial rate
428. For the reaction system

$2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}_2(\text{g})$  if the volume of the reaction vessel is reduced to one-third of its original volume, what will be the order of the reaction?

- a) Diminished to one fourth of its initial value      b) Diminished to one twenty seven of its initial value  
 c) Increase to twenty seven times of its initial value      d) Increase to four times of its initial value
429. The rate constant of a second order reaction is  $10^{-2} \text{M s}^{-1}$ . The rate constant expressed in  $\text{cc. molecule}^{-1} \text{min}^{-1}$  is:  
 a)  $9.96 \times 10^{-22}$       b)  $9.96 \times 10^{-23}$       c)  $9.96 \times 10^{-21}$       d)  $1.004 \times 10^{-24}$
430. Radioactive decay is a  
 a) First order reaction      b) Zero order reaction  
 c) Second order reaction      d) Third order reaction
431. For the decomposition of  $\text{N}_2\text{O}_5$  at a particular temperature, according to the equations  
 $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$   
 $\text{N}_2\text{O}_5 \rightarrow 2\text{NO}_2 + \frac{1}{2}\text{O}_2$   
 The activation energies are  $E_1$  and  $E_2$  respectively then  
 a)  $E_1 > E_2$       b)  $E_1 < E_2$       c)  $E_1 = 2E_2$       d)  $E_1 = E_2$
432. The rate of a gaseous reaction is equal to  $k[A][B]$ . The volume of the reaction vessel containing these gases is reduced by one-fourth of the initial volume. The rate of the reaction would be  
 a)  $\frac{1}{16}$       b)  $\frac{16}{1}$       c)  $\frac{1}{8}$       d)  $\frac{8}{1}$
433. The rate law of the reaction,  
 $A + 2B \rightarrow \text{Product}$ .  
 Product is given by  $\frac{d[P]}{dt} = K[A]^2 \cdot [B]$ . If  $A$  is taken in large excess, the order of the reaction will be:  
 a) Zero      b) 1      c) 2      d) 3
434. A first order reaction has a rate constant  $1.15 \times 10^{-3} \text{s}^{-1}$ . How long will 5g of this reactant take to reduce to 3 g?  
 a) 444 s      b) 402 s      c) 442 s      d) None of these
435. For a certain reaction a plot of  $\frac{[c_0 - c]}{c}$  against time  $t$ , yields a straight line.  $c_0$  and  $c$  are concentrations of reactant at  $t = 0$  and  $t = t$  respectively. The rate of reaction is:  
 a) 3      b) 0      c) 1      d) 2
436. The rate constant is doubled when temperature increases from  $27^\circ\text{C}$  to  $37^\circ\text{C}$ . Activation energy in kJ is  
 a) 34      b) 54      c) 100      d) 53
437. If the concentration of reactants is increased by ' $x$ ' then rate constant  $k$  becomes  
 a)  $\ln \frac{k}{x}$       b)  $\frac{k}{x}$       c)  $k + x$       d)  $k$