

GPLUS EDUCATION






Date :
Time :
Marks :

CHEMISTRY

HYDROCARBONS


Single Correct Answer Type

1. Thermal decomposition of




2. Which of the following is not a petroleum product?
a) Petrol b) Paraffin wax c) Bees wax d) Kerosene
3. A knocking sound is produced more in the engine when the fuel contains mainly:
a) *n*-alkanes b) CO₂ c) CO d) Lubricating oil
4. Reaction of HBr with propene in presence of peroxides gives:
a) Isopropyl bromide b) 3-bromopropane c) Allyl bromide d) *n*-propyl bromide
5. The next higher homologue of C₆H₁₄ is:
a) C₇H₁₄ b) C₇H₁₆ c) C₇H₁₀ d) C₇H₁₂
6. The reaction conditions used for converting 1,2-dibromopropane to propylene are
a) KOH, alcohol/Δ b) KOH, water/Δ c) Zn, alcohol/Δ d) Na, alcohol/Δ
7. A gas formed by the action of alcoholic KOH on ethyl iodide, decolourises alkaline KMnO₄. The gas is
a) C₂H₆ b) CH₄ c) C₂H₂ d) C₂H₄
8. Alkyne, C₇H₁₂, when reacted with alkaline KMnO₄ followed by acidification with HCl gives a mixture of (CH₃)₂CHCOOH + CH₃CH₂COOH, The alkyne C₇H₁₂ is
a) 3-hexyne b) 2-methyl-2-hexene c) 2-methyl-3-hexene d) 3-methyl-2-hexyne
9. The relationship between acetylene and benzene is comparable to the relationship between propyne and
a) Dimethyl benzene b) Neoprene c) Propyl benzene d) Mesitylene
10. Complete oxidation of one mole of an alkane forms 3 moles of CO₂. The alkane is
a) CH₄ b) C₂H₆ c) C₃H₈ d) C₆H₁₄
11. The ozonolysis of ethylene, acetylene and propylene respectively gives:
a) HCHO, CHO—CHO and CH₃CHO + HCHO
b) CHO—CHO, HCHO and CH₃CHO
c) HCHO + CH₃CHO, CHO—CHO and HCHO
d) CHO—CHO, CH₃CHO + HCHO and HCHO
12. The reaction, CH₂ = CH₂ + CH₃COCl $\xrightarrow{\text{AlCl}_3}$ gives the product:
a) CH₃COCH₂CH₂Cl
b) CH₃.CH₂.CH₂Cl
c) CH₃COCH₂.CH₂COCH₃
d) ClCH₂CH₂Cl
13. Alkyl halides react with dialkyl copper reagents to give
a) Alkenyl halides b) Alkanes
c) Alkyl copper halides d) Alkenes

14. The gas which is used for the artificial ripening of fruits is:
 a) C_2H_6 b) C_2H_2 c) C_2H_4 d) Marsh gas
15. $CH_3-C \equiv CH$ reacts with HCl to give:
 a) 2,2-dichloropropane b) 1,1-dichloropropane c) 1,2-dichloropropane d) 1-chloropropene
16. $CH_3CH_3 + HNO_3 \xrightarrow{675\text{ K}} ?$
 a) $CH_3CH_2NO_2$ b) $CH_3CH_2NO_2 + CH_3NO_2$
 c) $2CH_3NO_2$ d) $CH_2 = CH_2$
17. Which of the following is produced when coal is subjected to destructive distillation?
 a) Methane b) Ethane c) Acetylene d) Coal gas
18. The product of the following reaction are:
 $CH_3C \equiv C \cdot CH_2CH_3 \xrightarrow[\text{(ii) Hydrolysis}]{\text{(i) } O_3} ?$
 a) $CH_3COOH + CH_3COCH_3$
 b) $CH_3COOH + CH_3CH_2COOH$
 c) $CH_3CHO + CH_3CH_2CHO$
 d) $CH_3COOH + CO_2$
19. Methyl bromide heated with zinc in closed tube produces:
 a) Methane b) Ethane c) Ethylene d) Methanol
20. Aqueous solution of an organic compound, 'A' on electrolysis liberates acetylene and CO_2 at a node. 'A' is
 a) Potassium acetate b) Potassium succinate
 c) Potassium citrate d) Potassium maleate
21. The reaction of alkanes with halogen is explosive in the case of:
 a) F_2 b) Cl_2 c) I_2 d) Br_2
22. Which of the following is unsymmetrical alkene?
 a) 1-butene b) 2-hexene c) 1-pentene d) All of these
23. Which of the statement is wrong for alkanes?
 a) Most of the alkanes are soluble in water
 b) Their density is always less than water
 c) At room temperature some alkanes are liquid, some solid and other are gases
 d) All alkanes burn
24. Propane cannot be prepared from which reaction?
 a) $CH_3 - CH = CH_2 \xrightarrow[\text{OH}^-]{B_2H_6}$ b) $CH_3CH_2CH_2I \xrightarrow[P]{HI}$
 c) $CH_3CH_2CH_2COONa \xrightarrow{NaOH/CaO, \Delta}$ d) None of the above
25. Nitrating mixture is
 a) Fuming nitric acid
 b) Mixture of conc. H_2SO_4 and conc. HNO_3
 c) Mixture of nitric acid and anhydrous zinc chloride
 d) None of the above
26. Cyclohexene on reaction with OsO_4 followed by reaction with $NaHSO_3$ gives
 a) *cis* - diol b) *trans* - diol c) Epoxy d) Alcohol
27. Al_4C_3 on hydrolysis yields
 a) Nitrogen gas b) Methane gas c) Hydrogen gas d) Carbon dioxide
28. The compounds P, Q and S



where separately subjected to nitration using $\text{HNO}_3/\text{H}_2\text{SO}_4$ mixture. The major product formed in each case respectively, is





29. Which of the following is not a mixture of hydrocarbons?


- a) Candle wax b) Kerosene c) Vegetable oils d) Paraffin oil

30. C_8H_{10} (A) $\xrightarrow{\text{O}_3/\text{H}_2\text{O}}$ acid (B)





$\text{C}_3\text{H}_5\text{MgBr}$ (C) $\xrightarrow{\text{CO}_2, \text{H}_3\text{O}^+}$ acid B

Identify A, B and C





31. Which of the following has the maximum heat of hydrogenation?



32. $\text{CH}_3\text{CH}_2\text{CH}_3 \xrightarrow{400-600^\circ\text{C}} X + Y, X \text{ and } Y \text{ are}$

a) Hydrogen and methane

b) Hydrogen and ethylene

c) Ethylene and methane

d) Any of these

33. Position of double bond in alkenes is identified by


a) Ozonolysis

b) Bromine water

c) Ammonical silver nitrate

d) None of these

34. Consider the following reaction



I. $\text{H}_2/\text{Ni}_2\text{B}$

II. $\text{H}_2/\text{Pd} - \text{CaCO}_3$ in quinoline

III. Na/NH_3 or LiAlH_4

This reaction takes place by

a) I or II

b) I or III

c) II or III

d) I, II or III

35. Which of the following reagent can distinguish between 1-butyne and 2-butyne?

a) Aqueous NaOH

b) Bromine water

c) Fehling's solution

d) Ammoniacal AgNO_3

36. CH_4 is formed when:

a) Sodium acetate is heated with soda lime

b) Iodo methane is reduced

c) Aluminium carbide reacts with water

d) All of the above

37. Reaction of HBr with propene in the presence of peroxide gives






a) *iso*-propyl bromide

b) 3-bromo propane

c) Allyl bromide

d) *n*-propyl bromide

38. Predict structure of X in following reaction



39. The middle oil fraction of coal-tar distillation contains:

a) Benzene

b) Anthracene

c) Naphthalene

d) Xylene

40. On halogenation, an alkane (C_5H_{12}) gives only one monohalogenated product. The alkane is

a) *n*-pentane

b) 2-methyl butane

c) 2, 2-dimethyl propane

d) Cyclopentane

41. Acrylic emulsion in paints is a polymer of:

a) $\text{CH}_2 = \text{CH} - \text{COOCH}_3$





b) $\text{CH}_3 - \text{CH} = \text{CH} - \text{COOCH}_3$

- c) $\text{CH}_2 = \text{CH} - \text{COOH}$
 d) $\text{CH}_2 = \text{C}(\text{CH}_3) - \text{COOCH}_3$
42. A hydrocarbon X adds on one mole of hydrogen to give another hydrocarbon and decolourised bromine water. X react with KMnO_4 in presence of acid to give two mole of the same carboxylic acid. The structure of X is:
- a) $\text{CH}_3\text{CH} = \text{CHCH}_2\text{CH}_2\text{CH}_3$
 b) $\text{CH}_3\text{CH}_2\text{CH} = \text{CHCH}_2\text{CH}_3$
 c) $\text{CH}_3\text{CH}_2\text{CH}_2 - \text{CH} = \text{CHCH}_3$
 d) $\text{CH}_2 = \text{CH} - \text{CH}_2\text{CH}_2\text{CH}_3$
43. An anaesthetic narcylene is commercial name of:
- a) C_2H_4 b) C_2H_2 c) CHCl_3 d) ether
44. By which one of the following compounds both CH_4 and $\text{CH}_3 - \text{CH}_3$ can be prepared in one step?
- a) CH_3I b) CH_3OH c) $\text{CH}_3\text{CH}_2\text{I}$ d) $\text{C}_2\text{H}_5\text{OH}$
45. What volume of methane (NTP) is formed from 8.2 g of sodium acetate by fusion with sodalime?
- a) 10 litre b) 11.2 litre c) 5.6 litre d) 2.24 litre
46. When methyl iodide is treated with sodium in ethereal solution, it gives
- a) Methane b) Ethane
 c) Methyl sodium iodide d) Sodium methoxide
47. 2-methylpentene 2 on ozonolysis will give:
- a) Only propanal
 b) Propanal and ethanal
 c) Propanone-2 and ethanal
 d) Propanone-2 and propanal
48. The reaction,

$$2\text{RC}\equiv\text{CCu} \xrightarrow[\text{Pyridine}]{(\text{CH}_3\text{COO})_2\text{Cu}} \text{R}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{R}$$
- a) Eglinton's reaction
 b) Glaser reaction
 c) Gomberg-Beckmann's reaction
 d) Leuckart reaction
49. 2-Hexyne gives *trans*-2-hexene on treatment with:
- a) Li/NH_3 b) Pd/BaSO_4 c) LiAlH_4 d) Pt/H_2
50. Which of the following will give three mono-bromo derivatives?
- a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_3$ b) $\text{CH}_3\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_3$
 c) $\text{CH}_3\text{CH}_3(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}_3$ d) All the above can give
51. The reagent for the following conversion
 $\text{Br}-\text{C}\equiv\text{C}-\text{Br} \rightarrow \text{H}-\text{C}\equiv\text{C}-\text{H}$ is/are :
- a) Alc. KOH b) Alc. KOH followed by NaNH_2 c) Aqueous KOH followed by NaNH_2 d) $\text{Zn}/\text{CH}_3\text{OH}$
52. In a reaction if half of the double bond is broken and two new bonds are formed, this is a case of:
- a) Elimination b) Addition c) Displacement d) Rearrangement
53. Which represents a cyclic alkane?
- a) C_3H_6 b) C_3H_8 c) C_8H_{10} d) C_8H_{12}
54. $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_3 \xrightarrow[\text{HBr}]{\text{AlCl}_3}$ Product
 Product in the above reaction is
- a) $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{CH}_3 \\ | \\ \text{Br} \end{array}$ b) $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$
 c) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2$ d) All of these




55. According to Huckel's rule an aromatic compound must possess
 a) $(4n + 1)\pi$ -electrons
 b) $(4n + 2)\pi$ -electrons
 c) $4n \pi$ -electrons
 d) $(4n + 3)\pi$ -electrons
56. Acetylene gives:
 a) White ppt. with AgNO_3 and red ppt. with Cu_2Cl_2
 b) White ppt. with Cu_2Cl_2 and red ppt. with AgNO_3
 c) White ppt. with both
 d) Red ppt. with both
57. 1,1,2,2-tetrabromoethane on heating with Zn powder in alcohol finally gives:
 a) Methane
 b) Ethane
 c) Ethyne
 d) Ethene
58. The carbide which reacts with water to form ethyne is
 a) CaC_2
 b) SiC
 c) Mg_2C_3
 d) Al_4C_3
59. What is the product when 2-butyne is treated with liquid NH_3 in presence of lithium?
 a) *n*-butane
 b) *cis*-2-butene
 c) *trans*-2-butene
 d) 1-butene
60. $\text{Ph}-\text{C} \equiv \text{C}-\text{CH}_3 \xrightarrow{\text{Hg}^{2+}/\text{H}^+} \text{A}$. A is:






61. 1-butyne on reaction with hot alkaline KMnO_4 gives:
 a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$
 b) $\text{CH}_3\text{CH}_2\text{COOH} + \text{CO}_2$
 c) $\text{CH}_3\text{CH}_2\text{COOH}$
 d) $\text{CH}_3\text{CH}_2\text{COOH} + \text{HCOOH}$
62. Which statement is not correct in case of ethane?
 a) It can be catalytically hydrogenated
 b) When burnt produces CO_2 and H_2O
 c) It is homologue of isobutane
 d) It can be chlorinated with chlorine
63. CH_3COCH_3 can be converted to $\text{CH}_3\text{CH}_2\text{CH}_3$ by the action of
 a) HNO_3
 b) HIO_3
 c) H_3PO_3
 d) HI
64. When ethyl chloride and alcoholic KOH are heated, the compound obtained is
 a) C_2H_4
 b) C_2H_2
 c) C_6H_6
 d) C_2H_6
65. Which of the following will react with sodium metal?
 a) Ethene
 b) Propyne
 c) But-2-yne
 d) Ethane
66. When the boiling point of the first ten normal alkanes are plotted, the graph looks like:
- a)
- b)
- c)
- d)
67. Which is generally used as reducing agent in organic chemistry?
 a) $\text{Zn} + \text{HCl}$
 b) $\text{Zn} + \text{CH}_3\text{COOH}$
 c) $\text{Zn}/\text{Hg} + \text{HCl}$
 d) $\text{Na} + \text{C}_2\text{H}_5\text{OH}$
68. Alkynes can be reduced to alkenes by hydrogenation in presence of:
 a) Raney Ni
 b) Anhy. AlCl_3
 c) Pd
 d) Lindlar's catalyst
69. Which reagent distinguishes ethylene from acetylene?

- a) Aqueous alkaline permanganate
 b) Chlorine dissolved in carbon tetrachloride
 c) Ammoniacal cuprous chloride
 d) Concentrated sulphuric acid
70. By heating tetraethyl ammonium hydroxide, the product formed are:
 a) C_2H_4
 b) $(C_2H_5)_3N$
 c) H_2O
 d) All of these
71. Addition of ICl on propene gives the product:
 a) $CH_3CHClCH_3$ b) CH_3CHICH_2Cl c) $CH_3CHClCH_2I$ d) $CH_3CHClCH_2Cl$
72. Which of the following alkenes gives on acetaldehyde on ozonolysis?
 a) Ethene b) Propene c) 1-butene d) 2-butene
73. In the following sequence of reactions, the alkene affords the compound 'B'
 $CH_3CH = CHCH_3 \xrightarrow{O_3} A \xrightarrow{H_2O} B$
 The compound B is
 a) CH_3CH_2CHO b) CH_3COCH_3 c) $CH_3CH_2COCH_3$ d) CH_3CHO
74. $CH_3CH = CH - CH_3 + CH_2N_2 \rightarrow A$; A is
 a) $\begin{array}{c} CH_3CH-CH-CH_3 \\ | \quad | \\ CH_3 \quad N_2 \end{array}$ b) $\begin{array}{c} CH_3-CH-CH-CH_3 \\ \quad \quad \quad \diagdown \quad / \\ \quad \quad \quad \quad \quad CH_2 \end{array}$
 c) Both (a) and (b) d) None of these
75. Direct fluorination of alkanes is not made because:
 a) Reaction does not occur
 b) Alkane fluorides are not formed
 c) Reaction occurs violently
 d) None of the above
76. On monochlorination of *n*-pentane, the number of isomers formed is:
 a) 4 b) 3 c) 2 d) 1
77. Which of the following is the predominant product in the reaction of $HOBr$ with propene?
 a) 2-bromo-1-propanol b) 3-bromo-1-propanol
 c) 2-bromo-2-propanol d) 1-bromo-2-propanol
78. Acetylene is prepared industrially by passing electric discharge through graphite electrodes in the atmosphere of:
 a) Air b) N_2 c) H_2 d) CO_2
79. The reaction of an aromatic halogen compound with an alkyl halide in presence of sodium in ether is called
 a) Sandmeyer's reaction b) Wurtz reaction
 c) Kolbe reaction d) Wurtz-Fittig reaction
80. How many isomeric forms of pentane exist?
 a) 3 b) 2 c) 5 d) 6
81. Alkanes mainly show reactions involving:
 a) Carbonium formation
 b) Ionic elimination
 c) Ionic formation
 d) Heat/photochemical substitution
82. Ozonolysis of an organic compound A produces acetone and propionaldehyde in equimolar mixture. Identify A from the following compounds.


- a) 2-methyl-1-pentene
c) 2-pentene
- b) 1-pentene
d) 2-methyl-2-pentene
83. Using anhydrous AlCl_3 as catalyst, which one of the following reactions produce ethylbenzene(PhEt)?
a) $\text{H}_3\text{C} - \text{CH}_2\text{OH} + \text{C}_6\text{H}_6$
b) $\text{CH}_3 - \text{CH} = \text{CH}_2 + \text{C}_6\text{H}_6$
c) $\text{H}_2\text{C} = \text{CH}_2 + \text{C}_6\text{H}_6$
d) $\text{H}_3\text{C} - \text{CH}_3 + \text{C}_6\text{H}_6$
84. On vigorous oxidation by alkaline permanganate solution $(\text{CH}_3)_2\text{C} = \text{CH} - \text{CH}_2\text{CHO}$ gives:
a) $(\text{CH}_3)_2\text{C}(\text{OH}) - \text{CH}(\text{OH}) - \text{CH}_2\text{CH}_3$
b) $\text{CH}_3 \text{---} \text{CO} + \text{CH}_3\text{CH}_2\text{COOH}$
c) $\text{CH}_3 \text{---} \text{CHOH} + \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
d) $\text{CH}_3 \text{---} \text{CO} + \text{CH}_3\text{CH}_2\text{CHO}$
85. The compound that is most reactive towards electrophilic nitration is
a) toluene
b) benzene
c) benzoic acid
d) nitrobenzene
86. One mole of a symmetrical alkene on ozonolysis gives two moles of an aldehyde having a molecular mass of 44 u. The alkene is
a) Propene
b) 1-butene
c) 2-butene
d) Ethene
87. The conversion of propene to propanol is ... type of reaction.
a) Hydrogenation
b) Hydration
c) hydrolysis
d) Dehydrogenation
88. When *n*-hexane/*n*-heptane is passed through Cr_2O_3 supported over alumina at 600°C gives:
a) Hexane
b) Hexyne
c) Benzene, Toluene
d) None of these
89. If 20cm^3 of methane (CH_4) is burnt using 50cm^3 of oxygen. The volume of the gases left after cooling to room temperature will be:
a) 60cm^3
b) 70cm^3
c) 30cm^3
d) 50cm^3
90. An alkane of mol. weight 72 gives on monochlorination only one product. Name the alkane:
a) 2-methylbutane
b) *n*-pentane
c) 2,2-dimethylpropane
d) None of these
91. The number of disubstituted products of benzene is
a) 2
b) 3
c) 4
d) 5
92. The treatment of $\text{R}'\text{MgX}$ with $\text{RC} \equiv \text{CH}$ produces
a) RH
b) $\text{R}'\text{H}$
c) $\text{R} - \text{R}$
d) $\text{R} - \text{R}'$
93. Electrolysis of an aqueous solution of sodium acetate, yields
a) Ethane
b) Ethene
c) Ethyne
d) Propane
94. Propyne on passing through red hot copper tube forms
a) benzene
b) Toluene
c) Mesitylene
d) None of these
95. Among the following, the compound that be most readily sulphonated is
a) Benzene
b) Nitrobenzene
c) toluene
d) chlorobenzene
96. Propylene on hydrolysis with sulphuric acid forms
a) *n*-propyl alcohol
b) Isopropyl alcohol
c) Ethyl alcohol
d) Butyl alcohol
97. What is the product formed when acetylene reacts with hypochlorous acid?
a) CH_3COCl
b) ClCH_2CHO
c) Cl_2CHCHO
d) ClCH_2COOH
98. When CaC_2 was hydrolysed a gas was obtained. It had a garlic odour due to phosgene present as impurity. The gas was passed through ammoniacal solution of Cu_2Cl_2 , a red ppt. was obtained. The gas was:
a) Ethylene
b) Propyne
c) Acetylene
d) Ethane

99. Alkenes undergo
- Addition reactions
 - Substitution reactions
 - Both (a) and (b)
 - None of these
100. Aromatic compound among other things should have a π -electron cloud containing $(4n + 2)\pi$ electrons where, n cannot be
- $\frac{1}{2}$
 - 3
 - 2
 - 1
101. Polymer of propylene is:
- Polyethylene
 - Polythene
 - Benzene
 - Mesitylene
102. Which of the following has the least octane number?
- Octane
 - Cetane
 - 2,2,4-trimethylpentane
 - n*-heptane
103. Name the reaction $C_3H_6 \rightarrow C_3H_8$:
- Alkylation
 - Cracking
 - Hydrogenation
 - Dehydrogenation
104. The tar which is used to make roads is a solid known as:
- Pitch
 - Paraffin wax
 - Coal
 - None of these
105. Thermal decomposition of alkanes in the absence of air is called
- Cracking
 - Oxidation
 - Combustion
 - Hydrogenation
106. The conditions for aromaticity is
- Molecule must have clouds of delocalised π -electrons
 - Molecule must contain $(4n + 2)\pi$ -electrons
 - Both (a) and (b)
 - None of the above
107. C_2-C_3 bond length in but-1,3-diene is:
- 1.46Å
 - 1.20Å
 - 1.39Å
 - 1.34Å
108. For synthesis of 1-butene, CH_3MgI should be treated with
- Propene
 - 2-chloropropene
 - Allyl chloride
 - Ethyl chloride
109. The highest boiling point is expected for
- n*-butane
 - iso*-octane
 - n*-octane
 - 2,2,3,3-tetramethyl butane
110. When butane-1 is mixed with excess of bromine, the expected reaction product is:
- Hydrogen bromide
 - Butylene gas
 - 1,2-dibromobutane
 - Perbromobutane
111. An alkene having molecular formula C_9H_{18} on ozonolysis gives 2, 2-dimethyl propanal and 2-butanone. The alkene is
- 2,2,2-trimethyl-3-hexene
 - 2,2,6-trimethyl-3-hexene
 - 2,3,4-trimethyl-2-hexene
 - 2,2,4-trimethyl-3-hexene
112. Propene on reaction with diazomethane in presence of UV radiations gives:
- Cyclopropane
 - Methyl cyclopropane
 - Butane
 - Butene
113. Both methane and ethane may be obtained by a suitable one-step reaction from
- CH_3I
 - C_2H_5I
 - CH_3OH
 - C_2H_5OH
114. The product (s) obtained *via* oxymercuration ($HgSO_4 + H_2SO_4$) of but-1-yne would be
- $CH_3CH_2COCH_3$
 - $CH_3CH_2CH_2CHO$
 - $CH_3CH_2CHO + HCHO$
 - $CH_3CH_2COOH + HCOOH$
115. Alkene-1 on hydroboration followed with action of H_2O_2 gives:
- Alkanol-2
 - Alkanol-1
 - Alkanal
 - Alkanone
116.  on ozonolysis gives



d) None of these

117. The compound 'C' in the following reaction is



a) *o*-bromotoluene

b) *m*-bromotoluene

c) *p*-bromotoluene

d) 3-bromo-2,4,6-trichlorotoluene

118. Iodination of alkane is made in presence of:

a) $KMnO_4$


b) HgO or HIO_3

c) $K_2Cr_2O_7$

d) None of these

119. Pick out the wrong statement.

a) Toluene shows resonance



is non-aromatic.

c) The hybrid state of carbon in carbonyl group is sp^2 .

d) The hyperconjugative effect is known as no bond resonance.

120. An alkene on vigorous oxidation with $KMnO_4$ gives only acetic acid. The alkene is

a) $CH_3CH_2CH = CH_2$

b) $CH_3CH = CHCH_3$

c) $(CH_3)_2C = CH_2$

d) $CH_3CH = CH_2$

121. A hydrocarbon reacts with hypochlorous acid to give 2-chloroethanol. The hydrocarbon is:

a) Methane

b) Ethylene

c) Acetylene

d) Ethane

122. The angle strain in cyclobutane is

a) $24^\circ 44'$

b) $29^\circ 16'$

c) $19^\circ 22'$

d) $9^\circ 44'$

123. During chlorination of methane usually a mixture of all the chlorinated products, *i. e.*, methyl chloride, methylene dichloride, chloroform and carbon tetrachloride are obtained. What will happen, if we use excess of Cl_2 in this reaction?

a) Only methyl chloride will be formed

b) Only chloroform will be formed

c) Only CCl_4 will be formed

d) Only methylene dichloride will be formed

124. Aromatization of *n*-heptane and *n*-octane gives respectively:

a) Toluene, ethyl benzene

b) Ethyl benzene, toluene

c) Toluene, benzene

d) Benzene, ethyl benzene

125. Which of the following organic compounds exhibit acidic character?

a) $H_3C - C \equiv CH$

b) $H_3C - C \equiv C - CH_3$

c) $H_2C = CH_2$

d) $H_3C - CH_3$

126. Sodium formate on heating with soda lime gives:





a) CH_4

b) CO_2

c) H_2

d) All of these

127. Which of the following can be used for preparation of propane?



128. The marsh gas detector used by miners works on the principle of:

a) Difference in the rates of diffusion of gases

b) Avogadro's hypothesis


c) Gay-Lussac's law of gaseous volumes

d) Berzelius hypothesis

129. The compound with highest boiling point.

- a) *n*-hexane
 b) *n*-pentene
 c) 2,2-dimethyl propane
 d) 2-methyl butane

130. The most stable conformation of chlorocyclohexane at room temperature is:



131. Acetylene is not used in making:

- a) Textile yarn
 b) PVC
 c) Glucose
 d) Drugs

132. An aromatic compound 'X' with molecular formula C_8H_{10} produces on nitration one mononitro derivative and three dinitro derivatives. Compound 'X' would be

- a) Ethyl benzene
 b) *m*-xylene
 c) *o*-xylene
 d) *p*-xylene

133. That acetylene is a linear molecule is shown by

- a) Its $C \equiv C$ bond distance being 1.21 \AA
 b) Its $C - H$ bond distance being 1.08 \AA
 c) Its $H - C - C$ bond angle being 180°
 d) All of the above

134. Benzene on treatment with a mixture of conc. HNO_3 and conc. H_2SO_4 at $100^\circ C$ gives

- a) Nitrobenzene
 b) *m*-dinitrobenzene
 c) *p*-dinitrobenzene
 d) *o*-dinitrobenzene

135. Which of the following differs with the other three?

- a) Naphthalene
 b) Ethylene
 c) Toluene
 d) Xylene

136. A saturated hydrocarbon is shown by C_nH_{10} . The value of carbon atom 'n' in this compound is:

- a) 2
 b) 4
 c) 5
 d) 6

137. Which of the following reactions will yield, 2, 2-dibromopropane?

- a) $CH_3 - C \equiv CH + 2HBr \rightarrow$
 b) $CH_3CH = CHBr + HBr \rightarrow$
 c) $CH \equiv CH + 2HBr \rightarrow$
 d) $CH_3 - CH = CH_2 + HBr \rightarrow$

138. $CH_2 = CH_2$ reacts with HCl to form:

- a) $CH_2CH_2Cl_2$
 b) CH_2ClCH_3
 c) CH_2ClCH_2Cl
 d) CH_3CHCl_2

139. Reduction of carbonyl compounds to alkanes with $NH_2 - NH_2$ and NaOH is called:

- a) Clemmensen reduction
 b) Wolff-Kishner reduction
 c) Wurtz's reaction
 d) Pondrof Verley reduction

140. The compound which cannot decolourise alkaline $KMnO_4$:

- a) Acetylene
 b) Ethanol
 c) Ethanal
 d) Ethane

141. Which one of the following can distinguish propyne from propene?

- a) Br_2 water
 b) Ammoniacal $AgNO_3$
 c) Aq. $KMnO_4$
 d) Dil. H_2SO_4

142. The reaction of ethene with oxygen in presence of a silver catalyst gives:



- a) Ethylene glycol
 b) Ethylene epoxide
 c) Glyoxal
 d) Acetaldehyde

143. 4-nitrotoluene $\xrightarrow[H_2SO_4]{K_2Cr_2O_7}$ product. The product in the reaction is

- a) Benzoic acid
 b) 4-nitrobenzene
 c) 4-nitrobenzoic acid
 d) 2-nitrobenzoic acid


144. Which of the following is Wurtz-Fittig reaction?









145. Ozonolysis can be used to detect:






- 1-butene and 2-butene
- Branched alkene from unbranched alkene
- Location of double bond/triple bond in carbon chain
- All are correct



Product X is obtain by reaction R . X and R are




147.



148. In which of the following will Kharasch effect operate?


- $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2 + \text{HCl}$
- $\text{CH}_3\text{CH}_2-\text{CH}=\text{CH}_2 + \text{HBr}$
- $\text{CH}_3\text{CH}=\text{CH}-\text{CH}_3 + \text{HBr}$
- $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2 + \text{HI}$

149. In the following reaction, A and B , respectively are






- C_2H_4 , alc. KOH/Δ
- $\text{C}_2\text{H}_5\text{Cl}$, aq. KOH/Δ
- CH_3OH , aq. KOH/Δ
- C_2H_5 , PBr_3

150. Addition of HBr on:



Separately gives:



d) None of the above

151. Compound C_6H_{12} is an:





- Aliphatic saturated compound
- Alicyclic compound
- Aromatic compound

d) Heterocyclic compound

152. A lead compound known as....is used as anti-knock in petroleum industry to increase the efficiency of fuel consumption

- a) $(C_2H_5)_4Pb$ b) $Pb(CH_3COO)_2$ c) $(C_2H_5)_2Pb$ d) $PbCO_3$

153. Which of the following form alkynide?



154. Which of the following reagents when heated with ethyl chloride, forms ethylene?

- a) Aqueous KOH b) Zn/HCl c) Alcoholic KOH d) HI

155. Reduction of 2-methyl-1-bromopropane with metal and acid gives:


- a) Butyl bromide b) *n*-butane c) Isobutene d) None of these

156. Dehydration of 2-butanol yield

- a) 1-butene b) 2-butene c) 2-butyne d) Both (a) and (b)

157. Which statement is correct?

- a) Knocking decreases the efficiency of an internal combustion engine
 b) Knocking cannot be eliminated completely by adding anti-knock compounds
 c) The higher the octane number, the better is the quality of fuel
 d) All of the above

158. The treatment of  with $NaIO_4$ or boiling $KMnO_4$ produces $KMnO_4$ produces

- a) $CH_3COCH_3 + CH_3COOH$ b) $CH_3COCH_3 + CH_3CHO$
 c) $CH_3CHO + CO_2$ d) CH_3COCH_3 only

159. Which of the following reagents will be able to distinguish between 1-butyne and 2-butyne?

- a) $NaNH_2$ b) HCl c) O_2 d) Br_2

160. 2-chloro-3-methylbutane is treated with sodium in etheral solution, then it will give

- a) 2,4-dimethylhexane b) 3,5-dimethylhexane
 c) 2,3,4,5-tetramethylhexane d) 2,6-dimethyloctane

161. The hydrocarbon which can react with sodium in liquid ammonia is

- a) $CH_3CH_2CH_2C \equiv CCH_2CH_2CH_3$ b) $CH_3CH_2C \equiv CH$
 c) $CH_3CH = CHCH_3$ d) $CH_3CH_2C \equiv CCH_2CH_3$

162. Which of the following is incorrect? The members of the homologous series of alkanes?

- a) Are all straight chain compounds
 b) Have the general formula C_nH_{2n+2}
 c) Show a regular gradation in physical properties

176. A compound (*X*) on ozonolysis followed by reduction gives an aldehyde C_2H_4O and 2-butanone, compound (*X*) is

- a) 3-methyl pentene-2 b) 3-methyl pentene-3 c) 3-methyl hexene-3 d) 3-ethyl pentene-3

177. An octane number 100 is given to:

- a) *n*-hexane b) Iso-octane c) Neopentane d) Neo-octane

178. When butene-1 is mixed with HBr, the major reaction product is:

- a) 1,2-dibromobutane b) 1-bromobutane c) 2-bromobutane d) None of these


179. Which cycloalkane has the lowest heat of combustion per CH_2 group?

- a) Cyclopropane b) Cyclobutane c) Cyclopentane d) Cyclohexane

180. The order of appearance of the following with rising temperature during the refining of crude oil is:





- a) Kerosene, gasoline, diesel
b) Diesel, gasoline, kerosene
c) Gasoline, diesel, kerosene
d) Gasoline, kerosene, diesel

181. $CH_3-C \equiv C-CH_3 \xrightarrow{NaNH_2} X$; what is *X*?


- a) $CH_3-CH_2CH_2CH_3$ b) $CH_3CH_2C \equiv CH$ c)  d) $CH_2 = C = CH-CH_3$

182. $H_3C-\underset{\substack{| \\ CH_3}}{CH}-CH=CH_2 + HBr \rightarrow A$

A (Predominantly) is:



183. The reagent *X* in the reactions



- a) H_3O^+ b) $(CH_3COO)_2Hg$ c) OH^- d) $HCOOH$

184. Cetane number of diesel fuel increases with the addition of:


- a) Decane b) Hexadecane c) Pentane d) Methyl naphthalene



185. Distillation of acetone with concentrated sulphuric acid gives

- a) Diacetone alcohol b) Mesityl oxide c) Mesitylene d) Propene-2-ol

186.

Ozonolysis of  will give:





d) None of the above

187. Soda lime is used extensively in decarboxylation reaction to obtain alkanes. Soda lime is:

- a) NaOH b) NaOH and CaO c) CaO d) Na_2CO_3

188. Incomplete combustion of petrol or diesel oil in automobile engines can be best detected by testing fuel gases for the presence of:

- a) Carbon dioxide and water vapour
b) Carbon monoxide
c) Nitrogen oxide
d) Sulphur dioxide

189. A compound with molecular formula C_4H_6 may contain:

- a) A double bond
b) Two triple bonds
c) All single bonds
d) Two double bonds or a triple bond

190. Mustard gas is a

- a) Oil gas b) Poisonous gas c) Fuel gas d) Life gas

191. Which of the following is not true?

- a) Acetylene has a linear structure
b) Alkynes undergo electrophilic addition, but not nucleophilic addition reactions
c) Alkenes show geometrical isomerism
d) There is sp^3 -hybridisation in propane

192. Pure CH_4 can be obtained by:

- a) $\text{CH}_3\text{COONa} + \text{BaO}$ b) $\text{HCOONa} + \text{NaOH}$ c) $\text{CH}_3\text{COONa} + \text{Sodalime}$ d) Electrolysis of $\text{HCOONa}(aq.)$

193. Viscosity coefficients of some liquids are given below,

Liquid	η in millipoise at 30°C
$\text{CH}_3(\text{CH}_2)_3\text{CH}_3$	2.11
$\text{CH}_3(\text{CH}_2)_4\text{CH}_3$	2.89
$\text{CH}_3(\text{CH}_2)_5\text{CH}_3$	3.68

The order of viscosity coefficient of the liquids,

(A) $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$

(B) $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3-\text{CH}_2-\text{CH}-\text{CH}_2-\text{CH}_3 \end{array}$

(C) $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3-\text{CH}-\text{CH}-\text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$


is:

- a) The same b) $(A) > (B) > (C)$ c) $(A) < (B) < (C)$ d) $(A) > (B) = (C)$

194. Action of $RMgX$ with vinyl chloride gives:

- a) Alkane b) Alkyne c) Alkene d) All of these

195. The following reaction is called



- a) Michael addition reaction b) Diels-alder reaction
c) Wolff-Kishner reaction d) None of the above

196. Which branched chain isomer of the hydrocarbon with molecular mass 72u gives only one isomer of mono substituted alkyl halide?

- a) Neopentane
b) Isohexane
c) Neohexane
d) *Tertiary*-butyl chloride

197. A *meta* directing functional group is

- a) $-COOH$ b) $-OH$ c) $-CH_3$ d) $-Br$

198. Which one of the following compounds is prepared in the laboratory from benzene by a substitution reaction?

- a) Glyoxal b) Cyclohexane
c) Acetophenone d) Hexabromocyclohexane



199. Only two isomeric monochloro derivatives are possible for:

- a) *n*-pentane b) 2,4-dimethylpentane c) Benzene d) 2-methylpropane

200. Butene-1 may be converted to butane by reaction with

- a) $Zn - HCl$ b) $Sn - HCl$ c) $Zn - Hg$ d) Pd/H_2

201. Identify 'B' in the following reaction,







- a) CH_4 b) C_2H_6 c) C_2H_5Cl d) C_2H_5OH

202. The reaction of toluene with chlorine in presence of ferric chloride gives predominantly

- a) benzoyl chloride b) *m*-chlorotoluene
c) Benzyl chloride d) *o*-and *p*-chlorotoluene

203. $Ph - C \equiv C - CH_3 \xrightarrow{Hg^{2+}/H^+} A$

The product A is



204. During Wurtz reaction, which of the following is sometimes also obtained because of decomposition of free radicals?


- a) Alkynes b) Alkenes c) CO_2 d) Alkyl halide

205. Which of the following reagents cannot be used to locate the position of triple bond in $CH_3 - C \equiv C - CH_3$?





- a) Br_2 b) O_3 c) Cu^+ d) $KMnO_4$

206. Decarboxylation of malonic acid gives:




- a) CH_4 b) C_2H_6 c) C_3H_8 d) None of these







The product of reaction is:




208. Which compound will react with an aqueous solution of $\text{Ag}(\text{NH}_3)_2^+\text{OH}^-$?






209. Reactivity of tertiary H, secondary H and primary H towards elimination is:




210. 1-butyne on hydration gives



211. The hydration of propyne in the presence of $\text{HgSO}_4/\text{H}_2\text{SO}_4$ produces







212. The most reactive halogen in the halogenation of alkanes is:




213. A gas decolourised by KMnO_4 solution but gives no precipitate with ammoniacal cuprous chloride is




214. Indane is:







215. Which reacts with ammoniacal AgNO_3 ?




216. The conversion





Can be effected using



217. Which of the following cycloalkane gives open chain compound, when reacts with bromine?



218. The addition of HBr to an alkene in the presence of peroxide is the example of






219. On mixing a certain alkane with chlorine and irradiating it with UV light, it form one monochloro alkane.




The alkane could be




220. Which of the following statements is true for ethane, ethene and acetylene?




- d) Ethene is the strongest acid and has the shortest C – H bond distance
221. On cracking petrol we get:
- CH₄
 - C₃H₆
 - Both of the above
 - CH₃ + CH₄ + C₂H₆ + alcohols
222. Methyl bromide is converted into ethane by heating it in ether medium with
- Al
 - Zn
 - Na
 - Cu
223. The addition of oxygen gas to reaction mixture of chlorine and methane (photochemical chlorination):
- Accelerates the reaction
 - Retards the reaction for sometime
 - Has no effect on the rate of reaction
 - May accelerate or retard the reaction depending upon the amount of oxygen
224. Order of reactivity of C₂H₆, C₂H₄ and C₂H₂ is
- C₂H₆ > C₂H₄ > C₂H₂
 - C₂H₂ > C₂H₆ > C₂H₄
 - C₂H₄ > C₂H₂ > C₂H₆
 - All are equally reactive
225. Bacterial decomposition of cellulose material present in sewage water gives:
- H₂
 - CH₄
 - O₂
 - N₂
226. The reaction, CH₃Br + Na → Product, is called
- Perkin reaction
 - Levit reaction
 - Wurtz reaction
 - Aldol condensation
227. *Meso*-dibromobutane on debromination gives
- trans*-2-butene
 - cis*-2-butene
 - 1-butene
 - 1-butyne
228. CH ≡ CH + HBr → X, product X is
- Ethylene bromide
 - Vinyl bromide
 - Bromo ethane
 - Ethyledine bromide
229. Kolbe's synthesis of sodium salt of butanoic acid gives:
- n*-hexane
 - Isobutane
 - Butane-1
 - Ethylene
230. The compound formed when silver powder is heated with chloroform:
- CH₄
 - C₂H₂
 - C₂H₄
 - C₂H₆
231. The reaction of toluene with chlorine in the presence of ferric chloride gives predominantly
- m*-chlorotoluene
 - Benzyl chloride
 - Benzoyl chloride
 - o* and *p*-chlorotoluene
232. Which of the following will yield *trans* product from butyne?
- LiAlH₄
 - Na/Liq. NH₃
 - NaBH₄
 - Ni catalyst
233. A hydrocarbon of molecular formula C₆H₁₀ reacts with sodamide and the same on ozonolysis followed by hydrogen peroxide oxidation gives two molecules of carboxylic acids, one being optically active. Then, the hydrocarbon may be
- 1-hexyne
 - 3-hexyne
 - 3-methyl-1-pentyne
 - 3,3-dimethyl-1-butyne
234. Which of the following is not correct about the reaction,
- $$\text{CH}_2 = \text{CH}_2 + \text{Br}_2 \xrightarrow{\text{NaI(aq)}} ?$$
- The products formed are CH₂BrCH₂Br and CH₂BrCH₂I
 - The reaction follows polar mechanism
 - The reaction occurs readily in solution and is catalysed by inorganic halides
 - CH₂ICH₂I is formed only
235. During ozonolysis of CH₂ = CH₂ if hydrolysis is made in absence of Zn dust the products formed are:
- HCHO
 - HCOOH
 - CH₃OH
 - CH₂OHCH₂OH
236. The formation of butane on heating C₂H₅I with Na in presence of ether is contaminated with impurities of:



- a) C_2H_4 b) C_3H_6 c) CH_4 d) None of these
237. When sodium propionate is heated with soda-lime, the product formed is
 a) Methane b) Ethane c) Ethene d) Ethyne
238. Isopropyl bromide on Wurtz reaction gives
 a) Hexane b) Propane
 c) 2,3-dimethyl butane d) *neo*-hexane
239. Which one of the following has the minimum boiling point?
 a) *n*-butane b) 1-butyne c) 1-butene d) *Iso*-butene
240. The substance that would not at all be formed during the reaction of methane and chlorine in the presence of sunlight is:
 a) CH_3Cl b) $CHCl_3$ c) CH_3CH_3 d) $CH_3CH_2CH_3$
241. When isopropyl magnesium iodide is treated with water, the product is:
 a) Propane b) *n*-butane c) Isobutene d) Isobutyl alcohol
242. The monosodium salt of acetylene on treating with dry CO_2 forms:
 a) $CH \equiv CCOOH$ b) $CH \equiv CCOONa$ c) $CH \equiv CCONa$ d) None of these
243. Propyne on passing through red hot iron tube gives
 a)  b)  c)  d) None of these
244. $(CH_3)_3CMgCl$ on reaction with D_2O produces
 a) $(CH_3)_3COD$ b) $(CD_3)_3CH$ c) $(CH_3)_3CD$ d) $(CD_3)_3CD$
245. *n*-hexadecane (cetane) has cetane number:
 a) 100 b) Zero c) 90 d) 110
246. Acetylene does not react with
 a) Na b) ammoniacal $AgNO_3$ c) HCl d) NaOH
247. What volume of CH_4 at NTP is formed when 20.5 g of CH_3COONa is treated with sodalime?
 a) 4.4 litre b) 2.2 litre c) 3.2 litre d) 5.6 litre
248. The hydrocarbon which decolourizes alkaline $KMnO_4$ solution, but does not give any precipitate with ammoniacal silver nitrate is:
 a) Benzene b) Acetylene c) Propyne d) Butyne-2
249. What is the molecular formula of the product formed when benzene is reacted with ethyl chloride in presence of anhydrous aluminium chloride?
 a) C_8H_{10} b) C_6H_6 c) C_8H_8 d) C_6H_5Cl
250. Which will give $CH_2 = C = CH_2$?
 a) $CH_2Br - CBr = CH_2 \xrightarrow{Zn/CH_3OH}$
 b) $CH \equiv C - CH_2 - COOH \xrightarrow{K_2CO_3(aq)}$
 c) $2CH_2 = CH - CH_2I \xrightarrow{2Na}$
 d) None of the above
251. A dibromo derivative of an alkane reacts with sodium metal to form an alicyclic hydrocarbon. The derivative is
 a) 1,1-dibromopropane b) 2,2-dibromopropane
 c) 1,2-dibromoethane d) 1,4-dibromobutane
252. By coaltar distillation which is not obtained?
 a) Light oil b) Middle oil c) Heavy oil d) Mobil oil
253. In the following reaction:




The major product is:



254. The treatment of ethane with cold alkaline potassium permanganate produces
 a) Ethylene glycol b) Formaldehyde
 c) Formic acid d) Carbon dioxide and water
255. As compared to melting points of even carbon chain isomers, the melting points of odd carbon chain alkanes are:
 a) Lower
 b) Higher
 c) Same
 d) Not depend upon branching





257. In the reactions,



A and *B* are geometrical isomers. Then,

- a) *A* is *cis* and *B* is *trans* b) *A* is *trans* and *B* is *cis*
 c) *A* and *B* are *cis* d) *A* and *B* are *trans*

258. Identify 'A' in the reaction:



259. Choose the correct statement

- a) Acetylene is more reactive than ethylene to an electrophilic attack
 b) Acetylene and ethylene show similar reactivities towards an electrophilic attack with different rates
 c) The reactivities of acetylene and ethylene towards an electrophilic attack depend on the electrophilic reagent
 d) Acetylene is less reactive than ethylene to an electrophilic attack
260. $\text{C}_6\text{H}_5\text{CH}_3 \xrightarrow{\text{CrO}_2\text{Cl}_2} \text{Z}$

In the given sequence, Z is





- a) Benzaldehyde b) Toluic acid c) Phenyl acetic acid d) Benzoic acid

261. 2-hexyne can be converted to *trans*-2-hexene by the action of:
 a) $H_2 - Pd/BaSO_4$ b) $Li/Liq. NH_3$ c) $H_2 - Pt O_2$ d) $NaBH_4$
262. In the following reaction,
 $RCH_2CH = CH_2 + ICl \rightarrow [A]$
 Markownikoff's product [A] is
- a)
$$\begin{array}{c} RCH_2CH - CH_2I \\ | \\ Cl \end{array}$$
- b)
$$\begin{array}{c} RCH_2CH - CH_2Cl \\ | \\ I \end{array}$$
- c)
$$\begin{array}{c} RCH_2 - C = CH_2 \\ | \\ I \end{array}$$
- d) $RCH = CH - CH_2I$
263. Which of the following will not produce ethane?
 a) Reduction of CH_3COOH with HI and red P
 b) Reduction of CH_3COCH_3 with HI and red P
 c) Sodalime decarboxylation of sodium probionate
 d) Hydrogenation of ethane in presence of Raney-Ni
264. Which will not react with acetylene?
 a) NaOH b) Na c) HCl d) Amm. $AgNO_3$
265. Ozonolysis of an organic compounds gives formaldehyde as one of the products. This confirms the presence of
 a) Two ethylenic double bonds b) A vinyl group
 c) An *iso*-propyl group d) An acetylenic triple bond
266. Among the paraffins it is generally found that with an increase in the molecular weight:
 a) The freezing point decreases
 b) The boiling point decreases
 c) The boiling point increases
 d) The vapour density decreases
267. Which of the following reactions can be used to prepare methane?
 a) Clemmensen reduction
 b) Wurtz reaction
 c) Reduction of $CH_2 = CH_2$ by $LiAlH_4$
 d) Reduction of methyl iodine by using a zinc-copper couple
268. Ethylene reacts with dil. H_2SO_4 in presence of $HgSO_4$ to give:
 a) Ethanal b) Ethanol c) Ethane d) Ethene
269. Household gas or liquefied petroleum gas (L.P.G.) mainly contains:
 a) Methane and ethane
 b) Liquefied butane and isobutene
 c) Ethylene and CO
 d) C_2H_2 and H_2
270. Which one of the following gives, on ozonolysis, both aldehydes and ketones?
 a) $Me_2C = CHMe$ b) $Me_2C = CMe_2$
 c) $MeCH_2 - C(Me) = CMe_2$ d) $MeCH(Me) - CH = CHMe$
271. Which among the following give alkanes on reduction?
 a) Aldehydes b) Ketones c) Carboxylic acids d) All are correct
272. Lewisite (a war gas) is an.....compound.
 a) Organosulphur b) Organoarsenic c) Organoantimony d) Organophosphorus
273. In the following reaction,


$$C_2H_2 \xrightarrow[HgSO_4/H_2SO_4]{H_2O} X \rightleftharpoons CH_3CHO$$
. What is X?

- a) $\text{CH}_3\text{CH}_2\text{OH}$ b) $\text{CH}_3 - \text{O} - \text{CH}_3$ c) $\text{CH}_3\text{CH}_2\text{CHO}$ d) $\text{CH}_2 = \text{CHOH}$

274. Compound (A) on oxidation with $\text{OsO}_4/\text{NaIO}_4$ gives Hexanedinal. Structure of compound. (A) will be



275. Major product of the following reaction is:



- a) Butene-1 b) Butene-2 c) Butane d) Butyne-1

276. The compound formed as a result of oxidation of ethyl benzene by KMnO_4 is

- a) Benzophenone b) Acetophenone c) Benzoic acid d) Benzyl alcohol

277. Methane reacts with conc. HNO_3 at high temperature to yield:

- a) CO_2 and H_2O b) HCHO c) HCOOH d) CH_3NO_2

278. Butyne-1 and butyne-2 can be distinguished by:

- a) $\text{Br}_2, \text{CCl}_4$
b) H_2 , Lindler catalyst
c) Dilute $\text{H}_2\text{SO}_4, \text{HgSO}_4$
d) Ammoniacal cuprous chloride

279. An isolated alkadiene is:

- a) Penta-1,4-diene b) Penta-1,3-diene c) Penta-1,2-diene d) None of these

280. $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_3 \xrightarrow{\text{Lindlar's catalyst}} \text{A}$, the compound A is

- a) *cis*-2-butene b) *trans*-2-butene c) *iso*-butene d) 1-butene

281. If a halogen compound contains OH group, will it be possible to carry out the Wurtz reaction?

- a) Yes b) No c) - d) -

282. Reduction of 2-butyne with Na in liquid NH_3 gives predominantly:

- a) *n*-butane b) *Trans*-2-butene c) No reaction d) *Cis*-2-butene

283. Phenyl magnesium bromide reacts with methanol to give

- a) A mixture of anisol and $\text{Mg}(\text{OH})\text{Br}$ b) A mixture of benzene and $\text{Mg}(\text{OMe})\text{Br}$
c) A mixture of toluene and $\text{Mg}(\text{OH})\text{Br}$ d) A mixture of phenol and $\text{Mg}(\text{Me})\text{Br}$

284. Iso-octane is added to petrol:

- a) To precipitate inorganic material
b) To prevent freezing of petrol
c) To increase the boiling point of petrol
d) To increase octane number


285. When cyclohexane is poured on water, it floats, because:

- a) Cyclohexane is in 'boat' form
b) Cyclohexane is in 'chair' form
c) Cyclohexane is in 'crown' form
d) Cyclohexane is less dense than water

286. Ethylene reacts with 1% cold alkaline KMnO_4 (Baeyer's reagent) to form:


- a) Oxalic acid b) Acetic acid c) Glycerol d) Glycol

287. The reagent that would effect the following transformation is:



- a) NaBH_4 in alcohol b) $[(\text{C}_6\text{H}_5)_3\text{P}]_3\text{RhCl}/\text{H}_2$ c) LiAlH_4 d) All of these

288. How many primary and tertiary carbon atoms are present in




- a) 3p, 1t b) 2p, 2t c) 1p, 3t d) None of these

289. Which of these will not react with acetylene?

- a) NaOH b) Amm. AgNO₃ c) Na d) HCl

290. The catalytic hydrogenation is more easier in case of which alkene?



291. Addition of hydrogen on C=C is called hydrogenation. Addition of halogen on C=C is called:

- a) Halogenation
b) Dehalogenation
c) Elimination of halogen
d) None of these


292. The synthetic gas is:

- a) CH₄ b) C₂H₂ c) CO + 3H₂ d) NH₃

293. Toluene on treatment with CrO₃ and (CH₃CO)₂O followed by hydrolysis with dil. HCl gives

- a) Benzaldehyde b) Benzoic acid c) Phenol d) Phenylacetaldehyde

294. Identify the product (P) in the reaction:



- a) No reaction b) R₃C—CR₃ c) R₃C—OH d) R₃C—O—CR₃

295. Gem dihalides on treatment with alcoholic KOH give

- a) Alkyne b) Alkene c) Alkane d) All of these

296. The presence of Ag⁺ ion increases the solubility of alkenes due to the formation of

- a) dπ — dσ bonding b) pσ — pπ bonding c) pπ — dπ bonding d) pπ — dσ bonding

297. Acetylene and HCHO react in presence of copper acetylide catalyst to form

- a) 1-butyne-1,4-diol b) 2-butyne-1,2-diol c) 2-butyne-1,4-diol d) None of these

298. Decarboxylation of isobutyric acid leads to:

- a) Isobutene b) Propane c) Butane d) None of these

299. In the addition of HBr to propene in the absence of peroxides the first step involves the addition of:

- a) H⁺ b) Br⁻ c) H^o d) Br


300. The IUPAC name of —C ≡ C—CH₃ group is:

- a) Prop-2-ynyl b) Prop-2-enyl c) Prop-1-ynyl d) None of these

301. Pure methane can be produced by

- a) Wurtz reaction b) Kolbe's electrolytic method
c) Soda lime decarboxylation d) reduction with H₂

302. What are X and Y respectively, in the following reaction?



- a) Na/NH₃(liq.) and Pd/BaSO₄ + H₂ b) Ni/140°C and Pd/BaSO₄ + H₂
c) Ni/140°C and Na/NH₃(liq.) d) Pd/BaSO₄ + H₂ and Na/NH₃(liq.)

303. When a mixture of methane and oxygen is passed through heated molybdenum oxide, the main product formed is

- a) Methanoic acid b) Ethanal c) Methanol d) Methanal

304. Propyne and propene can be distinguished by





- a) conc. H₂SO₄ b) Br₂ in CCl₄ c) alk. KMnO₄ d) AgNO₃ in NH₃

305. Conformation in molecules is due to:

- a) Rotation about a single bond

- b) Change in direction of light
 c) Structural changes
 d) Restricted rotation about a double bond

306. The non-aromatic compound among the following is



307. Kerosene is a mixture of:

- a) Alkenes b) Alkanes c) Alkynes d) Arenes

308. Which of the following alkenes is most stable?

- a) $R_2C = CR_2$ b) $R-CH = CH-R$ c) $RCH = CH_2$ d) $CH_2 = CH_2$

309. What is obtained when chlorine is passed in boiling toluene and product is hydrolysed?

- a) *o*-cresol b) *p*-cresol
 c) 2,4-dihydroxytoluene d) Benzyl alcohol

310. It is necessary to use.....in the iodination of alkane.

- a) Alcohol b) Oxidant c) Benzene d) Reductant

311. Ozonolysis of propyne gives:

- a) CH_3CHO b) CH_3COCHO c) $HCHO$ d) $CHOCHO-$

312. Reactivity of alkenes towards HX decreases in the order:

- a) Butene > propene > ethene
 b) Butene > ethene > propene
 c) Ethene > propene > butene
 d) None of the above






313. Propyne on oxidation with SeO_2 gives:

- a) $CHOCHO$ b) CH_3CH_2CHO c) CH_3COCHO d) $CHOCH_2CHO$

314. 2-methylbutane on reacting with bromine in the presence of sunlight gives mainly

- a) 1-bromo 3-methylbutane b) 2-bromo 3-methylbutane
 c) 2-bromo 2-methylbutane d) 1-bromo 2-methylbutane

315. The product of following reaction is,



316. Which statement is correct?





- a) Alkanes are called paraffins because of their little chemical affinity
 b) Alkanes have only sigma bonds
 c) Most abundant alkane is CH_4
 d) All are correct

317. An activating group

- a) activates only *ortho* and *para* positions b) Deactivates *meta* position
 c) activates *ortho* and *para* more than *meta* d) Deactivates *meta* more than *ortho* and *para*

318. An alkyl bromide, RBr of molecular weight 151 is the exclusive product of bromination of which hydrocarbon?

- a) Dodecane b) 2, 2-dimethylpropane


- c) 2, 2-dimethylhexane
 319. The conversion of liquid hydrocarbon into a mixture of gaseous compounds by heat alone is known as:
 a) Hydrolysis b) Reduction c) Oxidation d) Cracking
320. Ethyl benzene cannot be prepared by
 a) Wurtz reaction b) Wurtz-Fittig reaction
 c) Friedel-Craft's reaction d) Clemmensen reduction
321. Silver acetylide when heated with HCl gives:
 a) C_2H_2 b) H_2 c) C_2H_4 d) C_6H_6
322. The addition of HCl to 3, 3, 3-trichloropropene gives
 a) $Cl_3CCH_2CH_2Cl$ b) $Cl_3CCH_2CHCl_2$ c) $Cl_2CHCH_2CHCl_2$ d) $Cl_2CHCH(Cl)CH_2Cl$
323. Sodium ethoxide is specific reagent for:
 a) Dehydration
 b) Dehydrohalogenation
 c) Dehydrogenation
 d) Dehalogenation
324. A fuel contains 25% *n*-heptane and 75% iso-octane. Its octane number is:
 a) 50 b) 75 c) 100 d) 25
325. The greatest strain is involved in cycloalkane, when the bond angle is:
 a) 60° b) 90° c) 120° d) 108°
326. Which of the following will be obtained by the bromination of ethylbenzene in the presence of light?
 a)  b)  c)  d) 
327. On passing electric discharge through graphite in presence of H_2 the compound formed is:
 a) CH_4 b) C_2H_6 c) C_2H_2 d) All of these
328. Propene reacts with Cl_2 at $400-600^\circ C$ to give:
 a) 1,2-dichloropropane b) Allyl chloride c) No reaction d) Polyvinyl chloride
329. Methane reacts with oxygen at 100 atm and $300^\circ C$ in presence of Cu to give:
 a) Acetaldehyde b) Methyl alcohol c) Acetic acid d) Ethyl alcohol
330. Ethylene is used in making:
 a) Anti-freeze b) Solvent c) Fumigant d) All of these
331. The main constituent of light oil fraction is:
 a) Benzene b) Toluene c) Phenol d) Naphthalene
332. The major product in the acid catalysed dehydration of 2-pentanol is:
 a) 4-pentene b) 3-pentene c) 2-pentene d) 1-pentene
333. Which gas is commonly used in welding?
 a) C_2H_4 b) C_2H_2 c) CH_4 d) C_2H_6
334. The synthesis of 3-octyne is achieved by adding a bromoalkane into a mixture of sodium amide and an alkyne. The bromoalkane and alkyne respectively are
 a) $BrCH_2CH_2CH_2CH_2CH_3$ and $CH_3CH_2C \equiv CH$ b) $BrCH_2CH_2CH_3$ and $CH_3CH_2CH_2C \equiv CH$
 c) $BrCH_2CH_2CH_2CH_2CH_3$ and $CH_3C \equiv CH$ d) $BrCH_2CH_2CH_2CH_3$ and $CH_3CH_2C \equiv CH$
335. Which is most acidic of the following?
 a) Methane b) Acetylene c) 1-butene d) *Neo*-pentane
336. Addition of HI on double bond of propene yields isopropyl iodide and not *n*-propyl iodide as the major product, because addition proceeds through:
 a) A more stable carbonium ion
 b) A more stable carbanion
 c) A more stable free radical
 d) None of the above

337. Correct statement about 1,3-dibutene




- a) Conjugated double bonds are present
 b) Reacts with HBr
 c) Forms polymer
 d) All of the above

338. Preparation of ethane by electrolysis of aqueous solution of potassium acetate is called






- a) Wurtz reaction
 b) Kolbe's synthesis
 c) Grignard reaction
 d) Sabatier-Sendersen's reaction



Thus, A and B are




340. The major product P in the following reaction is



341. $CH_3CH = CHCHO$ is oxidized to $CH_3CH = CHCOOH$ using:

- a) Alkaline permanganate
 b) Ammoniacal silver nitrate
 c) Selenium dioxide
 d) Osmium tetroxide



- a) C_2H_4 b) C_2H_2 c) C_2H_6 d) $Ca(OH)_2$

343. Acetylene reacts with hypochlorous acid to form

- a) $Cl_2CH.CHO$ b) $ClCH_2COOH$ c) CH_3COCl d) $ClCH_2CHO$

344. Dehydrohalogenation of 1,2-dibromobutane with alc. KOH gives:


- a) 1-butyne b) 2-butene c) 1-butene d) 1-bromo-1-butene

345. Water can be added across a triple bond in the presence of



- a) Acidic medium b) Alkaline medium c) Neutral medium d) Acid and $HgSO_4$



346. Both methane and ethane may be obtained in one step reaction from:

- a) CH_3COONa b) CH_3I c) Both (a) and (b) d) None of these







The reaction of with HBr gives predominantly










348. The product formed when acetylene is passed through red hot tube is:
 a) Benzene b) Cyclohexane c) Neoprene d) Ethane
349. The product formed when toluene is heated in light with Cl_2 and in absence of halogen carrier is
 a) Chlorobenzene b) Gammexane c) Benzotrichloride d) DDT
350. Among the following statement on the nitration of aromatic compounds, the false one is
 a) The rate of nitration of benzene is almost the same as that of hexadeuterobenzene
 b) The rate of nitration of toluene is greater than that of benzene
 c) The rate of nitration of benzene is greater than that of hexadeuterbenzene
 d) Nitration is an electrophilic substitution reaction
351. Reaction of one molecule of HBr with one molecule of 1, 3-butadiene at $40^\circ C$ gives predominantly
 a) 1-bromo-2-butene under kinetically controlled conditions
 b) 3-bromobutene under thermodynamically controlled conditions
 c) 1-bromo-2-butene under thermodynamically controlled conditions
 d) 3-bromobutene under kinetically controlled conditions
352. Which of the following compound is aromatic?



353. Ethylene reacts with 1% alkaline $KMnO_4$ to form
 a) Oxalic acid b) Ethylene glycol c) Ethyl alcohol d) HCHO
354. To prepare a pure sample of *n*-hexane using sodium metal as one reactant, the other reactant or reactants will be:
 a) Ethyl chloride and *n*-butyl chloride
 b) Methyl bromide and *n*-pentyl bromide
 c) *n*-propyl bromide
 d) Ethyl bromide and *n*-butyl bromide
355. Friedel-Craft acylation can be given by



X is



356. A mixture of CS_2 and H_2S on passing over heated Cu gives:
 a) C_2H_6 b) CH_4 c) C_3H_8 d) None of these
357. Photochemical chlorination of alkane is initiated by a process of:
 a) Pyrolysis b) Substitution c) Homolysis d) Peroxidation
358. Under which one of the following conditions, does the reaction,

$\text{CH} \equiv \text{CH} + \text{CH}_3\text{OH} \xrightarrow{?} \text{CH}_3\text{O}-\text{CH}=\text{CH}_2$ take place?


a) $\text{NH}_4\text{OH}/80^\circ\text{C}$

b) $\text{Conc. H}_2\text{SO}_4/160^\circ\text{C}$

c) Anhydrous $\text{ZnCl}_2/150^\circ\text{C}$

d) $\text{CH}_3\text{OK}/160-200^\circ\text{C}$

359. Which one is correct for the given change?



a) The product formed is *trans*-2-methyl-1-cyclopentanol

b) The product formed is

The diagram shows a skeletal structure of trans-2-methylcyclopentanol, which is a five-membered ring with a methyl group (CH_3) and a hydrogen atom (H) on one carbon, and a hydrogen atom (H) and a hydroxyl group (OH) on the adjacent carbon. The methyl and hydroxyl groups are on opposite sides of the ring.

c) The addition is syn addition

d) All of the above

360. The electrolysis of aqueous solution of potassium succinate produces

a) Methyl alcohol

b) ethyl alcohol

c) ethene





d) ethane

361. Ozonolysis products of an olefin are

CHO CH_2CHO





and
 CHO CH_2CHO

Olefin is



362. $\text{CH}_3\text{C} \equiv \text{CH} \xrightarrow[(2) \text{CH}_3\text{CH}_2\text{Br}]{(1) \text{NaNH}_2} \text{A} \xrightarrow[\text{Lindlar's catalyst}]{\text{H}_2} \text{B}$

What is B in the above reaction?



363. The gas believed to be the cause of explosion in coal-mines or fire damp is:

a) Methane

b) Ethane

c) C_3H_8

d) CO

364. Addition of HBr to propylene in presence of benzoyl peroxide, follows

- a) Markownioff's rule
 c) Carbanion mechanism
365. 2-phenyl propene on acidic hydration gives,
 a) 2-phenyl-2-propanol
 c) 3-phenyl-1-propanol
366. $\text{CH}_2 = \text{CH}_2$ is also called a:
 a) Monomer b) Polymer
367. Halogenation of alkanes is an example of:
 a) Electrophilic substitution
 b) Nucleophilic substitution
 c) Free radical substitution
 d) Oxidation
368. The most stable isomer of 1,2-dichloroethane is:
 a) Staggered b) Gauche

- b) Baeyer's rule
 d) *anti*-Markownioff's rule

- b) 2-phenyl-1-propanol
 d) 1-phenyl-2-propanol

- c) Isomer d) Equimer

369. Which does not decolourize Br_2 water?



- c) Eclipsed d) Partially eclipsed

- c) $\text{CH}_3\text{C} \equiv \text{CH}$ d) $\text{CH}_2 = \text{CHCH}_3$

370. Grignard's reagent gives alkane with:

- a) H_2O b) $\text{C}_2\text{H}_5\text{OH}$

- c) $\text{C}_2\text{H}_5\text{NH}_2$ d) All of these

371. The carbon-carbon bond length in benzene is

- a) In between C_2H_6 and C_2H_4
 c) In between C_2H_6 and C_2H_2
 b) Same as in C_2H_4
 d) In between C_2H_4 and C_2H_2





372. Electrolysis of a concentrated solution of sodium fumarate gives:

- a) Fumaric acid b) Ethylene c) Ethane d) Acetylene

373. In order to overcome angle strain, cyclohexane acquires:

- a) Square planar structure
 b) Planar structure
 c) Puckered ring structure
 d) Pyramidal structure


374. *o*-toluic acid on reaction with $\text{Br}_2 + \text{Fe}$ gives








375. The reaction, $\text{CH}_2 = \text{CH}_2 + \text{H}_2 \xrightarrow[250-300^\circ\text{C}]{\text{Ni}} \text{CH}_3 - \text{CH}_3$

is called:





- a) Wurtz's reaction b) Kolbe's reaction c) Sabatier and Senderens reaction
 d) Carbylamines reaction



Identify the X in the above reaction



Identify the adduct



378. Ethane can be freed (isolated) from the impurity of ethylene by washing with:

a) HCl

b) HNO_3

c) H_2SO_4

d) water

379. Poisonous gases are:

a) Phosgene

b) Lewisite

c) Mustard gas

d) All of these

380. A chlorohydrocarbon, named chlorodane is used especially as:

a) Insecticide

b) Anti-worm

c) Fungicide

d) Anti-termite

381. The highest boiling point is expected for

a) *iso*-octane

c) 2, 2, 3, 3-tetramethyl butane

b) *n*-octane

d) *n*-butane

382. The addition of tetraethyl lead of petrol:



a) Lowers its octane number

b) Raises its octane number

c) May raise or lower the octane number

d) Has no effect on octane number

383. Which of the following reactions will give an alkyne?



d) All of the above

384. Which one among the following is assigned an octane number of zero?

a) *Iso*-octane

b) *n*-heptane

c) *Iso*heptane

d) 2-methyloctane

385. The process where straight run gasoline is cracked in order to increase octane number is called:

a) Aromatization

b) Rearrangement

c) Substitution

d) Reforming

386. The treatment of aluminum carbide with water or dilute acid produces





a) acetylene

b) ethene


c) methane

d) ethane





387. When acetylene is passed through red hot iron tube, compound X is formed. Which one of the following reactions will yield X as the major product?







388. In the reaction






Identify C is

- a) C_6H_5OH b) C_6H_6 c) C_6H_5COONa d) C_6H_5ONa
389. The major product formed when a 3, 3-dimethyl butan-2-ol is heated with concentrated sulphuric acid, is
- a) 2,3-dimethyl-2-butene
 b) 2,3-dimethyl-1-butene
 c) 3,3-dimethyl-1-butene
 d) *cis* and *trans* isomers of 2,3-dimethyl-1-butene
390. Most of the hydrocarbons from petroleum are obtained by:
- a) Fractional distillation
 b) Fractional crystallization
 c) Vaporisation
 d) Polymerization
391. Cyclopentadienyl anion is
- a) Aromatic b) Non-aromatic c) Non-planar d) Aliphatic
392. Ozonolysis of buta-1,3-diene gives:
- a) HCHO and glyoxal
 b) CH_3CHO and glyoxal
 c) CO_2 and glyoxal
 d) HCHO+glyoxal+ CH_3CHO
393. Which is not true in the case of natural gas?
- a) It is a fuel
 b) It is used in the manufacture of fertilizer
 c) It is a mixture of CO_2 and H_2
 d) It is a mixture of gaseous hydrocarbons
394. Wurtz reaction using bromoethane yields:
- a) 2-bromobutane b) *n*-butane c) Isobutene d) Ethane
395. Which of the following compounds is not aromatic?
- a)  b)  c)  d) 
396. Which products are formed during the addition of Br_2 on ethylene in presence of aqueous $NaNO_3$ solution?
- a) CH_2Br . CH_2ONO_2
 b) CH_2Br . CH_2Br
 c) $CH_2(ONO_2)$. CH_2ONO_2
 d) Both (a) and (b)
397. Alkanes containing.....carbon atoms are converted into an aromatic hydrocarbon, when heated in presence of Cr_2O_3 on Al_2O_3
- a) 6 to 10 b) 4 to 8 c) 3 to 6 d) 5 to 6
398. Chlorination of toluene in the presence of light and heat followed by treatment with aqueous NaOH solution gives
- a) *o*-cresol b) *p*-cresol
 c) Benzoic acid d) 2,4-dihydroxytoluene
399. Toluene can be converted into benzaldehyde by oxidation with
- a) $KMnO_4$ /alkali b) CrO_2Cl_2 c) $K_2Cr_2O_7/H_2SO_4$ d) O_2/V_2O_5
400. $CH_3 - CH_2 - C \equiv CH \xrightarrow[H_2SO_4]{HgSO_4} A$
- The compound A is
- a) O b) $CH_3 - CH_2 - CH_2 - CHO$




413. Which molecule will undergo radical formation oxidation reaction most readily?



414. Which of the following is expected to be aromatic?




415. Aniline is treated with a mixture of sodium nitrite and hypophosphorus acid, the product formed is






416. Hexachloroethane is also called





417. In presence of nickel cyanide, acetylene gives



418. Conjugated double bonds are present in:







419. Normal alkanes can undergo sulphonation if they contain:




In the above reaction, X is







421. The dehydrohalogenation of neopentyl bromide with alcoholic KOH gives mostly:






422. What is obtained, when ammoniacal AgNO_3 reacts with acetylene?




423. Which of the following liberates methane on treatment with water?







424. Which statement is correct?



425. The most stable conformational isomer of cyclohexane is:



426. In the following reaction sequences,



the termination step is:



427. Which of the following is elimination reaction?

- a) $\text{CH}_3\text{CH}_2\text{OH} \rightarrow \text{CH}_2 = \text{CH}_2 + \text{H}_2\text{O}$
 b) $\text{CH}_3\text{CH}_2\text{Br} \rightarrow \text{CH}_2 = \text{CH}_2 + \text{HBr}$
 c) $\text{Br}-\text{CH}_2-\text{CH}_2-\text{Br} \xrightarrow{\text{Zn}} \text{CH}_2 = \text{CH}_2 + \text{ZnBr}_2$
 d) All of the above are correct

428. $\text{CH} \equiv \text{CH} \xrightarrow{\text{O}_3/\text{NaOH}} \text{X} \xrightarrow{\text{Zn}/\text{CH}_3\text{COOH}} \text{Y}$. Y is:

- a) $\text{CH}_2\text{OH}-\text{CH}_2\text{OH}$ b) $\text{CH}_3\text{CH}_2\text{OH}$ c) CH_3COOH d) CH_3OH


429. Baeyer's reagent is used in the laboratory for:

- a) Detection of double bonds b) Detection of glucose c) Reduction d) Oxidation

430. Product formed on electrolysis of potassium salt of fumaric and maleic acid is

- a) Ethane b) Ethene c) Ethyne d) Methane

431. The product 'X' in the following reaction is




a) $\text{CH}_3\text{Br}-\text{CH}=\text{CH}_2$


b) $\begin{array}{c} \text{Br} \\ | \\ \text{CH}_3-\text{C}=\text{CH}_2 \end{array}$


c) $\text{CH}_3\text{CH}=\text{CHBr}$



d) None of the above



432. Observe the following reactions and predict the nature of A and B.



a) A and B both are 

b) A and B both are 

c) A is  and B is 

d) A is  and B is 

433. HBr is added to $\text{CH}_3-\text{CH}=\text{CH}_2$ in presence of peroxides. The resultant compound is:

- a) $\text{CH}_3\text{CHBrCH}_3$ b) $\text{C}_2\text{H}_5\text{CH}_2\text{Br}$ c) $\text{CH}_2 = \text{CH}_2\text{CH}_2\text{Br}$ d) None of these

434. Amount of Br_2 required to react with 5 g pentene to form monobromo derivative is:

- a) 11.11 g b) 11.43 g c) 5.55 g d) None of these

435. The compound (i) decolourises KMnO_4 (ii) forms ozonide with ozone and (iii) undergoes polymerization. It will be:




- a) C_6H_6 b) C_3H_8 c) C_2H_4 d) C_2H_6

436. The strongest *ortho/para* directing group is

- a) $-\text{NH}_2$ b) $-\text{CH}_3$ c) $-\text{Cl}$ d) $-\text{C}_2\text{H}_5$

437. Which of the following species will be aromatic?

d) None of these



438. When chlorine is passed through warm benzene in presence of the sunlight, the product obtained is

- a) Benzotrichloride b) Chlorobenzene c) Gammexane d) DDT

439. The C = C bond distance in an organic compound is 1.34 Å. It can be

- a) Butene b) Hexatriene c) Cyclohexatriene d) Any of these

440. The lowest possible alkane with ethyl group as substituents possesses mol. mass equal to:

- a) 16 b) 72 c) 84 d) 128

441. The reagent(s) for the following conversion, cis/are

- a) Alcoholic KOH b) Alcoholic KOH followed by NaNH₂
c) Aqueous KOH followed by NaNH₂ d) Zn/CH₃OH

442. Aqueous H₂SO₄ reacts with 2-methyl-1-butene to give predominantly:

- a) Isopentyl hydrogen sulphate
b) 2-methyl-3-butene
c) 2-methyl-1-butene
d) Secondary butyl hydrogen sulphate

443. The number of conformation(s) for ethane is/are:

- a) 1 b) 2 c) 3 d) Infinite

444. The test for unsaturation is confirmed by the decolourisation of which of the following?

- a) Iodine water b) CuSO₄ solution c) Bromine water d) All of these

445. Which does not react with chlorine in dark?

- a) CH₄ b) C₂H₂ c) C₂H₄ d) CH₃CHO


446. The ozonolysis of isobutene gives:

- a) CH₃CHO b) CH₃COCH₃ and HCHO c) CH₃CH₂OH d) CH₃OH

447. Which compound on reductive ozonolysis forms only glyoxal?

- a) Ethyne b) Ethene c) Ethane d) 1,3-butadiene

448. The reaction,




is the example of:

- a) Sulphonation b) Dehydration c) Alkylation d) Decomposition

449. The catalyst used in the manufacture of polythene by Ziegler method is:

- a) Titanium tetrachloride and triphenyl aluminium
b) Titanium tetrachloride and trimethyl aluminium
c) Titanium dioxide
d) Titanium isopropoxide



On reductive ozonolysis yields

- a) 6-oxoheptanal b) 6-oxoheptanoic acid
c) 6-hydroxyheptanal d) 3-hydroxypentanal

451. The treatment of CH_3MgX with $\text{CH}_3\text{C} \equiv \text{C} - \text{H}$ produces

- a) $\text{CH}_3 - \text{CH} = \text{CH}_2$ b) $\text{CH}_3\text{C} \equiv \text{C} - \text{CH}_3$ c) $\begin{array}{cc} \text{H} & \text{H} \\ | & | \\ \text{CH}_3 - \text{C} = & \text{C} - \text{CH}_3 \end{array}$ d) CH_4

452. 1,3-butadiene has:

- a) Only sp -hybridised C-atoms
 b) Only sp^2 -hybridised C-atoms
 c) sp , sp^2 and sp^3 -hybridised C-atoms
 d) Sp and sp^2 -hybridised C-atoms


453. Chloroform, on warming with Ag powder gives

- a) C_2H_6 b) C_2H_6 c) C_2H_4 d) C_2H_2

454. By Wurtz reaction, a mixture of methyl iodide and ethyl iodide gives

- a) Butane b) Ethane
 c) Propane d) A mixture of the above three

455. The following reaction is an example of,



- a) An addition reaction
 b) A substitution reaction
 c) An elimination reaction
 d) None of the above

456. Acetylene on passing into excess of HOCl solution forms:


- a) Ethylene chlorohydrin
 b) Acetaldehyde
 c) Dichloroacetaldehyde
 d) Methyl chloride

457. Ethylene forms ethylene chlorohydrin by the action of:

- a) Dry chlorine gas
 b) Dry hydrogen chloride gas
 c) Solution of chlorine gas in water
 d) Dilute hydrochloric acid

458. Which one of the following has the smallest heat of hydrogenation per mole?

- a) 1-butene b) *Trans*-but-2-ene c) *Cis*-but-2-ene d) Buta-1, 3-diene

459.  $\xrightarrow[\text{H}_2\text{O}_2]{\text{OsO}_4}$ A, A is

- a) *meso* diol b) Racemic diol c) Both (a) and (b) d) None of these

460. Which of the following characteristic apply both to ethane and ethyne?

- a) Explode when mixed with chlorine
 b) Decolourise Baeyer's reagent giving brown precipitate
 c) Rapidly absorbed by cold conc. H_2SO_4
 d) Form white precipitate with AgNO_3 solution

461. Conjugated double bond is present in:

- a) Propylene b) Isobutylene c) 1,3-butadiene d) Butylene



462. The reactivities of ethane, ethylene and acetylene are of the order

- a) Ethane < ethene < ethyne b) Ethane < ethyne < ethene
 c) Ethyne = ethene > ethane d) Any of the above


463. Which is not linked with methane?

- a) Marsh gas b) Natural gas c) Producer gas d) Coal gas




464. Acetylene can be obtained by the reaction?



- a) Br, Br₂ b) Br₂, NBS c) NBS, NBS d) NBS, Br₂
476. During ozonolysis of CH₂ = CH₂ if reduction is carried out by LiAlH₄ the products formed are:
a) HCHO b) HCOOH c) CH₃OH d) CH₂OHCH₂OH
477. Ethyl hydrogen sulphate is obtained by reaction of H₂SO₄ on:
a) Ethylene b) Ethane c) Ethyl chloride d) Ethanal
478. When HCl gas is passed through propene in the presence of benzoyl peroxide, it gives:
a) n-propyl chloride b) 2-chloropropane c) Allyl chloride d) No reaction
479. Hydrocarbon which is liquid at room temperature is
a) Pentane b) Butane c) Propane d) Ethane
480. Which of the following reactions are not expected to give




In yields of more than 50%?




d) None of the above

481. Incorrect name of an alkyne is:
a) Propyne b) But-2-yne c) Pent-3-yne d) But-1-yne
482. The alkyne which gives pyruvic acid (CH₃COCOOH) on oxidation with alk. KMnO₄ is:
a) CH ≡ CH b) CH₃C ≡ CH c) CH₃C ≡ C—CH₃ d) CH₃—CH₂—C ≡ CH
483. A hydrocarbon of formula C₆H₁₀ absorbs only one molecule of H₂ upon catalytic hydrogenation. Upon ozonolysis the hydrocarbon yields,




The hydrocarbon is :

- a) Cyclohexane b) Benzene c) Cyclohexene d) Cyclobutane
484. Alkyl halides react with dialkyl copper reagents to give
a) Alkenyl halides b) Alkanes
c) Alkyl copper halides d) Alkenes




Thus, A is

- a) (CH₃)₂CHCH = CHCH = CH₂ b) (CH₃)₂C = CH — CH = CH₂
c) CH₃CH = CH — CH = CH — CH = CH₂ d) none of the above
486. In the series,







The compound Z is

- a) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ b) CH_3COCH_3 c) CH_3CHO d) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$
487. Paraffin dissolves in:
 a) Distilled water b) Benzene c) Methanol d) Salt water
488. Which cannot be prepared by Kolbe's electrolytic reaction using single salt?
 a) CH_4 b) C_2H_6 c) C_4H_{10} d) H_2
489. Which will react with NaBH_4 ?
 a) Benzoic acid b) Benzamide c) Cyclohexanone d) Acetic acid
490. When methane is made to react with a halogen (X_2), halides are formed, the order of reactivity is:
 a) $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$ b) $\text{Cl}_2 > \text{F}_2 > \text{Br}_2 > \text{I}_2$ c) $\text{I}_2 > \text{Br}_2 > \text{Cl}_2 > \text{F}_2$ d) $\text{Cl}_2 > \text{F}_2 > \text{I}_2 > \text{Br}_2$
491. Which of these does not follow *anti* -Markownikoff's rule?
 a) 2-butene b) 1-butene c) 2-pentene d) 2-hexene
492. Acetylene reacts with HCN in the presence of $\text{Ba}(\text{CN})_2$ to yield
 a) 1,1-dicyanoethane b) 1,2-dicyanoethane c) Vinyl cyanide d) None of these
493. An alkyl bromide (X) reacts with Na to form 4, 5-diethyl octane. Compound (X) is:
 a) $\text{CH}_3(\text{CH}_2)_3\text{Br}$
 b) $\text{CH}_3(\text{CH}_2)_5\text{Br}$
 c) $\text{CH}_3(\text{CH}_2)_3\text{CHBr}.\text{CH}_3$
 d) $\text{CH}_3(\text{CH}_2)_2\text{CHBrCH}_2\text{CH}_3$
494. To avoid lead pollution, a new anti-knock compound is used. It is:
 a)  b) Cyclopentadienyl manganese carbonyl
 c) AK-33-X d) All of the above
495. Identify B and D in the following sequence of reactions.
- ```

 graph TD
 A[CH2=CH2] -- "Conc. H2SO4" --> B[A]
 B -- "H2O, Δ" --> C[B]
 C -- "PBr3" --> D[C]
 D -- "D" --> A





```
- a) Methanol and bromoethane    b) Ethyl hydrogen sulphate and alcoholic KOH  
 c) Ethyl hydrogen sulphate and aqueous KOH    d) Ethanol and alcoholic KOH
496. Angle strain in cyclopropane is  
 a)  $24^\circ 44'$     b)  $9^\circ 44'$     c)  $44'$     d)  $-5^\circ 16'$
497. When propyne react with  $\text{H}_2\text{O}$  in presence of dil.  $\text{H}_2\text{SO}_4$  and  $\text{HgSO}_4$  product formed is  
 a) Acetone    b) Acetaldehyde    c) Acetic acid    d) Ethyl alcohol
498. Which of the following compounds cannot be prepared singly by the Wurtz reaction?  
 a)  $\text{C}_2\text{H}_6$     b)  $(\text{CH}_3)_2\text{CHCH}_3$     c)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$     d) All can be prepared
499. The olefin which on ozonolysis gives  $\text{CH}_3\text{CH}_2\text{CHO}$  and  $\text{CH}_3\text{CHO}$  is:  
 a) 1-butene    b) 2-butene    c) 1-pentene    d) 2-pentene
500. Which statement is false?  
 a) Peroxide effect is applicable only for  $\text{HBr}$  and not for the other halogen halides




- b) Meta directing groups are deactivating groups  
 c) Chlorination of methane follows an ionic mechanism  
 d) In benzene the C atoms are  $sp^2$ -hybridized
501. The presence of unsaturation (olefinic or acetylinic bond) in an organic compound can be tested with:  
 a) Schiff's reagent      b) Tollen's reagent      c) Fehling's solution      d) Baeyer's reagent
502. An alkene on reductive ozonolysis gives 2-molecules of  $\text{CH}_2(\text{CHO})_2$ . The alkene is  
 a) 2,4-hexadiene      b) 1,3-cyclohexadiene  
 c) 1,4-cyclohexadiene      d) 1-methyl-1, 3-cyclopentadiene
503. A mixture of ethyl iodide and *n*-propyl iodide is subjected to Wurtz reaction. The hydrocarbon that will not be formed is:  
 a) *n*-butane      b) *n*-propane      c) *n*-pentane      d) *n*-hexane
504. Which of the following reacts with benzene in presence of anhydrous aluminium chloride and forms acetophenone?  
 a)  $\text{CH}_3\text{Cl}$       b)  $\text{CH}_3\text{COOH}$       c)  $\text{CH}_3\text{CHO}$       d)  $\text{CH}_3\text{COCl}$
505. Oxidation of 1-butene with hot  $\text{KMnO}_4$  solution produces  
 a)  $\text{CH}_3\text{CH}_2\text{COOH} + \text{HCOOH}$       b)  $\text{CH}_3\text{CH}_2\text{COOH} + \text{CO}_2$   
 c)  $\text{CH}_3\text{COOH} + \text{CO}_2$       d)  $(\text{CH}_3)_2\text{C} = \text{O} + \text{CO}_2$
506. Action of  $\text{Br}_2$  on cyclopentene gives:  
 a) 1,2-dibromo cyclopentane  
 b) Cyclopentyl bromide  
 c) Cyclopentyl dibromide  
 d) No reaction
507. Which of the following species is aromatic?  
 a)       b)       c)       d) 
508. Propene,  $\text{CH}_3-\text{CH}=\text{CH}_2$  can be converted into 1-propanol by oxidation. Which set of reagents among the following is ideal to effect the conversion?  
 a) Alkaline  $\text{KMnO}_4$       b)  $\text{B}_2\text{H}_6$  and alk.  $\text{H}_2\text{O}_2$       c)  $\text{O}_3/\text{zinc dust}$       d)  $\text{OsO}_4/\text{CHCl}_3$
509. Compound which gives acetone on ozonolysis  
 a)  $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_3$       b)  $(\text{CH}_3)_2\text{C}=\text{C}(\text{CH}_3)_2$   
 c)  $\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$       d)  $\text{CH}_3\text{CH}=\text{CH}_2$
510. Toluene, when treated with  $\text{Br}_2/\text{Fe}$ , gives *p*-bromotoluene as the major product because the  $-\text{CH}_3$  group  
 a) Is *meta* directing      b) deactivates the ring  
 c) activates the ring by hyperconjugation      d) None of the above
511. Alkynes occur in nature in the:  
 a) Free state      b) Partially free state      c) Not in the free state      d) None of the above
512. Which of the following will have least hindered rotation about carbon-carbon bond?  
 a) Ethane      b) Ethylene      c) Acetylene      d) Hexachloroethane
513. Identify Z in the series,  

$$\text{CH}_2 = \text{CH}_2 \xrightarrow{\text{HBr}} \text{X} \xrightarrow{\text{aq.KOH}} \text{Y} \xrightarrow[\text{I}_2 \text{ excess}]{\text{NaCO}_3} \text{Z}$$
 a)  $\text{C}_2\text{H}_5\text{I}$       b)  $\text{C}_2\text{H}_5\text{OH}$       c)  $\text{CHI}_3$       d)  $\text{CH}_3\text{CHO}$
514. Action of  $\text{NH}_3$  over  $\text{C}_2\text{H}_2$  at high temperature gives:  
 a) Amine      b) Furan      c) Thiophene      d) Pyrrole
515. Wurtz reaction converts alkyl halide into alkane when it is made to react with  
 a) Na in alcohol      b) Na in dry ether      c) Zn in alcohol      d) Zn in dry ether
516. Polyethylene is a resin obtained by polymerization of:

- a) Butadiene                      b) Ethylene                      c) Isoprene                      d) Styrene
517. Cyclohexane ( $C_6H_{12}$ ) a hydrocarbon, floats on water because:
- It is immiscible with water
  - Its density is less than that of water
  - It is a non-polar substance
  - It is immiscible and lighter than water
518. Which of the following are produced from coaltar?
- Synthetic dyes
  - Drugs
  - Perfumes
  - All of these
519. The reduction of an alkyne to alkene using lithium metal in liquid ammonia as solvent results into
- cis* addition of hydrogen atoms
  - trans* addition of hydrogen atoms
  - Both *cis* and *trans* additions of hydrogen atoms. The relative amounts of the two depends on temperature
  - Both *cis* and *trans* additions of hydrogen atoms. The relative amounts depend on the nature of alkyne
520. Propene on reaction with hypochlorous acid to give







521. A mixture of nitrogen and acetylene, on passing electric spark through it gives:
- Hydrogen and carbon
  - Hydrogen cyanide
  - Nitromethane
  - Nitroethane
522. In the sequence of reactions,



Compound Z is

- N-methyl ethanamine
  - N-propylamine
  - N, N-dimethylamine
  - Ethyl cyanide
523. Which one of these is not true for benzene?
- It forms only one type of monosubstituted product.
  - There are three carbon-carbon single bonds and three carbon-carbon double bonds
  - The heat of hydrogenation of benzene is less than the theoretical value.
  - The bond angle between the carbon-carbon bonds is  $120^\circ$ .
524. Presence of a nitro group in a benzene ring
- Activates the ring towards electrophilic substitution
  - Renders the ring basic
  - Deactivates the ring towards nucleophilic substitution
  - Deactivates the ring towards electrophilic substitution

525. The major product in the reaction of 2-butyne with  $Li/liq. NH_3$  is



526. Hydrocarbon liquid at STP is:

- a) Ethane                                      b) Propane                                      c) Butane                                      d) Pentane
527. Chlorination of benzene is not possible in the following reaction
- a)  $C_6H_6 + Cl_2 \xrightarrow{FeCl_3}$                       b)  $C_6H_6 + HOCl \xrightarrow{H^+}$                       c)  $C_6H_6 + I-Cl \xrightarrow{ZnCl_2}$                       d)  $C_6H_6 + Cl_2 \xrightarrow{AlCl_3}$

528. In the series, ethane, ethene and ethyne, the C-H bond energy is
- a) Same in all the three compounds                      b) Greatest in ethane
- c) Greatest in ethene                      d) Greatest in ethyne


529. The shape of 2-butene is:
- a) Linear                                      b) Planar                                      c) Tetrahedral                                      d) Pyramidal

530. The substance used as an anti-knock compound is:
- a) Tetraethyl lead                      b) Lead tetrachloride                      c) Lead acetate                      d) Ethyl acetate


531. Petroleum refining is:
- a) Obtaining aromatic compounds from aliphatic compounds in petroleum                      b) Cracking of petroleum to get gaseous hydrocarbons                      c) Purification of petroleum                      d) Distillation of petroleum to get different fractions

532. Zinc-copper couple that can be used as a reducing agent is obtained by:
- a) Mixing zinc dust and copper gauze
- b) Zinc coated with copper
- c) Copper coated with zinc
- d) Zinc and copper wires welded together

533. Which of the following hydrocarbons has the lowest dipole moment?




534.  $(A) \xrightarrow[\Delta]{KMnO_4} \text{CH}_3COCH_2CH_2CH_2COOH$  is



535. A solution of sodium salt of fatty acid was electrolysed during Kolbe's reaction. The solution left after electrolysis is:


- a) Richer in NaOH                      b) Richer in  $H_2SO_4$                       c) Richer in sodium salt                      d) All of these

536. Sample of 2,3-dibromo-3-methylpentane is heated with zinc dust. The resulting product is isolated and heated with HI in the presence of phosphorus. Indicate which is the structure that represents the final organic product in the reaction?



537. Which compound does not give precipitate with ammoniacal silver nitrate solution?


- a)  $C_2H_5 - C \equiv CH$                       b)  $CH_3 - C \equiv C - CH_3$
- c)  $CH_3$                       d)  $Ph - CH_2 - C \equiv CH$




538. Hydroxylation of propyne in the presence of  $\text{HgSO}_4/\text{H}_2\text{SO}_4$  is initiated by the attack of:  
 a) Carbene                      b) Free radical                      c) Electrophile                      d) Nucleophile
539. Benzene vapour mixed with air when passed over  $\text{V}_2\text{O}_5$  catalyst at 775 K gives  
 a) Glyoxal                      b) Oxalic acid                      c) Maleic anhydride                      d) Fumaric acid
540. Kolbe's synthesis on electrolysis of sodium salt of butanoic acid gives :  
 a) *n*-hexane                      b) Isobutene                      c) Butane                      d) Ethene
541. Which among the following is aromatic?




542. The neutral wax called Ozokerite found near petroleum well is a mixture of:  
 a) Solid halides                      b) Solid hydrocarbons                      c) Solid alcohols                      d) None of these



Identify A:



544. Which among the following are used as catalyst in cracking?  
 a) Oxides of Al                      b) Oxides of Cr, Mo                      c) Oxides of V                      d) All of these
545. The general formula of a cycloalkane is  
 a)  $\text{C}_n\text{H}_n$                       b)  $\text{C}_n\text{H}_{2n}$                       c)  $\text{C}_n\text{H}_{2n-2}$                       d)  $\text{C}_n\text{H}_{2n+2}$
546. Toluene reacts, with excess of  $\text{Cl}_2$  in presence of sunlight to give a product, which on hydrolysis followed by reaction with  $\text{NaOH}$  gives







547. Which of the following alkanes can be easily sulphonated?  
 a) *n*-butane                      b) Isobutene                      c) *n*-pentane                      d) *n*-hexane
548. When propionic acid is treated with aqueous sodium bicarbonate,  $\text{CO}_2$  is liberated. The 'C' of  $\text{CO}_2$  comes from:  
 a) Methyl group                      b) Carboxylic group                      c) Methylene group                      d) Bicarbonate
549. 10mL of a certain hydrocarbon require 25mL of oxygen for complete combustion and the volume of  $\text{CO}_2$  produced is 20mL. what is the formula of hydrocarbon?  
 a)  $\text{C}_2\text{H}_2$                       b)  $\text{C}_2\text{H}_4$                       c)  $\text{CH}_4$                       d)  $\text{C}_2\text{H}_6$
550. Which of the following compounds is the most stable?



551. The octane number of any fuel increases with:  
 a) Increase in *n*-heptane  
 b) Decrease in 2,2,4-trimethylpentane

- c) Increase in 2,2,4-trimethylpentane  
d) None of the above


552. 3-methyl-2-pentene on reaction with HOCl gives:



553. The reaction of propene with HOCl proceeds *via* the addition of

- a)  $\text{Cl}^+$  and  $\text{OH}^-$  in a single step  
b)  $\text{Cl}^+$  in the first step  
c)  $\text{H}^+$  in the first step  
d)  $\text{OH}^-$  in the first step

554. Select the reagent for the following reaction,




- a)  $\text{SeO}_2$   
b)  $\text{O}_3, \text{Zn}/\text{H}_2\text{O}$   
c)  $\text{O}_3, \text{H}_2\text{O}_2 - \text{CH}_3\text{COOH}$   
d) PCC

555. The chemical reactivity of ethylene is due to:

- a) Short carbon to carbon bond distance  
b) High double bond energy  
c) Trigonal planar structure  
d) Presence of  $\pi$ -electrons

556. Which of the following species could be expected to exhibit aromatic character?



Select the correct answer from the following

- a) I and IV  
b) II and IV  
c) I and III  
d) II and III

557. Product formed when 1-butene is subjected to HBr in the presence of peroxide:

- a) 1-bromobutane  
b) 2-bromobutane  
c) 1,1-dibromobutane  
d) 1,2-dibromobutane



558. Nitrobenzene can be prepared from benzene by using a mixture of concentrated  $\text{HNO}_3$  and concentrated  $\text{H}_2\text{SO}_4$ . In the nitrating mixture,  $\text{HNO}_3$  acts as

- a) Base  
b) Acid  
c) Reducing agent  
d) Catalyst

559. In the reaction sequence,








- a) 1 and 4                      b) 2 and 3                      c) 1 and 2                      d) 1 and 3

582. A mixture of ethane, ethene and ethyne is passed through ammoniacal  $\text{AgNO}_3$  solution. The gases which remain unreacted are:


- a) Ethane and ethene              b) Ethane and ethyne              c) Ethene and ethyne              d) Ethane only

583. In the reaction,



The product C is

- a)  $\text{C}_6\text{H}_5\text{OH}$                       b)  $\text{C}_6\text{H}_6$                       c)  $\text{C}_6\text{H}_5\text{COONa}$                       d)  $\text{C}_6\text{H}_5\text{ONa}$



Identify A and B

- a)  $\text{CH}_3\text{CHO}, \text{CH}_3\text{COCH}_3$                       b)  $\text{CH}_3\text{CH}_2\text{CHO}, \text{CH}_3\text{COCH}_3$   
 c)  $\text{CH}_3\text{CH}_2\text{CHO}, \text{CH}_3\text{COCH}_2\text{CH}_3$                       d)  $\text{HCHO}, \text{CH}_3\text{COCH}_3$

585. Cyclobutadiene is said to be

- a) aromatic                      b) aliphatic                      c) non-aromatic                      d) None of these

586. Acetylene reacts with hypochlorous acid to form

- a)  $\text{Cl}_2\text{CHCHO}$                       b)  $\text{ClCH}_2\text{COOH}$                       c)  $\text{Cl}_3\text{COCl}$                       d)  $\text{ClCH}_2\text{CHO}$

587. To enable easy detection of gas leakage from cylinders, the substance added to L.P.G. is:

- a) Glycols                      b) Phenols                      c) Thioalcohols                      d) Glycerols

588. Octane no. of 2,3,3-trimethylbutane has been assumed to be:

- a) 100                      b) -45                      c) 124                      d) Zero

589.  $\text{C}_4\text{H}_6$  may contain

- a) One double bond                      b) Two double bond                      c) One triple bond                      d) Both (b) and (c)

590. Which of the following compounds can form metallic derivatives?

- a) Ethane                      b) Propyne                      c) 2-butyne                      d) 2-butene

591. Increasing order of volatility of  $\text{C}_2\text{H}_6, \text{C}_2\text{H}_4, \text{C}_2\text{H}_2$  and  $\text{C}_6\text{H}_6$  is:

- a)  $\text{C}_6\text{H}_6, \text{C}_2\text{H}_6, \text{C}_2\text{H}_4, \text{C}_2\text{H}_2$       b)  $\text{C}_2\text{H}_2, \text{C}_2\text{H}_4, \text{C}_2\text{H}_6, \text{C}_6\text{H}_6$       c)  $\text{C}_6\text{H}_6, \text{C}_2\text{H}_2, \text{C}_2\text{H}_4, \text{C}_2\text{H}_6$       d)  $\text{C}_2\text{H}_2, \text{C}_2\text{H}_6, \text{C}_2\text{H}_4, \text{C}_6\text{H}_6$

592. Octane no. of a fuel can be increased by:

- a) Isomerism                      b) Alkylation                      c) Reforming                      d) All of these

593. 1-propanol on dehydration with  $\text{H}_2\text{SO}_4$  produces:

- a)  $\text{CH}_3 - \text{CH} = \text{CH}_2$   
 b)  $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_3$   
 c)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{CH}_3$   
 d)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH} = \text{CH}_2$

594. Propadiene,  $\text{C}_3\text{H}_4$  molecule contains:

- a) Two  $sp^2$  and one  $sp$ -hybrid carbon  
 b) One  $sp^2$  and two  $sp$ -hybrid carbons  
 c) One  $sp^2$  and three  $sp$ -hybrid carbons  
 d) None of the above



595. Catalyst used in dimerization of acetylene to prepare chloroprene is:

- a)  $\text{HgSO}_4 + \text{H}_2\text{SO}_4$                       b)  $\text{Cu}_2\text{Cl}_2$                       c)  $\text{Cu}_2\text{Cl}_2 + \text{NH}_4\text{Cl}$                       d)  $\text{Cu}_2\text{Cl}_2 + \text{NH}_4\text{OH}$

596. Cyclopentene on treatment with alkaline  $\text{KMnO}_4$  gives:

- a) Cyclopentanol




- b) *trans*-1,2-cyclopentanediol  
 c) *cis*-1,2-cyclopentanediol  
 d) 1 : 1 mixture of *cis*- and *trans*-1,2-cyclopentanediol
597.  $C_7H_8 \xrightarrow{3Cl_2, Heat} A \xrightarrow{Fe/Br_2} B \xrightarrow{Zn/HCl} C$   
 Here, the compound *C* is  
 a) 3-bromo 2,4,5,6-trichlorotoluene  
 b) *o*-bromo toluene  
 c) *p*-bromo toluene  
 d) *m*-bromo toluene
598. Naphalene is an example of  
 a) Polynuclear hydrocarbon  
 b) alicyclic compound  
 c) heterocyclic compound  
 d) aliphatic compound
599. Which of the following will give *trans*-diols?  
 a)  $\text{>C=C<} \xrightarrow[2. H_2O]{1. KMnO_4}$   
 b)  $\text{>C=C<} \xrightarrow[2. Na_2SO_3]{1. OsO_4}$   
 c)   $\xrightarrow[2. Na_2SO_3]{1. OsO_4, 25^\circ C}$   
 d)   $\xrightarrow[HCO_2H, 25^\circ C]{35\% H_2O_2}$
600. Benzene can react with  
 a) Bromine water  
 b)  $HNO_3$   
 c)  $H_2O$   
 d)  $CH_3OH$
601. A mixture of methane and steam when passes over nickel supported on alumina catalyst at  $725^\circ C$  gives:  
 a)  $CH_3OH$   
 b)  $CO_2$  and  $H_2$   
 c)  $CO$  and  $H_2$   
 d) None of these
602. In which reaction addition takes place according to Markownikoff's rule?  
 a)  $CH_3CH = CH_2 + Br_2 \rightarrow$   
 b)  $CH_3CH = CH_2 + HBr \rightarrow$   
 c)  $CH_2 = CH_2 + HBr \rightarrow$   
 d)  $CH_3CH = CHCH_3 + Br_2 \rightarrow$
603. Paraffin wax is:  
 a) Ester  
 b) Alcohol  
 c) Unsaturated hydrocarbon  
 d) Saturated hydrocarbon
604. Propyne when passed through a hot iron tube at  $400^\circ C$  produces  
 a) Benzene  
 b) Methyl benzene  
 c) Dimethyl benzene  
 d) Trimethyl benzene
605. Which of the following is called Marsh gas?  
 a)  $C_2H_4$   
 b)  $C_2H_6$   
 c)  $C_2H_2$   
 d)  $CH_4$
606. Which can be easily oxidized?  
 a) Alkene  
 b) 1-alkyne  
 c) Alkane  
 d) Benzene
607. *n*-butane and isobutene, which have same number of hydrogen and carbon atoms in their molecules, boil at different temperatures because:  
 a) *n*-butane is much hotter  
 b) Their volumes are different  
 c) Isobutene is an alkene  
 d) Their atoms are not having the same carbon chain
608. Common oxidizing agents used in organic chemistry are:  
 a) Fenton's reagent  
 b) Osmium tetroxide  
 c) Acidified  $KMnO_4$   
 d) Alkaline  $KMnO_4$
609. Acetylenic hydrocarbons are acidic because:  
 a) Sigma electron density of C—H bond in acetylene is nearer a carbon which has 50% *s*-character  
 b) Acetylene has only one hydrogen atom at each carbon atom  
 c) Acetylene contains least number of hydrogen atoms among the possible

- d) Acetylene belongs to the class of alkynes with formula  $C_nH_{2n-2}$
610. Butene -1 may be converted to butane by the reaction with:  
 a) Zn-Hg                      b) Pd-H<sub>2</sub>                      c) Zn-HCl                      d) Sn-HCl
611. Number of acidic hydrogen atom in butyne-1 is:  
 a) 2                              b) 3                              c) 1                              d) 4
612. Propene on reaction with methylene iodide in presence of Zn-Cu couple gives:  
 a) Cyclopropane              b) Cyclopropene              c) Methyl cyclopropane      d) Cyclobutene
613. Addition of O<sub>2</sub> on ethylene in presence of Ag at 200°C forms:  
 a) Epoxy ethane              b) Oxiranes                      c) Cyclic ethers              d) All of these
614. The carbon-carbon bond distance in benzene is  
 a) Longer than a C - C single bond                      b) Longer than a C = C double bond  
 c) Shorter than a C = C double bond                      d) Shorter than a C ≡ C triple bond
615. Method of converting high boiling hydrocarbons into low boiling hydrocarbons is called:  
 a) Polymerisation              b) Isomerisation              c) Cracking                      d) Condensation
616. The mechanism of Wurtz reaction involves:  
 a) Free radical                      b) Carbocation                      c) Carbanion                      d) None of these
617. The most important energy yielding constituent in biogas is:  
 a) C<sub>2</sub>H<sub>4</sub>                              b) C<sub>2</sub>H<sub>2</sub>                              c) CH<sub>4</sub>                              d) H<sub>2</sub>S
618. PVC is a polymer of:  
 a) CH<sub>2</sub> = CH<sub>2</sub>                      b) ClCH<sub>2</sub>—CH<sub>2</sub>Cl              c) CH<sub>2</sub>—CHCl                      d) Cl—C = C—Cl
619. Cyclohexene on ozonolysis followed by reaction with zinc dust and water gives compound *E*. Compound *E* on further treatment with aqueous KOH yields compound *F*. Compound *F* is




620. The flash point in India is fixed at:  
 a) 44°C                              b) 35°C                              c) 22.8°C                              d) 30°C
621. Lindlar's catalyst is:  
 a) Pd- CaCO<sub>3</sub> deactivated by lead acetate  
 b) Pd - BaSO<sub>4</sub>  
 c) Pd  
 d) None of the above
622. The energy of π-bond in kcal is about :  
 a) 36                              b) 50                              c) 74                              d) 140
623. Ozonolysis (O<sub>3</sub>, H<sub>2</sub>O) of,  

$$\begin{array}{c} \text{CH}_3-\text{CH}-\text{C}\equiv\text{C}-\text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$$
 gives:  
 a) 
$$\begin{array}{c} \text{CH}_3-\text{CHCOOH} + \text{CH}_3\text{COOH} \\ | \\ \text{CH}_3 \end{array}$$
  
 b) 
$$\begin{array}{c} \text{CH}_3-\text{CHCHO} + \text{CH}_3\text{CHO} \\ | \\ \text{CH}_3 \end{array}$$
  
 c) 
$$\begin{array}{c} \text{CH}_3-\text{CHCHO} + \text{CH}_3\text{COOH} \\ | \\ \text{CH}_3 \end{array}$$
  
 d) None of the above
624. What is the end product of the following sequences of operations?




- a) Methyl alcohol                      b) Acetaldehyde                      c)  $\text{C}_2\text{H}_5\text{OH}$                       d)  $\text{C}_2\text{H}_4$
625. The order of relative acidic strengths of water, ethyne and propyne is:  
 a) Water > propyne > ethyne  
 b) Propyne > ethyne > water  
 c) Water > ethyne > propyne  
 d) Ethyne > water > propyne
626. Reaction of *trans*-2-phenyl-1-bromocyclopentane on reaction with alcoholic KOH produces:  
 a) 4-phenylcyclopentene  
 b) 2-phenylcyclopentene  
 c) 1-phenylcyclopentene  
 d) 3-phenylcyclopentene
627. Ethylene reacts with sulphur monochloride to give:  
 a) Phosgene                      b) Mustard gas                      c) Ethylene chloride                      d) None of these
628. The dihalogen derivative 'X' of a hydrocarbon with three carbon atoms reacts with alcoholic KOH and produces another hydrocarbon which forms a red precipitate with ammoniacal  $\text{Cu}_2\text{Cl}_2$ . 'X' gives an aldehyde on reaction with aqueous KOH. The compound 'X' is  
 a) 1,3-dichloropropane                      b) 1,2-dichloropropane  
 c) 2,2-dichloropropane                      d) 1,1-dichloropropane
629. Ethylene may be prepared by the dehydration of:  
 a) Ethyl alcohol                      b) Methyl alcohol                      c) Acetic acid                      d) Oxalic acid
630. Petroleum is formed by the chemical changes in:  
 a) Inorganic matter                      b) Vegetable matter                      c) Animal matter                      d) Both (b) and (c)
631. Common dehydrating agents for alkanes are:  
 a)  $\text{H}_2\text{SO}_4$                       b)  $\text{Al}_2\text{O}_3$                       c)  $\text{ZnCl}_2$                       d) All of the above
632. The most stable conformation of butane is:  
 a) Skew                      b) Staggered                      c) Gauche                      d) Eclipsed
633. A cyclic hydrocarbon molecule has all the carbon and hydrogen in a single plane. All the carbon-carbon bonds are of same length, less than  $1.54\text{\AA}$ , but more than  $1.34\text{\AA}$ . The C-c bond angle will be  
 a)  $109^\circ 28'$                       b)  $100^\circ$                       c)  $180^\circ$                       d)  $120^\circ$
634. The product of acid catalysed hydration of 2-phenyl propene is:  
 a) 3-phenyl-2-propanol                      b) 1-phenyl-2-propanol                      c) 2-phenyl-2-propanol                      d) 2-phenyl-1-propanol
635. When  $\text{C}_2\text{H}_5$ ,  $\text{CH}_4$  and  $\text{C}_2\text{H}_4$  passes through a test tube which have ammoniacal  $\text{Cu}_2\text{Cl}_2$ , find out which gas comes out unaffected from test tube?  
 a)  $\text{C}_2\text{H}_2$  and  $\text{CH}_4$                       b)  $\text{C}_2\text{H}_2$  and  $\text{C}_2\text{H}_4$                       c)  $\text{C}_2\text{H}_4$  and  $\text{CH}_4$                       d)  $\text{C}_2\text{H}_2$
636. Benzene reacts with chlorine in sunlight to give a final product  
 a)  $\text{CCl}_4$                       b)  $\text{C}_6\text{H}_6\text{Cl}_6$                       c)  $\text{C}_6\text{Cl}_6$                       d)  $\text{C}_6\text{H}_5\text{Cl}$
637. When 2-butyne is treated with Pd –  $\text{BaSO}_4$ ; the product formed will be  
 a) *cis*-2-butene                      b) *trans*-2-butene                      c) 1-butene                      d) 2-hydroxy butane
638. The overlapping of orbitals in benzene is of the type  
 a)  $sp - sp$                       b)  $p - p$                       c)  $sp^2 - sp^2$                       d)  $sp^3 sp^3$
639. The product obtained when methyl magnesium bromide reacts with methyl alcohol is:  
 a) Acetone                      b) Alcohol                      c) Methane                      d) Ethane
640. The treatment of benzene with benzoyl chloride in the presence of  $\text{AlCl}_3$  gives  
 a) Benzaldehyde                      b) Benzophenone                      c) Diphenyl                      d) Cyclohexane
641. Which of the following have delocalised electron?

- a) Benzene                      b) Cyclohexane                      c) CH<sub>4</sub>                      d) C<sub>2</sub>H<sub>6</sub>
642. The IUPAC name of CH<sub>2</sub> = CH—CH<sub>2</sub>—group is:  
 a) Allyl                      b) Propyl                      c) Prop-2-enyl                      d) Prop-1-enyl
643. Which statement is correct?  
 a) Low chemical reactivity of alkanes is due to strong C—C and C—H bonds  
 b) Alkanes show characteristic substitution reactions because they are saturated  
 c) Reaction of alkanes with fluorine is explosive even in dark  
 d) All of the above
644. Ease of sulphonation of alkanes is:  
 a) 3° > 2° > 1°                      b) 1° > 2° > 3°                      c) 2° > 3° > 1°                      d) 3° > 1° > 2°
645. Arrange the following in order of decreasing boiling point



- a) I > II > III > IV                      b) IV > III > II > I                      c) I > III > IV > II                      d) II > III > I > IV

646. The product *B* is:



- a)  $\text{CH}_3 \cdot \text{CH}_2 \cdot \text{CH}_2 \cdot \overset{\text{I}}{\underset{\text{Cl}}{\text{C}}} - \text{H}$                       b)  $\text{CH}_3 \cdot \text{CH}_2 \cdot \overset{\text{I}}{\underset{\text{Cl}}{\text{C}}} \cdot \text{CH}_2$                       c) CH<sub>3</sub>CH<sub>2</sub>C≡CH                      d) CH<sub>3</sub>CH=CHCH<sub>3</sub>

647. *n*-propyl bromide on treating with alcoholic KOH produces

- a) Propane                      b) Propene                      c) Propyne                      d) Propanol

648. An unsaturated hydrocarbon upon ozonolysis gives one mole each of formaldehyde, acetaldehyde and methylglyoxal(CH<sub>3</sub>COCHO). The structure of the hydrocarbon is

- a) CH<sub>2</sub> = CH — CH<sub>2</sub> — CH = CH<sub>2</sub>                      b) CH<sub>2</sub> = CH — C(CH<sub>3</sub>) = CH — CH<sub>3</sub>  
 c) (CH<sub>3</sub>)<sub>2</sub>C = CH — CH<sub>3</sub>                      d) CH<sub>3</sub> — CH = C(CH<sub>3</sub>) — CH<sub>3</sub>

649. Fischer-Tropsch process is used in the manufacture of:

- a) Synthetic petrol                      b) Ethanol                      c) Benzene                      d) Ethanoic acid

650. 2-methylpropene is isomeric with butane-1. They can be distinguished by:

- a) Baeyer's reagent                      b) Ammoniacal AgNO<sub>3</sub>                      c) Br<sub>2</sub> solution                      d) O<sub>3</sub>, Zn/H<sub>2</sub>O

651. Acetylene reacts with 42% H<sub>2</sub>SO<sub>4</sub> containing 1% HgSO<sub>4</sub> to give:

- a) C<sub>2</sub>H<sub>5</sub>HSO<sub>4</sub>                      b) CH<sub>3</sub>CHO                      c) HCHO                      d) CH<sub>2</sub> = CH<sub>2</sub>


652. The simplest alkyne is:

- a) CH                      b) CH<sub>2</sub>                      c) C<sub>2</sub>H<sub>2</sub>                      d) C<sub>2</sub>H<sub>4</sub>

653. A Friedel-Crafts reaction of benzene with chloroform produces

- a) C<sub>6</sub>H<sub>5</sub>CHCl<sub>2</sub>                      b)  $\text{C}_6\text{H}_5 - \overset{\text{Cl}}{\underset{\text{H}}{\text{C}}} - \text{C}_6\text{H}_5$                       c)  $\text{C}_6\text{H}_5 - \overset{\text{C}_6\text{H}_5}{\underset{\text{H}}{\text{C}}} - \text{C}_6\text{H}_5$                       d) All of these

654. An alkene, obtained by the dehydration of an alcohol (*A*), on ozonolysis gives two molecules of acetaldehyde for every molecule of alkene. The alcohol (*A*) is:



- a)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$       b)  $\text{CH}_3\text{CH}_2\text{OH}$       c)  $\text{CH}_3\text{CH} = \text{CHCH}_2\text{OH}$       d)  $\text{CH}_3\text{CH}_2\text{CHCH}_3$   
|  
OH

655. Which of the following annulenes is *anti*-aromatic?

- a) Benzene      b) Cyclobutadiene      c) Cyclodecapentene      d) Cyclooctatetraene

656. The number of possible isomers of alkane with formula  $\text{C}_6\text{H}_{14}$  is:

- a) 2      b) 3      c) 4      d) 5


657. Which statement is correct?

- a) Alkanes from  $\text{CH}_4$  to  $\text{C}_4\text{H}_{10}$  are colourless odourless gases  
b) Alkanes from  $\text{C}_5\text{H}_{12}$  to  $\text{C}_{17}\text{H}_{36}$  are colourless liquids  
c) All alkanes are lighter than water  
d) Melting point of alkanes increases with increase in the number of carbon atoms

658. Which compound does not decolourize bromine dissolved in carbon tetrachloride?

- a)  $\text{C}_2\text{H}_2$       b)  $\text{C}_3\text{H}_6$       c)  $\text{C}_6\text{H}_6$       d)  $\text{C}_2\text{H}_4$

659. The principal organic product formed in the reaction,



- a)  $\text{CH}_3 - \text{CHBr}(\text{CH}_2)_8\text{COOH}$   
b)  $\text{CH}_2 = \text{CH}(\text{CH}_2)_8\text{COBr}$   
c)  $\text{CH}_2\text{BrCH}_2(\text{CH}_2)_8\text{COOH}$   
d)  $\text{CH}_2 = \text{CH}(\text{CH}_2)_7\text{CHBrCOOH}$

660. What would be the product formed when 1-bromo-3-chlorocyclobutane reacts with two equivalents of metallic sodium in ether?

- a)       b)       c)       d) 

661.  $[A] \xleftarrow[\text{catalyst}]{\text{Lindlar's}} \text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_3 \xrightarrow[\text{liq. NH}_3]{\text{Na in}} [B]$

- [A] and [B] are respectively  
a) *cis, trans*-2-butene      b) Both *trans*-2-butene  
c) *trans, cis*-2-butene      d) Both *cis*-2-butene

662. Which of the following reacts with  $\text{KMnO}_4$  but does not react with  $\text{AgNO}_3$ ?

- a)  $\text{C}_2\text{H}_6$       b)  $\text{CH}_4$       c)  $\text{C}_2\text{H}_4$       d)  $\text{C}_2\text{H}_2$

663. Octane number 116 is given for:

- a) 2,2,2-trimethyl pentane  
b) 2,3,4-trimethyl pentane  
c) 2,2,3-trimethyl butane  
d) 2,2,4-trimethyl butane

664. Which of the following statements is incorrect?

- a) Acetylene is explosive above 2 atm  
b) It is transported by dissolving in acetone  
c) It has unpleasant garlic odour  
d) It is used in the manufacture of Lewisite

665. Formation of ethylene from ethyl bromide is a case of:


- a) Addition reaction  
b) Substitution reaction  
c) Elimination reaction  
d) Rearrangement reaction

666. The most stable alkene is,





- a)  $\text{R}_2\text{C} = \text{CR}_2$       b)  $\text{RCH} = \text{CHR}$       c)  $\text{CH}_2 = \text{CH}_2$       d)  $\text{RCH} = \text{CR}_2$

667. Ethylene can be prepared by electrolysis of an aqueous solution of:

- a) Sodium acetate      b) Sodium succinate      c) Sodium fumarate      d) Sodium propionate
668. HBr reacts with  $\text{CH}_2 = \text{CH} - \text{OCH}_3$  under anhydrous conditions at room temperature to give
- a)  $\text{CH}_3\text{CHO}$  and  $\text{CH}_3\text{Br}$       b)  $\text{BrCH}_2\text{CHO}$  and  $\text{CH}_3\text{OH}$   
 c)  $\text{BrCH}_2 - \text{CH}_2 - \text{OCH}_3$       d)  $\text{H}_3\text{C} - \text{CHBr} - \text{OCH}_3$
669. Identify Z in the following series?
- $$\text{CH}_2 = \text{CH}_2 \xrightarrow{\text{HBr}} X \xrightarrow{\text{Hydrolysis}} Y \xrightarrow[\text{I}_2 \text{ excess}]{\text{Na}_2\text{CO}_3} Z$$
- a)  $\text{C}_2\text{H}_5\text{I}$       b)  $\text{CHI}_3$       c)  $\text{CH}_3\text{CHO}$       d)  $\text{C}_2\text{H}_5\text{OH}$
670. Reactive species in halogenation of benzene in cold and dark
- a)  $\text{Cl}^\bullet$       b)  $\text{Cl}^+$       c)  $\text{Cl}^-$       d) None of these
671. An organic alkadiene on reductive ozonolysis produces
- (i)acetaldehyde  
 (ii)acetone  
 (iii)2-methylpropane-1, 3-dial
- The formula of alkadiene will be
- |                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                      |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>a) <math display="block">\begin{array}{c} \text{CH}_3\text{C} = \text{CHCHCH} = \text{CHCH}_3 \\   \quad   \\ \text{CH}_3 \quad \text{CH}_3 \end{array}</math></p> <p>c) <math display="block">\begin{array}{c} \text{CH}_3\text{C} = \text{CHCHC} = \text{CHCH}_3 \\   \quad   \\ \text{CH}_3 \quad \text{CH}_3 \end{array}</math></p> | <p>b) <math display="block">\begin{array}{c} \text{CH}_3\text{CHCH} = \text{CCH} = \text{CHCH}_3 \\   \quad   \\ \text{CH}_3 \quad \text{CH}_3 \end{array}</math></p> <p>d) <math display="block">\begin{array}{c} \text{CH}_3\text{CH}_2\text{CHCH} = \text{CHC} = \text{CH}_2 \\   \quad   \\ \text{CH}_3 \quad \text{CH}_3 \end{array}</math></p> |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
672. Synthetic petrol and kerosene can be obtained by passing.....under heat and pressure over coal.
- a)  $\text{O}_2$       b)  $\text{H}_2$       c)  $\text{N}_2$       d)  $\text{CO}_2$
673. A hydrocarbon containing 2 carbon atoms give Sabatier and Senderen's reaction but does not give precipitate with ammoniacal silver nitrate solution. The hydrocarbon in question is:
- a) Ethane      b) Acetylene      c) Ethylene      d) None of these
674. Acetylene can be converted to higher alkyne using the following sequence of reactions:
- a)  $\text{Na}, \text{RX}$       b)  $\text{RMgX}, \text{R}'\text{X}$       c) Either of these two      d) None of these
675. At low temperature, the slow addition of molecular bromine to  $\text{H}_2\text{C} = \text{CH} - \text{CH}_2 - \text{C} \equiv \text{CH}$  gives:
- a)  $\text{CH}_2 = \text{CH} - \text{CH}_2 - \text{CBr} = \text{CHBr}$   
 b)  $\text{BrCH}_2 - \text{CHBr} - \text{CH}_2 - \text{C} \equiv \text{CH}$   
 c)  $\text{H}_2\text{C} = \text{CH} - \text{CH}_2 - \text{CH}_2 - \text{CBr}_3$   
 d)  $\text{CH}_3 - \text{CBr}_2 - \text{CH}_2 - \text{C} \equiv \text{CH}$
676. Which of the following statement is correct?
- a) Benzene has a tetrahedral geometry like an alkane  
 b) Benzene is aromatic while naphthalene is not  
 c) Benzene and Cyclohexane are both aromatic  
 d) Benzene behaves more like and alkane than an alkene
677.  $\text{CaC}_2 + \text{H}_2\text{O} \rightarrow A \xrightarrow{\text{H}_2\text{SO}_4/\text{HgSO}_4} B$   
 Identify A and B in the given reaction
- a)  $\text{C}_2\text{H}_2$  and  $\text{CH}_3\text{CHO}$       b)  $\text{CH}_4$  and  $\text{HCOOH}$   
 c)  $\text{C}_2\text{H}_4$  and  $\text{CH}_3\text{COOH}$       d)  $\text{C}_2\text{H}_2$  and  $\text{CH}_3\text{COOH}$
678. The correct boiling point order for corresponding hydrocarbons is:
- a) Alkyne>alkane>alkene  
 b) Alkane>alkene>alkyne  
 c) Alkyne>alkene>alkane  
 d) Alkene>alkyne>alkane







Identify A and B



680. Electrolysis of cold concentrated aqueous solution of potassium methyl succinate yields:

- a) Ethane                      b) Ethyne                      c) Propene                      d) Ethane-1,2-diol

681. An alkene gives two moles of HCHO, one mole of CO<sub>2</sub> and one mole of CH<sub>3</sub>COCHO on ozonolysis. What is its structure?





682. Alkyl halides get converted to alkenes through:

- a) Electrophilic substitution                      b) Nucleophilic addition                      c) Elimination reaction                      d) Hydrolysis

683. In the complete combustion of C<sub>n</sub>H<sub>2n+2</sub>, the number of oxygen moles required is:

- a)  $\left(\frac{n}{2}\right) O_2$                       b)  $\left(\frac{n+1}{2}\right) O_2$                       c)  $\left(\frac{3n+1}{2}\right) O_2$                       d)  $\left(\frac{n+2}{2}\right) O_2$

684. When CH<sub>3</sub>CH<sub>2</sub>CHCl<sub>2</sub> is treated with NaNH<sub>2</sub> the product formed is:

- a) CH<sub>3</sub>CH = CH<sub>2</sub>                      b) CH<sub>3</sub> - C ≡ CH                      c)                       d) 

685. Cycloalkanes are isomeric with


- a) Alkanes                      b) Alkenes                      c) Alkynes                      d) Arenes

686. Which gives only one monosubstitution product on chlorination?

- a) n-pentane                      b) Neopentane                      c) Isopentane                      d) n-butane

687. The products obtained via oxymercuration (HgSO<sub>4</sub> + H<sub>2</sub>SO<sub>4</sub>) of 1-butyne would be:

- a) CH<sub>3</sub>CH<sub>2</sub>COCH<sub>3</sub>  
b) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CHO  
c) CH<sub>3</sub>CH<sub>2</sub>CHO + HCHO  
d) CH<sub>3</sub>CH<sub>2</sub>COOH + HCOOH



- a) Wolff-Kishner reaction                      b) Clemmensen reduction                      c) Red P+HI at 200°C                      d) Wurtz reaction

689. The presence of the chlorine atom on benzene ring makes the second substituent enter at a position

- a) ortho                      b) meta                      c) para                      d) ortho/para

690. Two organic compounds (A) and (B) both containing only carbon and hydrogen, on quantitative analysis gave the same percentage composition by weight

$$C = \left(\frac{12}{12}\right) \times 100\%, \quad H = \left(\frac{1}{13}\right) \times 100\%$$

*A* decolourises bromine water but *B* does not. *A* and *B* respectively are

- a)  $C_2H_2$  and  $C_6H_6$       b)  $C_6H_6$  and  $C_2H_2$       c)  $C_2H_4$  and  $C_2H_6$       d)  $C_2H_2$  and  $C_2H_6$
691. Which of the following compounds react with, an aqueous solution of  $Ag(NH_2)_2OH$ ?
- a) ethane      b) Ethene      c) 1-butyne      d) 2-butyne
692. Aromatisation of *n*-heptane by passing over  $(Al_2O_3 + Cr_2O_3)$  catalyst at 773 K gives
- a) Benzene      b) Toluene      c) Mixture of both      d) Heptylene
693. In a mixture of *n*-hexadecane and  $\alpha$ -methyl-naphthalene the percentage of the latter is 10. The value of cetane number is:
- a) 110      b) 90      c) 10      d) Zero
694. Addition of bromine to 1,3-butadiene gives:
- a) 1,2-addition product only  
b) 1,4-addition product only  
c) Both 1,2 and 1,4-addition products  
d) No reaction
695.  $R-COOH \rightarrow RCH_2OH$ . This mode of reduction can be effected only by:
- a)  $NaBH_4$       b)  $Na + Alcohol$       c)  $LiAlH_4$       d) All of these
696. A Wittig reaction with an aldehyde gives
- a) Ketone compound      b) A long chain fatty acid  
c) Olefin compound      d) Epoxide
697. Ethylene di bromide on heating with metallic sodium in ether solution yields
- a) Ethene      b) Ethyne      c) 2-butene      d) 1-butene
698. When alcoholic solution of ethylene dibromide is heated with granulated zinc, the compound formed is:
- a) Ethane      b) Ethylene      c) Butane      d) Isobutene
699. Octane number is:
- a) Number of carbon atoms in octane  
b) Number of molecules of octane formed in cracking of 1.0g of gasoline  
c) Number of hydrogen atoms in octane  
d) Number for representing standard rating of fuel
700. When an aqueous solution containing sodium acetate and sodium propionate is electrolysed we get:
- a) Ethane      b) Propane      c) Butane      d) All of these
701. Which one of the following methods is neither meant for the synthesis nor for separation of amines?
- a) Curtius reaction      b) Wurtz reaction      c) Hofmann method      d) Hinsberg method
702. *Vic*-dihalide on treatment with zinc dust gives:
- a) Alkane      b) Alkene      c) Alkyne      d) All of these
703. Identify the substitute group, that acts as *ortho* – *para* director, during electrophilic substitution in aromatic compounds.
- a)  $-NH_2$       b)  $-NO_2$       c)  $-SO_3H$       d)  $N_2$
704. Order of acidity of  $H_2O$ ,  $NH_3$  and acetylene is:
- a)  $NH_3 > CH \equiv CH > H_2O$   
b)  $H_2O > NH_3 > CH \equiv CH$   
c)  $H_2O > CH \equiv CH > NH_3$   
d)  $NH_3 > H_2O > CH \equiv CH$
705.  $C_2H_5I + C_5H_{11}I + 2Na \xrightarrow{\text{Ether}} C_2H_5-C_5H_{11} + 2NaI$   
The above equation represents:
- a) Hofmann's reaction  
b) Dow's reaction



- c) Wurtz synthesis  
 d) Reimer-Tiemann's reaction
706. Identify Z in the sequence,  

$$\text{CH}_3 - \text{CH}_2 - \text{CH} = \text{CH}_2 \xrightarrow{\text{HBr}/\text{H}_2\text{O}_2} \text{Y} \xrightarrow{\text{C}_2\text{H}_5\text{O}^- - \text{Na}^+} \text{Z} :$$
- a) 
$$\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{O} - \text{CH}_2 - \text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$$
- b) 
$$\begin{array}{c} \text{CH}_3 - \text{CH}_2 - \text{CH} - \text{O} - \text{CH}_2 - \text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$$
- c)  $\text{CH}_3 - (\text{CH}_2)_3 - \text{O} - \text{CH}_2 - \text{CH}_3$   
 d)  $\text{CH}_3 - (\text{CH}_2)_4 - \text{O} - \text{CH}_3$
707. Which will give cyclooctyne when treated with base?  
 a) 1,2-dibromocyclobutane  
 b) 1,1-dibromocyclobutane  
 c) 1,1-dibromocyclooctane  
 d) 1,2-dibromocyclopropane
708. The final product in following sequence of reaction is  

$$\text{CH} \equiv \text{CH} \xrightarrow{\text{NaNH}_2} \text{A} \xrightarrow{\text{CH}_3\text{Br}} \text{B}$$
- a)  $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$   
 b)  $\text{HC} \equiv \text{C} - \text{CH}_3$   
 c)  $\text{CH}_2 = \text{CH} - \text{CH}_3$   
 d)  $\text{CH}_3 - \text{CH}_2 - \text{CH}_3$
709. What are the products obtained upon the ozonolysis of pent-2-ene?  
 a)  $\text{CH}_3\text{CH}_2\text{CHO}$   
 b)  $\text{CH}_3\text{CHO}$   
 c)  $\text{CH}_3\text{COCH}_3$   
 d) Both (a) and (b)
710. Addition of halogen acid occurs at slowest rate in:  
 a)  $\text{CH}_2 = \text{CHCl}$   
 b)  $\text{CH}_2 = \text{CH}_2$   
 c)  $\text{CH}_3 - \text{CH} = \text{CH}_2$   
 d)  $(\text{CH}_3)_2\text{C} = \text{CH}_2$
711. Benzyl chloride ( $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ ) can be prepared from toluene by chlorination with  
 a)  $\text{SO}_2\text{Cl}_2$   
 b)  $\text{SOCl}_2$   
 c)  $\text{Cl}_2$   
 d)  $\text{NaOCl}$
712. The Markownikoff's rule is the best applicable to the reaction between  
 a)  $\text{C}_2\text{H}_4 + \text{HCl}$   
 b)  $\text{C}_3\text{H}_6 + \text{Br}_2$   
 c)  $\text{C}_3\text{H}_6 + \text{HBr}$   
 d)  $\text{C}_3\text{H}_8 + \text{Cl}_2$
713. Which of the following acid reacts to reverse the Markownikoff's rule?  
 a)  $\text{HCl}$   
 b)  $\text{HBr}$   
 c)  $\text{HF}$   
 d)  $\text{HI}$
714. The addition of  $\text{HOCl}$  on alkenes in presence of strong acids to form halohydrins proceeds via formation of:  
 a) Chloronium ion  
 b) Carbocation  
 c) Chloro carbocation  
 d) None of these
715. On treatment with chlorine in presence of sunlight, toluene gives the product  
 a) *o*-chloro toluene  
 b) 2,5-dichloro toluene  
 c) *p*-chloro toluene  
 d) Benzyl chloride
716. The most oxidized form of hydrocarbon  $\text{RCH}_3$  is:  
 a)  $\text{CO}_2$   
 b)  $\text{RCHO}$   
 c)  $\text{RCOOH}$   
 d)  $\text{RCOCOOH}$
717. Ethylene is used for:  
 a) Ripening of food  
 b) Preparing ethylene oxide  
 c) For preparing ethylene chloride  
 d) All are correct