

GPLUS EDUCATION

Date :
Time :
Marks :

CHEMISTRY

ELECTROCHEMISTRY

Single Correct Answer Type

- The desired amount of charge for obtaining one mole of Al from Al^{3+} is
a) 96500 C b) 2×96500 C c) 3×96500 C d) $\frac{96500}{2}$ C
- A certain current liberates 0.504 g of hydrogen in 2 hr. How many gram of copper can be liberated by the same current flowing for the same time in CuSO_4 solution?
a) 12.7 b) 16 c) 31.8 d) 63.5
- If the E_{cell}° for a given reaction has a negative value, then which of the following gives the correct relationships for the value of ΔG° and K_{eq} ?
a) $\Delta G^\circ > 0$; $K_{\text{eq}} < 1$ b) $\Delta G^\circ > 0$; $K_{\text{eq}} > 1$ c) $\Delta G^\circ < 0$; $K_{\text{eq}} > 1$ d) $\Delta G^\circ < 0$; $K_{\text{eq}} < 1$
- The Edison storage cell is represented as :
 $\text{Fe}(s) + \text{FeO}(s) | \text{KOH}(aq) | \text{Ni}_2\text{O}_3(s) | \text{Ni}_2\text{O}_3(s) | \text{Ni}(s)$
The half reactions are $\text{Ni}_2\text{O}_3(s) + \text{H}_2\text{O}(l) + 2e^- \rightarrow 2\text{NiO}(s) + 2\text{OH}^-$; $E^\circ = +0.40$ V
 $\text{FeO}(s) + \text{H}_2\text{O}(l) + 2e^- \rightarrow \text{Fe}(s) + 2\text{OH}^-$; $E^\circ = -0.87$ V
Choose the incorrect statement
a) E_{anode} increases with increase in concentration of OH^-
b) E_{cathode} decreases with increase in concentration of OH^-
c) $E_{\text{cell}}^\circ = 1.27$ V
d) E_{cell} increases with increase in concentration of FeO
- Standard reduction potentials of the half reactions are given below :
 $\text{F}_2(g) + 2e^- \rightarrow 2\text{F}^-(aq)$; $E^\circ = +2.85$ V
 $\text{Cl}_2(g) + 2e^- \rightarrow 2\text{Cl}^-(aq)$; $E^\circ = +1.36$ V
 $\text{Br}_2(l) + 2e^- \rightarrow 2\text{Br}^-(aq)$; $E^\circ = +1.06$ V
 $\text{I}_2(s) + 2e^- \rightarrow 2\text{I}^-(aq)$; $E^\circ = +0.53$ V
The strongest oxidising and reducing agents respectively are :
a) F_2 and I^- b) Br_2 and Cl^- c) Cl_2 and Br^- d) Cl_2 and I_2
- The standard reduction potential for $\text{Fe}^{2+} | \text{Fe}$ and $\text{Sn}^{2+} | \text{Sn}$ electrodes are -0.44 V and -0.14 V respectively. For the cell reaction, $\text{Fe}^{2+} + \text{Sn} \rightarrow \text{Fe} + \text{Sn}^{2+}$, the standard e.m.f. is:
a) $+0.30$ V b) 0.58 V c) $+0.58$ V d) -0.30 V
- Electrolytes when dissolved in water dissociates into ions because
a) They are unstable
b) The water dissolves it
c) The force of repulsion increases
d) The force of electrostatic attraction are broken down by water
- Which ion has exceptionally higher Λ^∞ values?
a) H^+ b) K^+ c) NH_2^- d) OH^-
- Limiting molar ionic conductivities of a uni-univalent electrolyte are 57 and 73. The limiting molar conductivity of the solution will be :
a) $130 \text{ S cm}^2 \text{ mol}^{-1}$ b) $65 \text{ S cm}^2 \text{ mol}^{-1}$ c) $260 \text{ S cm}^2 \text{ mol}^{-1}$ d) $187 \text{ S cm}^2 \text{ mol}^{-1}$
- Molten NaCl conducts electricity due to the presence of :
a) Free electrons b) Free molecules c) Free ions d) Atoms of Na and Cl

11. The emf of the cell, ($E_{Zn^{2+}/Zn} = -0.76 V$)
 $Zn / Zn^{2+} (1 M) || Cu^{2+} (1 M) | Cu$
($E_{Cu^{2+}/Cu} = +0.34 V$) will be
a) +1.10 V b) -1.10 V c) +0.42 V d) -0.42 V
12. Which represents a concentration cell?
a) $PtH_2 | HCl (c_1) || HCl (c_2) | PtH_2$ b) $PtH_2 | HCl (c_1) || Cl_2 | Pt$ c) $Zn | Zn^{2+} || Cu^{2+} | Cu$ d) $Fe | Fe^{2+} || Cu^{2+} | Cu$
13. In electrolysis of aqueous copper sulphate, the gas at anode and cathode are
a) O_2 and H_2 b) H_2 and O_2 c) SO_2 and H_2 d) SO_3 and O_2
14. Consider the reaction, $M^{n+}(aq) + ne \rightarrow M^0(s)$. The standard reduction potential values of the metals M_1, M_2 and M_3 are $-0.34 V, -3.05 V$ and $-1.66 V$ respectively. The order of their reducing power will be :
a) $M_1 > M_2 > M_3$ b) $M_3 > M_2 > M_1$ c) $M_1 > M_3 > M_2$ d) $M_2 > M_3 > M_1$
15. The charge required to liberate one gram equivalent of an element is
a) 96500 F b) 1 F c) 1 C d) None of these
16. What will be pH of aqueous solution of electrolyte in electrolytic cell during electrolysis of $CuSO_4(aq)$ between graphite electrodes?
a) pH = 14.0 b) pH > 7.0 c) pH < 7.0 d) pH = 7.0
17. In an electrolytic cell, the anode and cathode are respectively represented as :
a) Positive electrode, negative electrode
b) Negative electrode, positive electrode
c) Both positive and negative electrode
d) None of the above
18. The cell reaction is spontaneous, when
a) E_{red}° is negative b) E_{red}° is positive c) ΔG° is negative d) ΔG° is positive
19. The emf of the cell $Mg | Mg^{2+} (0.01 M) || Sn^{2+} (0.1 M) | Sn$ at 298 K is (Given, $E_{Mg^{2+}, Mg}^\circ = -2.34 V, -2.34 V, E_{Sn^{2+}, Sn}^\circ = -0.14 V$)
a) 2.23 V b) 1.86 V c) 1.56 V d) 3.26 V
20. When an aqueous solution of lithium chloride is electrolysed using graphite electrodes :
a) pH of the resulting solution increases
b) pH of the resulting solution decreases
c) As the current flows, pH of the solution around the cathode increases
d) None of the above
21. In electrolytic purification, which of the following is made of impure metal?
a) Anode b) Cathode c) Both (a) and (b) d) None of these
22. The specific conductivity of 0.1 N KCl solution is $0.0129 \Omega^{-1} cm^{-1}$. The resistance of the solution in the cell 100Ω . The cell constant of the cell will be
a) 1.10 b) 1.29 c) 0.56 d) 2.80
23. Which graph correctly correlates E_{Cell} as a function of concentrations for the cell (for different values of M and M')?
 $Zn(s) + Cu^{2+}(M) \rightarrow Zn^{2+}(M') + Cu(s)$;
 $E_{Cell}^\circ = 1.10 V$
X - axis : $\log_{10} \frac{[Zn^{2+}]}{[Cu^{2+}]}$, Y - axis : E_{Cell}
- a)

b)

c)

d)
24. In acidic medium MnO_4^- is converted to Mn^{2+} . The quantity of electricity in faraday required to reduce 0.5

- mole of MnO_4^- to Mn^{2+} would be
 a) 2.5 b) 5 c) 1 d) 0.5
25. In electrolysis, oxidation takes place at:
 a) Anode
 b) Cathode
 c) The anode as well as cathode
 d) The surface of electrolyte solution
26. A depolariser used in dry cell batteries is :
 a) Ammonium chloride b) Manganese dioxide c) Potassium hydroxide d) Sodium phosphate
27. The $E^\circ_{M^{3+}/M^{2+}}$ values for Cr, Mn, Fe and Co are $-0.41, +1.57, +0.77$ and $+1.97$ V respectively. For which one of these metals, the change in oxidation state from +2 to +3 is easiest?
 a) Fe b) Mn c) Co d) Cr
28. The standard reduction electrode potential values of the elements A, B and C are $+0.68, -2.50$ and -0.50 V respectively. The order of their reducing power is :
 a) $A > B > C$ b) $A > C > B$ c) $C > B > A$ d) $B > C > A$
29. The number of electrons involved in the reaction when a faraday of electricity is passed through an electrolyte in solution is :
 a) 12×10^{46} b) 96500 c) 8×10^{16} d) 6.02×10^{23}
30. The electrolysis of a solution resulted in the formation of H_2 at the cathode and Cl_2 at the anode. The liquid is:
 a) Pure water
 b) H_2SO_4 solution
 c) NaCl solution in water
 d) CuCl_2 solution in water
31. The passage of electricity in the Daniell cell when Zn and Cu electrodes are connected:
 a) From Cu to Zn inside the cell
 b) From Cu to Zn outside the cell
 c) From Zn to Cu outside the cell
 d) None of the above
32. $\text{Ni} / \text{Ni}^{2+} [1.0 \text{ M}] || \text{Au}^{3+} [1.0 \text{ M}] / \text{Au}$ where E° for $\text{Ni}^{2+} / \text{Ni}$ is -0.250 V; and E° for $\text{Au}^{3+} / \text{Au}$ is 0.150 V. The emf of the cell is
 a) $+1.25$ V b) -1.75 V c) $+1.75$ V d) $+0.4$ V
33. The product obtained at anode when 50% H_2SO_4 aqueous solution is electrolysed using platinum electrodes is
 a) H_2SO_3 b) $\text{H}_2\text{S}_2\text{O}_8$ c) O_2 d) H_2
34. The approximate e.m.f. of a dry cell is :
 a) 2.0 V b) 1.2 V c) 6 V d) 1.5 V
35. $E_1, E_2,$ and E_3 are the emfs of the following three galvanic cells respectively
 I. $\text{Zn} (\text{s}) | \text{Zn}^{2+} (0.1 \text{ M}) || \text{Cu}^{2+} (1 \text{ M}) | \text{Cu} (\text{s})$
 II. $\text{Zn} (\text{s}) | \text{Zn}^{2+} (1 \text{ M}) || \text{Cu}^{2+} (1 \text{ M}) | \text{Cu} (\text{s})$
 III. $\text{Zn} (\text{s}) | \text{Zn}^{2+} (1 \text{ M}) || \text{Cu}^{2+} (0.1 \text{ M}) | \text{Cu} (\text{s})$
 Which one of the following is true?
 a) $E_2 > E_1 > E_3$ b) $E_1 > E_2 > E_3$ c) $E_3 > E_1 > E_2$ d) $E_3 > E_2 > E_1$
36. The fraction of the total current carried by an ion is known as:
 a) Transport number of that ion
 b) Conductance of that ion
 c) Both(a) and (b)
 d) None of the above
37. In a galvanic cell, which is wrong?

- a) Anode has negative polarity
 b) Cathode has positive polarity
 c) Reduction takes place at anode
 d) Reduction takes place at cathode
38. The rusting of iron takes place as follows

$$2\text{H}^+ + 2\text{e}^- + \frac{1}{2}\text{O}_2 \rightarrow \text{H}_2\text{O} (l);$$

$$E^\circ = +1.23\text{ V}$$

$$\text{Fe}^{2+} + 2\text{e}^- \rightarrow \text{Fe} (s); \quad E^\circ = -0.44\text{ V}$$
 Calculate ΔG° for the net process.
 a) -322 kJ mol^{-1} b) -161 kJ mol^{-1} c) -152 kJ mol^{-1} d) -76 kJ mol^{-1}
39. What weight of copper will be deposited by passing 2 faraday of electricity through a solution of Cu(II) salt?
 a) 35.6 g b) 63.5 g c) 6.35 g d) 3.56 g
40. Chlorine cannot displace :
 a) Fluorine from NaF b) Iodine from NaI c) Bromine from NaBr d) None of these
41. For Acell reaction involving Atwo-electron change, the standard emf of the cell is found to be 0.295 V at 25°C. The equilibrium constant of the reaction at 25 °C will be
 a) 1×10^{-10} b) 29.5×10^{-2} c) 10 d) 1×10^{10}
42. The resistance of a decinormal solution of a salt occupying a volume between two platinum electrodes 1.80 cm apart and 5.4 cm² in area was formed to be 32 ohm. The specific and equivalent conductivity respectively in their proper units are :
 a) 104.1 and 0.0104 b) 208.2 and 0.0208 c) 0.0104 and 104.0 d) None of these
43. The value of equilibrium constant for a feasible cell reaction is :
 a) < 1 b) Zero c) = 1 d) > 1
44. At 25°C, the standard e.m.f. of cell having reactions involving a two electron change is found to be 0.295 V. The equilibrium constant of the reaction is :
 a) 29.5×10^{-2} b) 10 c) 10^{10} d) 29.5×10^{10}
45. E° for $\text{Fe}^{2+} + 2\text{e}^- \rightarrow \text{Fe}$ is -0.44 V and E° for $\text{Zn}^{2+} + 2\text{e}^- \rightarrow \text{Zn}$ is -0.76 V thus
 a) Zn is more electropositive than Fe b) Zn is more electronegative than Fe
 c) Fe is more electropositive than Zn d) None of the above
46. A certain quantity of electricity is passed through aqueous solution of AgNO_3 and CuSO_4 connected in series. If Ag (at.wt.108) deposited at the cathode is 1.08 g then Cu deposited at the cathode is (at. wt. of Cu is 63.53):
 a) 6.354 g b) 0.317 g c) 0.6354 g d) 3.177 g
47. $\text{I}_2(s)|\text{I}^- (0.1\text{ M})$ half-cell is connected to a $\text{H}^+(aq)|\text{H}_2(1\text{ bar})|\text{Pt}$ half-cell and emf is found to be 0.7714 V. If $E^\circ_{\text{I}_2/\text{I}^-} = 0.535\text{ V}$, find the pH of H^+/H_2 half-cell
 a) 1 b) 2 c) 3 d) 5
48. The $E^\circ_{M^{3+}/M^{2+}}$ values for Cr, Mn, Fe and Co are -0.41 V , $+1.57\text{ V}$, $+0.77\text{ V}$ and $+1.97\text{ V}$ respectively. For which one of these metals the change in oxidation state from +2 to +3 is easiest?
 a) Cr b) Mn c) Fe d) Co
49. In which cell, liquid junction potential need to be eliminated?
 a) $\text{Pt}/\text{H}_2(\text{P}_1)|\text{HCl}|\text{Pt}/\text{H}_2(\text{P}_2)$
 b) $\text{Pt}/\text{H}_2|_{c_1}\text{HCl}|\text{HCl}|_{c_2}\text{Pt}/\text{H}_2$
 c) Nicad cell
 d) Lead storage battery
50. Which one of the following nitrates will leave behind Ametal on strong heating?
 a) Ferric nitrate b) Copper nitrate c) Manganese nitrate d) Silver nitrate

51. $E_{Cu}^{\circ} = 0.34 V$, $E_{Zn}^{\circ} = 0.76 V$. A Daniel cell contains 0.1 M $ZnSO_4$ solution and 0.01 M $CuSO_4$ solution at its electrodes. EMF of the cell is
 a) 1.10 V b) 1.04 V c) 1.16 V d) 1.07 V
52. The E° of Fe^{2+} / Fe and Sn^{2+} / Sn are -0.44 V and -0.14 V respectively. If cell reaction is $Fe + Sn^{2+} \rightarrow Fe^{2+} + Sn$ then emf of the cell is
 a) +0.30 V b) -0.58 V c) +0.58 V d) -0.30 V
53. Electrolysis rules of Faraday's states that mass deposited on electrode is proportional to
 a) Q b) Q^2 c) I^2 d) None of these
54. A silver cup is plated with silver by passing 965 C of electricity. The amount of Ag deposited is
 a) 107.89 g b) 9.89 g c) 1.0002 g d) 1.08 g
55. The molecular conductivity and equivalent conductivity are same for the solution of :
 a) 1 M NaCl b) 1 M $Ba(NO_3)_2$ c) 1 M $La(NO_3)_3$ d) 1 M $Th(NO_3)_4$
56. Dipping iron article into a strongly alkaline solution of sodium phosphate
 a) Does not affect the article b) Forms $Fe_2O_3 \cdot xH_2O$ on the surface
 c) Forms iron phosphate film d) Forms ferric hydroxide
57. When an electric current is passed through an aqueous solution of sodium chloride :
 a) H_2 is evolved at the anode
 b) Oxygen is evolved at the cathode
 c) Its pH progressively decreases
 d) Its pH progressively increases
58. The cell reaction of the galvanic cell $Cu(s) | Cu^{2+}(aq) || Hg^{2+}(aq) | Hg(l)$ is
 a) $Hg + Cu^{2+} \rightarrow Hg^{2+} + Cu$ b) $Hg + Cu^{2+} \rightarrow Cu + Hg^{2+}$
 c) $Cu + Hg \rightarrow CuHg$ d) $Cu + Hg^{2+} \rightarrow Cu^{2+} + Hg$
59. Calculate the volume of hydrogen at NTP obtained by passing a current of 0.4 ampere through acidified water for 30 minute :
 a) 0.0836 litre b) 0.1672 litre c) 0.0432 litre d) 0.836 litre
60. The standard emf of a cell involving one electron change is found to be 0.591 V and 25 °C. The equilibrium constant of the reaction is ($F = 96500 C mol^{-1}$)
 a) 1.0×10^1 b) 1.0×10^5 c) 1.0×10^{10} d) 1.0×10^{30}
61. The relationship between Gibbs' free energy change (ΔG) and emf (E) of a reversible electrochemical cell is given by
 a) $\Delta G = nFE$ b) $\Delta G = nF/E$ c) $\Delta G = -nFE$ d) $\Delta G = E/nF$
62. The reduction electrode potential, E of 0.1 M solution of M^+ ions ($E_{RP} = -2.36 V$) is
 a) -4.82 V b) -2.41 V c) +2.41 V d) None of these
63. Passage of 1 faraday of electricity through a solution of $CuSO_4$, deposits :
 a) 1 mole of Cu b) 1 g-atom of Cu c) 1 molecule of Cu d) 1 g equivalent of Cu
64. The conductivity of $N/50$ solution of KCl in a cell at 25°C is 0.002765 mho cm^{-1} . If the resistance of a cell containing this solution is 400 ohm, the cell constant is :
 a) 1.106 cm b) 1.106 cm^{-1} c) 1 cm d) 1 cm^{-1}
65. The equilibrium constant for the reaction given below at 298 K is :
 $Zn(s) + Fe^{2+}(aq) \rightarrow Zn^{2+}(aq) + Fe(s)$;
 $E_{cell}^{\circ} = 2905 V$ at 298 K
 a) $e^{0.32/0.0295}$ b) $10^{0.595/0.76}$ c) $10^{0.0250/0.32}$ d) $10^{0.32/0.295}$
66. When the sample of copper with zinc impurity is to be purified by electrolysis, the appropriate electrodes are

Cathode	Anode
a) Pure zinc	pure copper

- b) Impure sample pure copper
 c) Impure zinc impure sample
 d) Pure copper impure sample
67. A current of 12 A is passed through an electrolytic cell containing aqueous NiSO_4 solution. Both Ni and H_2 gas are formed at the cathode. The current efficiency is 60%. What is the mass of nickel deposited on the cathode per hour?
 a) 7.883 g b) 3.941 g c) 5.91 g d) 2.645 g
68. 10^{-2} g atom of Ag can be oxidised to Ag^+ during the electrolysis of AgNO_3 solution using silver electrode by :
 a) 965 coulomb b) 96500 coulomb c) 9650 coulomb d) 96.500 coulomb
69. A gas X at 1 atm is bubbled through a solution containing a mixture of $1 \text{ M } y^-$ and $1 \text{ M } z^-$ at 25°C . If the order of reduction potential is $z > y > x$ then
 a) y will oxidize x and not z b) y will oxidize x and z
 c) y will oxidize z and not x d) y will reduce both x and z
70. Which one of the following solutions will have highest conductivity?
 a) 0.1 M CH_3COOH b) 0.1 M NaCl c) 0.1 M KNO_3 d) 0.1 M HCl
71. A current of strength 2.5 A was passed through CuSO_4 solution for 6 min 26 s. The amount of copper deposited is (At. Wt. of Cu = 63.5, $1F = 96500 \text{ C}$)
 a) 0.3175 g b) 3.175 g c) 0.635 g d) 6.35 g
72. A student made the following observations in the laboratory,
 i) Clean copper metal did not react with 1 molar $\text{Pb}(\text{NO}_3)_2$ solution.
 ii) Clean lead metal dissolved in a 1 molar AgNO_3 solution and crystals of Ag metal appeared.
 iii) Clean silver metal did not react with 1 molar $\text{Cu}(\text{NO}_3)_2$ solution.
 The order of decreasing reducing character of the three metals is :
 a) Cu, Pb, Ag b) Cu, Ag, Pb c) Pb, Cu, Ag d) Pb, Ag, Cu
73. The e. m. f. of the cell $\text{Zn} | \text{Zn}^{2+} (1 \text{ M}) || \text{Cu}^{2+} | \text{Cu} (1 \text{ M})$ is 1.1 volt. If the standard reduction potential of $\text{Zn}^{2+} | \text{Zn}$ is -0.78 volt, what is the oxidation potential of $\text{Cu} | \text{Cu}^{2+}$?
 a) $+1.86 \text{ V}$ b) 0.32 V c) -0.32 V d) -1.86 V
74. Standard reduction electrode potentials of three metals A, B and C are respectively $+0.5 \text{ V}$, -3.0 V and -1.2 V . The reducing powers of these metals are
 a) $A > B > C$ b) $C > B > A$ c) $A > C > B$ d) $B > C > A$
75. Quantity of charge is measured in :
 a) ampere-sec. b) ampere c) ampere sec^{-1} . d) ampere $^{-1}$ sec.
76. Which of the following will form a cell with the highest voltage?
 a) 0.1 M Ag^+ , 2 M Co^{2+} b) 2 M Ag^+ , 2 M Co^{2+} c) 1 M Ag^+ , 1 M Co^{2+} d) 2 M Ag^+ , 0.1 M Co^{2+}
77. When electric current is passed through acidified water for 1930 s, 1120 mL of H_2 gas is collected (at STP) at the cathode. What is the current passed in amperes?
 a) 0.05 b) 0.50 c) 5.0 d) 50
78. In which of the following pairs, the constants/ quantities are not mathematically related to each other?
 a) Gibbs free energy and standard cell potential
 b) Equilibrium constant and standard cell potential
 c) Rate constant and activation energy
 d) Rate constant and standard cell potential
79. The charge required for reduction of 1 mole of $\text{Cr}_2\text{O}_7^{2-}$ ions to Cr^{3+} is
 a) 96500 C b) $2 \times 96500 \text{ C}$ c) $3 \times 96500 \text{ C}$ d) $6 \times 96500 \text{ C}$
80. Cell constant has the unit:
 a) cm b) cm^{-1} c) cm^2 d) cm sec^{-1}
81. The resistance of 0.01 N solution of an electrolyte was found to be 210 ohm at 298 K, using a conductivity cell of cell constant 0.66 cm^{-1} . The equivalent conductivity of solution is :
 a) $314.28 \text{ mho cm}^2\text{eq.}^{-1}$ b) $3.14 \text{ mho cm}^2\text{eq.}^{-1}$ c) $314.28 \text{ mho}^{-1} \text{ cm}^2\text{eq.}^{-1}$ d) $3.14 \text{ mho}^{-1} \text{ cm}^2\text{eq.}^{-1}$


82. IV. $\text{Cu} + 2\text{HCl} \rightarrow \text{CuCl}_2 + \text{H}_2(\text{g})$
 $[E_{\text{Cu}^{2+}/\text{Cu}}^\circ = +0.34 \text{ V}]$
 V. $\text{Zn} + 2\text{HCl} \rightarrow \text{ZnCl}_2 + \text{H}_2(\text{g})$
 $[E_{\text{Zn}^{2+}/\text{Zn}}^\circ = -0.76 \text{ V}]$
 VI. $\text{Ag} + 2\text{HCl} \rightarrow \text{AgCl} + \frac{1}{2}\text{H}_2(\text{g})$
 $[E_{\text{Ag}^+/\text{Ag}}^\circ = +0.80 \text{ V}]$

Which of the following reaction is feasible ?

- a) (ii) b) (i) c) (iii) d) All of these
83. The standard potential at 25°C for the following half-reactions are given against them $\text{Zn}^{2+} + 2e^- \rightarrow \text{Zn}$, $E^\circ = -0.762 \text{ V}$
 $\text{Mg}^{2+} + 2e^- \rightarrow \text{Mg}$, $E^\circ = -2.37 \text{ V}$
 When zinc dust is added to the solution of MgCl_2 ,
 a) ZnCl_2 is formed b) Mg is precipitated
 c) Zn dissolves in the solution d) No reaction takes place
84. 1 coulomb of charge passes through solution of AgNO_3 and CuSO_4 connected in series and the concentration of two solution being in the ratio 1 : 2. The ratio of amount of Ag and Cu deposited on Pt electrode is :
 a) 107.9 : 63.54 b) 54 : 31.77 c) 107.9 : 31.77 d) 54 : 63.54
85. When lead accumulator is charged, it is :
 a) An electrolytic cell b) A galvanic cell c) A Daniell cell d) None of these
86. If the ΔG of a cell reaction $\text{AgCl} + e^- \rightarrow \text{Ag} + \text{Cl}^-$ is -21.20 kJ , the standard emf of cell is
 a) 0.239 V b) 0.220 V c) -0.320 V d) -0.110 V
87. $\Lambda_{\text{ClCH}_2\text{COONa}}^\infty = 224 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
 $\Lambda_{\text{NaCl}}^\infty = 38.5 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
 $\Lambda_{\text{HCl}}^\infty = 203 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
 What is the value of $\lambda_{\text{ClCH}_2\text{COOH}} = ?$
 a) $288.5 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$ b) $289.5 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
 c) $388.5 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$ d) $59.5 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
88. When a copper wire is immersed in a solution of AgNO_3 , the colour of the solution becomes blue because copper :
 a) Forms a soluble complex with AgNO_3
 b) Is oxidised to Cu^{2+}
 c) Is reduced to Cu^{2-}
 d) Splits up into atomic form and dissolves
89. The electrode potential of a glass electrode depends upon :
 a) Concentration of chloride ions
 b) Concentration of hydrogen ions
 c) Concentration of KCl solution
 d) None of the above
90. 0.04 N solution of a weak acid has conductivity $4.23 \times 10^{-4} \text{ mho cm}^{-1}$. If the degree of dissociation of acid at this dilution is 0.0612, then equivalent conductivity at infinite dilution ismho $\text{cm}^2 \text{ eq.}^{-1}$:
 a) 172.8 b) 180 c) 190 d) 160
91. The highest electrical conductivity of the following aqueous solutions is of
 a) 0.1 M difluoroacetic acid b) 0.1 M fluoroacetic acid
 c) 0.1 M chloroacetic acid d) 0.1 M acetic acid
92. Which one is correct?
 a) Ni displaces zinc from its solution
 b) Zn displaces iron from its solution


- c) Ag displaces copper from its solution
 d) Cu displaces nickel from its solution
93. In an electrolytic cell of $\text{Ag} | \text{AgNO}_3 | \text{Ag}$, when current is passed, the concentration of AgNO_3 :
 a) Increases b) Decreases c) Remains same d) None of these
94. The resistance of 1N solution of acetic is 250Ω , when measured in a cell having a cell constant of 1.15cm^{-1} . The equivalent conduction (in $\text{ohm}^{-1} \text{cm}^2 \text{equiv}^{-1}$) of 1N acetic acid is
 a) 2.3 b) 4.6 c) 9.2 d) 18.4
95. The standard reduction potential E° for the half reactions are as
 $\text{Zn} \rightarrow \text{Zn}^{2+} + 2e^-$, $E^\circ = 0.76 \text{ V}$
 $\text{Cu} \rightarrow \text{Cu}^{2+} + 2e^-$, $E^\circ = 0.34 \text{ V}$
 The emf for the cell reaction,
 $\text{Zn} + \text{Cu}^{2+} \rightarrow \text{Zn}^{2+} + \text{Cu}$
 a) 0.42 V b) -0.42 V c) -1.1 V d) 1.1 V
96. Ionic mobility is equal to:
 a) Speed of ions
 b) Speed of ion under potential difference of 1 volt
 c) Speed of ions under unit potential gradient
 d) None of the above

97. At $\text{pH} = 2$, $E^\circ_{\text{Quinhydrone}} = 1.30 \text{ V}$, $E_{\text{Quinhydrone}}$ will be :



- a) 1.36 V b) 1.30 V c) 1.42 V d) 1.20 V
98. The equilibrium constant for the reaction : $\text{Cu} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{Ag}$; $E^\circ = 0.46 \text{ V}$ at 298 K is :
 a) 2.0×10^{10} b) 4.0×10^{10} c) 4.0×10^{15} d) 2.4×10^{10}
99. For a given cell reaction; $\text{Cr} + 3\text{H}_2\text{O} + \text{OCl}^- \rightarrow \text{Cr}^{3+} + 3\text{Cl}^- + 6\text{OH}^-$, the species undergoing reduction is :
 a) Cr b) Cr^{6+} c) OCl^- d) Cl^-
100. If the H^+ concentration is decreased from 1 M to 10^{-4} M at 25°C for the couple $\text{MnO}_4^- / \text{Mn}^{2+}$, then the oxidising power of the $\text{MnO}_4^- / \text{Mn}^{2+}$ couple decreases by
 a) -0.18 V b) 0.18 V c) 0.38 V d) -0.38 V
101. The standard e.m.f. of a galvanic cell involving the cell reaction with $n = 2$ is found to be 0.295 V at 25°C . The equilibrium constant of the reaction is :
 a) 2.0×10^{11} b) 4.0×10^{12} c) 1.0×10^2 d) 1.0×10^{10}
102. If an iron rod is dipped in CuSO_4 solution, then :
 a) Blue colour of the solution turns red
 b) Brown layer is deposited on iron rod
 c) No change occurs in the colour of the solution
 d) None of the above
103. Which of the following liberates hydrogen on reaction with dilute H_2SO_4 ?
 a) Al b) Fe c) Cu d) Hg
104. A galvanic cell with electrode potential of 'A' = +2.23 V and 'B' = -1.43 V. The value of E°_{cell} is
 a) 3.66 V b) 0.80 V c) -0.80 V d) -3.66 V
105. A galvanic cell is composed of two hydrogen electrodes, one of which is a standard one. In which of the following solutions should the other electrode be immersed to get maximum e. m. f.?
 a) 0.1 M HCl b) 0.1 M CH_3COOH c) 0.1 M H_3PO_4 d) 0.1 M H_2SO_4
106. Which metal does not give the following reaction?
 $M + \text{water or steam} \rightarrow \text{oxide} + \text{H}_2 \uparrow$

- a) Iron b) Sodium c) Mercury d) Magnesium
107. 4.5 g of Al (at. mass 27 amu) is deposited at cathode from Al^{3+} solution by a certain quantity of charge. The volume of H_2 produced at STP from H^+ ions in solution by the same quantity of charge will be :
- a) 11.2 L b) 44.8 L c) 5.6 L d) 22.4 L
108. In the electrolysis of acidulated water, it is desired to obtain 1.12 cc of hydrogen per second under STP condition. The current to be passed is
- a) 1.93 A b) 9.65 A c) 19.3 A d) 0.965 A
109. The speed of ions during passage of current depends upon :
- a) Nature of ion b) Potential gradient c) Dilution of solution d) All of these
110. The best way to prevent rusting of iron is
- a) Making it cathode b) Putting in saline water
c) Both (a) and (b) d) None of these
111. The hydrogen electrode is dipped in a solution of $\text{pH} = 3$ at 25°C . The reduction potential of the cell would be :
- a) 0.177 V b) -0.177 V c) 0.087 V d) 0.059 V
112. Conductivity (unit Siemen) is directly proportional to area of the vessel and the concentration of the solution in it and is inversely proportional to the length of the vessel. Then, the units of the constant of proportionality is
- a) $\text{S}^2 \text{m}^2 \text{mol}$ b) $\text{S}^2 \text{m}^2 \text{mol}^{-2}$ c) $\text{S m}^2 \text{mol}^{-1}$ d) S m mol^{-1}
113. The metal that cannot be produced on reduction of its oxide by aluminium is :
- a) K b) Mn c) Cr d) Fe
114. In the concentration cells, the electrical energy is produced due to :
- a) Oxidation of fuel
b) Heat energy
c) Chemical reaction
d) Transfer of a substance from one concentration to other
115. How many faraday are needed to reduce a mole of MnO_4^- of Mn^{2+} ?
- a) 4 b) 5 c) 3 d) 2
116. For the cell,
 $\text{Tl} \mid \text{Tl}^+ (0.001 \text{ M}) \parallel \text{Cu}^{2+} (0.1 \text{ M}) \mid \text{Cu}$
 E_{cell} at 25°C is 0.83 V. E_{cell} can be increased
- a) By decreasing $[\text{Cu}^{2+}]$ b) By increasing $[\text{Cu}^{2+}]$
c) By increasing $[\text{Tl}^+]$ d) None of these
117. In an aqueous solution, hydrogen (H_2) will not reduce :
- a) Fe^{3+} b) Cu^{2+} c) Zn^{2+} d) Ag^+
118. How many faradays of electricity are required to electrolyse 1 mole CuCl_2 to copper metal and chlorine gas?
- a) 1 F b) 2 F c) 3 F d) 4 F
119. Which statement is not correct?
- a) Conductance of an electrolytic solution increases with dilution
b) Conductance of an electrolytic solution decreases with dilution
c) Specific conductance of an electrolytic solution decreases with dilution
d) Equivalent conductance of an electrolytic solution increase with dilution
120. The correct value of e.m.f. of cell is given by :
- i) $E_{\text{cell}} = E_{\text{OP anode}} - E_{\text{RP cathode}}$
ii) $E_{\text{cell}} = E_{\text{OP anode}} + E_{\text{RP cathode}}$
iii) $E_{\text{cell}} = E_{\text{RP anode}} + E_{\text{RP cathode}}$
iv) $E_{\text{cell}} = E_{\text{OP anode}} - E_{\text{OP cathode}}$
- a) (iii) and (i) b) (i) and (ii) c) (iii) and (iv) d) (ii) and (iv)
121. $\text{Zn}^{2+} \rightarrow \text{Zn (s)}; E^\circ = -0.76 \text{ V}$



Which of the following is spontaneous?

- a) $\text{Zn}^{2+} + \text{Cu} \rightarrow \text{Zn} + \text{Cu}^{2+}$
 b) $\text{Cu}^{2+} + \text{Zn} \rightarrow \text{Cu} + \text{Zn}^{2+}$
 c) $\text{Zn}^{2+} + \text{Cu}^{2+} \rightarrow \text{Zn} + \text{Cu}$
 d) None of the above

122. Reduction potentials of A, B, C, and D are 0.8 V, 0.79 V, 0.34 V and -2.37 V respectively. Which element displaces all the other three elements?

- a) B b) A c) D d) C

123. Given,

$$E^\circ_{\text{Cr}^{3+}/\text{Cr}} = 0.72 \text{ V}, E^\circ_{\text{Fe}^{2+}/\text{Fe}} = 0.42 \text{ V.}$$

The potential for the cell



- a) 0.26 V b) 0.399 V c) -0.339 V d) -0.26 V

124. The electroplating with chromium is undertaken because :

- a) Electrolysis of chromium is easier
 b) Chromium can form alloys with other metals
 c) Chromium gives a protective and decorative coating to the base metal
 d) Of high reactivity of chromium metal



125. Which of the following is not correct?

- a) Aqueous solution of NaCl is an electrolyte.
 b) The units of electrochemical equivalent are g-coulomb.
 c) In the Nernst equation, n represents the number of electrons transferred in the electrode reaction.
 d) Standard reduction potential of hydrogen electrode is zero volt.

126. H_2 cannot be displaced by

- a) Li^+ b) Sr^{2+} c) Al^{3+} d) Ag^+

127. The standard reduction potential of Zn and Ag in water at 298 K are,



- a) $\text{Zn}^{2+} (\text{aq}) + 2\text{Ag} (\text{s}) \rightarrow 2\text{Ag}^+ (\text{aq}) + \text{Zn} (\text{s})$ b) $\text{Zn} (\text{s}) + 2\text{Ag}^+ (\text{aq}) \rightarrow \text{Zn}^{2+} (\text{aq}) + 2\text{Ag} (\text{s})$
 c) $\text{Zn}^{2+} (\text{aq}) + \text{Ag}^+ (\text{aq}) \rightarrow \text{Zn} (\text{s}) + \text{Ag} (\text{s})$ d) $\text{Zn} (\text{s}) + \text{Ag} (\text{s}) \rightarrow \text{Zn}^{2+} (\text{aq}) + \text{Ag}^+ (\text{aq})$

128. The amount of an ion discharged during electrolysis is not dependent of :

- a) Resistance of solution
 b) Time
 c) Current strength
 d) Electrochemical equivalent of the element

129. The conductivity of a 0.1 N KCl solution at 23°C is $0.012 \text{ ohm}^{-1} \text{ cm}^{-1}$. The resistance of the cell containing the solution at the same temperature was found to be 55 ohm. The cell constant will be :

- a) 0.918 cm^{-1} b) 0.66 cm^{-1} c) 1.142 cm^{-1} d) 1.12 cm^{-1}

130. Reduction potential of four elements P, Q, R, S is -2.90, +0.34, +1.20 and -0.76. Reactivity decreases in the order

- a) $P > Q > R > S$ b) $S > R > Q > P$ c) $P > S > Q > R$ d) $Q > S > R > P$

131. Which of the following statements are correct concerning redox properties?

I Ametal M for which E° for the half reaction



II The oxidizing power of the halogens decreases from chlorine to iodine.

III The reducing power of hydrogen halides increases from hydrogen chloride to hydrogen iodide.

- a) I, II and III b) I and II c) I only d) II and III only

132. A cell with two electrodes, one of grey tin and the other white tin, both dipping in solution of $(\text{NH}_4)_2\text{SnCl}_6$ showed zero e.m.f. at 18°C. What conclusion may be drawn from this?

- a) The e.m.f. developed at the electrode-solution phase boundary cancels the normal e.m.f.

- b) Grey tin being non-metallic ceases to provide a reversible electrode reaction
 c) Electrode surface develops a protective layer and the cell develops a very large internal resistance
 d) The standard Gibbs energy change of the cell becomes zero
133. Aluminium displaces hydrogen from dilute HCl whereas silver does not. The emf of Acell prepared by combining Al/ Al³⁺ and Ag / Ag⁺ is 2.46 V. The reduction potential of silver electrode is +0.80 V. The reduction potential of aluminium electrode is
 a) +1.66 V b) -3.26 V c) 3.26 V d) -1.66 V
134. For $I_2 + 2e^- \rightarrow 2I^-$, standard reduction potential = + 0.54 volt. For $2Br^- \rightarrow Br_2 + 2e^-$, standard oxidation potential = - 1.09 volt. For $Fe \rightarrow Fe^{2+} + 2e^-$, standard oxidation potential = + 0.44 volt. Which of the following reactions is non-spontaneous?
 a) $Br_2 + 2I^- \rightarrow 2Br^- + I_2$
 b) $Fe + Br_2 \rightarrow Fe^{2+} + 2Br^-$
 c) $Fe + I_2 \rightarrow Fe^{2+} + 2I^-$
 d) $I_2 + 2Br^- \rightarrow 2I^- + Br_2$
135. When KMnO₄ acts as an oxidizing agent and ultimately forms MnO₄²⁻, MnO₂, Mn₂O₃ and Mn²⁺ then the number of electrons transferred in each case respectively, are
 a) 4, 3, 1, 5 b) 1, 5, 3, 7 c) 1, 3, 4, 5 d) 3, 5, 7, 1
136. For a cell reaction involving a two electron change, the standard emf of the cell is found to be 0.295 V at 25°C. The equilibrium constant of the reaction, at 25°C, will be
 a) 10 b) 1×10^{10} c) 1×10^{-10} d) 10×10^{-2}
137. Which one of the following has the highest molar conductivity?
 a) Diaminedichloroplatinum (III) b) Tetraaminedichlorocobalt (III) chloride
 c) Potassium hexacyanoferrate (II) d) Hexaaquochromium (III) bromide
138. Electrode potential of Zn²⁺/Zn is - 0.76 V and that of Cu²⁺/Cu is + 0.34 V. The emf of the cell constructed between these two electrodes is
 a) 1.10 V b) - 1.10 V c) 2.20 V d) - 2.20 V
139. The standard reduction potentials at 298 K for the following half-cell reactions are given
 $Zn^{2+} (aq) + 2e^- \rightleftharpoons Zn (s); -0.762 V$
 $Cr^{3+} (aq) + 3e^- \rightleftharpoons Cr (s); -0.74 V$
 $2H^+ (aq) + 2e^- \rightleftharpoons H_2(g); +0.00 V$
 $Fe^{3+} (aq) + e^- \rightleftharpoons Fe^{2+} (aq); +0.77 V$
 Which one of the following is the strongest reducing agent?
 a) Zn (s) b) Cr(s) c) H₂ (s) d) Fe²⁺ (aq)
140. How long (in hours) must a current of 5.0 A be maintained to electroplate 60 g of calcium from molten CaCl₂?
 a) 27 h b) 8.3 h c) 11 h d) 16 h
141. Use of electrolysis is
 a) Electrorefining b) Electroplating c) Both (a) and (b) d) None of these
142. What is the cell reaction occurring in Daniel cell (Galvanic cell)?
 a) $Cu(s) + ZnSO_4(aq) \rightarrow CuSO_4(aq) + Zn(s)$
 b) $Zn(s) + CuSO_4(aq) \rightarrow Cu(s) + ZnSO_4(aq)$
 c) $Ni(s) + ZnSO_4(aq) \rightarrow NiSO_4(aq) + Zn(s)$
 d) $2Na(s) + CdSO_4(aq) \rightarrow Na_2SO_4(aq) + Cd (s)$

143.

Electrolyte	KCl	KNO ₃	HCl	NaOAc	NaCl
$\Lambda^\infty (S cm^2 mol^{-1})$	149.9	145.0	426.2	91.0	126.5

Calculate Λ_{HOAc}^∞ using appropriate molar conductances of the electrolytes listed above at infinite dilution in H₂O at 25°C.

- a) 217.5 b) 390.7 c) 552.7 d) 517.2
144. Is the reaction, $2\text{Al} + 3\text{Fe}^{2+} \rightarrow 2\text{Al}^{3+} + 3\text{Fe}$ possible?
 a) No, because standard oxidation potential of Al < Fe
 b) Yes, because standard oxidation potential of Al > Fe
 c) Neither (a) nor (b)
 d) Data are unpredictable
145. What will be the electrode potential of that hydrogen electrode is filled with HCl solution of pH value 1.0?
 a) -59.15 V b) +59.15 c) +59.15 mV d) -59.15 mV
146. The conductivity of a 0.01 N solution is found to be $0.005 \text{ ohm}^{-1} \text{ cm}^{-1}$. The equivalent conductivity of the solution will be
 a) $5 \times 10^{-2} \text{ ohm}^{-1} \text{ cm}^2 \text{ equiv}^{-1}$ b) $5.00 \times 10^{-3} \text{ ohm}^{-1} \text{ cm}^2$
 c) $500 \text{ ohm}^{-1} \text{ cm}^{-2} \text{ equiv}^{-1}$ d) $0.5 \text{ ohm}^{-1} \text{ cm}^2 \text{ equiv}^{-1}$
147. A correct electrochemical series can be obtained from K, Ca, Na, Al, Mg, Zn, Fe, Pb, H, Cu, Hg, Ag, Au by interchanging :
 a) Al and Mg b) Zn and Fe c) Zn and Pb d) Pb and H
148. The emf of the cell $\text{Zn} | \text{Zn}^{2+} (0.01 \text{ M}) || \text{Fe}^{2+} (0.001 \text{ M}) | \text{Fe}$ at 298 K is 0.2905. The value of equilibrium constant for the cell reaction is
 a) $10^{\frac{0.32}{0.0298}}$ b) $e^{\frac{0.32}{0.0295}}$ c) $10^{\frac{0.32}{0.0591}}$ d) $10^{\frac{0.26}{0.0295}}$
149. When Alcad storage battery is discharged
 a) Lead sulphate is consumed b) SO_2 is evolved
 c) Lead is formed d) Sulphuric acid is consumed
150. EMF of hydrogen electrode in term of pH is (at 1 atm pressure)
 a) $E_{\text{H}_2} = \frac{RT}{F} \times \text{pH}$ b) $E_{\text{H}_2} = \frac{RT}{F} \cdot \frac{1}{\text{pH}}$
 c) $E_{\text{H}_2} = \frac{2.303RT}{F} \cdot \text{pH}$ d) $E_{\text{H}_2} = -0.0591 \text{ pH}$
151. If $E_{\text{Fe}^{2+}/\text{Fe}}^\circ = -0.441 \text{ V}$ and $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ = 0.771 \text{ V}$, the standard e.m.f. of the reaction $\text{Fe} + 2\text{Fe}^{3+} \rightarrow 3\text{Fe}^{2+}$ will be :
 a) 1.212 V b) 0.111 V c) 0.330 V d) 1.653 V
152. When Zn piece is kept in CuSO_4 solution, copper gets precipitated because:
 a) Standard reduction potential of zinc is more than copper
 b) Standard reduction potential of zinc is less than copper
 c) Atomic number of zinc is larger than copper
 d) Atomic number of zinc is lower than copper
153. Ionic mobility of of electricity is 1 M solution of :
 a) CH_3COOH b) H_2SO_4 c) H_3PO_4 d) Boric acid
154. The equivalent conductivity of 0.1 M weak acid is 100 times less than that at infinite dilution. The degree of dissociation of weak electrolyte at 0.1 M is :
 a) 100 b) 10 c) 0.01 d) 0.001
155. Standard electrode potential of cell $\text{H}_2 | \text{H}^+ || \text{Ag}^+ | \text{Ag}$ is (Given, $E_{\text{Ag}^+/\text{Ag}}^\circ = 0.80 \text{ V}$)
 a) 0.4 V b) 0.8 V c) 1.4 V d) 1.8 V
156. If the current is passed into the solution of an electrolyte:
 a) Anions move towards anode, cations towards cathode
 b) Anions and cations both move towards anode
 c) Anions move towards cathode, cations towards anode
 d) No movement of ions takes place
157. The element that is easiest to be reduced is :
 a) Fe b) Cu c) Ag d) Sn
158. Standard reduction potential for, $\text{Li}^+ | \text{Li}, \text{Zn}^{2+} | \text{Zn}, \text{H}^+ | \text{H}_2$ and $\text{Ag}^+ | \text{Ag}$ is $-3.05, -0.762, 0.00$ and +


- 80 V. Which has highest reducing capacity?
 a) Ag b) H₂ c) Zn d) Li
159. What is the quantity of electricity (in Coulombs) required to deposit all the silver from 250mL of 1 MAgNO₃ solution?
 a) 2412.5 b) 24125 c) 4825.0 d) 48250
160. When 1 faraday of electricity is passed through CuSO₄ solution, number of atoms formed is :
 a) 6.02×10^{23} b) 3.01×10^{23} c) 2 d) 6.02×10^{23}
161. Hydrogen gas is not liberated when the following metal is added to dil. HCl
 a) Ag b) Zn c) Mg d) Sn
162. In a hydrogen-oxygen fuel cell, combustion of hydrogen occurs to
 a) Generate heat
 b) Create potential difference between the two electrodes
 c) Produce high purity water
 d) Remove adsorbed oxygen from electrode surfaces.
163. The sum of the two transport number of ions for an electrolyte is always equal to :
 a) 1 b) 2 c) 1/2 d) None of these
164. On passing 0.5 F electricity through molten sodium chloride, sodium deposited at cathode will be
 a) 29.25 g b) 11.50 g c) 58.50 g d) 0.00 g
165. A solution of CuSO₄ is electrolysed for 10 min with a current of 1.5 A. What is the mass of copper deposited at the cathode?
 a) 2.096 g b) 0.296 g c) 3.029 g d) 2.906 g
166. The correct order of molar conductivity at infinite dilution of LiCl, NaCl and KCl is
 a) LiCl > KCl > NaCl b) KCl > NaCl > LiCl c) LiCl > NaCl > KCl d) NaCl > KCl > LiCl
167. Salts of A (atomic weight 7), B (atomic weight 27) and C (atomic weight 48) were electrolyzed under identical conditions using the same quantity of electricity. It was found that when 2.1 g of A was deposited, the weights of B and C deposited were 2.7 g and 7.2 g. The valencies of A, B and C are respectively :
 a) 3, 1 and 2 b) 1, 3 and 2 c) 3, 1 and 3 d) 2, 3 and 2
168. Indicator electrode is :
 a) SHE
 b) Calomel electrode
 c) Ag/AgCl electrode
 d) Quinhydrone electrode
169. Molar conductance of electrolytic solution Λ_m is
 a) $\propto l$ b) $\propto (1/A)$ c) $\propto (1/C)$ d) $\propto (\sqrt{C})$
170. Which metal is most readily corroded in moist air?
 a) Copper b) Iron c) Silver d) Nickel
171. Electrolysis of dilute aqueous NaCl solution was carried out by passing 10mA current. The time required to liberate 0.01 mole of H₂ gas at the cathode is (1F = 96500 C mol⁻¹)
 a) 9.65×10^4 s b) 19.3×10^4 s c) 28.95×10^4 s d) 38.6×10^4 s
172. The oxidation potential of Mg and Al are + 2.37 and + 1.66 volt respectively. The Mg in chemical reactions :
 a) Will be replaced by Al
 b) Will replace Al
 c) Will not be able to replace Al
 d) None of the above
173. The weight of silver (eq. wt. = 108) displaced by that quantity of current which displaced 5600 mL of hydrogen at STP is :
 a) 54 g b) 108 g c) 5.4 g d) None of these
174. When during electrolysis of a solution of a AgNO₃, 9650 C of charge pass through the electroplating bath, the mass of silver deposited on the cathode will be

- a) 1.08 g b) 10.8 g c) 21.6 g d) 108 g
175. The standard oxidation potentials of the electrodes $\text{Ag} | \text{Ag}^+$, $\text{Sn} | \text{Sn}^{2+}$, $\text{Ca} | \text{Ca}^{2+}$, $\text{Pb} | \text{Pb}^{2+}$ are -0.8 , 0.136 , 2.866 and 0.126 V respectively. The most powerful oxidising agent among these metal ions is :
 a) Pb^{2+} b) Ca^{2+} c) Sn^{2+} d) Ag^+
176. Pure water does not conduct electricity because it
 a) Is neutral b) Is readily decomposed
 c) Is almost totally unionized d) Has a low boiling point
177. The minimum equivalent conductance in fused state is shown by :
 a) MgCl_2 b) BeCl_2 c) CaCl_2 d) SrCl_2
178. A cell necessarily does not contain :
 a) An anode
 b) A cathode
 c) An electrolyte or a fuel
 d) A porous diaphragm
179. The standard redox potentials for the reactions
 $\text{Mn}^{2+} + 2e^- \rightarrow$ and $\text{Mn}^{3+} + e^- \rightarrow \text{M}^{2+}$ are -1.18 V and 1.51 V respectively. What is the redox potential for the reaction
 $\text{Mn}^{3+} + 3e^- \rightarrow \text{Mn}$?
 a) 0.33 V b) 1.69 V c) -0.28 V d) -0.85
180. During electrolysis of fused CaH_2 , H_2 is liberated at :
 a) Anode b) Cathode c) Either electrode d) Not at all
181. Total charge on 1 mole of a monovalent metal ion is equal to :
 a) 6.28×10^{18} coulomb b) 1.6×10^{-19} coulomb c) 9.65×10^4 coulomb d) None of these
182. For which case Λ values $vs\sqrt{c}$ show a straight line?
 a) KCl b) HCOOH c) CH_3NH_2 d) CH_3COOH
183. Which is not true for a standard hydrogen electrode?
 a) The hydrogen ion concentration is $1M$
 b) Temperature is 25°C
 c) Pressure of hydrogen is 1 atmosphere
 d) It contains a metallic conductor which does not adsorb hydrogen
184. The laws of electrolysis were proposed by
 a) Kohlrausch b) Faraday c) Haber d) Bergius
185. The metal that cannot be obtained by electrolysis of the aqueous solution of its salts is :
 a) Ag b) Cr c) Cu d) Al
186. A certain current liberated 0.504 g of hydrogen in 2 h. How many grams of copper can be liberated by the same current flowing for the same time in a copper sulphate solution?
 a) 12.9 g b) 15.9 g c) 31.7 g d) 36.9 g
187. If mercury is used as cathode in the electrolysis of aqueous NaCl solution, the ions discharged at cathode are :
 a) H^+ b) Na^+ c) OH^- d) Cl^-
188. Specific conductivity of a solution
 a) Increases with dilution b) Decreases with dilution
 c) Remains unchanged with dilution d) Depends on mass of electrolyte
189. When an electrolytic solution conducts electricity, current is carried out by :
 a) Electrons b) Cations and anions c) Neutral atoms d) None of these
190. e.m.f. of a cell in terms of reduction potential of its left and right electrode is :
 a) $E = E_L + E_R$ b) $E = E_L - E_R$ c) $E = E_R - E_L$ d) $E = -[E_R + E_L]$
191. Which defines the standard reduction electrode potential of Zn^{2+} ions?
 a) $\text{Zn}^{2+}(\text{aq}) + 2e^- \rightarrow \text{Zn}(\text{s}); [\text{Zn}^{2+}] = 1M$
 b) $\text{Zn}(\text{g}) \rightarrow \text{Zn}^{2+} + 2e^-; [\text{Zn}^{2+}] = 1M$

- c) $\text{Zn}^{2+}(\text{aq}) \rightarrow \text{Zn}(\text{s}) + 2\text{e}^-$; $[\text{Zn}^{2+}] = 1\text{M}$
 d) $\text{Zn}^{2+}(\text{g}) \rightarrow \text{Zn}(\text{s}) - 2\text{e}^-$; $[\text{Zn}^{2+}] = 1\text{M}$
192. Given, the data at 25 °C ,
 $\text{Ag} + \text{I}^- \rightarrow \text{AgI} + \text{e}^-$; $E^\circ = 0.152\text{V}$
 $\text{Ag} \rightarrow \text{Ag}^+ + \text{e}^-$; $E^\circ = -0.800\text{V}$
 What is the value of $\log K_{\text{sp}}$ for AgI?
 $\left(2.303 \frac{RT}{F} = 0.059\text{V}\right)$
- a) -8.12 b) +8.612 c) -37.83 d) -16.13
193. The molar conductivity of HCl, NaCl and CH_3COONa are 425, 188, 96 $\text{S cm}^2 \text{mol}^{-1}$ at 298 K. The molar conductivity of CH_3COOH at the same temperature is $\text{S cm}^2 \text{mol}^{-1}$.
- a) 333 b) 451 c) 325 d) 550
194. In the electrolysis of CuCl_2 solution using Cu electrodes the mass of cathode increases by 3.18 g. What happened at the other electrode?
- a) 0.05 mole of Cu^{2+} ions passed into solution
 b) 0.112 litre of Cl_2 was liberated
 c) 0.56 litre O_2 was liberated
 d) 0.1 mole of Cu^{2+} ions passed into the solution
195. When a quantity of electricity is passed through CuSO_4 solution, 0.16 g of copper gets deposited. If the same quantity of electricity is passed through acidulated water, then the volume of H_2 liberated at STP will be [Given , atomic weight of Cu = 64]
- a) 4.0 cm^3 b) 56 cm^3 c) 604 cm^3 d) 8.0 cm^3
196. Faraday's laws hold good at :
- a) All pressures b) Only at 298 K c) In different solvents d) All of these
197. The standard reduction potentials at 25°C of $\text{Li} + |\text{Li}, \text{Ba}^{2+}|\text{Ba}, \text{Na}^+|\text{Na}$ and $\text{Mg}^{2+}|\text{Mg}$ are -3.05, -2.73, -2.71 and -2.37 V respectively. Which is strongest reducing agent?
- a) Li b) Ba c) Na d) Mg
198. In which cell, electrical energy is converted into chemical energy?
- a) Water voltameter b) Silver voltameter c) Coulometer d) Either of these
199. Passage of 96500 coulomb of electricity liberateslitre of O_2 at NTP during electrolysis.
- a) 5.6 b) 6.5 c) 22.2 d) 11.2
200. The number of coulombs required for the deposition of 107.870 g silver is
- a) 96500 b) 48250 c) 1 d) 10000
201. The units of equivalent conductance, are
- a) $\Omega \text{ cm}^2 \text{equiv}^{-1}$ b) $\Omega \text{ cm}^2 \text{equiv}$ c) $\Omega^{-1} \text{ cm}^2 \text{equiv}^{-1}$ d) $\Omega \text{ cm}^2 \text{equiv}$
202. For strong electrolytes the plot of molar conductance vs \sqrt{C} is
- a) Parabolic b) Linear c) Sinusoidal d) Circular
203. The value of $\Lambda_{\text{eq}}^\infty$ for NH_4Cl , NaOH and NaCl are respectively, 149.74, 248.1 and $126.4 \Omega^{-1} \text{ cm}^2 \text{equiv}^{-1}$. The value of $\Lambda_{\text{eq}}^\infty$ of NH_4OH is
- a) 371.44 b) 271.44
 c) 71.44 d) Cannot be predicted from given data
204. The standard electrode potentials of Ag^+ / Ag is +0.80 V and Cu^+ / Cu is +0.34 V. These electrodes are connected through Asalt bridge and if
- a) Copper electrode acts as Acathode then E_{cell}° is +0.46 V
 b) Silver electrode acts as anode then E_{cell}° is -0.34 V
 c) Copper electrode acts as anode then E_{cell}° is +0.46 V
 d) Silver electrode acts as Acathode then E_{cell}° is -0.34 V
205. e.m.f. of cell $\text{Ni}|\text{Ni}^{2+}(0.1\text{M})||\text{Au}^{3+}(1.0\text{M})|\text{Au}$ is, if E° for $\text{Ni}^{2+}|\text{Ni}$ is -0.25 V, E° for $\text{Au}^{3+}|\text{Au}$ is 1.50 V.
- a) +1.25 V b) -1.75 V c) +1.75 V d) +4.0 V

206. The position of some metals in the electrochemical series in decreasing electropositive character is given as $Mg > Al > Zn > Cu > Ag$. What will happen, if a copper spoon is used to stir a solution of aluminium nitrate?
- The spoon will get coated with aluminium
 - An alloy of copper and aluminium is formed
 - The solution becomes blue
 - There is no reaction
207. Which of the following statements is correct? Galvanic cell converts
- Chemical energy into electrical energy
 - Electrical energy into chemical energy
 - Metal from its elemental state to the combined state
 - Electrolyte into individual ions
208. For cell reaction,
 $Zn + Cu^{2+} \rightarrow Zn^{2+} + Cu$
 Cell representation is
- $Zn | Zn^{2+} || Cu^{2+} | Cu$
 - $Cu | Cu^{2+} || Zn^{2+} | Zn$
 - $Cu | Zn^{2+} || Zn | Cu^{2+}$
 - $Cu^{2+} | Zn || Zn^{2+} | Cu$
209. By passing 9.65 A current for 16 min 40 s, the volume of O_2 liberated at STP will be
- 280 mL
 - 560 mL
 - 1120 mL
 - 2240 mL
210. Consider the following disproportionation
 $2ClO_3^- \rightleftharpoons ClO_2^- + ClO_4^-$
 If the initial concentration of perchlorate ion is 0.1 M what it would be at equilibrium at 298 K?
 $(E_{ClO_4^- / ClO_3^-}^\circ = 0.36 \text{ V and } E_{ClO_3^- / ClO_2^-}^\circ = 0.33 \text{ V})$
- 0.1 M
 - 0.05 M
 - 0.07 M
 - 0.19 M
211. When Cu reacts with $AgNO_3$ solution, the reaction takes place is
- Oxidation of Cu
 - Reduction of Cu
 - Oxidation of Ag
 - Reduction of NO_3^-
212. E° for $F_{2+} + 2e = 2F^-$ is 2.8 V, E° for $1/2 F_2 + e = F^-$ is:
- 2.8 V
 - 1.4 V
 - 2.8 V
 - 1.4 V
213. Which one of the following solutions has highest conductance power?
- 0.1 M CH_3COOH
 - 0.1 M NaCl
 - 0.1 M KNO_3
 - 0.1 M HCl
214. Standard electrode potentials of $Fe^{2+} + 2e \rightarrow Fe$ and $Fe^{3+} + 3e \rightarrow Fe$ are - 440 V and -0.036 V respectively. The standard electrode potential (E°) for $Fe^{3+} + e \rightarrow Fe^{2+}$ is:
- 0.476 V
 - 0.404 V
 - + 0.404 V
 - + 0.772 V
215. Stainless steel does not rust because
- Chromium and nickel combine with iron
 - Chromium forms an oxide layer and protects iron from rusting
 - Nickel present in it, does not rust
 - Iron forms Ahard chemical compound with chromium present in it
216. Cu(II) sulphate solution is treated separately with KCl and KI. In which case, Cu^{2+} be reduced to Cu^+ ?
- With KCl
 - With KI
 - With both (a) and (b)
 - None of these
217. The main function of the salt bridge is :
- To allow ions to go from one cell to another
 - To provide link between two half cells
 - To keep the e.m.f. of the cell positive
 - To maintain electrical neutrality of the solution in two half cells
218. When 9.65 C of electricity is passed through a solution of silver nitrate (atomic weight of Ag = 107.87 taking as 108), the amount of silver deposited is
- 5.8 mg
 - 10.8 mg
 - 15.8 mg
 - 20.8 mg
219. The oxidation number of S in $Na_2S_4O_6$ is
- 2.5 for each S atom

233. The factors which influence the conductance of solution.
- Solute-solute interaction
 - Solute-solvent interaction
 - Temperature
 - All of the above
234. In a cell containing zinc electrode and standard hydrogen electrode(SHE), the zinc electrode acts as :
- Anode
 - Cathode
 - Neither cathode nor anode
 - Both anode and cathode
235. The best conductor of electricity is a 0.1 M solution of:
- Boric acid
 - Sulphuric acid
 - Acetic acid
 - Propionic acid
236. Electrode potential of hydrogen electrode is volt.
- 0
 - +1
 - 1
 - None of these
237. Which aqueous solution will conduct an electric current quite well?
- Glycerol
 - Sugar
 - Hydrochloric acid
 - Pure water
238. Use of electrolysis is not done in
- Production of Na
 - Production of water
 - Purification of metals
 - Production of KOH
239. Beryllium is placed above magnesium in the II group. Beryllium dust, therefore, when added to $MgCl_2$ solution will :
- Have no effect
 - Precipitate Mg metal
 - Precipitate MgO
 - Lead to dissolution of Be metal
240. When electric current is passed through an ionic hydride in molten state
- Hydrogen is liberated at anode
 - Hydrogen is liberated at cathode
 - No change takes place
 - Hydride ions migrates towards cathode
241. Which of the following electrolytic solutions has the least specific conductance?
- 0.02 N
 - 0.2 N
 - 2 N
 - 0.002 N
242. During the electrolysis of an electrolyte, the number of ions produced, is directly proportional to the
- Time consumed
 - Mass of electrons
 - Quantity of electricity passed
 - Electrochemical equivalent of electrolytes
243. 1.8 g of metal were deposited by a current of 3 amperes for 50 minute. The equivalent wt. of metal is :
- 20.5
 - 25.8
 - 19.3
 - 30.7
244. Which substance is obtained in the solution on electrolysis of aqueous $CuSO_4$ solution using graphite electrodes?
- H_2O
 - H_2SO_4
 - Na_2SO_4
 - $Cu(OH)_2$
245. During the electrolysis of fused NaCl, which reaction occurs at anode?
- Chloride ions are oxidized
 - Sodium ions are oxidized
 - Chloride ions are reduced
 - Sodium ions are reduced
246. Which one of the following condition will increase the voltage of the cell represented by the equation?
- $$Cu(s) + 2Ag^+(aq) \rightleftharpoons Cu^{2+}(aq) + 2Ag(s)$$
- Increase in the dimension of Cu electrode
 - Increase in the dimension of Ag electrode
 - Increase in the concentration of Cu^{2+} ions
 - Increase in the concentration of Ag^+ ions
247. Which will reduce zinc oxide to zinc?
- Mg
 - Pb
 - Cu
 - Fe
248. The unit of electrochemical equivalent is :
- gram
 - Gram/ampere
 - Kg/coulomb
 - Coulomb/gram
249. $Sn^{4+} + 3e^- \rightarrow Sn^{2+}$, $E^\circ = 0.13 V$



Calculate K_{eq} for the cell reaction for the cell formed by two electrodes.

- a) 10^{41} b) 10^{32} c) 10^{-32} d) 10^{-42}
250. SI unit of conductivity is :
 a) $\text{ohm}^{-1} \text{ cm}^{-1}$ b) $\text{ohm}^{-1} \text{ cm}^{-1}$ or Sm^{-1} c) ohm m^{-1} d) ohm cm^{-1}
251. Ionic mobility of Ag^+ is
 ($\lambda_{\text{Ag}^+} = 5 \times 10^{-1} \Omega^{-1} \text{ cm}^2 \text{ equiv}^{-1}$)
 a) 5.2×10^{-9} b) 2.4×10^{-9} c) 1.52×10^{-9} d) 8.25×10^{-9}
252. $E_{\text{Fe}^{3+}/\text{Fe}}^\circ = -0.036 \text{ V}$, $E_{\text{Fe}^{2+}/\text{Fe}}^\circ = -0.439 \text{ V}$. The value of standard electrode potential for the charge,
 $\text{Fe}^{3+}(\text{aq}) + e^- \rightarrow \text{Fe}^{2+}(\text{aq})$ will be
 a) -0.072 V b) 0.385 V c) 0.770 V d) -0.270 V
253. Whether tin can displace lead from aqueous lead bromide solution?
 a) No
 b) Yes, because standard reduction potential of $\text{Sn} < \text{Pb}$
 c) Yes, because standard reduction potential of $\text{Sn} > \text{Pb}$
 d) None of the above
254. Given the standard reduction potentials
 $\text{Zn}^{2+}/\text{Zn} = -0.74 \text{ V}$, $\text{Cl}_2/\text{Cl}^- = 1.36 \text{ V}$
 $\text{H}^+/\frac{1}{2}\text{H}_2 = 0 \text{ V}$ and $\text{Fe}^{2+}/\text{Fe}^{3+} = 0.77 \text{ V}$
 The order of increasing strength as reducing agent is
 a) Cl^- , Zn , H_2 , Fe^{2+} b) H_2 , Zn , Fe^{2+} , Cl^- c) Cl^- , Fe^{2+} , Zn , H_2 d) Cl^- , Fe^{2+} , H_2 , Zn
255. Molar conductivities (Λ_m) at infinite dilution of NaCl , HCl and CH_3COONa are 126.4, 425.9 and 91.0 $\text{S cm}^2 \text{ mol}^{-1}$ respectively. Λ_m for CH_3COOH will be :
 a) $425.5 \text{ S cm}^2 \text{ mol}^{-1}$ b) $180.5 \text{ S cm}^2 \text{ mol}^{-1}$ c) $290.8 \text{ S cm}^2 \text{ mol}^{-1}$ d) $390.5 \text{ S cm}^2 \text{ mol}^{-1}$
256. $\text{KCl}(\text{aq})$ cannot be used as a salt bridge for the cell $\text{Cu}(\text{s})|\text{CuSO}_4(\text{aq})||\text{AgNO}_3(\text{aq})|\text{Ag}(\text{s})$ because :
 a) CuCl_2 is precipitated b) Cl_2 gas is given out c) AgCl is precipitated d) All of these
257. The ionic conductance of Ba^{2+} and Cl^- are respectively 127 and $76 \Omega^{-1} \text{ cm}^2$ at infinite dilution. The equivalent conductance (in $\Omega^{-1} \text{ cm}^2$) of BaCl_2 at infinite dilution will be
 a) 139.5 b) 203 c) 279 d) 101.5
258. The Gibbs energy for the decomposition of Al_2O_3 at 500°C is as follows :
 $\frac{2}{3}\text{Al}_2\text{O}_3 \rightarrow \frac{4}{3}\text{Al} + \text{O}_2, \Delta_r G = +966 \text{ kJ mol}^{-1}$.
 The potential difference needed for electrolytic reduction of Al_2O_3 at 500°C is atleast :
 a) 5.0 V b) 4.5 V c) 3.0 V d) 2.5 V
259. Which of the following statements (or equation) is correct?
 a) The units of cell emf are V. cm^{-1}
 b) $\Delta G = -\frac{nF}{E_{\text{cell}}}$
 c) In galvanic cell, chemical energy is transformed into electrical energy.
 d) Oxidation state of Mn in potassium permanganate is +6
260. Faraday's law of electrolysis fails when :
 a) Temperature is increased
 b) Inert electrodes are used
 c) A mixture of electrolytes is used
 d) In none of the above cases
261. Conductance 'C' (in S) is directly proportional to the area of the electrode and concentration and inversely proportional to length of separation of electrode, the unit of constant of proportionality is:
 a) S m mol^{-1} b) $\text{S m}^2 \text{ mol}^{-1}$ c) $\text{S}^{-2} \text{ m}^2 \text{ mol}$ d) $\text{S}^2 \text{ m}^2 \text{ mol}^2$
262. A certain metal fails to liberate H_2 gas from a moderately conc. HCl solution. However, it displaces Ag from

- AgNO₃ solution. Which among the followings may it be?
 a) Mg b) Fe c) Cu d) Cd
263. For the cell reaction $\text{Fe} + 2\text{Fe}^{3+} = 3\text{Fe}^{2+}$, which is not possible?
 a) One cell can be constructed
 b) Three different cells with different E_{cell}° are possible
 c) Three different cells with different number of electrons used in redox reaction are possible
 d) Three different cells with same ΔG° value are possible
264. Copper sulphate solution does not react with
 a) Zinc b) Iron c) Silver d) All of these
265. Rust is
 a) FeO + Fe(OH)₂ b) Fe₂O₃
 c) Fe₂O₃ + Fe(OH)₂ d) Fe₂O₃ and Fe(OH)₃
266. The conductivity of strong electrolyte
 a) Increases on dilution slightly b) Decreases on dilution
 c) Does not change with dilution d) Depends upon density of electrolyte itself
267. An electric current of c ampere was passed through a solution of an electrolyte for ' t ' second depositing P g of the metal M on the cathode. The equivalent weight E of the metal will be :
 a) $E = \frac{c \times t}{P \times 96500}$ b) $E = \frac{c \times P}{t \times 96500}$ c) $E = \frac{96500 \times P}{c \times t}$ d) $E = \frac{c \times t \times 9650}{P}$
268. Cu⁺ ion is not stable in aqueous solution because of disproportionation reaction. E° value for disproportionation of Cu⁺ is (given, $E_{\text{Cu}^{2+}/\text{Cu}^{+}}^{\circ} = 0.15$, $E_{\text{Cu}^{2+}/\text{Cu}}^{\circ} = 0.34$ V)
 a) + 0.38 V b) - 0.38 V c) + 0.49 V d) - 0.49 V
269. The molar conductivities $\Lambda_{\text{NaOAc}}^{\circ}$ and $\Lambda_{\text{HCl}}^{\circ}$ at infinite dilution in water at 25°C are 91.0 and 426.2 S cm²/mol respectively. To calculate $\Lambda_{\text{HOAc}}^{\circ}$, the additional value required is
 a) $\Lambda_{\text{H}_2\text{O}}^{\circ}$ b) $\Lambda_{\text{KCl}}^{\circ}$ c) $\Lambda_{\text{NaOH}}^{\circ}$ d) $\Lambda_{\text{NaCl}}^{\circ}$
270. The molar conductivity of NaCl, HCl and CH₃COONa at infinite dilution are 126.45, 426.16 and 91 ohm⁻¹ cm² mol⁻¹ respectively. The molar conductivity of CH₃COOH at infinite dilution is :
 a) 201.28 ohm⁻¹ cm² mol⁻¹
 b) 698.28 ohm⁻¹ cm² mol⁻¹
 c) 390.71 ohm⁻¹ cm² mol⁻¹
 d) 540.48 ohm⁻¹ cm² mol⁻¹
271. If the electrolyte used in problem 4 is Ba(NO₃)₂, then molecular conductivity of solution is :
 a) 628.56 mho⁻¹ cm² mol⁻¹ b) 628.56 mho cm² mol⁻¹ c) 6.28 mho cm² mol⁻¹ d) 6.28 mho⁻¹ cm² mol⁻¹
272. The equivalent conductivity of KCl at infinite dilution is 130 mho cm² eq⁻¹. The transport number of Cl⁻ ion in KCl at the same temperature is 0.505. The transport number of K⁺ ion is :
 a) 0.495 b) 0.505 c) 0.0495 d) Cannot be predicted
273. A cell in which electric current is produced by net oxidation and reduction process is called :
 a) Voltaic cell b) Electrolytic cell c) Concentration cell d) None of these
274. Electrolysis of aq. Solution of LiCl shows :
 a) pH < 7 b) pH = 7 c) pH > 7 d) No change
275. On the basis of the following E° values, the strongest oxidizing agent is :
 $[\text{Fe}(\text{CN})_6]^{4-} \rightarrow [\text{Fe}(\text{CN})_6]^{3-} + e^{-}; E^{\circ} = -0.35$ V
 $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^{-}; E^{\circ} = 0.77$ V
 a) Fe²⁺ b) Fe³⁺ c) $[\text{Fe}(\text{CN})_6]^{3-}$ d) $[\text{Fe}(\text{CN})_6]^{4-}$
276. The specific conductance of 0.1 N KCl solution at 23°C is 0.012 ohm⁻¹ cm⁻¹. The resistance of cell containing the solution at the same temperature was found to be 55 ohm. The cell constant will be
 a) 0.66 cm⁻¹ b) 1.12 cm⁻¹ c) 0.918 cm⁻¹ d) 1.66 cm⁻¹
277. 20 g of chlorine are evolved in 6 hour from sodium chloride solution by the current of :
 a) 5 ampere b) 10 ampere c) 2.5 ampere d) 50 ampere

278. For the electrochemical cell, $M | M^+ || X^- | X$,
 $E^\circ (M^+ | M) = 0.44 \text{ V}$, $E^\circ (X | X^-) = 0.33 \text{ V}$. From this data one can deduce that
 a) $E^\circ_{\text{cell}} = -0.77 \text{ V}$
 b) $M^+ + X^- \rightarrow M + X$ is the spontaneous reaction
 c) $M + X \rightarrow M^+ + X^-$ is the spontaneous reaction
 d) $E^\circ_{\text{cell}} = 0.77 \text{ V}$
279. The standard reduction potential of the reaction,
 $\text{H}_2\text{O} + e^- \rightarrow 1/2 \text{H}_2 + \text{OH}^-$ at 298 K is
 a) $E^\circ = \frac{RT}{F} \ln K_w$
 b) $E^\circ = -\frac{RT}{F} \ln [p_{\text{H}_2}]^{1/2} [\text{OH}^-]$
 c) $E^\circ = -\frac{RT}{F} \ln \frac{[p_{\text{H}_2}]^{1/2}}{[\text{H}^+]}$
 d) $E^\circ = -\frac{RT}{F} \ln K_w$
280. The correct order $E^\circ_{M^{2+}/M}$ values with negative sign for the four successive elements Cr, Mn, Fe and Co is :
 a) $\text{Cr} > \text{Mn} > \text{Fe} > \text{Co}$ b) $\text{Mn} > \text{Cr} > \text{Fe} > \text{Co}$ c) $\text{Cr} > \text{Fe} > \text{Mn} > \text{Co}$ d) $\text{Fe} > \text{Mn} > \text{Cr} > \text{Co}$
281. The increase in equivalent conductivity of a weak electrolyte solution with dilution is attributed to :
 a) Increase in degree of dissociation
 b) Increase in ionic mobility
 c) Both (a) and (b)
 d) None of the above
282. Resistance of 0.2 M solution of an electrolyte is 50Ω . The specific conductance of the solution is 1.3 S m^{-1} .
 If resistance of the 0.4M solution of the same electrolyte is 260Ω , its molar conductivity is
 a) $6250 \text{ Sm}^2 \text{ mol}^{-1}$
 b) $6.25 \times 10^{-4} \text{ Sm}^2 \text{ mol}^{-1}$
 c) $625 \times 10^{-4} \text{ Sm}^2 \text{ mol}^{-1}$
 d) $62.5 \text{ Sm}^2 \text{ mol}^{-1}$
283. When electric current is passed through a cell having an electrolyte, the positive ions move towards the cathode and the negative ions towards the anode. If the cathode is pulled out of the solution :
 a) The positive and the negative ions both will move towards the anode
 b) The positive ions will start moving towards the anode; the negative ions will stop moving
 c) The negative ions will continue to move towards the anode; the positive ions will stop moving
 d) The positive ions and the negative ions will start moving randomly
284. $\text{Cu}^+ (aq)$ is unstable in solution and undergoes simultaneous oxidation and reduction, according to the reaction
 $2\text{Cu}^+ (aq) \rightleftharpoons \text{Cu}^{2+} (aq) + \text{Cu} (s)$
 choose correct E° for the above reaction if
 $E^\circ_{\text{Cu}^{2+}/\text{Cu}} = 0.34 \text{ V}$, $E^\circ_{\text{Cu}^{2+}/\text{Cu}^+} = 0.15 \text{ V}$
 a) -0.38 V b) $+0.49 \text{ V}$ c) $+0.38 \text{ V}$ d) -0.19 V
285. Standard electrode potential of NHE at 298 K is
 a) 0.05 V b) 0.10 V c) 0.50 V d) 0.00 V
286. A galvanic cell is constructed using the redox reaction,
 $\frac{1}{2} \text{H}_2 (g) + \text{AgCl} (s) \rightleftharpoons \text{H}^+ (aq) + \text{Cl}^- (aq) + \text{Ag} (s)$
 It is represented as
 a) $\text{Pt} | \text{H}_2 (g) | \text{HCl solution} || \text{AgNO}_3 \text{ solution} | \text{Ag}$
 b) $\text{Ag} | \text{AgCl} (s) | \text{KCl solution} | | \text{HCl solution} | \text{H}_2 (g) | \text{Pt}$
 c) $\text{Pt} | \text{H}_2 (g) | \text{KCl solution} || \text{AgCl} (s) | \text{Ag}$
 d) $\text{Pt} | \text{H}_2 (g), \text{HCl solution} || \text{AgCl} (s) | \text{Ag}$
287. $\text{Zn} | \text{Zn}^{2+} (A = 0.1 \text{ M}) || \text{Fe}^{2+} (A = 0.01 \text{ M}) | \text{Fe}$.
 The emf of the above cell is 0.2905 V . Equilibrium constant for the cell reaction is
 a) $10^{0.32/0.0591}$ b) $10^{0.32/0.0295}$ c) $10^{0.26/0.0295}$ d) $e^{0.32/0.0295}$

288. The conductance of a solution of an electrolyte is same as that of its conductivity. The cell used can be said to have cell constant equal to :
- a) 1 b) Zero c) 100 d) 10
289. A conductivity cell has two platinum electrodes of 1.2 cm^2 area, separated by a distance of 0.8 cm . The cell constant is :
- a) 0.66 cm^{-1} b) 1.5 cm^{-1} c) 0.96 cm^{-1} d) 0.66 cm
290. A current of i ampere was passed for t second through three cells P, Q and R connected in series. These contain respectively silver nitrate, mercuric nitrate and mercurous nitrate. At the cathode of the cell P , 0.216 g of Ag was deposited. The weights of mercury deposited in the cathode of Q and R respectively are :
- a) 0.4012 and 0.8024 g b) 0.4012 and 0.2006 g c) 0.2006 and 0.4012 g d) 0.1003 and 0.2006 g
291. Out of $\text{Cu}, \text{Al}, \text{Fe}$ and Zn , metal which can displace all others from their salt solution is
- a) Al b) Cu c) Zn d) Fe
292. The equivalent conductances of two strong electrolytes at infinite dilution in H_2O (where ions move freely through a solution) at 25°C are given below
 $\Lambda^\circ_{\text{CH}_3\text{COONa}} = 91.0 \text{ S cm}^2/\text{equiv}$
 $\Lambda^\circ_{\text{HCl}} = 426.2 \text{ S cm}^2/\text{equiv}$
 What additional information/quantity one needs to calculate Λ° of an aqueous solution of acetic acid?
- a) Λ° of NaCl
 b) Λ° of $\text{CH}_3 \text{COOK}$
 c) The limiting equivalent conductance of H^+ ($\Lambda^\circ_{\text{H}^+}$)
 d) Λ° of chloroacetic acid ($\text{ClCH}_2 \text{COOH}$)
293. The emf of the cell
 $\text{Ni} | \text{Ni}^{2+} (1.0 \text{ M}) || \text{Au}^{3+} (1.0 \text{ M}) | \text{Au}$
 is [$E^\circ (\text{Ni}^{2+} / \text{Ni}) = -0.25 \text{ V}$ and
 $E^\circ (\text{Au}^{3+} / \text{Au}) = +1.5 \text{ V}$]
- a) 2.00 V b) 1.25 V c) -1.25 V d) 1.75 V
294. The standard reduction potential for Fe^{2+}/Fe and Sn^{2+}/Sn electrodes are -0.44 and -0.14 V respectively. For the given cell reaction, $\text{Fe}^{2+} + \text{Sn} \rightarrow \text{Fe} + \text{Sn}^{2+}$, the standard emf is
- a) 0.42 V b) -0.42 V c) -0.30 V d) -1.10 V
295. In a cell that utilises the reaction,
 $\text{Zn} (s) + 2\text{H}^+ (aq) \rightarrow \text{Zn}^{2+} (aq) + \text{H}_2 (g)$
 addition of H_2SO_4 to cathode compartment, will
- a) Lower the E and shift the equilibrium to the left
 b) Lower the E and shift the equilibrium to the right
 c) Increase the E and shift the equilibrium to the right
 d) Increase the E and shift the equilibrium to the left
296. Which will increase the voltage of the cell $\text{Sn}(s) + 2\text{Ag}^+(aq) \rightarrow \text{Sn}^{2+}(aq) + 2\text{Ag}(s)$?
- a) Increase in size of the silver rod
 b) Increase in the concentration of Sn^{2+} ions
 c) Increase in the concentration of Ag^+ ions
 d) None of the above
297. Given standard electrode potentials
 $\text{Fe}^{2+} + 2e^- \rightarrow \text{Fe} \quad E^\circ = -0.440 \text{ V}$
 $\text{Fe}^{3+} + 3e^- \rightarrow \text{Fe} \quad E^\circ = -0.036 \text{ V}$
 The standard electrode potential (E°) for
 $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$ is
- a) $+0.772 \text{ V}$ b) -0.772 V c) $+0.417 \text{ V}$ d) -0.414 V
298. Normal $\text{Al} - \text{AlCl}_3$ coupled with normal hydrogen electrode gives an e.m.f. of 1.66 V . The standard oxidation electrode potential of aluminium is :
- a) -1.66 V b) $+1.66 \text{ V}$ c) -0.83 V d) $+0.83 \text{ V}$

299. Which of the following statements is true for fuel cells?
 a) They are more efficient
 b) They are free from pollution
 c) They run till reactants are active
 d) All of the above
300. The Λ^∞ of NH_4OH at infinite dilution is $\text{S cm}^2 \text{eq}^{-1}$. Given $\lambda_{\text{OH}^-}^\infty = 174$, $\lambda_{\text{Cl}^-}^\infty = 66$ and $\lambda_{\text{NH}_4\text{Cl}}^\infty = 130 \text{ S cm}^2 \text{eq}^{-1}$:
 a) 238
 b) 218
 c) 198
 d) 160
301. The metal that forms a self protecting film of oxide to prevent corrosion is
 a) Na
 b) Al
 c) Cu
 d) Au
302. The number of Faraday's needed to reduce 4 g-equivalents of Cu^{2+} to Cu metal will be
 a) 1
 b) 2
 c) 4
 d) 8
303. The atomic weight of Al is 27. When a current of 5 faraday is passed through a solution of Al^{3+} ions, the wt. of Al deposited is :
 a) 27 g
 b) 36 g
 c) 45 g
 d) 9 g
304. Which is correct representation for a cell at equilibrium?
 a) $\Delta G^\circ = -2.303 RT \log K_{eq}$.
 b) $E^\circ = \frac{2.303RT}{nF} \log K_{eq}$.
 c) $-\Delta G^\circ = RT \ln K_{eq}$.
 d) All of the above.
305. Consider the following E° values
 $E^\circ_{\text{Fe}^{3+}/\text{Fe}^{2+}} = +0.77 \text{ V}$
 $E^\circ_{\text{Sn}^{2+}/\text{Sn}} = -0.14 \text{ V}$
 Under standard conditions, the potential for the reaction
 $\text{Sn}(s) + 2\text{Fe}^{3+}(aq) \rightarrow 2\text{Fe}^{2+}(aq) + \text{Sn}^{2+}(aq)$ is
 a) -0.91 V
 b) $+0.91 \text{ V}$
 c) -0.41 V
 d) $+0.41 \text{ V}$
306. According to Kohlrausch's law the limiting value of equivalent conductivity of an electrolyte A_2B is given by :
 a) $\lambda_{\text{A}^+}^\infty + \lambda_{\text{B}^{2-}}^\infty$
 b) $\frac{1}{2}\lambda_{\text{A}^+}^\infty + \lambda_{\text{B}^{2-}}^\infty$
 c) $\lambda_{\text{A}^+}^\infty + \frac{1}{2}\lambda_{\text{B}^{2-}}^\infty$
 d) $2\lambda_{\text{A}^+}^\infty + \lambda_{\text{B}^{2-}}^\infty$
307. Standard reduction potential of an element is equal to :
 a) $+1 \times$ its reduction potential
 b) $-1 \times$ its standard oxidation potential
 c) 0.00 V
 d) $+1 \times$ its standard oxidation potential
308. Rusting of iron is catalysed by which of the following?
 a) Fe
 b) Zn
 c) O_2
 d) H^+
309. The equivalent conductivity of monobasic acid at infinite dilution is $348 \text{ ohm}^{-1} \text{ cm}^2 \text{eq}^{-1}$. If the resistivity of the solution containing 15 g acid (mol. wt. 49) in 1 litre is 18.5 ohm cm , what is the degree of dissociation of acid?
 a) 45.9%
 b) 40.2%
 c) 60.4%
 d) 50.7%
310. The standard reduction potential E° for half reactions are
 $\text{Zn} = \text{Zn}^{2+} + 2e^- \quad E^\circ = +0.76 \text{ V}$
 $\text{Fe} = \text{Fe}^{2+} + 2e^- \quad E^\circ = +0.41 \text{ V}$
 The emf of the cell reaction
 $\text{Fe}^{2+} + \text{Zn} = \text{Zn}^{2+} + \text{Fe}$ is
 a) -0.35 V
 b) $+0.35$
 c) $+1.17 \text{ V}$
 d) -1.17 V
311. If a salt bridge is removed from the two half cells, the voltage
 a) Drops to zero
 b) Does not change
 c) Increase gradually
 d) Increase rapidly
312. The standard oxidation potentials of Zn, Cu, Ag and Ni electrodes are $+0.76$, -0.34 , -0.80 and $+0.25 \text{ V}$ respectively. Which of the following reaction will provide maximum voltage?

- a) $\text{Cu} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{Ag}$ b) $\text{Zn} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + 2\text{Ag}$
 c) $\text{H}_2 + \text{Ni}^{2+}(\text{aq}) \rightarrow 2\text{H}^+(\text{aq}) + \text{Ni}$ d) $\text{Zn} + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + \text{Cu}$
313. An apparatus used for the measurement of quantity of electricity is known as a :
 a) Calorimeter b) Cathetometer c) Coulometer d) colorimeter
314. For the cell prepared from electrode *A* and *B*; Electrode *A* : $\text{Cr}_2\text{O}_7^{2-} | \text{Cr}^{3+}$, $E_{\text{red}}^\circ = +1.33 \text{ V}$ and Electrode *B* : $\text{Fe}^{3+} | \text{Fe}^{2+}$, $E_{\text{red}}^\circ = 0.77 \text{ V}$. Which of the following statement is correct?
 a) The electrons will flow from *B* to *A* when connection is made
 b) The e.m.f. of the cell will be 0.56 V
 c) *A* will be positive electrode
 d) All of the above
315. Which colourless gas evolves when NH_4Cl reacts with zinc in Adry cell battery?
 a) NH_3 b) N_2 c) H_2 d) Cl_2
316. The standard E_{red}° values of *A*, *B* and *C* are +0.68 V, - 2.54 V, - 0.50 V respectively. The order of their reducing power is
 a) $A > B > C$ b) $A > C > B$ c) $C > B > A$ d) $B > C > A$
317. Based on the data given below, the correct order of reducing power is :
 $\text{Fe}^{3+}(\text{aq}) + e \rightarrow \text{Fe}^{2+}(\text{aq}); E^\circ = +0.77 \text{ V}$
 $\text{Al}^{3+}(\text{aq}) + 3e \rightarrow \text{Al}(\text{s}); E^\circ = -1.66 \text{ V}$
 $\text{Br}_2(\text{aq}) + 2e \rightarrow 2\text{Br}^-(\text{aq}); E^\circ = +1.08 \text{ V}$
 a) $\text{Br}^- < \text{Fe}^{2+} < \text{Al}$ b) $\text{Fe}^{2+} < \text{Al} < \text{Br}^-$ c) $\text{Al} < \text{Br}^- < \text{Fe}^{2+}$ d) $\text{Al} < \text{Fe}^{2+} < \text{Br}^-$
318. Small quantities of solutions of compounds *TX*, *TY* and *TZ* are put into separate test tubes containing *X*, *Y* and *Z* solutions. *TX* does not react with any of these. *TY* reacts with both *X* and *Z*. *TZ* reacts with *X*. The decreasing order of ease of oxidation of the anions X^- , Y^- , Z^- is
 a) Y^-, Z^-, X^- b) Z^-, X^-, Y^- c) Y^-, X^-, Z^- d) X^-, Z^-, Y^-
319. What flows in the internal circuit of Agalvanic cell?
 a) Ions b) Electrons c) Electricity d) Atoms
320. The standard electrode potential of Zn^{2+}/Zn and Ag^+/Ag are -0.763 V and $+0.799 \text{ V}$ respectively. The standard potential of the cell is
 a) 1.56 V b) -1.56 V c) 0.036 V d) -0.036 V
321. Consider the following E° values
 $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ = +0.77 \text{ V}$
 $E_{\text{Sn}^{2+}/\text{Sn}}^\circ = -0.14 \text{ V}$
 Under standard conditions the potential for the reaction
 $\text{Sn}(\text{s}) + 2\text{Fe}^{3+}(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq})$ is
 a) 1.68 V b) 1.40 V c) 0.91 V d) 0.63 V
322. The standard electrode potential for the half - cell reactions are
 $\text{Zn}^{2+} + 2e^- \rightarrow \text{Zn}; E^\circ = -0.76 \text{ V}$
 $\text{Fe}^{2+} + 2e^- \rightarrow \text{Fe}; E^\circ = -0.44 \text{ V}$
 The emf of the cell reaction,
 $\text{Fe}^{2+} + \text{Zn} \rightarrow \text{Zn}^{2+} + \text{Fe}$ is
 a) -0.32 V b) -1.20 V c) $+1.20 \text{ V}$ d) $+0.32 \text{ V}$
323. The reduction potential of hydrogen half-cell will negative if
 a) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 2.0 \text{ M}$ b) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 1.0 \text{ M}$
 c) $p(\text{H}_2) = 2 \text{ atm}$ and $[\text{H}^+] = 1.0 \text{ M}$ d) $p(\text{H}_2) = 2 \text{ atm}$ and $[\text{H}^+] = 2.0 \text{ M}$
324. Give the products available on the cathode and the anode respectively during the electrolysis of an aqueous solution of MgSO_4 between inert electrodes.
 a) $\text{H}_2(\text{g})$ and $\text{O}_2(\text{g})$ b) $\text{O}_2(\text{g})$ and $\text{H}_2(\text{g})$ c) $\text{O}_2(\text{g})$ and $\text{Mg}(\text{s})$ d) $\text{O}_2(\text{g})$ and $\text{SO}_2(\text{g})$
325. Which of the following statements is not applicable to electrolytic conductors?
 a) A single stream of electrons flows from cathode to anode

- b) Show a positive temperature coefficient for conductance
 c) New products show up at the electrodes
 d) Ions are responsible for carrying the current
326. Which of the following expressions correctly represents the equivalent conductance at infinite dilution of $\text{Al}_2(\text{SO}_4)_3$. Given that $\Lambda_{\text{Al}^{3+}}^\infty$ and $\Lambda_{\text{SO}_4^{2-}}^\infty$ are the equivalent conductances at infinite dilution of the respective ions?
- a) $2\Lambda_{\text{Al}^{3+}}^\infty + 3\Lambda_{\text{SO}_4^{2-}}^\infty$ b) $2\Lambda_{\text{Al}^{3+}}^\infty + \Lambda_{\text{SO}_4^{2-}}^\infty$ c) $(\Lambda_{\text{Al}^{3+}}^\infty + \Lambda_{\text{SO}_4^{2-}}^\infty) \times 6$ d) $\frac{1}{3}\Lambda_{\text{Al}^{3+}}^\infty + \frac{1}{2}\Lambda_{\text{SO}_4^{2-}}^\infty$
327. Conductivity of a strong electrolyte
- a) Decreases on dilution b) Increases on dilution
 c) Does not change considerably on dilution d) Depends on density
328. Which of the following compounds will not undergo decomposition on passing electricity through aqueous solution?
- a) Sugar b) Sodium acetate c) Sodium chloride d) Sodium bromide
329. Which loses charge at cathode?
- a) Ions
 b) Cations
 c) Anions
 d) Both anions and cations
330. An electrochemical cell is set up as follows
 $\text{Pt}(\text{H}_2, 1 \text{ atm}) \mid 0.1 \text{ M HCl} \mid \mid 0.1 \text{ M acetic acid} \mid (\text{H}_2, 1 \text{ atm})\text{Pt}$ Emf of this cell will not be zero because
- a) The pH of 0.1 M HCl and 0.1 M acetic acid is not the same b) Acids used in two compartments are different
 c) Emf of a cell depends on the molarities of acids used d) The temperature is constant
331. Which of the following reactions cannot be a base for electrochemical cell?
- a) $\text{H}_2 + \text{O}_2 \rightarrow \text{H}_2\text{O}$
 b) $\text{AgNO}_3 + \text{Zn} \rightarrow \text{Zn}(\text{NO}_3)_2 + \text{Ag}$
 c) $\text{AgNO}_3 + \text{NaCl} \rightarrow \text{AgCl} \downarrow + \text{NaNO}_2$
 d) $\text{KMnO}_4 + \text{FeSO}_4 + \text{H}_2\text{SO}_4 \rightarrow \text{K}_2\text{SO}_4 + \text{Fe}_2(\text{SO}_4)_3 + \text{MnSO}_4 + \text{H}_2\text{O}$
332. The emf of a galvanic cell, with electrode potentials of silver = + 0.80 V and that of copper = + 0.34 V, is
- a) + 0.46 V b) + 0.66 V c) + 0.86 V d) - 0.66 V
333. The standard oxidation potentials of Zn and Ag in water at 25°C are,
 $\text{Zn}(s) \rightarrow \text{Zn}^{2+} + 2e; E^\circ = 0.76 \text{ V}$
 $\text{Ag}(s) \rightarrow \text{Ag}^+ + e; E^\circ = -0.80 \text{ V}$
 Which reaction actually takes place?
- a) $\text{Zn}(s) + 2\text{Ag}^+(aq) \rightarrow \text{Zn}^{2+} + 2\text{Ag}(s)$
 b) $\text{Zn}^{2+} + 2\text{Ag}^+(s) \rightarrow 2\text{Ag}^+(aq) + \text{Zn}(s)$
 c) $\text{Zn}(s) + 2\text{Ag}(s) \rightarrow \text{Zn}^{2+}(aq) + 2\text{Ag}^+(aq)$
 d) $\text{Zn}^{2+}(aq) + \text{Ag}^+(aq) \rightarrow \text{Zn}(s) + \text{Ag}(s)$
334. Amount of electricity that can deposit 108 g of silver from AgNO_3 solution is
- a) 1 F b) 1 A c) 1 C d) None of these
335. Also the $[\text{H}^+]$ for problem 9 using the same data is :
- a) 0.00133 M b) 0.133 M c) 0.0133 M d) None of these
336. A hydrogen electrode placed in a buffer solution of CH_3COONa and acetic acid in the ratio's $x : y$ and $y : x$ has electrode potential values E_1 volt and E_2 volt respectively at 25°C. The pK_a values of acetic acid is (E_1 and E_2 are oxidation potential) :


- a) $\frac{E_1 + E_2}{0.118}$
 b) $\frac{E_2 - E_1}{0.118}$
 c) $\frac{E_1 + E_2}{0.118}$
 d) $\frac{E_1 - E_2}{0.118}$

337. When an electric current is passed through acidulated water, 112 mL of hydrogen gas at STP collects at the cathode in 965 second. The current passed, in ampere is :
 a) 1.0 b) 0.5 c) 0.1 d) 2.0
338. The hydrogen electrode is dipped in A solution of pH 3 at 25°C. The potential would be (the value of $\frac{2.303RT}{F}$ is 0.059 V)
 a) 0.177 V b) 0.087 V c) 0.059 V d) -0.177 V
339. The potential of the following cell is 0.34 V at 25°C. Calculate the standard reduction potential of the copper half-cell.
 Pt | H₂(1 atm) | H⁺ (1 M) || Cu²⁺ (1 M) | Cu
 a) -3.4 V b) +3.4 V c) -0.34 V d) +0.34 V
340. The hydrogen electrode is dipped in a solution of pH = 3 at 25°C. The potential of the cell would be (the value of $\frac{2.303RT}{T}$ is 0.059 V)
 a) 0.059 V b) 0.088 V c) 0.178 V d) -0.177 V
341. The apparatus in which electrical energy is converted into chemical energy is known as :
 a) Voltmeter b) Coulometer c) Both (a) and (b) d) None of these
342. When 1 F of electricity is passed through acidulated water, O₂ evolved is
 a) 1.0 dm³ b) 5.6 dm³ c) 11.2 dm³ d) 22.4 dm³
343. A current of strength 2.5 A was passed through CuSO₄ solution for 6 min 26 s. The amount of copper deposited is:
 (Atomic weight of Cu = 63.5) (1 F = 96500 C)
 a) 0.3175 g b) 3.0175 g c) 7.0135 g d) 6.0275 g
344. The potential of the cell for the reaction,
 $M(s) + 2H^+(1M) \rightarrow H_2(g)(1\text{ atm}) + M^{2+}(0.1M)$
 is 1.500 V. The standard reduction potential for M²⁺/M(s) couple is
 a) 0.1470 V b) 1.470 V c) 14.70 V d) None of these
345. For the cell Zn|Zn²⁺||Cu²⁺|Cu, if the concentration of Zn²⁺ and Cu²⁺ ions is doubled, the e.m.f. of the cell :
 a) Doubles b) Reduces to half c) Remains same d) Becomes zero
346. The ionic conductivity of H⁺ and OH⁻ at 298 K are 349.8 and 198.5 mho cm²eq⁻¹ respectively. The equivalent conductivity of H₂O at infinite dilution is :
 a) 548.3 b) 151.3 c) 699.6 d) 54.83
347. The equivalent conductivity of a solution containing 2.54 g of CuSO₄ per L is 91.0 Ω⁻¹ cm² eq⁻¹. Its conductivity would be
 a) $2.9 \times 10^{-3} \Omega^{-1} \text{ cm}^2$ b) $1.8 \times 10^{-2} \Omega^{-1} \text{ cm}^2$ c) $2.4 \times 10^{-4} \Omega^{-1} \text{ cm}^2$ d) $3.6 \times 10^{-3} \Omega^{-1} \text{ cm}^2$
348. Standard electrode potential data are useful for understanding the suitability of an oxidant in a redox titration. Some half cell reactions and their standard potentials are given below ;
 $\text{MnO}_4^-(aq) + 8\text{H}^+(aq) + 5e^- \rightarrow \text{Mn}^{2+}(aq) + 4\text{H}_2\text{O}; E^\circ = 1.51 \text{ V}$
 $\text{Cr}_2\text{O}_7^{2-}(aq) + 14\text{H}^+(aq) + 6e^- \rightarrow 2\text{Cr}^{3+}(aq) + 7\text{H}_2\text{O}; E^\circ = 1.38 \text{ V}$
 $\text{Fe}^{3+}(aq) + e^- \rightarrow \text{Fe}^{2+}(aq); E^\circ = 0.77 \text{ V}$
 $\text{Cl}_2(g) + 2e^- \rightarrow 2\text{Cl}^-(aq); E^\circ = 1.40 \text{ V}$
 Identify the only incorrect statement regarding the quantitative estimation of aqueous Fe(NO₃)₂:

- a) MnO_4^- can be used in aqueous HCl
 b) $\text{Cr}_2\text{O}_7^{2-}$ can be used in aqueous HCl
 c) MnO_4^- can be used in aqueous H_2SO_4
 d) $\text{Cr}_2\text{O}_7^{2-}$ can be used in aqueous H_2SO_4
349. The standard H electrode is written as :
 a) Pt, H_2 , H^+ ($a = 1$)
 b) PtH_2/H^+ ($a = 1$)
 c) $\text{PtH}_2(\text{g})(1 \text{ atm})/\text{H}^+$ ($a = 1$)
 d) None of the above
350. Standard electrode potential of cell $\text{H}_2 | \text{H}^+ || \text{Ag}^+ | \text{Ag}$ is
 a) 0.8 V b) -0.8 V c) -1.2 V d) 1.2 V
351. A dilute solution of Li_2SO_4 is electrolyzed. The products formed at the anode and cathode, respectively are :
 a) S and Li b) O_2 and Li c) SO_2 and O_2 d) O_2 and H_2
352. 3 F electricity was passed through an aqueous solution of iron (II) bromide. The weight of iron metal (at. Wt. = 56) deposited at the cathode (in g) is
 a) 65 b) 84 c) 112 d) 168
353. 5 A is passed through a solution of zinc sulphate for 40 min. Find the amount of zinc deposited at the cathode
 a) 4.065 g b) 8.065 g c) 16.065 g d) 32.065 g
354. Saturated solution of KNO_3 is used to make 'salt-bridge' because
 a) Velocities of both K^+ and NO_3^- are nearly the same
 b) Velocity of K^+ is greater than that of NO_3^-
 c) Velocity of NO_3^- is greater than that of K^+
 d) KNO_3 is highly soluble in water
355. The calomel electrode is a :
 a) Standard hydrogen electrode
 b) Reference electrode
 c) Platinum electrode
 d) Mercury electrode
356. Calculate the emf of the cell
 $\text{Cu}(s) | \text{Cu}^{2+}(aq) || \text{Ag}^+(aq) | \text{Ag}(s)$
 Given,
 $E_{\text{Cu}^{2+}/\text{Cu}}^\circ = +0.34 \text{ V}$, $E_{\text{Ag}^+/\text{Ag}}^\circ = 0.80 \text{ V}$,
 a) +0.46 V b) +1.14 V c) +0.57 V d) -0.46 V
357. The electrolytic conductance is a direct measure of
 a) Resistance b) Potential c) Dissociation d) Concentration
358. On the basis of electrochemical theory of aqueous corrosion, the reaction occurring at the cathode is
 a) $\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}(\text{l})$ b) $\text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + 2\text{e}^-$
 c) $\text{Fe}(\text{s}) \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}^-$ d) $\text{Fe}^{2+}(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + \text{e}^-$
359. The resistance of $\frac{\text{N}}{10}$ solution is found to be $2.5 \times 10^3 \Omega$. The equivalent conductance of the solution is (cell constant = 1.25 cm^{-1})
 a) $2.5 \Omega^{-1} \text{ cm}^2 \text{ equiv}^{-1}$ b) $5.0 \Omega^{-1} \text{ cm}^2 \text{ equiv}^{-1}$
 c) $2.5 \Omega^{-1} \text{ cm}^{-2} \text{ equiv}^{-1}$ d) $5.0 \Omega^{-1} \text{ cm}^{-2} \text{ equiv}^{-1}$
360. In a concentration cell :
 a) Two electrodes are of different elements
 b) Two electrolytic solutions of the same electrolyte but having different concentrations are used
 c) Electrolyte of one strength but electrodes of two different concentrations are used
 d) Both (b) and (c)

361. Using the following data, for the electrode potentials calculate ΔG° , in kJ, for the indicated reaction
 $5\text{Ce}^{4+}(\text{aq}) + \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l}) \rightarrow 5\text{Ce}^{3+}(\text{aq}) + \text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq})$
 $\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightarrow \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l}) E^\circ = +1.51 \text{ V}$
 $\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq}) E^\circ = +1.61 \text{ V}$
 a) -36.24 b) -48.25 c) -31.54 d) -19.65
362. During electrolysis of an aqueous solution of Cu^{2+} sulphate, 0.635 g of copper was deposited at cathode. The amount of electricity consumed in coulomb is :
 a) 1930 b) 3860 c) 9650 d) 4825
363. Each of the three metals X, Y and Z were put in turn into aqueous solution of the other two. $X + \text{Salt of } Y \text{ (or } Z) = Y \text{ (or } Z) + \text{Salt of } X$. Which observation is probably incorrect?
 a) $Y + \text{Salt of } X = \text{No action observed}$
 b) $Y + \text{Salt of } Z = \text{Nothing can be decided}$
 c) $Z + \text{Salt of } X = X + \text{Salt of } Z$
 d) $Z + \text{Salt of } X = \text{No action observed}$
364. During the charging of lead storage battery, the reaction at anode is represented by :
 a) $\text{Pb}^{2+} + \text{SO}_4^{2-} \rightarrow \text{PbSO}_4$
 b) $\text{PbSO}_4 + \text{H}_2\text{O} \rightarrow \text{PbO}_2 + \text{SO}_4^{2-} + 2\text{H}^+$
 c) $\text{Pb} \rightarrow \text{Pb}^{2+} + 2\text{e}^-$
 d) $\text{Pb}^{2+} + 2\text{e}^- \rightarrow \text{Pb}$
365. Which of the formula given below is correct?
 a) $\kappa = \frac{1}{R} \times \frac{1}{a}$ b) $\kappa = C \times \frac{1}{a}$ c) $\Lambda_{eq} = \kappa \times V_{\text{in mL}}$ d) All of these
366. The number of faraday required to generate 1 g-atom of Mg from MgCl_2 is :
 a) 1 b) 2 c) 3 d) 4
367. During the electrolysis of molten NaCl solution, 230 g of sodium metal is deposited on the cathode, then how many moles of chlorine will be obtained at anode?
 a) 10.0 b) 5.0 c) 35.5 d) 17.0
368. 1.05 g of lead ore containing impurity of Ag was dissolved in HNO_3 and the volume was made 350 mL. An Ag electrode was dipped in the solution. $\text{Pt}(\text{H}_2) | \text{H}^+(1 \text{ M}) || \text{Ag}^+ | \text{Ag}$
 The E_{cell} is 0.503 V at 298 K. The percent of lead in the ore is ($E_{\text{Ag}^+ | \text{Ag}}^\circ = 0.80 \text{ V}$)
 a) 0.033% b) 0.050% c) 0.066% d) 0.13%
369. The equivalent conductivity of 0.05 N solution of a monobasic acid is $15.8 \text{ mho cm}^2 \text{ eq}^{-1}$. If equivalent conductivity of the acid at infinite dilution is $350 \text{ mho cm}^2 \text{ eq}^{-1}$ at the same temperature. What is its degree of dissociation?
 a) 0.04514 b) 0.4514 c) 4.514 d) 0.004514
370. What is the potential of the cell containing two hydrogen electrodes as represented ahead,
 $\text{Pt}; \frac{1}{2} \text{H}_2(\text{g}) | \text{H}^+(10^{-8} \text{ M}) || \text{H}^+(0.001 \text{ M}) | 1/2 \text{H}_2(\text{g}) \text{Pt}$?
 a) -0.295 V b) -0.0591 V c) 0.295 V d) 0.0591 V
371. The standard emf for the given cell reaction, $\text{Zn} + \text{Cu}^{2+} \rightarrow \text{Cu} + \text{Zn}^{2+}$ is 1.10 V at 25°C . The emf for the cell reaction, when 0.1 M Cu^{2+} and 0.1 M Zn^{2+} solutions are used, at 25°C , is
 a) 1.10 V b) -1.10 V c) 2.20 V d) -2.20 V
372. Four colourless salt solutions are placed in separate test tubes and a strip of copper is placed in each. Which solution finally turns blue?
 a) $\text{Pb}(\text{NO}_3)_2$ b) $\text{Zn}(\text{NO}_3)_2$ c) AgNO_3 d) $\text{Cd}(\text{NO}_3)_2$
373. The same amount of electricity was passed through two separate electrolytic cells containing solutions of nickel nitrate and chromium nitrate respectively. If 0.3 g of nickel was deposited in the first cell, the amount of chromium deposited is :
 (at. wt. Ni = 59, Cr = 52)
 a) 0.1 g b) 0.17 g c) 0.3 g d) 0.6 g

374. The molar conductivities of KCl, NaCl and KNO_3 are 152, 128 and $111 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. What is the molar conductivity of NaNO_3 ?
 a) $101 \text{ S cm}^2 \text{ mol}^{-1}$ b) $87 \text{ S cm}^2 \text{ mol}^{-1}$ c) $-101 \text{ S cm}^2 \text{ mol}^{-1}$ d) $-391 \text{ S cm}^2 \text{ mol}^{-1}$
375. The degree of ionisation of weak electrolytes is influenced by :
 a) Temperature
 b) Concentration of electrolyte
 c) Nature of solvent
 d) All of the above
376. At 25°C temperature, the cell potential of a given electrochemical cell is 1.92 V. Find the value of x .
 $\text{Mg}(s) | \text{Mg}^{2+}(aq) \ x \ \text{M} || \text{Fe}^{2+}(aq) \ 0.01 \ \text{M} | \text{Fe}(s)$
 $E^\circ \text{Mg}/\text{Mg}^{2+}(aq) = 2.37 \ \text{V}$ $E^\circ \text{Fe}/\text{Fe}^{2+}(aq) = 0.45 \ \text{V}$
 a) $x = 0.01 \ \text{M}$ b) $x < 0.01 \ \text{M}$
 c) $x > 0.01 \ \text{M}$ d) x cannot be predicted
377. The corrosion of iron object is favoured by :
 a) Presence of H^+ ion
 b) Presence of moisture in air
 c) Presence of impurities in iron object
 d) All of the above
378. The cathodic reaction of a dry cell is represented by
 $2\text{MnO}_2(s) + \text{Zn}^{2+} + 2e^- \rightarrow \text{ZnMn}_2\text{O}_4(s)$
 If, there are 8 g of MnO_2 in the cathodic compartment then the time for which the dry cell will continue to give a current of 2 mA is
 a) 25.675 day b) 51.35 day c) 12.8 day d) 6.423 day
379. The standard emf of the cell,
 $\text{Cd}(s) | \text{CdCl}_2(aq) \ (0.1 \ \text{M}) || \text{AgCl}(s) | \text{Ag}(s)$
 In which the cell reaction is
 $\text{Cd}(s) + 2\text{AgCl}(s) \rightarrow 2\text{Ag}(s) + \text{Cd}^{2+}(aq) + 2\text{Cl}^-(aq)$
 is 0.6915 V at 0°C and 0.6753 V at 25°C . The enthalpy change of the reaction at 25°C is
 a) -176 kJ b) -234.7 kJ c) +123.5 kJ d) -167.26 kJ
380. The factor which is not affecting the conductivity of any solution is
 a) Temperature b) Dilution c) Nature of electrolyte d) None of these
381. The standard reduction potential for the half-cell having reaction
 $\text{NO}_3^-(aq) + 2\text{H}^+(aq) + e^- \rightarrow \text{NO}_2(g) + \text{H}_2\text{O}$
 Is 0.78 V. What will be the reduction potential of the half-cell is a neutral solution?
 a) 0.78 V b) 0.89 V c) 0.36 V d) 0.59 V
382. Two different electrolytic cells filled with molten $\text{Cu}(\text{NO}_3)_2$ and molten $\text{Al}(\text{NO}_3)_3$ respectively are connected in series. When electricity is passed 2.7 g Al is deposited on electrode. Calculate the weight of Cu deposited on cathode.
 $[\text{Cu} = 63.5; \text{Al} = 27.0 \ \text{g mol}^{-1}]$
 a) 190.5 g b) 9.525 g c) 63.5 g d) 31.75 g
383. 1 volt coulomb is :
 a) Equal to 1 joule b) Equal to 10^7 erg c) An unit of energy d) All of these
384. $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}$; $\log[\text{Cu}^{2+}]$ vs. E_{red} graph is of the type as shown in figure where $OA = 0.34 \ \text{V}$, then electrode potential of the half cell of $\text{Cu} | \text{Cu}^{2+}(0.1 \ \text{M})$ will be :



- a) $-0.34 + \frac{0.0591}{2} V$ b) $0.34 + 0.0591 V$ c) $0.34 V$ d) None of these
385. If ϕ denotes standard reduction potential, which is true:
 a) $E_{\text{cell}}^{\circ} = \phi_R - \phi_L$ b) $E_{\text{cell}}^{\circ} = \phi_L + \phi_R$ c) $E_{\text{cell}}^{\circ} = \phi_L - \phi_R$ d) $E_{\text{cell}}^{\circ} = (\phi_L + \phi_R)$
386. A substance that will reduce Ag^+ to Ag but will not reduce Ni^{2+} to Ni is :
 a) Zn b) Pb c) Mg d) Al
387. The correct order of the mobility of the alkali metal ions in aqueous solution is :
 a) $\text{K}^+ > \text{Rb}^+ > \text{Na}^+ > \text{Li}^+$
 b) $\text{Rb}^+ > \text{K}^+ > \text{Na}^+ > \text{Li}^+$
 c) $\text{Li}^+ > \text{Na}^+ > \text{K}^+ > \text{Rb}^+$
 d) $\text{Na}^+ > \text{K}^+ > \text{Rb}^+ > \text{Li}^+$
388. Calculate the volume of H_2 gas at NTP obtained by passing 4 A through acidified H_2O for 30 min is
 a) 0.0836 L b) 0.0432 L c) 0.1672 L d) 0.836 L
389. Resistance of a conductivity cell filled with a solution of an electrolyte of concentration 0.1 M is 100Ω . The conductivity of this solution is 1.29 S m^{-1} . Resistance of the same cell when filled with 0.2 M of the same solution is 520Ω . The molar conductivity of 0.02 M solution of the electrolyte will be
 a) $124 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$ b) $1240 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 c) $1.24 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$ d) $12.4 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
390. Ionic mobility of Ag^+ at infinite dilution is :
 ($\lambda_{\text{Ag}^+}^0 = 5 \times 10^{-3} \text{ S m}^2 \text{ eq}^{-1}$)
 a) 5.2×10^{-8} b) 2.4×10^{-8} c) 1.52×10^{-8} d) 8.25×10^{-8}
391. The number of electrons passing per second through a cross-section of copper wire carrying 10^{-6} ampere :
 a) 6.2×10^{23} b) 6.2×10^{12} c) 6.2×10^{10} d) None of these
392. The amount of substance deposited by the passage of 1 A of current for 1 s is equal to
 a) Equivalent mass b) Molecular mass
 c) Electrochemical equivalent d) Specific equivalent
393. 9.65 C electric current is passed through fused anhydrous MgCl_2 . The magnesium metal thus obtained is completely converted into a Grignard reagent. The number of moles of Grignard reagent obtained is
 a) 5×10^{-4} b) 1×10^{-4} c) 5×10^{-5} d) 1×10^{-5}
394. Which one is correct relation :
 a) $\Delta S = \left(\frac{\partial E}{\partial T}\right)_P \times nF$
 b) $\left(\frac{\partial E}{\partial T}\right)_P = \frac{\Delta G - \Delta H}{T}$
 c) $\left(\frac{\partial E}{\partial T}\right)_P = \frac{\partial(\Delta S)}{\partial T}$
 d) $-\Delta S = \left(\frac{\partial E}{\partial T}\right)_P \times nF$
395. A current is passed through two voltameters connected in series. The first voltameter contains $\text{XSO}_4 (aq)$ while the second voltameter contains $\text{Y}_2\text{SO}_4 (aq)$. The relative atomic masses of X and Y are in the ratio of 2 : 1. The ratio of the mass of X liberated to the mass of Y liberated is :

- a) 1 : 1 b) 1 : 2 c) 2 : 1 d) None of these
396. Given, standard electrode potentials
 $\text{Fe}^{2+} + 2e^- \rightarrow \text{Fe}, E^\circ = -0.440 \text{ V}$
 $\text{Fe}^{3+} + 3e^- \rightarrow \text{Fe}, E^\circ = -0.036 \text{ V}$
 The standard electrode potential (E°) for
 $\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$ is
 a) + 0.771 V b) - 0.771 V c) + 0.417 V d) - 0.417 V
397. The standard cell potential for the cell is : $\text{Zn} | \text{Zn}^{2+}(1M) || \text{Cu}^{2+}(1M) | \text{Cu}$
 $[E^\circ \text{ for } \text{Zn}^{2+}/\text{Zn} = -0.76; E^\circ \text{ for } \text{Cu}^{2+}/\text{Cu} = +0.34]$
 a) $-0.76 + 0.34 = -0.42 \text{ V}$
 b) $-0.34 - (-0.76) = +0.42 \text{ V}$
 c) $0.34 - (-0.76) = +1.10 \text{ V}$
 d) $-0.76 - (+0.34) = -1.10 \text{ V}$
398. The speed of migration of Ag^+ ion and NO_3^- ion are $0.00057 \text{ cm sec}^{-1}$ and $0.00063 \text{ cm sec}^{-1}$ at infinite dilution. The equivalent conductivity of AgNO_3 at infinite dilution is:
 a) 140.2 b) 130.1 c) 120.8 d) 115.8
399. In electrochemical corrosion of metals, the metal undergoing corrosion :
 a) Acts as anode b) Acts as cathode c) Undergoes reduction d) None of these
400. Which does not get oxidised by bromine water?
 a) Fe^{2+} to Fe^{3+} b) Cu^+ to Cu^{2+} c) Mn^{2+} to MnO_4^- d) Sn^{2+} to Sn^{4+}
401. 3 faraday of electricity is passed through molten Al_2O_3 , aqueous solution of CuSO_4 and molten NaCl taken in three different electrolytic cells. The amount of Al, Cu and Na deposited at the cathodes will be in the ratio of :
 a) 1 mole : 2 mole : 3 mole
 b) 3 mole : 2 mole : 1 mole
 c) 1 mole : 1.5 mole : 3 mole
 d) 1.5 mole : 2 mole : 3 mole
402. In a galvanic cell, the electrons flow from
 a) Anode to cathode through the solution b) Cathode to anode through the solution
 c) Anode to cathode through the external circuit d) Cathode to anode through the external circuit
403. Which of the following reactions is correct for a given electrochemical cell at 25°C ?
 $\text{Pt} | \text{Br}_2(\text{g}) | \text{Br}^-(\text{aq}) || \text{Cl}^-(\text{aq}) | \text{Cl}_2(\text{g}) | \text{Pt}$
 a) $2\text{Br}^-(\text{aq}) + \text{Cl}_2(\text{g}) \rightarrow 2\text{Cl}^-(\text{aq}) + \text{Br}_2(\text{g})$ b) $\text{Br}_2(\text{g}) + 2\text{Cl}^-(\text{aq}) \rightarrow 2\text{Br}^-(\text{aq}) + \text{Cl}_2(\text{g})$
 c) $\text{Br}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow 2\text{Br}^-(\text{aq}) + 2\text{Cl}^-(\text{aq})$ d) $2\text{Br}^-(\text{aq}) + 2\text{Cl}^-(\text{aq}) \rightarrow \text{Br}_2(\text{g}) + \text{Cl}_2(\text{g})$
404. Which gains electrons more easily?
 a) H^+ b) Na^+ c) K^+ d) Mg^{2+}
405. Two electrolytic cells, one containing acidified ferrous chloride and another acidified ferric chloride are connected in series. The ratio of iron deposited at cathodes in the two cells when electricity is passed through the cells will be :
 a) 3 : 1 b) 2 : 1 c) 1 : 1 d) 3 : 2
406. Limiting molar conductivity of NH_4OH , i.e., $\Lambda_m(\text{NH}_4\text{OH})$ is equal to :
 a) $\Lambda_m(\text{NH}_4\text{OH}) + \Lambda_m(\text{NH}_4\text{Cl}) - \Lambda_m(\text{HCl})$
 b) $\Lambda_m(\text{NH}_4\text{Cl}) + \Lambda_m(\text{NaOH}) - \Lambda_m(\text{NaCl})$
 c) $\Lambda_m(\text{NH}_4\text{Cl}) + \Lambda_m(\text{NaCl}) - \Lambda_m(\text{NaOH})$
 d) $\Lambda_m(\text{NaOH}) + \Lambda_m(\text{NaCl}) - \Lambda_m(\text{NH}_4\text{Cl})$
407. Given :
 i) $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}, E^\circ = 0.337 \text{ V}$
 ii) $\text{Cu}^{2+} + e^- \rightarrow \text{Cu}^+, E^\circ = 0.153 \text{ V}$
 Electrode potential, E° for the reaction,

$\text{Cu}^+ + e^- \rightarrow \text{Cu}$, will be :

- a) 0.38 V b) 0.52 V c) 0.90 V d) 0.30 V

408. The reaction taking place at anode when an aqueous solution of CuSO_4 is electrolysed using inert Pt electrode :

- a) $2\text{SO}_4^{2-} \rightarrow \text{S}_2\text{O}_8^{2-} + 2e^-$
 b) $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}$
 c) $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4e^-$
 d) $2\text{H}^+ + 2e^- \rightarrow \text{H}_2$

409. Deduce from the following E° values of half cells, what combination of two half cells would result in a cell with the largest potential?

- i) $\text{A}^{3-} \rightarrow \text{A}^{2-} + e^-; \quad E^\circ = 1.5 \text{ V}$
 ii) $\text{B}^{2+} + e^- \rightarrow \text{B}^+; \quad E^\circ = -2.1 \text{ V}$
 iii) $\text{C}^{2+} + e^- \rightarrow \text{C}^+; \quad E^\circ = +0.5 \text{ V}$
 iv) $\text{D} \rightarrow \text{D}^{2+} + 2e^-; \quad E^\circ = -1.5 \text{ V}$

- a) (i) and (ii) b) (i) and (iv) c) (ii) and (iv) d) (iii) and (iv)

410. An ion is reduced to the element when it absorbs 6×10^{20} electrons. The number of equivalents of the ion is :

- a) 0.10 b) 0.01 c) 0.001 d) 0.0001

411. The standard e.m.f. of a galvanic cell can be calculated from :

- a) The size of the electrode
 b) The pH of the solution
 c) The amount of metal in the anode
 d) The E° values of the two half cells

412. The charge in coulomb on Cu^{2+} ion is :

- a) 3.2×10^{-19} b) 2.3×10^{-12} c) 0.23×10^{-19} d) 0.32×10^{-19}

413. $\text{Zn}^{2+} + 2e^- \rightarrow \text{Zn}(s), E^\circ = -0.76$

$\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}, E^\circ = -0.77$

$\text{Cr}^{3+} + 3e^- \rightarrow \text{Cr}, E^\circ = -0.79$

$\text{H}^+ + e^- \rightarrow \frac{1}{2} \text{H}_2, E^\circ = 0.00$

Strongest reducing agent is

- a) H_2 b) Zn c) Fe^{2+} d) Cr

414. The standard reduction potentials at 298 K for the following half reactions are given against each

$\text{Zn}^{2+}(\text{aq}) + 2e^- \rightarrow \text{Zn}(s); \quad E^\circ = -0.762 \text{ V}$

$\text{Cr}^{3+}(\text{aq}) + 3e^- \rightarrow \text{Cr}(s); \quad E^\circ = -0.740 \text{ V}$

$2\text{H}^+(\text{aq}) + 2e^- \rightarrow \text{H}_2(\text{g}); \quad E^\circ = 0.00 \text{ V}$

$\text{Fe}^{3+}(\text{aq}) + e^- \rightarrow \text{Fe}^{2+}(\text{aq}); \quad E^\circ = +0.762 \text{ V}$

The strongest reducing agent is

- a) Zn (s) b) Cr (s) c) $\text{H}_2(\text{g})$ d) $\text{Fe}^{2+}(\text{aq})$

415. Strong electrolytes are those which

- a) Conduct electricity b) Dissolve readily in water
 c) Dissociate into ions at high dilution d) Completely dissociate into ions at all dilutions

416. The cell reaction of Acell is

$\text{Mg}(s) + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Cu}(s) + \text{Mg}^{2+}(\text{aq})$.

If the standard reduction potentials of Mg and Cu are -2.37 and $+0.34 \text{ V}$ respectively. The emf of the cell is




- a) 2.03 V b) -2.03 V c) +2.71 V d) -2.71 V

417. Consider the following reaction :

$2\text{Ag}^+\text{C}_6\text{H}_{12}\text{O}_6 + \text{H}_2\text{O} \rightarrow 2\text{Ag}(s) + \text{C}_6\text{H}_{12}\text{O}_7 + 2\text{H}^+$

When ammonia is added to the solution, pH is raised to 11. Which half-cell reaction is affected by pH and

by how much?



a) E_{oxid}° will increase by a factor of 0.65 from E_{oxid}° . b) E_{oxid}° will decrease by a factor of 0.65 from E_{oxid}° .

c) E_{red}° will increase by a factor of 0.65 from E_{red}° . d) E_{red}° will decrease by a factor of 0.65 from E_{red}° .

418. Which process occurs in the electrolysis of aqueous solution of nickel chloride at nickel anode?



419. A solution containing one mole per litre of each $\text{Cu}(\text{NO}_3)_2$, AgNO_3 , $\text{Hg}_2(\text{NO}_3)_2$ and $\text{Mg}(\text{NO}_3)_2$ is being electrolysed by using inert electrodes. The values of standard electrode potentials in volts (reducing potentials) are $\text{Ag}/\text{Ag}^+ = +0.80$, $2\text{Hg}/\text{Hg}_2^{2+} = +0.79$, $\text{Cu}/\text{Cu}^{2+} = +0.34$, $\text{Mg}/\text{Mg}^{2+} = -2$. With increasing voltage, the sequence of deposition of metals on the cathode will be

a) Ag, Hg, Cu b) Cu, Hg, Ag c) Ag, Hg, Cu, Mg d) Mg, Cu, Hg, Ag

420. The metal that does not displace hydrogen from an acid is

a) Ca b) Al c) Zn d) Hg

421. For an electrolyte A_xB_y , the molar conductivity at infinite dilution can be given by :

a) $\Lambda_M^\circ = x\lambda^\circ \text{A}^{y+} + y\lambda^\circ \text{B}^{x-}$

b) $\Lambda_M^\circ = \frac{1}{x}\lambda^\circ \text{A}^{y+} + \frac{1}{y}\lambda^\circ \text{B}^{x-}$

c) $\Lambda_M^\circ = \frac{1}{y}\lambda^\circ \text{A}^{y+} + \frac{1}{x}\lambda^\circ \text{B}^{x-}$

d) $\Lambda_M^\circ = \lambda^\circ \text{A}^{y+} + \lambda^\circ \text{B}^{x-}$

422. $\text{Sn}^{4+}/\text{Sn}^{2+}$ couple is + 0.15 V and that for the Cr^{3+}/Cr couple is - 0.74 V. These two couples in their standard state are connected to make a cell. The cell potential will be :

a) + 1.83 V b) + 1.19 V c) + 0.89 V d) + 0.18 V

423. The standard reduction potentials for Zn^{2+}/Zn , Ni^{2+}/Ni and Fe^{2+}/Fe are - 0.76, -0.23 and -0.44 V respectively. The reaction $\text{X} + \text{Y}^{2+} \rightarrow \text{X}^{2+} + \text{Y}$ will be spontaneous when :

a) $\text{X} = \text{Ni}, \text{Y} = \text{Zn}$ b) $\text{X} = \text{Fe}, \text{Y} = \text{Zn}$ c) $\text{X} = \text{Zn}, \text{Y} = \text{Ni}$ d) $\text{X} = \text{Ni}, \text{Y} = \text{Fe}$

424. Given that $E_{\text{Fe}^{3+}|\text{Fe}}^\circ$ and $E_{\text{Fe}^{3+},\text{Fe}^{2+}|\text{Pt}}^\circ$ are -0.36 V and - 0.439 V, respectively. The value of $E_{\text{Fe}^{3+},\text{Fe}^{2+}|\text{Pt}}^\circ$ would be :


a) $(-36 - 0.439)\text{V}$

b) $[3(-0.36) + 2(-0.439)]\text{V}$

c) $(-0.36 + 0.439)\text{V}$

d) $[3(-0.36) - 2(-0.439)]\text{V}$

425. The standard emf for the cell reaction,



is +0.36 V at 298 K. The equilibrium constant of the reaction is

a) 5×10^6 b) 1.4×10^{12} c) 7.4×10^{12} d) 1.2×10^6





426. Electrolytic conduction is due to migration of :

a) Protons b) Electrons c) Ions d) All of these

427. the amount of sodium deposited by 5 ampere current for 10 minute from fused NaCl is :

a) 0.715 g b) 71.5 g c) 5.17 g d) 0.517 g


428. Which of the following reactions is used to make a fuel cell?



429. When 1 F of electricity is passed through acidulated water, O_2 evolved is

a) 11.2 dm³ b) 5.6 dm³ c) 22.4 dm³ d) 1.0 dm³

430. The Gibbs energy for the decomposition of Al_2O_3 at 500°C is as follows :



$$\Delta_r G = +960 \text{ kJ mol}^{-1}.$$

The potential difference needed for the electrolytic reduction of aluminium oxide (Al_2O_3) at 500°C is at least :

- a) 4.5 V b) 3.0 V c) 2.5 V d) 5.0 V

431. Which one will liberate Br_2 from KBr ?

- a) HI b) I_2 c) Cl_2 d) SO_2

432. In a galvanic cell, the electrons flow from

- a) Anode to cathode through the external circuit b) Anode to cathode through the solution
c) Cathode to anode through the external circuit d) Cathode to anode through the solution

433. The value of molar conductance of HCl is greater than that of NaCl at a particular temperature and dilution because :

- a) mol. wt. of $\text{HCl} <$ mol. wt. of NaCl
b) $u_{\text{H}^+} > u_{\text{Na}^+}$ (u is speed of ion)
c) HCl is acid
d) Ionisation of HCl is more than NaCl

434. Maximum number of mole of oxygen gas that can be obtained by the electrolytic decomposition of 90 g of water will be:

- a) 1 b) 2.5 c) 5 d) 9


435. More electropositive elements have :

- a) Positive reduction potential
b) Tendency to gain electrons
c) Negative reduction potential
d) Negative oxidation potential

436. Al_2O_3 is reduced by electrolysis at low potentials and high currents. If 4.0×10^4 amperes of current is passed through molten Al_2O_3 for 6 hours, what mass of aluminium is produced? (Assume 100% current efficiency. At mass of $\text{Al} = 27 \text{ g mol}^{-1}$)

- a) $1.3 \times 10^4 \text{ g}$ b) $9.0 \times 10^3 \text{ g}$ c) $8.05 \times 10^4 \text{ g}$ d) $2.4 \times 10^5 \text{ g}$

437. The reaction,



Occurs in the galvanic cell

- a) $\text{Pt}/\text{H}_2(\text{g}) \text{KCl}(\text{sol})||\text{AgCl}(\text{s})|\text{Ag}$ b) $\text{Pt}/\text{H}_2(\text{g}) \text{HCl}(\text{sol})||\text{AgNO}_3(\text{sol})|\text{Ag}$
c) $\text{Pt}/\text{H}_2(\text{g}) \text{HCl}(\text{sol})||\text{AgCl}(\text{s})|\text{Ag}$ d) $\text{Ag}/\text{AgCl}(\text{s})\text{KCl}(\text{sol})||\text{AgNO}_3|\text{Ag}$

438. Which of the following ions can be replaced by H^+ ions when H_2 gas is bubbled through the solutions containing these ions?

- a) Li^+ b) Ba^{2+} c) Cu^{2+} d) Be^{2+}

439. The cell reaction, $\text{Zn} + \text{Cu}^{2+} \rightarrow \text{Zn}^{2+} + \text{Cu}$ is best represented by :

- a) $\text{Cu}/\text{Cu}^{2+}||\text{Zn}^{2+}/\text{Zn}$ b) $\text{Zn}/\text{Zn}^{2+}||\text{Cu}^{2+}/\text{Cu}$ c) $\text{Cu}^{2+}/\text{Cu}||\text{Zn}/\text{Zn}^{2+}$ d) $\text{Pt}/\text{Zn}^{2+}||\text{Pt}/\text{Cu}^{2+}$

440. Ionic mobility (u^∞) of an ion at infinite dilution is related to its ionic conductance (λ_∞) by :

- a) $\lambda_\infty = u_\infty \times \text{Faraday}$ b) $u_\infty = \lambda_\infty \times \text{Faraday}$ c) $\text{Faraday} = u_\infty \times \lambda_\infty$ d) None of these

441. Coulomb is the quantity of current defined as :

- a) One ampere of current passing for 1 sec
b) One which deposits 0.001118 g of Ag on cathode
c) One which deposits electrochemical equivalence of metal
d) All of the above

442. The standard electrode potential is measured by

- a) Electrometer b) Voltmeter c) Pyrometer d) Galvanometer

443. Chlorine gas is passed into a solution containing KF , KI , and KBr and CHCl_3 is added. The initial colour in

CHCl₃ layer is :

- a) Violet due to formation of I₂
 b) Orange due to formation of Br₂
 c) Colourless due to formation of F₂
 d) No colour change due to no reaction
444. On passing 3 A of electricity for 50 min, 1.8 g metal deposits. The equivalent mass of metal is
 a) 9.3 b) 19.3 c) 38.3 d) 39.9
445. How many atoms of calcium will be deposited from a solution of CaCl₂ by a current 0.25 mA flowing for 60 s?
 a) 4.68×10^{18} b) 2.34×10^{18} c) 1.24×10^{18} d) 0.46×10^{18}
446. If 'F' is faraday and 'N' is Avogadro number, then charge of electron can be expressed as
 a) $F \times N$ b) $\frac{F}{N}$ c) $\frac{N}{F}$ d) $F^2 N$
447. By how much is the oxidizing power of Cr₂O₇²⁻/Cr³⁺ couple decreased if the H⁺ concentration is decreased from 1 M to 10⁻³ M at 25°C?
 a) 0.207 V b) 0.414 V c) 0.001 V d) 0.287 V
448. Which process involves corrosion?
 a) Brown deposits on iron articles
 b) Green deposits on battery terminals
 c) Black deposits on silver coin
 d) All of the above
449. The electric conduction of a salt solution in water depends on the
 a) Size of its molecules b) Shape of its molecules
 c) Size of solvent molecules d) Extent of its ionization
450. The electrode potentials for

$$\text{Cu}^{2+}(\text{aq}) + e^{-} \rightarrow \text{Cu}^{+}(\text{aq})$$
 and
$$\text{Cu}^{+}(\text{aq}) + e^{-} \rightarrow \text{Cu}(\text{s})$$
 are + 0.15 V and + 0.50 V respectively. The value of $E_{\text{Cu}^{2+}/\text{Cu}}^{\circ}$ will be :
 a) 0.150 V b) 0.500 V c) 0.325 V d) 0.650 V
451. By diluting a weak electrolyte , specific conductivity (K_c) and equivalent conductivity (λ_c) change as
 a) Both increase b) K_c increases , λ_c decreases
 c) K_c decreases , λ_c increases d) Both decrease
452. The cell reaction for the given cell is :


$$\text{Pt}(\text{H}_2) \mid \text{pH} = 2 \mid \mid \text{pH} = 3 \mid \text{Pt}(\text{H}_2)$$

$$P_1=1 \text{ atm} \qquad \qquad \qquad P_2=1 \text{ atm}$$
 a) Spontaneous b) Non-spontaneous c) In equilibrium d) Either of these
453. If the molar conductance value of Ca²⁺ and Cl⁻ at infinite dilution are respectively 118.88 × 10⁻⁴ m² mho mol⁻¹ and 77.33 × 10⁻⁴ m² mho mol⁻¹ then that of CaCl₂ is (in m² mho mol⁻¹)
 a) 118.88×10^{-4} b) 154.66×10^{-4} c) 273.54×10^{-4} d) 196.21×10^{-4}
454. During electrolysis, the species discharged at cathode are
 a) Anion b) Cation c) Ions d) All of these
455. In the electrolysis of which solution, OH⁻ ions are discharged in preference to Cl⁻ ions?
 a) Dilute NaCl b) Very dilute NaCl c) Fused NaCl d) Solid NaCl
456. A cell constructed by coupling a standard copper electrode and a standard magnesium electrode has emf of 2.7 V. If the standard reduction potential of copper electrode is +0.34 V then that of the magnesium electrode is
 a) + 2.36 V b) - 2.36 V c) + 3.26 V d) - 3.26 V
457. Variation of equivalent conductivity with concentration of strong electrolyte is given by Hückel-Onsager equation expressed as :
 a) $\Lambda_M = \Lambda^{\infty} - b\sqrt{c}$ b) $\Lambda_{\infty} = \Lambda M - b\sqrt{c}$ c) $\Lambda_M = b\sqrt{c} - \Lambda^{\infty}$ d) None of these

458. An electric current is passed through silver nitrate solution using silver electrodes. 10.79 g of silver was found to be deposited on the cathode. If the same amount of electricity is passed through copper sulphate solution using copper electrodes, the weight of copper deposited on the cathode is
 a) 1.6 g b) 2.3 g c) 3.2 g d) 6.4 g
459. The amount of energy expanded during the passage of one ampere current for 100 second under a potential of 115 V is :
 a) 20 kJ b) 11.5 kJ c) 115 kJ d) 0.115 kJ
460. If a strip of copper metal is placed in a solution of ferrous sulphate :
 a) Copper will precipitate out
 b) Iron will precipitate out
 c) Both copper and iron will be dissolved
 d) No reaction will take place
461. The process of zinc-plating on iron sheet is known as
 a) Annealing b) Roasting c) Galvanisation d) Smelting
462. For the following cell with hydrogen electrodes at two different pressures p_1 and p_2
 $\text{Pt}(\text{H}_2) | \text{H}^+(aq) | \text{Pt}(\text{H}_2)$
 $p_1 \quad 1 \text{ M} \quad p_2$
 emf is given by
 a) $\frac{RT}{F} \log_e \frac{p_1}{p_2}$ b) $\frac{RT}{2F} \log_e \frac{p_1}{p_2}$ c) $\frac{RT}{F} \log_e \frac{p_2}{p_1}$ d) $\frac{RT}{2F} \log_e \frac{p_2}{p_1}$
463. During the electrolysis of a solution of AgNO_3 , 9650 C of charge is passed through the electroplating bath. The mass of silver deposited at the cathode will be
 a) 108 g b) 10.8 g c) 1.08 g d) 18.10 g
464. What is the time (in sec) required for depositing all the silver present in 125mL of 1 M AgNO_3 solution by passing a current of 241.25 A ? ($1F = 96500 \text{ C}$)
 a) 10 b) 50 c) 1000 d) 100
465. For the redox reaction,
 $\text{Zn}(s) + \text{Cu}^{2+}(0.1 \text{ M}) \rightarrow \text{Zn}^{2+}(1 \text{ M}) + \text{Cu}(s)$
 taking place in a cell, E_{cell}° is 1.10 V. E_{cell}° for the cell will be
 $\left(2.303 \frac{RT}{F} = 0.0591\right)$
 a) 2.14 V b) 1.80 V c) 1.07 V d) 0.82 V
466. The limiting molar conductivities Λ° for NaCl, KBr and KCl are 126, 152 and 150 $\text{S cm}^2 \text{ mol}^{-1}$ respectively. The Λ° for NaBr is
 a) 128 $\text{S cm}^2 \text{ mol}^{-1}$ b) 248 $\text{S cm}^2 \text{ mol}^{-1}$ c) 328 $\text{S cm}^2 \text{ mol}^{-1}$ d) 348 $\text{S cm}^2 \text{ mol}^{-1}$
467. The emf of the cell,
 $\text{Ag} | \text{Ag}^+(0.1 \text{ M}) || \text{Ag}^+(1 \text{ M}) | \text{Ag}$ at 298 K is
 a) 0.0059 V b) 0.059 V c) 5.9 V d) 0.59 V
468. A solution of sodium sulphate in water is electrolysed using inert electrodes. The products at the cathode and anode are respectively
 a) H_2, O_2 b) O_2, H_2 c) O_2, Na d) O_2, SO_2
469. The standard electrode potential for the change ;
 $\text{Sn}(s) + 2\text{Fe}^{3+}(aq) \rightarrow 2\text{Fe}^{2+}(aq) + \text{Sn}^{2+}(aq)$ is :
 (Given $E_{\text{Fe}^{3+}/\text{Fe}^{2+}}^\circ = 0.77 \text{ V}$ and $E_{\text{Sn}^{2+}/\text{Sn}}^\circ = -0.14 \text{ V}$)
 a) 0.63 V b) 1.40 V c) 0.91 V d) 1.68 V
470. Hydrogen-oxygen fuel cells are used in spacecraft to supply :
 a) Power for heat and light
 b) Power for pressure
 c) Oxygen

- d) None of the above
471. The resistance of 0.01 *N* solution of an electrolyte was found to be 210 ohm at 298 K. Its conductance is :
 a) 4.76×10^{-3} mho b) 4.76 mho c) 210 mho d) None of these
472. The amount of silver deposited on passing 2 F of electricity through aqueous solution of AgNO_3 is
 a) 54 g b) 108 g c) 216 g d) 324 g
473. Cell constant of a conductivity cell is usually derived by using a solution of :
 a) KCl b) NaCl c) NH_4Cl d) LiCl
474. Four successive members of the first series of the transition metals are listed below. For which one of them the standard potential ($E_{M^{2+}/M}^\circ$) value has a positive sign?
 a) Co (Z = 27) b) Ni (Z = 28) c) Cu (Z = 29) d) Fe (Z = 26)
475. When same quantity of electricity is passed through aqueous AgNO_3 and H_2SO_4 solutions connected in series, 5.04×10^{-2} g of H_2 is liberated. What is the mass of silver (in grams) deposited? (Eq. wts. of hydrogen = 1.008, silver = 108)
 a) 54 b) 0.54 c) 5.4 d) 10.8
476. The term infinite dilution refers when :
 a) $\alpha \rightarrow 1$, for weak electrolytes
 b) An electrolyte is 100% dissociated
 c) All interionic effects disappears
 d) All of the above
477. In the problem 15, the limiting mobility of K^+ ion is :
 a) 6.1×10^{-4} b) 6.67×10^{-4} c) 7.1×10^{-4} d) 7.67×10^{-4}
478. The standard reduction potential at 290 K for the following half reactions are,
 (i) $\text{Zn}^{2+} + 2e \rightarrow \text{Zn}(s); E^\circ = -0.762 \text{ V}$
 (ii) $\text{Cr}^{3+} + 3e \rightarrow \text{Cr}(s); E^\circ = -0.740 \text{ V}$
 (iii) $2\text{H}^+ + 2e \rightarrow \text{H}_2(\text{g}); E^\circ = -0.000 \text{ V}$
 (iv) $\text{Fe}^{3+} + e \rightarrow \text{Fe}^{2+}; E^\circ = +0.77 \text{ V}$
 Which is the strongest reducing agent?
 a) Zn b) Cr c) Fe^{2+} d) H_2
479. Daniel cell, anode and cathode are respectively
 a) $\text{Zn} | \text{Zn}^{2+}$ and $\text{Cu}^{2+} | \text{Cu}$ b) $\text{Cu} | \text{Cu}^{2+}$ and $\text{Zn}^{2+} | \text{Zn}$ c) $\text{Fe} | \text{Fe}^{2+}$ and $\text{Cu}^{2+} | \text{Cu}$ d) $\text{Cu} | \text{Cu}^{2+}$ and $\text{Fe}^{2+} | \text{Fe}$
480. Iron sheets are galvanized to :
 a) Prevent action of O_2 and H^+ on Fe
 b) Prevent oxidation of Fe
 c) Prevent rusting
 d) All of the above
481. The conductance of all the ions present in a solution containing 1 g equivalent in it is known as :
 a) Conductivity
 b) Equivalent conductivity
 c) Molecular conductivity
 d) None of the above
482. For the reduction of silver ions with copper metal, the standard cell potential is 0.46 V at 25°C. The value of standard Gibbs energy ΔG° will be :
 a) - 89.0 kJ b) - 89.0 J c) - 44.5 kJ d) - 98.0 kJ
483. E° of an electrode is :
 a) Extensive property b) Constitutive property c) Colligative property d) Intensive property
484. The one which decreases with dilution is
 a) Molar conductance b) Conductance
 c) Specific Conductance d) Equivalent conductance
485. Which of the following metal can replace zinc from ZnSO_4 solution?
 a) Cu b) Hg c) Fe d) Al

486. Several blocks of magnesium are fixed to the bottom of Aship to
 a) Keep away the sharks
 b) Make the ship lighter
 c) Prevent action of water and salt
 d) Prevent puncturing by under- seaArocks




$E^\circ_{\text{MnO}_4^-|\text{MnO}_2}$ is

- a) 1.70 V b) 0.91 V c) 1.37 V d) 0.548 V

488. In a salt bridge, KCl is used because :

- a) It is an electrolyte
 b) It is good conductor of electricity
 c) The transport number of K^+ and Cl^- ions are nearly same or both have same ionic mobility.
 d) It is ionic compound.

489. $\text{AgNO}_3 (aq)$ was added to an aqueous KCl solution gradually and the conductivity of the solution was measured. The plot of conductance (Λ) versus the volume of AgNO_3 is



- a) (P) b) (Q) c) (R) d) (S)

490. For the electrochemical cell, $M | M^+ || X^- | X, E^\circ(M^+ | M) = 0.44 \text{ V}$ and $E^\circ(X | X^-) = 0.33 \text{ V}$. From this data one can deduce that

- a) $E_{\text{cell}} = 0.77 \text{ V}$
 b) -0.77 V
 c) $M^+ + X^- \rightarrow M + X$ is the spontaneous reaction
 d) $M + X \rightarrow M^+ + X^-$ is the spontaneous reaction

491. The specific conductance (κ) of an electrolyte of 0.1 N concentration is related to equivalent conductance (Λ) by the following formula

- a) $\Lambda = \kappa$ b) $\Lambda = 10\kappa$ c) $\Lambda = 100\kappa$ d) $\Lambda = 10000\kappa$


492. Which is the correct representation for Nernst equation?

- a) $E_{RP} = E^\circ_{RP} + \frac{0.059}{n} \log \frac{[\text{oxidant}]}{[\text{reductant}]}$
 b) $E_{OP} = E^\circ_{OP} - \frac{0.059}{n} \log \frac{[\text{oxidant}]}{[\text{reductant}]}$
 c) $E_{OP} = E^\circ_{OP} + \frac{0.059}{n} \log \frac{[\text{reductant}]}{[\text{oxidant}]}$
 d) All of the above

493. The number of electrons required to deposit 1 g atom of Al(at. wt. = 27) from a solution of AlCl_3 are :

- a) 1 N b) 2 N c) 3 N d) 4 N

494. The standard reduction potential of some electrodes are,




The Strongest oxidant is :

- a) Copper b) Zinc c) Hydrogen d) Cu^{2+}

495. In the electrolytic cell, flow of electrons is from

- a) Cathode to anode in solution
 c) Cathode to anode through internal supply
- b) Cathode to anode through external supply
 d) Anode to cathode through internal supply
496. The weight ratio of Mg and Al deposited during the passage of same current through their molten salts :
 a) 12 : 9 b) 9 : 12 c) 6 : 2 d) 2 : 3
497. $2\text{Fe}^{3+} + 3\text{I}^{-} \rightleftharpoons 2\text{Fe}^{2+} + \text{I}_3^{-}$
 The standard reduction potentials in acidic conditions are 0.77 and 0.54 V respectively for $\text{Fe}^{3+} / \text{Fe}^{2+}$ and $\text{I}_3^{-} / \text{I}^{-}$ couples. The equilibrium constant for the reaction is
 a) 6.26×10^{-7} b) 5.33×10^{-4} c) 6.26×10^7 d) 5.33×10^4
498. In a cell that utilizes the reaction
 $\text{Zn}(s) + 2\text{H}^{+}(aq) \rightarrow \text{Zn}^{2+}(aq) + \text{H}_2(g)$
 addition of H_2SO_4 to cathode compartment will
 a) Lower the E and shift equilibrium to the right
 b) Lower the E and shift equilibrium to the left
 c) Increase the E and shift equilibrium to the right
 d) Increase the E and shift equilibrium to the left
499. On passing electricity through dilute H_2SO_4 solution the amount of substance liberated at the cathode and anode are in the ratio :
 a) 1 : 8 b) 8 : 1 c) 16 : 1 d) 1 : 16
500. The increase in equivalent conductivity of a strong electrolytic solution with dilution is attributed to :
 a) Increase in number of ions per unit volume
 b) Increase in molecular attraction
 c) Increase in degree of dissociation
 d) Increase in ionic mobility
501. The cell, $\text{Zn} | \text{Zn}^{2+} (1 \text{ M}) || \text{Cu}^{2+} (1 \text{ M}) | \text{Cu}$
 $(E_{\text{cell}}^{\circ} = 1.10 \text{ V})$, was allowed to be completely discharged at 298 K. The relative concentration of Zn^{2+} to Cu^{2+} $\left(\frac{[\text{Zn}^{2+}]}{[\text{Cu}^{2+}]}\right)$ is
 a) Antilog (24.08) b) Antilog (37.3) c) $10^{37.3}$ d) 9.65×10^4
502. The algebraic sum of potentials of two electrodes of a galvanic cell is called :
 a) Potential difference b) Ionic difference c) e.m.f. d) Electrode difference
503. The standard oxidation potentials, E° for the half reactions are ;
 $\text{Zn} \rightarrow \text{Zn}^{2+} + 2e; \quad E^{\circ} = +0.76 \text{ V}$
 $\text{Ag} \rightarrow \text{Ag}^{+} + e; \quad E^{\circ} = -0.77 \text{ V}$
 The standard e. m. f. of the cell,
 $\text{Ag}^{+} + \text{Zn} \rightarrow \text{Zn}^{2+} + \text{Ag}$ is :
 a) + 1.53 V b) - 1.53 V c) -0.01 V d) + 0.01 V
504. Rust is a mixture of :
 a) FeO and $\text{Fe}(\text{OH})_2$ b) FeO and $\text{Fe}(\text{OH})_3$ c) Fe_2O_3 and $\text{Fe}(\text{OH})_3$ d) Fe_3O_4 and $\text{Fe}(\text{OH})_3$
505. A metal having negative reduction potential when dipped in the solution of its own ions, has a tendency :
 a) To pass into the solution
 b) To be deposited from the solution
 c) To become electrically positive
 d) To remain neutral
506. The resistance of 0.5 N solution of an electrolyte in a conductivity cell was found to be 45 ohms. The equivalent conductivity of the same solution isif the electrodes in the cell are 2.2 cm apart and have an area of 3.8 cm^2 .
 a) 25.73 b) 30.75 c) 35.75 d) 15.75
507. The SI unit for ionic mobility is:
 a) $\text{m}^2\text{volt}^{-1}\text{sec}^{-1}$ b) $\text{cm}^2\text{volt}^{-1}\text{sec}^{-1}$ c) $\text{cm} \text{ volt}^{-1}\text{sec}^{-1}$ d) $\text{cm}^{-2}\text{volt}^{-1}\text{sec}^{-1}$
508. Which modifications are necessary to determine resistance of solution by usual method of measurements

- on Wheatstone bridge principle?
- A.C. should be used
 - A conductivity cell is used
 - Galvanometer is replaced by magic eye or head phone arrangement
 - All of above
509. The number of electrons passing per second through a cross-section of Cu wire carrying 10 ampere is :
- 6×10^{19}
 - 8×10^{19}
 - 1×10^{19}
 - 1.6×10^{19}
510. Electrolytic reduction of alumina to aluminium by Hall-Heroult process is carried out :
- In the presence of NaCl
 - In the presence of fluoride
 - In the presence of cryolite, which forms a melt with lower melting temperature
 - In the presence of cryolite, which forms a melt with higher melting temperature
511. Electrolytes, when dissolved in water, dissociate into their constituent ions. The degree of dissociation of a weak electrolyte increases with
- The presence of a substance yielding common ion
 - Decreasing temperature
 - Decreasing concentration of the electrolyte
 - Increasing concentration of the electrolyte
512. The electrolytic bath used in gold plating of copper articles contains :
- Molten gold
 - CuSO_4
 - AuCl_3
 - $\text{AuCl}_3 + \text{NaCN}$
513. Pure water does not conduct electricity because it is
- Basic
 - Almost not ionised
 - Decomposed easily
 - Acidic
514. Galvanic cell is Adevice in which
- Chemical energy is converted into electrical energy.
 - Electrical energy is converted into chemical energy.
 - Chemical energy is seen in the form of heat.
 - Thermal energy from an outside source is used to drive the cell reaction.
515. Astandard hydrogen electrode has zero electrode potential because
- Hydrogen is easier to oxidise
 - This electrode potential is assumed to be zero
 - Hydrogen atom has only one electron
 - Hydrogen is the lightest element
516. The molar conductivity at infinite dilution of AgNO_3 , NaCl and NaNO_3 are 116.5, 110.3 and 105.2 $\text{mho cm}^2\text{mol}^{-1}$ respectively. The molar conductivity of AgCl is :
- 121.6
 - 111.4
 - 130.6
 - 150.2
517. Which is correct about fuel cells?
- Cells continuously run as long as fuels are supplied
 - These are more efficient and free from pollution
 - These are used to provide power and drinking water to astronauts in space programme
 - All of the above
518. The value of electronic charge is equal to :
- $\frac{\text{Faraday}}{\text{Av. number}}$
 - $\text{Faraday} \times \text{Av. number}$
 - $\frac{\text{Av. number}}{\text{Faraday}}$
 - None of these
519. The formula $\alpha = \frac{\Lambda_v}{\Lambda_\infty}$ is valid for :
- Weak electrolytes
 - Strong electrolytes
 - Salts
 - None of these
520. A hypothetical electrochemical cell is shown below;



The e.m.f. measured is + 0.20 V. The cell reaction is :

- a) The cell reaction cannot be predicted
- b) $A + B^+ \rightarrow A^+ + B$
- c) $A^+ + B \rightarrow A + B^+$
- d) $A^+ + e^- \rightarrow A; B^+ + e^- \rightarrow B$

521. The laws of electrolysis were proposed by

- a) Kohlraush
- b) Faraday
- c) Nernst
- d) Berthelot

522. When X amperes of current is passed through molten $AlCl_3$ for 96.5 s. 0.09 g of aluminium is deposited. What is the value of X ?

- a) 10 A
- b) 20 A
- c) 30 A
- d) 40 A

523. It is impossible to measure the actual voltage of any half cell by itself because:

- a) Both half cell reactions takes place simultaneously
- b) Of resistance of wire
- c) A reaction does not take place on its own
- d) None of the above

524. The art of electroplating was given by :

- a) Faraday
- b) Edison
- c) Graham
- d) Brugan

525. If 1 faraday of charge is passed through a solution of $CuSO_4$, the amount of copper deposited will be equal to its :

- a) Gram equivalent weight
- b) Gram molecular weight
- c) Atomic weight
- d) Electrochemical equivalent

526. The oxidation potential values of A, B, C and D are -0.03, +0.108 V, -0.07 V and +0.1 V respectively. The non-spontaneous cell reaction takes place between

- a) A and B
- b) B and D
- c) D and A
- d) B and C

527. The conductivity of $N/10$ KCl solution at $20^\circ C$ is $0.0212 \text{ ohm}^{-1} \text{ cm}^{-1}$ and the resistance of cell containing this solution at $20^\circ C$ is 55 ohm. The cell constant is:

- a) 2.173 cm^{-1}
- b) 1.166 cm^{-1}
- c) 4.616 cm^{-1}
- d) 3.324 cm^{-1}

528. What is the value of E_{cell} ?



$$\text{Given, } E^\circ_{Cr^{3+}/Cr} = -0.74 V$$


$$\text{and } E^\circ_{Fe^{2+}/Fe} = -0.44 V$$

- a) +0.2941 V
- b) +0.5212 V
- c) +0.1308 V
- d) -0.2606 V

529. The E° for OCl^-/Cl^- and $Cl^-/\frac{1}{2}Cl_2$ are 0.94 V and $-1.36 V$; E° for $OCl^-/\frac{1}{2}Cl_2$ is :

- a) $-0.42 V$
- b) $-2.20 V$
- c) $0.52 V$
- d) $1.04 V$

530. The cell reaction for the given cell is spontaneous if :



- a) $P_1 > P_2$
- b) $P_1 < P_2$
- c) $P_1 = P_2$
- d) $P_1 = 1 \text{ atm}$


531. When an acid cell is charged, then



- a) Voltage of cell increases
- b) Resistance of cell increases
- c) Electrolyte of cell dilutes
- d) None of the above

532. An electrolytic cell contains a solution of Ag_2SO_4 and platinum electrodes. A current is passed until 1.6 g of O_2 has been liberated at anode. The amount of Ag deposited at cathode would be:

- a) 1.6 g
- b) 0.8 g
- c) 21.6 g
- d) 107.88 g

533. For Acell given below,





E°_{cell} is

a) $x + 2y$

b) $2x + y$

c) $y - x$

d) $y - 2x$

534. EMF of a cell in terms of reduction potential of its left and right electrodes is

a) $E = E_{\text{left}} - E_{\text{right}}$

b) $E = E_{\text{right}} - E_{\text{left}}$

c) $E = E_{\text{left}} + E_{\text{right}}$

d) None of these

535. At 18°C the conductance of H^+ and CH_3COO^- at infinite dilution are 315 and 35 mho $\text{cm}^2 \text{eq}^{-1}$ respectively. The equivalent conductivity of CH_3COOH at infinite dilution ismho $\text{cm}^2 \text{eq}^{-1}$:

a) 350

b) 280

c) 30

d) 315

536. An alloy of Pb-Ag weighing 1.08 g was dissolved in dilute HNO_3 and the volume made to 100 mL. A silver electrode was dipped in the solution and the emf of the cell set up



Was 0.62 V. If $E^\circ_{\text{cell}} = 0.80 \text{ V}$, what is the percentage of Ag in the alloy?

[At 25°C, $RT/F = 0.06$]

a) 25

b) 2.50

c) 10

d) 50

537. A lamp draws a current of 1.0 A. Find the charge in coulomb used by the lamp in 60 s.

a) 0.6 C

b) 60 C

c) 600 C

d) 0.006 C

538. During electrolysis of water the volume of O_2 liberated is 2.24 dm^3 . The volume of hydrogen liberated, under same conditions will be

a) 2.24 dm^3

b) 1.12 dm^3

c) 4.48 dm^3

d) 0.56 dm^3

539. The amount of electricity required to liberate 1 g-equiv of Cu is

a) 96500 F

b) 1 F

c) 1 C

d) 96500 A

540. Which of the following is correct?

a) Zinc acts as cathode in Daniell cell

b) In a Li – Zn couple, zinc acts as anode

c) Copper will displace iron in solution

d) Zinc displaces tin from its solution

541. The number of electrons involved in redox reactions when a faraday of electricity is passed through an electrolyte in solution is :

a) 6×10^{23}

b) 8×10^{19}

c) 69500

d) 6×10^{-23}

542. During electrolysis of fused sodium chloride, the reaction of the electrodes are:

Anode **Cathode**



543. Which one is correct about conductivity water?

a) The water whose own conductance is very small

b) The water obtained after 7-8 times distillation

c) Kohlrausch prepared the conductivity water for the first time

d) All of the above

544. Blocks of magnesium metal are often strapped to the steel hulls of ocean going ships in order to:

a) Provide cathodic protection

b) Protect oxidation of steel

c) Both (a) and (b) are correct

d) Neither (a) nor (b) is correct

545. Given the limiting molar conductivity as

$$\Lambda_m^0(\text{HCl}) = 425.9 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

$$\Lambda_m^0(\text{NaCl}) = 126.4 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

$$\Lambda_m^0(\text{CH}_3\text{COONa}) = 91 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

The molar conductivity, at infinite dilution, of acetic acid (in $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$) will be


- a) 481.5 b) 390.5 c) 299.5 d) 516.9

546. If the standard electrode potential of $\text{Cu}^{2+} / \text{Cu}$ electrode is 0.34 V, what is the electrode potential at 0.01 M concentration of Cu^{2+} ?

($T = 298 \text{ K}$)

- a) 0.399 V b) 0.281 V c) 0.222 V d) 0.176 V

547. If the ΔG° of Acell reaction,





The standard emf of the cell is

- a) 0.220 V b) -0.220 V c) 0.229 V d) -0.110 V

548. For the cell reaction, $\text{Cu}_2^{2+}(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{Zn}_2^{2+}(\text{aq}) + \text{Cu}(\text{s})$, the change in free energy (ΔG) at a given temperature is a function of :

- a) $\ln c_1$ b) $\ln (c_2/c_1)$ c) $\ln (c_1 + c_2)$ d) $\ln c_2$


549. Consider the following cell reaction



At $[\text{Fe}^{2+}] = 10^{-3} \text{ M}$, $P(\text{O}_2) = 0.1 \text{ atm}$ and $\text{pH} = 3$, the cell potential at 25°C is

- a) 1.47 V b) 1.77 V c) 1.87 V d) 1.57 V



550. Aluminium oxide may be electrolysed at 1000°C to furnish aluminium metal (atomic mass = 27 u; 1 F = 96500 C). The cathode reaction is



To prepare 5.12 kg of aluminium metal by this method would require

- a) $5.49 \times 10^1 \text{ C}$ of electricity b) $5.49 \times 10^4 \text{ C}$ of electricity
c) $1.83 \times 10^7 \text{ C}$ of electricity d) $5.49 \times 10^7 \text{ C}$ of electricity


551. The standard potentials at 25°C for the following half reactions are given ahead,



When zinc dust is added to the solution of MgCl_2 :

- a) ZnCl_2 is formed
b) Zinc dissolves in the solution
c) No reaction takes place
d) Mg is precipitated

552. The cell reaction for the given cell is spontaneous if :



- a) $P_1 > P_2$ b) $P_1 < P_2$ c) $P_1 = P_2$ d) $P_1 = 1 \text{ atm}$


553. Passage of three faraday of charge through aqueous solution of AgNO_3 , CuSO_4 , $\text{Al}(\text{NO}_3)_3$ and NaCl will deposit metals at the cathode in the molar ratio of:

- a) 1 : 2 : 3 : 1 b) 6 : 3 : 2 : 6 c) 6 : 3 : 0 : 0 d) 3 : 2 : 1 : 0

554. In the problem 15, ionic conductance of K^+ ion is :

- a) 64.35 b) 60.20 c) 262.26 d) 26.22

555. In the electrochemical reaction,





increasing the concentration of Fe^{2+}


- a) Increases cell emf b) Increases the current flow
c) Decreases the cell emf d) Alter the pH of the solution


556. How many electrons are there in one coulomb?

- a) 6.02×10^{21} b) 6.24×10^{18} c) 6.24×10^{15} d) 6.02×10^{16}
557. The element which can displace three other halogens from their compound is
a) F b) Cl c) Br d) I
558. The units of equivalent conductivity is
a) $S\text{ cm}^2$ b) $\text{ohm cm}^2(\text{g} - \text{equivalent})$
c) ohm cm d) $\text{ohm}^{-1}\text{ cm}^2(\text{g} - \text{equivalent})^{-1}$
559. Calculate the equilibrium constant for the reaction, at 25°C
 $\text{Cu}(s) + 2\text{Ag}^+(aq) \rightarrow \text{Cu}^{2+}(aq) + 2\text{Ag}(s)$
at 25°C , $E_{\text{cell}}^\circ = 0.47\text{ V}$, $R = 8.314\text{ JK}^{-1}$
 $F = 96500\text{ C}$ is
a) 1.8×10^{15} b) 8.5×10^{15} c) 1.8×10^{10} d) 85×10^{15}
560. The ratio of weights of hydrogen and magnesium deposited by the same amount of electricity from H_2SO_4 and MgSO_4 in aqueous solution are :
a) 1 : 8 b) 1 : 12 c) 1 : 16 d) None of these
561. The Λ^∞ of NH_4Cl , NaOH and NaCl are 129.8, 217.4 and $108.9\text{ ohm}^{-1}\text{ cm}^2\text{ eq.}^{-1}$ respectively. The λ_∞ of NH_4OH is $\text{ohm}^{-1}\text{ cm}^2\text{ eq.}^{-1}$.
a) 238.3 b) 218 c) 240 d) 260
562. The reaction at cathode during the electrolysis of aqueous solution of NaCl in Nelson cell is :
a) $2\text{Cl}^- \rightarrow \text{Cl}_2 + 2e$
b) $2\text{H}^+ + 2e \rightarrow \text{H}_2$
c) $2\text{OH}^- \rightarrow \text{H}_2 + \text{O}_2 + 2e$
d) $\text{Na}^+ + e \rightarrow \text{Na}$
563. Which of the following is an additive property?
a) Conductance b) Viscosity c) Surface tension d) None of these
564. The limiting molar conductivities of NaCl , KBr and KCl are 126, 152 and 150 $\text{S cm}^2\text{ mol}^{-1}$ respectively. The Λ for NaBr is :
a) $302\text{ S cm}^2\text{ mol}^{-1}$ b) $176\text{ S cm}^2\text{ mol}^{-1}$ c) $278\text{ S cm}^2\text{ mol}^{-1}$ d) $128\text{ S cm}^2\text{ mol}^{-1}$
565. The calomel electrode is reversible with respect to :
a) Hg_2^{2+} b) H^+ c) Hg^{2+} d) Cl^-
566. Reaction taking place at anode in dry cell is :
a) $\text{Zn}^{2+} + 2e \rightarrow \text{Zn}(s)$ b) $\text{Zn}(s) \rightarrow \text{Zn}^{2+} + 2e$ c) $\text{Mn}^{2+} + 2e \rightarrow \text{Mn}(s)$ d) $\text{Mn}(s) \rightarrow \text{Mn}^{2+} + 2e$
567. Number of faraday required to liberate 8 g of H_2 is :
a) 8 b) 16 c) 4 d) 2
568. The number of coulombs required to reduce 12.3 g of nitrobenzene to aniline, is
a) 96500 C b) 5790 C c) 95700 C d) 57900 C
569. On passing 0.1 F of electricity through aluminium metal deposited at cathode is ($\text{Al} = 27$)
a) 0.3 g b) 0.6 g c) 0.9 g d) 1.2 g
570. During electrolysis of H_2O , the molar ratio of H_2 and O_2 formed is :
a) 2 : 1 b) 1 : 2 c) 1 : 3 d) 1 : 1
571. At infinite dilution stage, the solution of CH_3COOH in water does not contain :
a) H^+ ion b) CH_3COO^- ion c) CH_3COOH molecule d) All of these
572. 1 faraday of electricity will liberate 1 g-atom of the metal from the solution of :
a) NaCl b) BaCl_2 c) CuSO_4 d) AlCl_3
573. The standard electrode potential of hydrogen electrode at 1 M concentration and hydrogen gas at 1 atm pressure is
a) 1 V b) 6 V c) 8 V d) 0 V
574. The emf of a Daniell cell at 298 K is E_1 , $\text{Zn} | \text{ZnSO}_4 || \text{CuSO}_4 | \text{Cu}$. When the concentration of (0.01 M) (1.0 M) ZnSO_4 is 1.0 M and that of CuSO_4 is 0.01 M, the emf changed to E_2 . What is the relationship between E_1 and E_2 ?

- a) $E_1 = E_2$ b) $E_1 > E_2$ c) $E_1 < E_2$ d) $E_2 = 0 \neq E_1$
575. The acid used in lead storage battery is
 a) H_2SO_4 b) H_3PO_4 c) HCl d) HNO_3
576. The conductance of 1 cm^3 of a solution is known as its :
 a) Resistance
 b) Conductivity
 c) Equivalent conductivity
 d) Molecular conductivity
577. The limiting molar conductivities Λ° for NaCl, KBr and KCl are 126, 152 and 150 $S\text{ cm}^2\text{mol}^{-1}$ respectively. The Λ° for NaBr is
 a) $128\text{ S cm}^2\text{mol}^{-1}$ b) $176\text{ S cm}^2\text{mol}^{-1}$ c) $278\text{ S cm}^2\text{mol}^{-1}$ d) $302\text{ S cm}^2\text{mol}^{-1}$
578. The variation of equivalent conductivity of weak electrolyte with $\sqrt{\text{concentration}}$ is correctly shown in figure :
- a) 

b) 

c) 

d) 
579. The electrode potential measures the :
 a) Tendency of the electrode to gain or lose electrons
 b) Tendency of the cell reaction to occur
 c) Difference in the ionisation potential of electrode and metal ion
 d) Current carried by an electrode
580. Metals can be prevented from rusting by :
 a) Connecting iron to more electropositive metal, *i. e.*, cathodic protection
 b) Connecting iron to more electropositive metal, *i. e.*, anodic protection
 c) Connecting iron to less electropositive metal, *i. e.*, anodic protection
 d) Connecting iron to less electropositive metal, *i. e.*, cathodic protection
581. The number of faraday required to liberate 1 mole of any element indicates :
 a) Weight of element
 b) Conductance of electrolyte
 c) Charge on the ion of that element
 d) None of the above
582. Lithium is generally used as an electrode in high energy density batteries. This is because:
 a) Lithium is the lightest element
 b) Lithium has quite high negative reduction potential
 c) Lithium is quite reactive
 d) Lithium does not corrode easily
583. Corrosion of iron is essentially an electrochemical phenomenon where the cell reactions are
 a) Fe is oxidised to Fe^{2+} and dissolved oxygen in water is reduced to OH^- b) Fe is oxidised to Fe^{3+} and H_2O is reduced to O_2^{2-}
 c) Fe is oxidised to Fe^{2+} and H_2O is reduced to O_2^- d) Fe is oxidised to Fe^{2+} and H_2O is reduced to O_2
584. In the electrodeposition of Ag, the silver ions are:
 a) Reduced at anode b) Reduced at cathode c) Oxidised at anode d) Oxidised at cathode
585. Standard E° of the half cell $Fe|Fe^{2+}$ is $+0.44\text{ V}$ and standard E° of half cell $Cu|Cu^{2+}$ is -0.32 V then :
 a) Cu oxidises Fe^{2+} ion b) Cu^{2+} oxidises Fe c) Cu reduces Fe^{2+} ion d) Cu^{2+} reduces Fe
586. Which of the following is displaced by Fe?
 a) Ag b) Zn c) Na d) All of these
587. The electrochemical cell stops working after sometimes because
 a) Electrode potential of both the electrodes becomes zero

- b) Electrode potential of both the electrodes becomes equal
 c) One of the electrodes is eaten away
 d) The cell reaction gets reversed
588. The resistance of 0.01 *N* solution of an electrolyte was found to be 210 ohm at 298 K, using a conductivity cell of cell constant 0.66 cm^{-1} . The conductivity of solution is :
 a) $3.14 \times 10^{-3} \text{ mho cm}^{-1}$
 b) $3.14 \times 10^{-3} \text{ mho}^{-1} \text{ cm}$
 c) 3.14 mho cm^{-1}
 d) $3.14 \text{ mho}^{-1} \text{ cm}^{-1}$
589. The molar conductivity of acetic acid at infinite dilution is 390.7 and for 0.1 *M* acetic acid solution is 5.2 $\text{mho cm}^2 \text{ mol}^{-1}$. The degree of dissociation of 0.1 *M* CH_3COOH solution is :
 a) 13.3% b) 0.0133% c) 1.33% d) 133%
590. When a lead storage battery is charged, it acts as
 a) A primary cell b) A galvanic cell c) A concentration cell d) An electrolytic cell
591. For gold plating, the electrolyte used is
 a) AuCl_3 b) HAuCl_4 c) $\text{K}[\text{Au}(\text{CN})_2]$ d) None of these
592. How many coulomb of electricity are consumed when 100 mA current is passed through a solution of AgNO_3 for 30 minute during an electrolysis experiment?
 a) 108 b) 18000 c) 180 d) 3000
593. How many kJ of energy is evolved, when a current of 2.00 A passes for 200 s under the potential of 230 V?
 a) 56 kJ b) 86 kJ c) 36 kJ d) 92 kJ
594. What will be the emf for the given cell $\text{Pt} | \text{H}_2(p_1) | \text{H}^+(aq) || \text{H}_2(p_2) | \text{Pt}$?
 a) $\frac{RT}{2F} \log \frac{p_1}{p_2}$ b) $\frac{RT}{F} \log \frac{p_1}{p_2}$ c) $\frac{RT}{F} \log \frac{p_2}{p_1}$ d) None of these
595. The time required to coat a metal surface of 80 cm^2 with $5 \times 10^{-3} \text{ cm}$ thick layer of silver (density 1.05 g cm^{-3} with the passage of 3A current through a silver nitrate solution is :
 a) 115 sec b) 125 sec c) 135 sec d) 145 sec
596. On electrolysing a solution of dilute H_2SO_4 between platinum electrodes, the gas evolved at the anode and cathode are respectively :
 a) SO_2 and O_2 b) SO_3 and H_2 c) O_2 and H_2 d) H_2 and O_2
597. The electrochemical equivalent of silver is 0.0011180 g. When an electric current of 0.5 ampere is passed through an aqueous silver nitrate solution of 200 sec, the amount of silver deposited is:
 a) 1.1180 g b) 0.11180 g c) 5.590 g d) 0.5590 g
598. Galvanised iron sheets have coating of :
 a) Cu b) Sn c) Zn d) Carbon
599. Ionisation depends upon
 a) Pressure b) Volume c) Dilution d) None of these
600. Standard free energies of formation (in kJ/mol) at 298 K are -237.2 , -394.4 and -8.2 for $\text{H}_2\text{O}(l)$, $\text{CO}_2(g)$ and pentane (g), respectively. The value of E°_{cell} for the pentane-oxygen fuel cell is :
 a) 2.0968 V b) 1.0968 V c) 0.0968 V d) 1.968 V
601. In the electrolysis of water, 1 F of electrical energy would evolve
 a) 1 mole of oxygen b) 1 g atom of oxygen c) 8 g of oxygen d) 22.4 L of oxygen
602. Given $l/a = 0.5 \text{ cm}^{-1}$, $R = 50 \text{ ohm}$, $N = 1.0$. The equivalent conductance of the electrolytic cell is
 a) $10 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$ b) $20 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
 c) $300 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$ d) $100 \Omega^{-1} \text{ cm}^2 \text{ g equiv}^{-1}$
603. If 3 F of electricity is passed through the solutions of AgNO_3 , CuSO_4 and AuCl_3 , the molar ratio of the cations deposited at the cathodes will be
 a) 1:1:1 b) 1:2:3 c) 3:2:1 d) 6:3:2
604. If $\text{Mg}^{2+} + 2e \rightarrow \text{Mg}(s)$; $E = -2.37 \text{ V}$,
 $\text{Cu}^{2+} + 2e \rightarrow \text{Cu}(s)$; $E = +0.34 \text{ V}$?

The e.m.f. of the cell $\text{Mg} | \text{Mg}^{2+} || \text{Cu}^{2+} | \text{Cu}$ is :

- a) 2.71 V b) 2.30 V c) 2.80 V d) 1.46 V

605. The standard reduction potentials of

$\text{Zn}^{2+} | \text{Zn}$, $\text{Cu}^{2+} | \text{Cu}$ and $\text{Ag}^+ | \text{Ag}$ are respectively - 0.76, 0.34 and 0.8 V. The following cells were constructed

I $\text{Zn} | \text{Zn}^{2+} || \text{Cu}^{2+} | \text{Cu}$

II $\text{Zn} | \text{Zn}^{2+} || \text{Ag}^+ | \text{Ag}$

III $\text{Cu} | \text{Cu}^{2+} || \text{Ag}^+ | \text{Ag}$

What is the correct order of E°_{cell} of these cells?

- a) II > III > I b) II > I > III c) I > II > III d) III > I > II

606. What is the effect of dilution on the equivalent conductance of strong electrolyte?

- a) Decreases on dilution b) Remains unchanged
c) Increases on dilution d) None of these

607. For which electrolyte the evaluation of A^∞ is not possible by extrapolation of Λ vs \sqrt{c} curves to zero concentration?

- a) KCl b) NH_4OH c) NaCl d) K_2SO_4

608. The standard reduction potential, E° for the half-reactions are as

$\text{Zn} \rightleftharpoons \text{Zn}^{2+} + 2e^-$, $E^\circ = \oplus 0.76 \text{ V}$

$\text{Fe} \rightleftharpoons \text{Fe}^{2+} + 2e^-$, $E^\circ = +0.41 \text{ V}$

The E°_{cell} for the cell formed by these two electrodes is

- a) -0.35 V b) -1.17 V c) +0.35 V d) +1.17 V

609. In the electrochemical cell, $\text{H}_2(\text{g}) | 1 \text{ atm} | \text{H}^+(1 \text{ M}) || \text{Cu}^{2+}(1 \text{ M}) | \text{Cu}(\text{s})$

Which one of the following statements is true?

- a) H_2 is anode, Cu is cathode b) Cu is anode, H_2 is cathode
c) Oxidation occurs at Cu electrode d) Reduction occurs at H_2 electrode

610. Which of the following does not conduct electricity?

- a) Fused NaCl b) Solid NaCl c) Brine solution d) Copper

611. The ionic mobility of alkali metal ions in aqueous solution is maximum for :

- a) K^+ b) Rb^+ c) Li^+ d) Na^+

612. The e.m.f. of the cell involving following changes,

$\text{Zn}(\text{s}) + \text{Ni}^{2+}(1\text{M}) \rightarrow \text{Zn}^{2+}(1\text{M}) + \text{Ni}(\text{s})$ is 0.5105 V. The standard e.m.f. of the cell is :

- a) 0.540 V b) 0.4810 V c) 0.5696 V d) 0.5105 V

613. The factor temperature coefficient of e.m.f. is :

- a) $(\partial E / \partial T)_P$ b) $(\partial E / \partial T)_T$ c) $(\partial E / \partial V)_T$ d) None of these

614. On passing 1 F of electricity through the electrolytic cells containing Ag^+ , Ni^{2+} and Cr^{3+} ions solution, the deposited Ag (at. wt. = 108), Ni (at. wt. = 59) and Cr (at. wt. = 52) is

- | | | | | | |
|----------|--------|--------|----------|---------|--------|
| Ag | Ni | Cr | | | |
| a) 108 g | 29.5 g | 17.3 g | b) 108 g | 59.5 g | 52.0 g |
| c) 108 g | 108 g | 108 g | d) 108 g | 117.5 g | 166 g |

615. Which of the following expression is correct?

- a) $\Delta G^\circ = -nFE^\circ_{\text{cell}}$ b) $\Delta G^\circ = +nFE^\circ_{\text{cell}}$
c) $\Delta G^\circ = -2.303 RT nFE^\circ_{\text{cell}}$ d) $\Delta G^\circ = -nF \log K_c$

616. For which cell e.m.f. is independent of the concentration of electrolytes used?

- a) $\text{Fe} | \text{FeO}(\text{s}) | \text{KOH}(\text{aq}) | \text{Nb}$ $\text{Pt}(\text{H}_2) | \text{HCl} | \text{Pt}(\text{Cl}_2)$ c) $\text{Zn} | \text{Zn}(\text{NO}_3)_2 || \text{CuSO}_4 | \text{Cd}$ $\text{Hg}, \text{HgCl}_2 | \text{KCl} || \text{AgNO}_3 |$

617. In the problem 13, the dissociation constant of acid is :

- a) 2.067×10^{-4} b) 1.02×10^{-4} c) 1.02×10^{-3} d) 1.02×10^{-5}

618. Which are used as secondary reference electrodes?

- a) Calomel electrode
b) Ag/AgCl electrode
c) $\text{Hg}/\text{Hg}_2\text{Cl}_2 - \text{KCl}$ electrode

- d) All of the above
619. The amount of electricity required to produce one mole of copper from copper sulphate solution will be
 a) 1 F b) 2.33 F c) 2 F d) 1.33 F
620. The weight ratio of Al and Ag deposited using the same quantity of current is :
 a) 9 : 108 b) 2 : 12 c) 108 : 9 d) 3 : 8
621. When same electric current is passed through the solution of different electrolytes in series the amounts of the element deposited on the electrode are in the ratio of their:
 a) At.no. b) At. wt. c) Sp. gravity d) Eq. wt.
622. The metal used to recover copper from a solution of CuSO_4 is
 a) Fe b) He c) Na d) Ag
623. If the half-cell reaction $A + e \rightarrow A^-$ has a large negative reduction potential, it follows that :
 a) A is readily reduced b) A is readily oxidized c) A^- is readily reduced d) A^- is readily oxidized
624. Same amount of electric current is passed through solutions of AgNO_3 and HCl. If 1.08 g of silver is obtained in the first case, the amount of hydrogen liberated at STP in the second case is
 a) 224 cm^3 b) 1.008 g c) 112 cm^3 d) 22400 cm^3
625. The standard emf of an galvanic cell involving cell reaction with $n = 2$ is found to be 0.295 V at 25 °C. The equilibrium constant of the reaction would be
 (Given, $F = 96500 \text{ C mol}^{-1}$, $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$)
 a) 2.0×10^{11} b) 4.0×10^{12} c) 1.0×10^2 d) 1.0×10^{10}
626. The correct order of chemical reactivity with water according to electrochemical series is:
 a) $\text{K} > \text{Mg} > \text{Zn} > \text{Cu}$ b) $\text{Mg} > \text{Zn} > \text{Cu} > \text{K}$ c) $\text{K} > \text{Zn} > \text{Mg} > \text{Cu}$ d) $\text{Cu} > \text{Zn} > \text{Mg} > \text{K}$
627. Calculate using appropriate molar conductance of the CH_3COOH from the molar conductances of electrolytes listed below at infinite dilution in H_2O at 25°C :
- | Electrode | KCl | NaCl | HCl | NaOAc | KNO_3 |
|----------------------------------|-------|-------|-------|-------|----------------|
| $\text{S cm}^2 \text{ mol}^{-1}$ | 149.9 | 126.5 | 426.2 | 91.0 | 145.0 |
- a) 51.2 b) 552.7 c) 390.7 d) 217.5
628. The E° for half-cell Fe/Fe^{2+} and Cu/Cu^{2+} are -0.44 V and $+0.32 \text{ V}$ respectively, then
 a) Cu^{2+} oxidises Fe b) Cu oxidises FeFe^{2+} c) Cu reduces Fe^{2+} d) Cu^{2+} oxidises Fe^{2+}
629. The same amount of electricity was passed through two cells containing molten Al_2O_3 and molten NaCl. If 1.8 g of Al were liberated in one cell, the amount of Na liberated in the other cell is :
 a) 4.6 g b) 2.3 g c) 6.4 g d) 3.2 g
630. 1 mole of Al is deposited by X coulomb of electricity passing through aluminium nitrate solution. The number of mole of silver deposited by X coulomb of electricity from silver nitrate solution is :
 a) 3 b) 4 c) 2 d) 1
631. The platinum electrodes were immersed in a solution of cupric sulphate and electric current was passed through the solution. After some time, it was found that colour of copper sulphate disappeared with evolution of gas at the electrode. The colourless solution contain
 a) Copper sulphate b) Copper hydroxide c) Platinum sulphate d) Sulphuric acid